

LIBRARY OF CONGRESS.

QD 31
Chap. 7 Copyright No.

Shelf A 88

UNITED STATES OF AMERICA.

CHEMISTRY:

GENERAL, MEDICAL, AND PHARMACEUTICAL,

INCLUDING

THE CHEMISTRY OF THE U. S. PHARMACOPŒIA.

A MANUAL

ON THE GENERAL PRINCIPLES OF THE SCIENCE, AND THEIR
APPLICATIONS IN MEDICINE AND PHARMACY.

BY

JOHN ATTFIELD, F.R.S.,

Ph.D. of the University of Tübingen, F. I. C.; F. C. S;

Professor of Practical Chemistry to the Pharmaceutical Society of Great Britain;

Formerly Demonstrator of Chemistry at St. Bartholomew's Hospital, London;

Honorary Member of the Pharmaceutical Societies of Great Britain, St. Petersburg, Austria,
Denmark, East Flanders, Switzerland, Australasia, New South Wales, and Queensland;

Honorary Member of the American Pharmaceutical Association;

Honorary Corresponding Member of the Society of Pharmacy of Paris;



Honorary Member of the Colleges of Pharmacy of Philadelphia, New York,
Massachusetts, Chicago, and Ontario, and of the Pharmaceutical Associations of New
Hampshire, Virginia, Liverpool, Manchester, Georgia, Maryland, and
the Province of Quebec;

Editor of the British Pharmacopœia;

Annual Reporter of the British Pharmacopœia to the Medical Council;

Author of a Handbook on Water and Water-Supplies.

FOURTEENTH EDITION.



PHILADELPHIA:

LEA BROTHERS & CO.

1894.

Entered according to Act of Congress, in the year 1894, by
LEA BROTHERS & CO.,
in the Office of the Librarian of Congress, at Washington. All rights reserved.

"But the greatest error of all is, mistaking the ultimate end of knowledge; for some men covet knowledge out of a natural curiosity and inquisitive temper; some to entertain the mind with variety and delight; some for ornament and reputation; some for victory and contention; many for lucre and a livelihood; and but few for employing the divine gift of reason to the use and benefit of mankind."—LORD BACON.

"I hold that the greatest friend to man is labor; that knowledge without toil, if possible, were worthless; that toil in pursuit of knowledge is the best knowledge we can attain; that the continuous effort for fame is nobler than fame itself; that it is not wealth suddenly acquired which is deserving of homage, but the virtues which a man exercises in the slow pursuit of wealth—the abilities so called forth, the self-denials so imposed; in a word, that Labor and Patience are the true schoolmasters on earth."—LORD LYTTON.

"I want to learn all that one human being can. It is awful to be buried alive in the coffin of one's own ignorance and helplessness."—GRAHAM TRAVERS.

Q1731
A 88

PREFACE.

70-978 mll

THE short title on the back of a book, and even the words on the title-page, are generally, and even necessarily, imperfect descriptions of the contents, and hence not unfrequently induce at the outset misconceptions in the minds of readers. The author of *Chemistry: General, Medical, and Pharmaceutical*, would at once state, therefore, that his chief aim is to teach the general truths of chemistry to medical and pharmaceutical pupils. So far as laws and principles are concerned, the book is a work on General Chemistry; but inasmuch as those laws and principles are elucidated and illustrated by that large portion of chemistry which is directly interesting to medical practitioners and pharmacists, the book may be said to be a work on Medical Chemistry and on Pharmaceutical Chemistry. Only in this conventional sense would the author speak of Medical and Pharmaceutical Chemistry; for the truths of chemistry are the same for all students—crystalline verities which cannot be expanded or compressed to suit any class of workers. The leading principles of the science, however, can as easily be illustrated by or deduced from those facts which have interest as from those which have little or no special interest to the followers of medicine and pharmacy. The grand and simple leading truths or laws of chemistry, the lesser truths or principles, and nearly all the interesting relationships of elements and compounds—in a word, the *science* of chemistry—can be taught to medical and pharmaceutical students with little other aid than that afforded by the materials which lie in rich abundance all around these workers. Such a mode of teaching what is stated on the title-page as “the general principles of the science and their applications in medicine and pharmacy” is adopted in this volume. It is a mode which greatly increases the usefulness of the science to the students

chiefly addressed, while it in no way diminishes the value of chemistry to them as an instrument of mental culture—an instrument which sharpens and expands the powers of observation, which enlarges and strengthens memory and imagination, which gives point to the perceptive faculties, and which develops and elaborates the powers of thought and of reason.

This Manual is intended, then, as a systematic exponent of the general truths of chemistry, but is written mainly for the pupils, assistants, and principals engaged in medicine and pharmacy. It is essentially a Manual of Applied Chemistry or Technical Chemistry, but is first of all a Manual of Chemistry.

The book will be found equally useful as a reading-book for students having no opportunities of attending lectures or performing experiments, or, on the other hand, as a text-book for college pupils; while its comprehensive Index, containing nearly ten thousand references, will fit the work for after-consultation in the course of business or professional practice.

From other chemical text-books it differs in three particulars: first, in the exclusion of matter relating to compounds which at present are only of interest to the scientific chemist; secondly, in containing more or less of the chemistry of every substance recognized officially or in general practice as a remedial agent; thirdly, in the paragraphs being so cast that the volume may be used as a guide in studying the science experimentally.

The order of subjects is that which, in the author's opinion, best meets the requirements of medical and pharmaceutical students in Great Britain, Ireland, America, India, and the English colonies. Introductory pages are devoted to a few leading properties of the elements. A review of the facts thus unfolded affords opportunity for stating the views of philosophers respecting the manner in which these elements influence each other as components of terrestrial matter. The consideration in detail of the relations of the elementary and compound radicals follows, synthetical and analytical bearings being pointed out, and attention frequently directed to connecting or underlying truths or general principles. The chem-

istry of substances met with in vegetables and animals, or of similar substances artificially produced (the so-called "organic chemistry"), is next considered. Chemical toxicology and the chemical as well as microscopical characters of morbid urine, urinary sediments, and calculi are then given. The concluding sections form a laboratory-guide to beginners in the chemical and physical study of quantitative analysis. In the Appendix is a long table of tests for impurities in medicinal preparations, a short one of the saturating-powers of acids and alkalies, designed for use in prescribing and dispensing, and an alphabetical list of elements and atomic weights.

In the course of the treatment outlined in the preceding paragraph it will be observed that the whole of the elements are first noticed very shortly, to give the pupil a general view of his course of study, and afterward at length and thoroughly; that the chemistry of the common metallic radicals precedes that of the rarer, and that the sections on the acidulous radicals are similarly divided; while the basylous radicals are arranged according to analytical relations, the common acidulous according to exchangeable value or quantivalence, and the rarer acidulous radicals alphabetically. By this plan the more important facts and principles are repeatedly brought under consideration, the points of view, however, differing according as interest is concentrated on physical, synthetical, analytical, or quantitative properties. This arrangement of matter was adopted, also, partly in the belief that the separate and general truths of chemistry never do or can enter the mind in the order of any scientific classification at present possible. Chemical facts are not yet united by any single, consistent theory. In the current state of chemical knowledge consistency in the methodical arrangement even of elements can only be carried out in one direction, and is necessarily accompanied by inconsistencies in other directions—a result most perplexing to learners, and hence totally subversive of the chief advantages of classification. For this reason the writer has preferred to lead up to, rather than follow, scientific classification—has allowed analogies and affinities to suggest,

rather than be suggested by, classification. Among the acidulous radicals, especially, any known system of classification would have given undue prominence to one set of relations, and undeserved obscurity to others. Then, by separating more important from less important matter, instruction is adapted to the wants of gentlemen whose opportunities of studying chemistry vary greatly, and are unavoidably insufficient to enable them to gain a knowledge of the whole area of the science. One great advantage of the mode of treatment is, that difficulties of nomenclature, notation, chemical constitution, and even those arising from conventionality of language, are explained as they arise, instead of being massed under the head of "Introductory Chapters," "Preliminary Considerations," or "General Remarks," which are not unfrequently too difficult to be understood by a beginner, too voluminous to be remembered except by the aid of subsequent lessons, and are consequently the cause of much trouble and confusion. For an illustration of the treatment thus adopted by the author the reader is referred to the various notes on chemical constitution. (See "Constitution," "Structure," etc. in the Index.) This plan has also admitted of greater prominence being given to "The General Principles of Chemical Philosophy," the only section to which the student is asked frequently to return until he finds himself naturally employing those principles in the interpretation of the phenomena obtained by experiment.

An elementary knowledge of the subjects of Gravitation, Heat, Light, Sound, Electricity, and Magnetism cannot be too strongly recommended to the student of chemistry. The first portion of this Manual would have been devoted to an exposition of these branches of physics, so far as they bear on chemistry, did not the many special books on Physics render such a course unnecessary. Quantitative chemical analysis frequently involving determinations of temperature, specific gravity, and atmospheric pressure, a few paragraphs on these subjects are made introductory to the section on quantitative operations.

The theories that matter consists of molecules, and that

molecules consist of atoms, are freely adopted in this book, the author believing that in the present state of knowledge and education philosophic conceptions concerning chemistry can only be induced in the minds of medical, pharmaceutical, and general students by the aid of those theories.

The chemical notation of the work is in accordance with modern views. Equations illustrative of pharmacopœial processes have in all important cases a name attached to each formula.

In the first edition of this Manual, in 1867, chemical nomenclature was modernized to the extent of defining the alkali-metal salts and the earthy compounds as those of potassium, sodium, ammonium, barium, calcium, magnesium, and aluminium, instead of potash, soda, ammonia, baryta, lime, magnesia, and alumina. The author confidently believes that this change, founded on views now adopted by all prominent writers on chemistry, will ultimately be accepted by all the various followers of medicine and pharmacy. It is a step in the direction of simplicity and consistency, and involves far less hypothesis than is contained in the old system. The name nitrate of potash, for example, was based on the pure assumption that nitre contained oxide of potassium or potash and nitric anhydride, then erroneously termed nitric acid. By the modern name, nitrate of potassium or potassium nitrate, all that is intended to be conveyed is that nitre contains the element common to all potassium compounds and the group of elements common to all nitrates. Under the old method students always experienced difficulty in distinguishing salts of the metal from salts of its oxide—salts of potassium, for instance, from salts of potash; under the new view no such difficulty arises. The contractions in Latin for names like “sulphate of magnesium,” *magnesi sulphas*, and “magnesium sulphate,” *magnesium sulphuricum*, are identical with the contractions for names resembling “sulphate of magnesia,” *magnesiæ sulphas*—an accidental circumstance that will facilitate the general introduction of either of the former names among the older medical practitioners and pharmacists, and a practical advantage that must determine the choice over the latter.

The author ventures to express some gratification that his use and advocacy of this system from 1867 onward resulted in its adoption in 1873 in the "Pharmacopœia of the United States," and in 1885 in the "British Pharmacopœia." Pharmacy amongst English-speaking nations will thus sooner or later, in the important matter of chemical nomenclature, be in accord with the teachings of chemical science.

Respecting chemical nomenclature generally, the one characteristic demanded by the best interests alike of pure and of applied chemistry should be permanence. The author therefore, whenever possible, has avoided the employment of names which include numeral syllables, any doubling or halving of atomic weights that the progress of research may necessitate at once rendering such names useless, nay, misleading. Instead of "platinum tetrachloride" he has written "platinic chloride" or "platinum perchloride," and so on.

The metric system of weights and measures (that which, doubtless, is destined to supersede all others) is alone used in the sections on quantitative analysis. In other parts of the Manual avoirdupois weights and imperial measures are employed, necessarily.

It is hoped that the numerous etymological references scattered throughout the following pages will be found useful. Words in Greek continue to be rendered, by special request, in English characters, letter for letter. The word "official" is used throughout for things recognized officially by the compilers of pharmacopœias; "officinal," in its original application to the officina, or shop—restrictions of meaning which have been advocated in every edition of this Manual since 1869, and which have been adopted in the recently-issued United States Pharmacopœia.

Chemical substances recognized in the United States Pharmacopœia, but not in the British Pharmacopœia, have, nevertheless, a certain amount of notice in the British editions of the Manual, and the chemical substances official in Great Britain are noticed in the American editions. The author hopes that this slight broadening of the horizon of his readers

and students, while enabling him more fully to illustrate chemical principles, will perhaps have influence in promoting extended as well as concurrent applications of chemistry to pharmacy in the great English-speaking countries.

Students are strongly recommended to test their progress by frequent examination. To this end appropriate questions are appended to each subject.

The author's ideal of a manual of chemistry for medical and pharmaceutical students is one in which not only the *science* of chemistry is taught, but in which the chemistry of every substance having interest for the followers of medicine and pharmacy is noticed at more or less length in proportion to its importance, and at least its position in relation to the leading principles of chemistry is set forth with all attainable exactness. The extent to which he has realized this ideal he leaves to others to decide. Such a work will doubtless in certain parts partake of the character of a dictionary; but this is by no means a fault, especially if a good index be appended, for the points of contact between pure and applied chemistry are thus multiplied, and abundant outlets supplied by which a lover of the science may pass into other chemical domains by aid of other guides, or even into the regions of original research. Among the rarer alkaloids, bitter bodies, glucosides, salts of organic radicals, solid fats, fixed oils, volatile oils, resins, oleo-resins, gum-resins, balsams, and coloring-matters mentioned in this volume, will be found many such points whence the ardent student may start for the well-known, the less-known, or the untrodden paths of scientific chemistry.

Within twenty-six years a demand has arisen for fifteen large editions of this Manual. The First, in 1867, was intended as a handbook of practical chemistry only, but the notes and remarks made respecting most of the experiments were found to be so useful by students that this portion of the volume was in the Second Edition (1869) sufficiently extended to render the book more fairly complete in itself. It and all subsequent editions included the chemistry of the British Pharmacopœia. In response to a call from pro-

professional friends in the United States in 1870, the work was revised by the author for the followers of Medicine and Pharmacy in America, descriptions of the Chemistry of the Preparations and Materia Medica of the United States Pharmacopœia being introduced, while those specially British were curtailed, and such other adaptations were included as to form a Third Edition. A Fourth was presented to English workers in 1872, and, founded on the fourth, a Fifth Edition for American students in 1873. A very large Sixth Edition was published in England in 1875, and in America a Seventh in 1876, and an Eighth in 1879, the Sixth and all following Editions containing notices of substances official in Waring's Indian Pharmacopœia. The Ninth, 1881, was an English Edition; the Tenth, 1883, an American; the Eleventh, 1885, English; the Twelfth, 1889, American; the Thirteenth, 1889, English.

The present, Fourteenth, Edition, and the concurrently produced Fifteenth, British Edition, contain such alterations and additions as seemed necessary for the demonstration of the latest developments of chemical principles and the latest applications of chemistry to pharmacy. Hence, even if compared with the immediately preceding editions, on nearly every page will be found some indication of the recent rapid growth of the science and the art of chemistry. On the other hand, synthetical and analytical processes which have become more or less superseded have been either omitted or described less fully. The whole work has thus been kept within the limits of a learner's manual, while the author has endeavored faithfully to portray the present relationship of chemistry to at least those areas of medicine and pharmacy with which students should become familiar in the days of their studentship. The voluminous Index will maintain the usefulness of the book to them afterward in the course of the practice of their respective callings.

The author is indebted to his son Dr. D. Harvey Attfield, M. A., D. P. H., Cantab., and to Mr. Charles S. Ellis, for much valuable help in preparing this edition for the press.

17 BLOOMSBURY SQUARE, LONDON.

ADVICE TO STUDENTS

RESPECTING THEIR OBJECT IN STUDYING.

AVOID studying chemistry, or indeed any subject, *merely* by way of "preparation for examination;" all ordinary "examinations" being, admittedly, inefficient tests of competency. Do not so mistake the means for the end. You are studying to fit yourself for your position in the world. Work diligently, study thoughtfully and deliberately; above all, be thorough, otherwise your knowledge will be inaccurate and transient, and will be unaccompanied by that enlightenment of the understanding, that mental training, mental discipline, and general elevation of the intellect, which constitute, in a word, education. When you are thus educated you will with ease and pleasure pass any examination in the knowledge you have thus acquired.

All authorities on education, whether statesmen, teachers, or examiners, regard "examinations," even by the most highly skilled "Board," with ample time at its disposal and a wide area from which to select questions, as but a partial test of knowledge and an imperfect test of education. It is the least unsatisfactory, however, that has been devised, and is especially useful when, following instead of leading education, it is restricted to the subjects of a well-defined, earnestly followed, compulsory public curriculum of study—a curriculum directed by a competent representative body, administered by properly qualified teachers, and followed by pupils who have had sound preliminary training.

Students! in all honor and in the highest self-interest take care that any inefficiencies inseparable from "examination" are abundantly compensated by the extent and precision of your knowledge and by the soundness and thoroughness of your whole education.

APPARATUS.

LIST OF APPARATUS FOR EXPERIMENTS IN ANALYSIS.

List of apparatus suitable for the three months' course of practical chemistry in the summer session of medical schools or for any similar series of lessons—including the preparation of elementary gases, analytical reactions of common metals and acidulous radicals, analysis of single salts, chemical toxicology, and the examination of urine, urinary sediments, and calculi:

One dozen test-tubes.	A 2-inch and a 3-inch evaporating-basin.
Test-tube stand.	Two porcelain crucibles.
Test-tube cleaning-brush.	Blowpipe.
A few pieces of glass tubing, eight to sixteen inches long, with a few inches of india-rubber tubing to fit.	Crucible tongs.
Small flask.	Round file.
Two small beakers.	Triangular file.
Two small funnels.	Small retort-stand.
Two watch-glasses.	Sand-tray.
Two or three glass rods.	Wire triangles.
Wash-bottle.	Platinum wire and foil.
Small pestle and mortar.	Test-papers.
A 2-pint earthenware basin.	Filter-paper.
	Towel.
	Two dozen corks.

(This set, packed in a case, can be obtained of any chemical-apparatus maker for about seven dollars.)

LIST OF APPARATUS FOR EXPERIMENTS IN SYNTHESIS AND ANALYSIS.

A larger set, suitable for the performance of most of the synthetic as well as qualitative analytical experiments described in this Manual:

A set of evaporating-basins, of the following sizes:	Two soup-plates.
Two 3-inch; one 4-inch; one 6½-inch; one 7½-inch; one 8½-inch.	One flat-plate.
One retort-stand and three rings.	Two spatula knives.
Two test-glasses.	One pair of scissors.
One half-pint flask.	One round file.
Half a quire of filter-paper.	One triangular file.
Two porcelain crucibles.	Half a pound of glass rod.
One measure-glass, 5 oz.	Half a pound of glass tubing.
Blowpipe, 8-inch.	One foot of small india-rubber tubing.
Two glass funnels.	Three dozen corks of various sizes.
One dozen test-tubes (hard glass).	Platinum wire and foil.
One test-tube brush.	Test-papers.
One pair of 8-inch brass crucible tongs.	A nest of three beakers.

(This set, packed in a case, can be obtained of any chemical-apparatus maker for about twelve dollars.)

A sponge, towels, and a note-book may be included.

LIST OF FURNITURE OF A CHEMICAL LABORATORY.

The following apparatus should be ready to hand for students following an extended course of practical chemistry, in a room set apart for the purpose:

A bench or table and stool.	Test-tube rack, two dozen holes.
Water-supply and waste-pipe.	Iron stand or cylinder for supporting large dishes.
A cupboard attached to a chimney with an outward draught.	Iron adapters for fitting dishes to cylinder.
A furnace fed with coke; tongs, hot-plate or sand-bath, etc.	Pestle and mortar, 5 or 6 inches.
A waste-box.	One 6-inch funnel.
Shelves for chemical and other materials in jars or bottles.	Brown pan, 1- or 2-gallon.
Gas-supply and lamp with flexible tube (or a spirit-lamp and spirit).	White jug, 1-gallon.
	Water-bottle, quart.
	Twenty-eight test-bottles, 6-oz.

Other articles, such as flasks, retorts, receivers, condensers, large evaporating-dishes, may be obtained as wanted. In Quantitative Analysis the apparatus described in the sections on that subject will be required.

LIST OF FLUID REAGENTS.

Certain chemical substances are used so frequently in analytical processes that it is desirable to have small quantities placed in bottles in front of the operator. (See p. 18.) As these reagents are generally employed in a state of solution, nearly all the solid salts may at once be dissolved (in distilled water). The bottles employed should be well stoppered, and of 5 or 6 ounces capacity. Common glass bottles of this size may be had for about one dollar and a quarter per dozen. The bottles should not be more than about three-fourths full; single drops, if required, can then be poured out with ease and precision. The following list of test-solutions is recommended; directions for methods of preparing the substances not readily purchasable will be found by referring to the Index:

Sulphuric Acid, strong.	Sol. of Potash, 5 per cent. or U.S.P.
Nitric Acid, strong.	“ Soda, 5 to 15 per cent.
Hydrochloric Acid, strong.	“ Ammon., 10 per cent. or U.S.P.
Acetic Acid, strong.	Lime-water, saturated.

The next nine may contain about 10 per cent. of solid salt:

Ammonium Carbonate, with a little solution of Ammonia added.	Ammonium Sulphhydrate.
Ammonium Chloride.	Barium Chloride.
Ammonium Phosphate or Arsenate.	Calcium Chloride.
	Sodium Phosphate.
	Neutral Chromate.

The succeeding seven may have a strength of about 5 per cent :

Potassium Ferrocyanide.	Iron Perchloride.
Potassium Ferricyanide.	Silver Nitrate.
Potassium Iodide.	Platinum Perchloride.
Ammonium Oxalate.	

LISTS OF SOLID CHEMICAL SUBSTANCES FOR STUDY.

List of chemical substances necessary for the practical study of the non-metallic elements mentioned on pp. 14 to 32. The quantities are sufficient for several experiments :

Potassium Chlorate	1 oz.	Phosphorus	$\frac{1}{2}$ oz.
Black Manganese Oxide	1 oz.	Hydrochloric Acid	1 oz.
Zinc	1 oz.	Sulphur	$\frac{1}{2}$ oz.
Oil of Vitriol	2 oz.	Iodine	$\frac{1}{4}$ oz.

List of chemical substances necessary for the *analytical* study of the important metallic and acidulous radicals (pp. 61 to 381). The quantities will depend on the frequency with which experiments are repeated or analyses performed ; those mentioned are sufficient for one or two students. The articles are given in the order in which they will be required. The eight substances mentioned in the above list are included :

The set of test-solutions described in the previous section :		Black Manganese Oxide	$\frac{1}{2}$ lb.
Potassium Carbonate	1 oz.	Manganese Chloride	$\frac{1}{4}$ oz.
Tartaric Acid	1 oz.	Cobalt Chloride	50 grs.
Litmus	$\frac{1}{4}$ oz.	Nickel Nitrate	$\frac{1}{4}$ oz.
Magnesium Sulphate	1 oz.	Chromium Chloride	$\frac{1}{4}$ oz.
Zinc Sulphate	1 oz.	Gold-leaves	2 or 3.
Alum	1 oz.	Cadmium Chloride	$\frac{1}{4}$ oz.
Ferrous Sulphide	1 lb.	Bismuth Nitrate	$\frac{1}{4}$ oz.
Oak-galls	1 oz.	Potassium Bromide	$\frac{1}{4}$ oz.
Potassium Sulphocyanate	$\frac{1}{4}$ oz.	Starch	1 oz.
White Arsenic	$\frac{1}{2}$ oz.	Potassium Nitrate	1 oz.
Zinc	$\frac{1}{2}$ lb.	Copper borings or turnings	1 oz.
Charcoal	$\frac{1}{2}$ lb.	Indigo	$\frac{1}{4}$ oz.
Ferrous Sulphate	1 oz.	Potassium Chlorate	1 oz.
Copper-foil	1 oz.	Iodine	$\frac{1}{2}$ oz.
Copper Sulphate	1 oz.	Spirit of Wine	1 oz.
Tartar Emetic	$\frac{1}{2}$ oz.	Sulphur	1 oz.
Mercury	1 oz.	Acid Potassium Oxalate	1 oz.
Corrosive Sublimate	$\frac{1}{2}$ oz.	Citric Acid	1 oz.
Calomel	$\frac{1}{2}$ oz.	Phosphorus	1 oz.
Tin	1 oz.	Borax	1 oz.
Potassium Bicarbonate	1 oz.	Turmeric	$\frac{1}{4}$ oz.
Lead Acetate	1 oz.	Benzoic Acid	50 grs.
Potassium Cyanide	$\frac{1}{2}$ oz.	Fluor Spar	1 oz.
Sodium Hyposulphite	1 oz.	Tannic Acid	50 grs.
A Lithium Salt	10 grs.	Gallic Acid	50 grs.
Strontium Nitrate	$\frac{1}{4}$ oz.	Pyrogallie Acid	50 grs.

The quantities of materials required for the study of chemistry *synthetically* will necessarily vary with the desires and tastes of the operator or according to the number and requirements of students working together.

The materials that will be needed for the home-study of *organic chemistry* will vary with the requirements of the student. By the time he has qualified himself for a preliminary experimental course in that section of the science he may trust largely to his own judgment as regards both materials and apparatus.

CONTRACTIONS USED IN THIS MANUAL.

B. P., British Pharmacopœia.	F., Fahrenheit.
U. S. P., United States Pharmacopœia.	gm., Gramme.
C., Centigrade.	mm., Millimètre.
cc., Cubic centimètres.	T. S., Test solution, U. S. P.
	V. S., Volumetric solution, U. S. P.

CONTENTS.

	PAGE
PREFACE	v
ADVICE TO STUDENTS	xiii
LISTS OF APPARATUS	xiv
LIST OF FURNITURE OF A CHEMICAL LABORATORY	xv
LIST OF FLUID REAGENTS	xv
LISTS OF SOLID CHEMICAL SUBSTANCES FOR STUDY	xvi
INTRODUCTION	13
GENERAL PROPERTIES OF THE NON-METALLIC ELEMENTS	15
SYMBOLS AND DERIVATION OF NAMES OF ELEMENTS	31
THE GENERAL PRINCIPLES OF CHEMICAL PHILOSOPHY	35
COMMON METALLIC ELEMENTS, THEIR OFFICIAL PREPARATIONS AND TESTS:	
Salts of Potassium, Sodium, Ammonium, Barium, Calcium, Magnesium, Zinc, Aluminium, Iron, Arsenum, Antimony, Copper, Mercury, Lead, Silver	60
ANALYTICAL CHARTS FOR ORDINARY METALS	224
RARER METALLIC ELEMENTS, THEIR OFFICIAL PREPARATIONS AND TESTS:	
Salts of Lithium, Strontium, Manganese, Cobalt, Nickel, Chromium, Tin, Gold, Platinum, Cadmium, Bismuth	226
ANALYTICAL CHARTS FOR ALL METALS	256
COMMON ACIDULOUS RADICALS, OFFICIAL ACIDS, AND TESTS:	
Chlorides, Bromides, Iodides, Cyanides, Nitrates, Chlorates, Acetates, Sulphides, Sulphites, Sulphates, Carbonates, Oxa- lates, Tartrates, Citrates, Phosphates, Borates	262
SALTS OF RARER ACIDULOUS RADICALS:	
Benzoates, Cyanates, Formates, Hippurates, Ferrocyanides, Ferri-cyanides, Fluorides, Hypophosphites, Hyposulphites, Lactates, Malates, Meconates, Metaphosphates, Nitrites, Phosphites, Pyrophosphates, Silicates, Sulphocyanates, Tannates, Gallates, Urates, Valerianates	335

	PAGE
ANALYTICAL CHART FOR ACIDULOUS RADICALS	369
SYSTEMATIC ANALYSIS	370
ORGANIC CHEMISTRY	383
CHEMICAL TOXICOLOGY	561
EXAMINATION OF MORBID URINE AND CALCULI	572
OFFICIAL GALENICAL PREPARATIONS	589
OFFICIAL CHEMICAL PREPARATIONS	591
QUANTITATIVE ANALYSIS:	
INTRODUCTORY REMARKS	591
DETERMINATION OF ATMOSPHERIC PRESSURE	593
DETERMINATION OF TEMPERATURE	594
DETERMINATION OF WEIGHT	600
WEIGHTS AND MEASURES	600
SPECIFIC GRAVITY	613
CORRECTION OF THE VOLUME OF GASES FOR PRESSURE AND TEMPERATURE	619
VOLUMETRIC QUANTITATIVE ANALYSIS	625
GRAVIMETRIC QUANTITATIVE ANALYSIS	659
DIALYSIS	704
APPENDIX:	
TABLE OF TESTS FOR IMPURITIES IN PREPARATIONS OF THE UNITED STATES PHARMACOPŒIA	707
SATURATION TABLES	724
THE ELEMENTS, THEIR SYMBOLS, AND ATOMIC WEIGHTS	725
INDEX	727

CHEMISTRY:

GENERAL, MEDICAL, AND PHARMACEUTICAL.

INTRODUCTION.*

MAN can neither create matter nor destroy matter; he can only alter its form—alter the relation of its elements to each other, or note similar alterations proceeding within and around him in Nature. To study those alterations in all their known length and breadth and depth is to study *Chemistry*.

The infinite varieties of solid, liquid, and gaseous matter of which our earth and atmosphere are composed may be resolved, with more or less difficulty, into a very few distinct substances appropriately termed “elements,” for by no known means can they be further decomposed. About seventy of these elements have been proved to exist. Some (such as gold) occur naturally in the uncombined state, but the greater number are combined in so subtle a manner as to conceal them from ordinary methods of observation. Thus, none of the common properties of water indicate that it is composed of two elements, both gases, but differing much from each other; nor can the senses of sight, touch, and taste, or other common means of examination, detect in their concealment the three elements of which sugar is composed. The art by which these and all other compound substances are resolved into their elements is termed *Chemistry*, a name derived possibly from the Arabic word *kamai*, to conceal.† The *art* of Chemistry also includes the

* Students using this book as a guide in following Chemistry practically should read the first four pages, and then commence work by preparing oxygen. All students should read the prefatory pages, especially the page of “Advice to Students.”

† The idea that common metals contained valuable metals concealed within them was the one seed from which mainly sprung chemical knowledge. The men who endeavored to find the secret of such concealment were appropriately termed *alchemists*, and their efforts spoken of as *alchemy* (*al kimia*, from *kamai*, to conceal). Their persistent labors, generation after generation, were unsuccessful so far as the transmutation of baser metals into gold was concerned, yet were invaluable to pos-

construction of compounds from elements, and the conversion of substances of one character into those of another. The general principles or leading truths relating to the elements, to the manner in which they severally combine, and to their properties when combined—that is, to the properties of the compound substances formed by their union—constitute the *science of Chemistry*.*

From these few words concerning the nature of the art and science of Chemistry it will be seen that in most of the occupations that engage the attention of man Chemistry plays an important part—in few more so than in the practice of therapeutics † and of pharmacy.‡

terity, for new substances were discovered and truths of nature unveiled : from these discoveries multiplication of discoveries resulted, and thus grew the still growing branch of knowledge called Chemistry.

* Persons who practise the art and science of Chemistry are known as chemists. Some two hundred or more years ago, and before Chemistry was a science, the only “chemists” were the makers and vendors of chemical substances, then only used as medicines. They were the successors of the alchemists. In Great Britain these chemists and the herb-dealers, otherwise drug-grocers, otherwise drug-gists, gradually associated to form the “chemist and druggist.” Between the “chemist and druggist” and the physician there existed the apothecary—the putter together of medicines or compounder of physicians’ prescriptions. The apothecary has since become a medical practitioner, prescriptions being now “made up” by the chemist and druggist. The latter in Great Britain, since the year 1868, has the title of *chemist and druggist*, his higher title being *pharmaceutical chemist*; these respective designations he legally assumes on passing the minor and major examinations conducted by the Pharmaceutical Society of Great Britain, in accordance with the provisions of the Pharmacy Acts of 1852 and 1868. The whole class is often spoken of as that of *pharmacists* or *pharmaceutists*, terms also used in the United States. Other classes of chemists are the analytical chemists, who give special attention to analysis; manufacturing chemists, who restrict their labors to the preparation of chemical substances; while others devote a portion of their knowledge and energies to chemical education or to chemical research, or are appealed to as consulting chemists by the persons, firms, corporations, or governments needing chemical advice respecting industrial processes, hygienic matters, etc. The callings of the consulting and analytical chemists are generally united, and the professional gentlemen who follow these conjoint avocations also not infrequently occupy professorial or other tutorial positions, sometimes adding to these labors more or less work at original chemical research. In England, Scotland, and Ireland, nearly all the leading professional chemists are Fellows of the Institute of Chemistry of Great Britain and Ireland.

† Therapeutics (θεραπευτικός, *therapeutikos*, from θεραπεύω, *therapeuō*, I nurse, serve, or cure) is that branch of medicine which treats of the application of remedies for diseases. The therapist also takes cognizance of *hygiene*—that department of medicine which respects the preservation of health—and of *dietetics*, the subject of diet or food. By *pharmacology* is understood the normal action of drugs upon the system as underlying the therapeutic action.—*Physics*, see p. 38.

‡ Pharmacy (from φάρμακον, *pharmakon*, a drug) is the generic name for

Air, water, food, drugs, and chemical substances—in short, all material things—are composed, as stated, of elements. An intimate knowledge of the properties of the more important elements, both in the free and in the combined state, and of the various substances they form when they have combined with each other, some knowledge or idea of the power or force (the chemical force or chemical affinity) by which the elements contained in the compounds are held together, and an application of such knowledge to pharmacy and medicine, must be the objects sought to be attained by the learner, for whom especially this book has been written.

The Elements.—Of the seventy or so known elements, about forty are of medical or pharmaceutical interest; of these, two-thirds are metals and one-third non-metals; the remainder* are so seldom met with in Nature as to have received no practical application either in medicine, art, or manufacture. Before intimately studying the elements it is desirable to acquire some general notions concerning them: such a procedure will also serve to introduce the practical student to his apparatus, and make him better acquainted with the various methods of manipulation.†

Metallic Elements.—With regard to the metallic elements it may safely be assumed that the reader has sufficient knowledge for present purposes; but little, therefore, need now be said respecting them. He has an idea of the appearance, relative weight, hardness, etc. of such metals as gold, silver, copper, lead, tin, zinc, and iron. If he has not a similar knowledge of mercury, antimony, arsenium, platinum, nickel, aluminium, magnesium, potassium, and sodium, he should embrace the ear-

the operations of preparing or compounding medicines, whether performed by the medical practitioner or by the chemist and druggist. It is also sometimes applied, like the corresponding term surgery, to the apartment in which the operations are conducted. *Pharmacognosy* is the study of the crude drugs of the vegetable and animal kingdom.

* A list of the elements will be found at the end of the volume.

† This allusion to apparatus need not discourage the youngest pupil. With the aid of a few phials, wine-glasses, or other similar vessels always at hand, he may, by studying the following pages, learn the chemical reactions which are constantly occurring in the course of making up medicine, understand the processes by which medicinal preparations are manufactured, and detect adulterations, impurities, or faults of manufacture. Among the substances used in medicine will be found nearly all the chemical materials required. If, in addition, a dozen test-tubes and a few feet of glass tubing be procured, many of the experiments described may be performed. For full lists of apparatus and chemical material see the introductory pages.

liest opportunity of seeing and handling specimens of each of these metals.

Non-Metallic Elements, wrongly termed Metalloids.—With regard to the non-metallic elements it is here supposed that the student has no general knowledge. He should commence his studies, therefore, by a series of operations as follows on eight of their number :

OXYGEN.

Preparation.—Oxygen is the most abundant element in nature, forming (in a state of combination) about one-half of the whole weight of our globe. To obtain it for experimental purposes all that is necessary is to apply heat—that force which will often be noticed as antagonistic, so to speak, to chemical union ; heat generally separating particles of matter farther from each other, while chemical attraction tends to bind them closer together—to heat certain compounds containing oxygen ; the latter is then evolved in its normal, natural gaseous condition. Several substances when heated yield oxygen ; but for convenience of students the crystalline body known as potassium chlorate is best fitted for the experiment. The size and form of the vessel in which to heat it will mainly depend on the quantity required, but for the purposes of the student the best is a *test-tube*, an instrument in constant requisition in studying practical chemistry. It is simply a tube of thin glass, a few inches in length and half or three-quarters of an inch in diameter, closed by fusion at one end. It is made of thin glass, in order that it may be rapidly heated or cooled without risk of fracture. (See Figs. 3 and 4.)

Outline of the Process.—Heat potassium chlorate (say as much as will lie on a shilling) in a test-tube by means of a spirit or gas flame ; gaseous oxygen is quickly evolved. Before applying heat, however, provision should be made for collecting the gas.

Collection of Gases (see Fig. 3).—Procure a piece of glass tubing about the thickness of a quill pen and a foot or eighteen inches long, and fit it to the test-tube by means of a cork in the following manner (longer tubes may be neatly cut to any size by smartly drawing the edge of a triangular file across the glass at the required point, then clasp the tube, the scratch being between the hands, and pulling the portions asunder, force being exerted in a *slightly* curved direction, so as to open out the crack which the file has commenced) : The tube is fixed in the cork through a round hole made by the aid of a red-hot wire, or, better, by a rat-tail file, or, best of all, by one of a set of cork-borers—pieces of brass tubing sharpened at one end and having a flat head at the other. Fit the cork and test-tube

to each other accurately and closely, but not so tightly as to break the test-tube. Setting aside the test-tube for a few minutes, proceed to bend the long piece of tubing to the most convenient shape for collecting the gas.

To Bend Glass Tubes.—Hold the part of the tube required to be bent in any gas or spirit flame (a fish-tail gas-jet, for example, Fig. 1), constantly rotating it, so that about an inch of the glass becomes heated. It will soon be felt to soften,

Fig. 1.




Softening and Bending Glass Tubes.

and will then, yielding to the *gentle* pressure of the fingers, assume any required angle. In the present case the tube should be heated at about four inches from the extremity to which the cork is attached, and bent to an angle of 90 degrees (Fig. 2).

Source of Heat.—The source of heat for the test-tube may be the flame of an ordinary spirit-lamp, or, still better where coal-gas is procurable, a mixture of the latter with air. Gas-lamps, especially constructed to burn a mixture of coal-gas and air, are sold by chemical-apparatus manufacturers. (See Figs. 3 and 7.)

Collection, etc. (continued).—Fit the cork and bent tube into the test-tube; the apparatus will then be ready for delivering gas at a convenient distance from the heated portion of the arrangement. To collect it, have ready three or four test-tubes (or small wide-mouthed bottles) filled with water, and inverted in a basin or other vessel, also containing water, taking care to keep the mouths of the still full tubes a little below the surface. Now apply heat to the chlorate contained in the test-tube, and so arrange the open end of the bent tube under the water, that the gas, which presently escapes with effervescence from the melted chlorate, may pass out from the free end of the tube, and may bubble into and gradually fill the previously

Fig. 2.



water-filled inverted test-tubes. The first tubeful may be rejected, as it probably consists of little more than the air

Fig. 3.



This engraving represents the preparation, collection, and storage of small quantities of oxygen gas. A test-tube and bent glass tube, joined together by a perforated cork, are supported by the arm of an iron stand. (The apparatus might be held by the fingers.) The tube is heated by a gas-lamp. (The spirit-lamp shown at back might be used instead.) Gas evolved from the heated substance in the test-tube is displacing water from an inverted test-tube. Spare tubes in a test-tube rack are at hand, and tubes already filled are set aside till wanted. A nest of cork-borers, a round file, a triangular file, and a test-tube cleaning brush are lying on the table or student's bench. Below are cupboards for apparatus; above are bottles containing testing liquids, etc.

originally in the apparatus, and which has been displaced by the oxygen. That which comes afterward will be pure oxygen.

As each tube or bottle becomes full, close its mouth (still under the surface of the water) by a cork, and then set it aside; or a little cup (such as a porcelain crucible or small gallipot) may be brought under the mouth, and the cup, with the mouth of the tube in it, be lifted out of the water and placed close by till wanted, the water remaining in the cup effectually preventing the gas from escaping.

On the large scale, oxygen may be made in the same way, larger vessels (glass flasks or iron bottles) being employed. Less heat also will be necessary if the potassium chlorate be previously mixed with very fine sand, or, still better, with about an equal weight of common black manganese oxide.

Note on the Collection and Storage of Gases.—It may be as well to state that nearly all gases, whether for experimental or practical purposes, are collected and stored in a similar manner. Even coal-gas is generated at gas-works in iron retorts very much the shape of test-tubes, only they are as many feet long as a test-tube is inches, and the well-known gigantic gas-holders may be viewed as inverted iron test-tubes of great diameter.

Properties.—Free oxygen is a colorless gas. Cailletet and Pictet succeeded in liquefying it, and Wroblewski and Obszewski have obtained it in some amount as a definite, colorless, transparent fluid. Air, which contains 20 per cent. of oxygen, has been solidified, but not yet the pure oxygen itself. Obviously, it is not very soluble in water, or the gas could not be collected by the aid of that liquid. Oxygen is soluble to a certain extent, however (about 3 volumes in 100 at common temperatures), or fishes could not breathe. Other noticeable negative features are its want of taste and smell.

To show the relation of oxygen to combustion, remove one of the tubes from the water by placing the thumb over its mouth, and apply for a second a lighted wood match to the orifice; the gas will be found to be incombustible. Extinguish the flame of the match, and then quickly introduce the still incandescent carbonaceous extremity of the wood halfway down the test-tube; the wood will at once burst into flame, owing to the extreme violence with which oxygen gas supports combustion. These tests of the presence of free oxygen may also be applied at the extremity of the delivery-tube whilst the gas is being evolved. (It is desirable to retain two tubes of the gas for use in subsequent experiments; also one tube in which only one-third of the water has been displaced by oxygen.)

Relation of Oxygen to Animal and Vegetable Life.—Not only the carbon at the end of a piece of charred wood, but any other sub-

stance that will burn in air (which, as will be seen presently, is diluted oxygen), will burn more brilliantly in pure oxygen. The warmth of the bodies of animals is kept up by the continuous burning of the tissues in the oxygen (of the air) drawn into the system through the lungs. The product of this combustion is exhaled into the air as a gaseous compound of carbon and oxygen termed carbonic acid gas—a gas which in sunlight is absorbed by and decomposed in the cells of plants, with fixation of the carbon and liberation of the oxygen; hence the atmosphere is kept constant in composition.

Memorandum.—At present it is not advisable that the reader should trouble himself with the consideration of the chemical action which occurs either in the elimination of oxygen from its compounds or in the separation of any of the following non-metallic elements from their combinations. It is to the properties of those elements themselves, especially in their free and least active condition, that he should at present restrict his attention. Working thus from simple to more complex facts, he will in due time find that the comprehension of such actions as occur in the preparation of these few elements will be easier than if he attempted their full study now.

HYDROGEN.

Preparation and Collection.—The element hydrogen is also, in the free state, a gas,* and is obtainable from its commonest compound, water (of which one-ninth by weight is hydrogen), by the agency of hot zinc or iron, but more conveniently by the action of either of those metals on cold diluted sulphuric acid. The apparatus used for making oxygen may be employed for this experiment, but no lamp is required. Place several pieces of thin zinc † in the generating-tube (Fig. 4), or in any common glass bottle (Fig. 5) or flask, and cover them with water. The collecting-tubes (these also may be wide-mouthed bottles) being ready, add strong sulphuric acid (oil of vitriol)

*Graham obtained alloys of hydrogen with palladium and other metals, compounds in which several hundred times its bulk of gas is retained by the metal *in vacuo* or even at a red heat. This was physical confirmation of the opinion long held by chemists, that hydrogen is a gaseous metal. Graham termed it *hydrogenium*, other chemists *hydrium*, and considered its relative weight in the solid state to be nearly three-fourths that of water. Cailletet and Pictet have since actually liquefied and solidified this element.

† The best form is *granulated zinc* (*Zincum*, U. S. P.), made by heating scraps of common sheet zinc in an iron ladle over a fire, and, immediately the metal is fused, pouring it, in a slow stream, into a pail of water from a height of eight or ten feet. Each drop of zinc thus yields a thin little bell, which, for its weight, presents a large surface to the action of the acid liquid. If the zinc is allowed to become hotter than necessary, the little bells will not be formed. A trace of iron in the zinc greatly increases the rate at which the hydrogen is evolved.

to the zinc and water, in the proportion of about 1 volume of acid to 5 of water, and fit on the delivery-tube; or pour the acid down such a funnel-tube* as is shown in Fig. 5; the

Fig. 4.



Fig. 5.



Preparation of Hydrogen.

hydrogen at once escapes with effervescence from the fluid. Having rejected the first portions (or having waited until the air originally in the bottle may be considered to be all expelled), collect four or five tubes of the gas in the manner described under Oxygen.

Note.—This process is similar to that of the “British Pharmacopœia.” In making larger quantities bottles of appropriate size may be employed. Other metals, notably potassium and sodium, liberate hydrogen the moment they come into contact with water, but the processes are not economical and the action is dangerously violent.

Properties.—Like oxygen, hydrogen gas is invisible, inodorous, and tasteless. If iron be used, the gas has a marked smell, but this is due to impurities derived from the iron.

Apply a flame to the mouth of the delivery-tube as soon as the operator's judgment tells him that the brisk effervescence of hydrogen must have resulted in the driving out of all air from the tube, for the *mixture* of hydrogen and air may explode. Ignition of the hydrogen gas ensues, showing that, unlike oxygen gas, it is combustible.

Plunge a lighted match well into a tube (or wide-mouthed bottle) containing free hydrogen; the gas is ignited, but the match becomes extinguished. This shows that hydrogen is not a supporter of combustion.

* Funnel-tubes may be purchased of the apparatus-maker, or, if the pupil has access to a table blowpipe and the advantage of a tutor to direct his operations, they may be made by himself.

Hydrogen gas in burning unites with the oxygen of the air and forms water, which may be condensed on a cool glass or other surface. Prove this by holding a glass vessel a few inches above a hydrogen flame. In burning the hydrogen contained in one of the tubes or bottles the flame is best seen when the tube is held mouth upward and water poured in so as to expel the gas gradually.

If, instead of this gradual combination of the two elements oxygen and hydrogen, they be mixed together in bulk in the right proportions and then ignited, they will rapidly combine, and explosion will result. Prepare a mixture of this kind by filling up with hydrogen a test-tube from which one-third of the water has been expelled by oxygen. Remove the tube from the water, placing a finger over the mouth, and, having a lighted match ready, apply the flame; explosion ensues, owing to the instantaneous combination of the whole bulk of the two elements and the expansive force of the highly heated steam produced. If anything larger than a test-tube is employed in this experiment, it should be a soda-water bottle or some such vessel equally strong.

Notes.—These gases thus unite at a temperature far higher than that of boiling water, 2 volumes of hydrogen and 1 of oxygen yielding 2 of gaseous water (true steam).

The noise of such explosions is caused by concussion between the suddenly expanded gaseous body and the air.

The force of the explosion—or, in other words, the force of the suddenly heated and therefore suddenly expanded steam—is below that necessary to break the test-tube. Some force, however, is exerted; and hence the necessity of the precaution previously suggested, of allowing all the air which may be in a hydrogen apparatus to escape before proceeding with the experiments. If a flame be applied to the delivery-tube before all the air is expelled, the probable result will be ignition of the mixture of hydrogen and oxygen (of the air) and consequent explosion. But even in this case the generating vessel is not often fractured, unless it be large and of thin glass, the ordinary effect being that the cork is blown out and the delivery-tube broken on falling to the ground.

Hydrogen is a constituent of all the substances used for producing artificial light, such as solid fats, oil, and coal-gas. The explosive force of large quantities, such as a roomful, of coal-gas and air, though vastly below that of an equal weight of gunpowder, is well known to suffice for blowing out that side of the room which offers least resistance.

The composition of water can be proved analytically as well as synthetically, a current of electricity decomposing it, by "electrolysis" (*λίσω, luo*, I loose, or I decompose), into its constituent gases, twice as much hydrogen as oxygen by volume being produced.


COMBUSTION (from *comburo*, I burn).—The experiments with hydrogen and oxygen illustrate the true character of combustion. Whenever chemical combination is sufficiently intense to be accompanied by heat and light, the materials are said to undergo combustion. Combustion only occurs at the line of contact of the combining bodies; a jet of oxygen will burn in an atmosphere of hydrogen quite as easily as a jet of hydrogen in oxygen. A jet of air (diluted oxygen) will burn as readily in a jar of coal-gas as a jet of coal-gas burns in air; each is combustible, each supports the combustion of the other. Hence the terms *combustible* and *supporter of combustion* are purely conventional, and only applicable so long as the circumstances under which they are applied remain the same. In the case of substances burning in air the conditions are, practically, always the same; hence no confusion arises from regarding air as the great supporter of combustion, and bodies which burn in it as being combustible.

Fig. 6.



Structure of Flames.

Fig. 7.



"Bunsen," or Air-Gas, Burner.

Structure of Flame.—A candle flame (Fig. 6) or oil flame is a jet of gas intensely heated; the central portion is unburnt gas; the next envelope is formed of partially burnt and very dense gaseous and solid particles sufficiently highly heated to give light; and the outer cone of completely burnt gases. In the figure the sharpness of limit of these cones is purposely somewhat exaggerated. Air made by any mechanical contrivance to mix with the gas in the interior of a flame at once burns up, or perhaps prevents the formation of, dense gases, giving a hotter but non-luminous jet. The air-gas lamps (Fig. 7), or "Bunsen" gas-burners, commonly used in chemical laboratories, are constructed on this principle: their flame has the additional advantage of not yielding a deposit of soot.

In the air-gas lamp coal-gas escaping from a small orifice draws rather more than twice its volume of air (supplied through adjacent holes) into its column, and the mixture of gas and air passes upward

along a pipe. It only burns at the end, and not within the pipe, partly because the metal of the burner, by conducting heat away, cools the mixture below the temperature at which it can ignite; partly because the velocity with which the mixture flows out is greater than the rate at which such a mixture ignites; and partly because the proportion of air to gas in the mixture is insufficient for perfect combustion, the external air contributing materially to the complete combustion of the jet of air-gas. The *Davy safety-lamp* acts on the principle first named: a wire-gauze cage surrounds an oil flame; an inflammable mixture of gas (fire-damp) and air can pass through the gauze and catch fire and burn inside; but the flame cannot, ordinarily, be communicated to the mixture outside, because the metal of the gauze and of the other parts cools down the gas below the temperature at which combustion can continue.

Properties (continued).—Gaseous hydrogen is the lightest substance known. It was formerly used for filling balloons, but was superseded by coal-gas, because coal-gas, though not lighter, is cheaper and more easily obtained. The lightness of hydrogen may be rendered evident by the following experiment: Fill two test-tubes with the gas, and hold one with its mouth downward and the other with its mouth upward. The hydrogen will have escaped from the latter in a few seconds, whereas the former will still contain the gas after the lapse of many seconds. This may be proved by applying a lighted match to the mouths.

The relative weight or specific gravity of oxygen is sixteen times that of hydrogen. A vessel holding 1 grain of hydrogen will hold 16 grains of oxygen. The relation of the weight of hydrogen to air is as 1 to 14.44, or as 0.0693 to 1.0. 1 grain of hydrogen by weight would measure about 27 fluidounces, and therefore would about fill a common wine-bottle. Such a bottle would, at ordinary temperatures, hold about $14\frac{1}{2}$ grains of air or 16 grains of oxygen.

Mem.—It is desirable to retain two tubes of hydrogen for use in subsequent experiments.

DIFFUSION OF GASES.—Hydrogen gas cannot be kept in such vessels as the inverted test-tube, for, though much lighter than air, it *diffuses* downward into the air, while the air, though much heavier, diffuses upward into the hydrogen. This power of *diffusion* is characteristic of all gases, and proceeds according to a fixed rate—namely, “in inverse proportion to the square root of the specific gravity of the gas” (*Graham*). Thus hydrogen diffuses four times faster than oxygen. This great and important property of diffusion strongly suggests that the particles of gases, at least, are always moving, never at rest; how otherwise *could* gases diffuse into each other as they do, notwithstanding the opposing influence of gravitation? Diffusion strongly supports this (*Clausius’s* kinetic (*κινῶ, kineō*, I move or put in motion) theory of the physical condition of gases.

PHOSPHORUS.

Appearance and Source.—Phosphorus (*Phosphorus*, U. S. P.) is a solid element, in appearance and consistence resembling white wax, but it gradually becomes yellow by exposure to light. It is a characteristic constituent of bones, and may be prepared from bones by a process which will be described subsequently.

Caution.—Phosphorus, on account of its great affinity for oxygen, takes fire very readily in the air, and should therefore be kept under water. When wanted for use it must be cut under water. It is employed in tipping lucifers, though *red* or *amorphous phosphorus* (*vide* Index) is less objectionable for this purpose.

Experiment.—Dry a piece about one-fourth the size of a pea by quickly and carefully pressing it between the folds of porous (filter- or blotting-) paper; place it on a plate, and ignite by touching it with a piece of warm wire or wood. The product of combustion is a dense white suffocating smoke, which must be confined at once by placing an inverted tumbler, or beaker, or other similar vessel over the phosphorus. The fumes rapidly aggregate, and fall in white flakes on the plate. When this has taken place and the phosphorus is no longer burning, moisten the powder with a drop or two of water, and observe that some of the water is converted into steam, an effect due to the intense affinity with which another portion of the water and the powder have combined, yielding heat.

The powder produced by the combustion of phosphorus is termed phosphoric anhydride; the combination of the latter with the elements of water produces a variety of phosphoric acid which dissolves in the water, forming, on standing, a dilute solution of ordinary phosphoric acid. The diluted phosphoric acid of the British and American Pharmacopœias is a similar solution, made in a somewhat different way and of definite strength.

NITROGEN.

Source.—The chief source of this gaseous element is the atmosphere, nearly four-fifths of which consists of nitrogen (the remaining fifth being almost entirely oxygen).

Preparation.—Burn a piece of dried phosphorus, the size of a pea, in a confined portion of air. The oxygen gas is thus removed, and the nitrogen gas remains. The readiest mode of performing this experiment is to fix a piece of earthenware (the lid of a small porcelain crucible answers very well) on a thin piece of cork, so that it may float in a dish of water (Fig. 8). Place the phosphorus on the lid, ignite with a warm rod, and then invert a tumbler or any glass vessel of about a half-pint capacity over the burning phosphorus, so that the


mouth of the glass may dip into the water. Let the arrangement rest for a short time, for the fumes of phosphoric anhydride to subside and dissolve in the water, and then decant the gas into test-tubes as indicated in Fig. 9, using a tub or other vessel of water of sufficient depth to admit of the glass containing the nitrogen gas being turned on one side without air gaining access.

Fig. 8.



Preparation of Nitrogen.

Fig. 9.



Decantation of Gases.

Larger quantities of nitrogen gas are obtained in the same way. Other combustibles, as sulphur or a candle, might be used to burn out the oxygen gas from the air, but none answers so quickly and completely as phosphorus; added to which the product of their combustion would not always be dissolved by water, but would remain with the nitrogen.

Mem.—The statement concerning the composition of the air is roughly confirmed in isolating nitrogen, about one-fifth of the volume of the air originally in the glass vessel having disappeared, its place being occupied by water.

Properties.—Like oxygen and hydrogen, nitrogen gas is invisible, tasteless, and inodorous. By pressure, Cailletet and Pictet condensed it to a liquid. Wroblewski and Oboszewski obtained it in some amount as a definite, colorless, transparent fluid, which congeals, by its own evaporation, to a white snow-like solid. It is only slightly soluble in water. *Free* nitrogen is distinguished from all other gases by the absence of any very characteristic or positive properties. Apply a flame to some contained in a tube; it will be found to be incombustible. Immerse a lighted match in the gas; the flame is extinguished, showing that nitrogen is a non-supporter of combustion.

Nitrogen is fourteen times as heavy as hydrogen.

The chief office of the free nitrogen in the air is to dilute the energetic oxygen, a mere *mechanical* mixture resulting.

The air is nearly fourteen and a half (14.44) times as heavy as hydrogen. It may be liquefied and, apparently, solidified. Its average composition, including minor constituents (which will be referred to subsequently), is as follows:

Composition of the Atmosphere.

	In 100 volumes.
Oxygen	20.61
Nitrogen	77.95
Carbonic acid gas	.04
Aqueous vapor	1.40
Nitric acid	} traces.
Ammonia	
Carburetted hydrogen	} traces in
Sulphuretted hydrogen	
Sulphurous acid	towns.

Pure dry air free from carbonic acid *invariably* contains, by weight, 23 parts of oxygen to 77 of nitrogen, or, by volume, 20.76 parts of oxygen to 79.24 of nitrogen. Ozone (*vide* Index) is said to be a normal constituent of air.

Free Nitrogen and Combined Nitrogen.

The comparative inactivity or negative character of nitrogen in its free condition—that is, when uncombined with other elements—contrasts strongly with its apparent influence in a state of combination. When its compounds with hydrogen come to be studied, it will be found to be, apparently, the chief or leading, or, in a sense, the most important, element of those compounds—the ammoniacal compounds. United with carbon, it gives the poisonous cyanic substances. With oxygen it yields quite a large group of bodies, amongst which are the common and important class of salts termed nitrates. With carbon, as well as hydrogen and some oxygen, it affords powerful agents termed alkaloids—near relatives of ammonia; while the same elements otherwise grouped, and sometimes a little sulphur or phosphorus, form the various albumenoid and gelatinoid matters characteristic of the tissues of animals and vegetables. In a perfect structure we should perhaps scarcely regard any one element or member as more important than another; still, such a conclusion almost forces itself upon us as we become acquainted with the chemical history of combined nitrogen. Free nitrogen is not, however, altogether inactive, for the nitrogen of the air appears to be absorbed and assimilated by plants, a given crop containing more nitrogen than the soil and manure whence it grew.

CHLORINE.

Source.—In the free state this element is a gas. Its chief source is common salt, more than half of which is chlorine.

Preparation.—About a quarter of an ounce of salt and the same amount of black manganese oxide are mixed, and placed in a test-tube with sufficient water to cover them; on adding a

small quantity of sulphuric acid the evolution of chlorine gas commences. For the mode of collection see the following paragraphs.

Another Process.—As the action of the sulphuric acid on the salt in the above process is mainly to give hydrochloric acid, the latter acid (about 4 parts) and the black manganese oxide (about 1 part) may be used in making the gas, instead of salt, sulphuric acid, and black manganese oxide. This, the usual process, is that adopted in the British and United States Pharmacopœias.

Collection and Properties.—Free chlorine is a suffocating gas. Care therefore must be observed in experimenting with this element. As soon as its penetrating odor indicates that it is escaping from the test-tube, the cork and delivery-tube (similar to that used in making oxygen) should be fitted on, and the gas passed to the bottom of another test-tube containing water (Fig. 10). When thirty or forty small bubbles have passed, their evolution being assisted by slightly heating the generating tube, the latter should be removed to the cupboard usually provided in laboratories for performing operations with noxious gases, or be dismantled and the contents carefully and rapidly

Fig. 10.





Fig. 11.



Preparation of Chlorine.

washed away. The water in the collecting-tube will now be found to smell of the gas, chlorine being, in fact, soluble in about half its bulk of water. Chlorine-water is official* in

* "The Pharmacopœia and all in it is official (*office*, Fr. from *L. officium*, an office). There are many things which in Pharmacy are official (Fr. from *L. officina*, a shop), but not official. To restrict the word *official* to the contents of a pharmacist's shop, and to that portion of the contents which is pharmacopœial, is radically wrong, and should be avoided."—J. A., second edition, 1869.

the British and United States Pharmacopœias (*Aqua Chlori*, U. S. P.).

Larger quantities may be made from the hydrochloric acid and black manganese oxide (4 to 1) in a Florence flask fitted with a delivery-tube, the flask being supported over a flame by the ring of a retort-stand or any similar mechanical contrivance (Fig. 11). A piece of cardboard on the neck of the collecting-bottle, as indicated in the figure, retards diffusion of the chlorine gas from the bottle during the process of collection.

Mem.—Flasks and similar glass vessels are less liable to fracture if protected from the direct action of the flame by being placed on a piece of wire gauze two to four inches square, or on a *sand-bath*; that is, a saucer-shaped tray of sheet iron on which a *thin* layer of sand is placed.

The *Vapor Chlori*, B. P., or inhalation of chlorine, is simply moist chlorinated lime so placed that some of the chlorine given off may be inhaled.

During these manipulations the operator will have noticed that chlorine is of a light yellowish-green color. The tint is observable when the gas is collected in large vessels. As it is soluble in water ($2\frac{1}{2}$ vols. in 1 vol. at 60° F.), it cannot be economically stored over that liquid. Being, however, nearly twice and a half as heavy as air, the gas may be collected by simply allowing the delivery-tube to pass to the bottom of a dry test-tube or dry bottle (Fig. 11).

A distinctive property of free chlorine is its bleaching power. Prepare some colored liquid by placing a few chips of logwood or other dyeing material in a test-tube half full of hot water. Pour off some of this red infusion into another tube and add a few drops of the chlorine-water; the red color is rapidly destroyed.

Free chlorine readily decomposes offensive effluvia; it is one of the most powerful of *deodorizers*. It also decomposes putrid and infectious matter; it is one of the best of *disinfectants*. (*Antiseptics* are substances which *prevent* putrefaction. See Index.)

"An *official* formula is one given under authority. An *official* formula is one made in obedience to the customary usage of the shop (*officina*). To state that any preparation under the sanction of the British Pharmacopœia is *official* is a misapprehension of the meaning of the word."—J. BROUGH.

"That is *official* which emanates from a recognized authority. That is *official* which is issued from an *officina* or workshop."—J. INCE.

"*Official* writings and orders are those issued by official persons. *Official* articles are such as are found in a shop."—J. F. STANFORD, M.A., F. R. S.

"It will be observed that the word '*official*' has been used in this edition of the Pharmacopœia in place of the word '*official*.' This change was made by a special vote of the committee at one of its first meetings in 1890."—UNITED STATES PHARMACOPEIA.

Combination of Hydrogen with Chlorine, forming Hydrochloric Acid.—If an opportunity occurs of generating chlorine in a closed chamber or in the open air, a test-tube, of the same size as one of those in which hydrogen has been retained from a previous operation, is filled with the gas. The hydrogen tube is then inverted over that containing the chlorine, the mouths being kept together by encircling them with a finger. After the gases have mixed, the mouths of the tubes are quickly in succession brought near a flame, when explosion occurs, and fumes of a compound of hydrochloric acid gas with the moisture of the air are formed. The hydrochloric acid of pharmacy (*Acidum Hydrochloricum*, U. S. P.) is a solution of the gas (made in a more economical way) in water.

The foregoing experiment affords evidence of the powerful affinity of chlorine and hydrogen for each other. Chlorine dissolved in water will, in sunlight, slowly remove hydrogen from some of the water and liberate oxygen. The bleaching power of chlorine is generally referred to this indirect oxidizing effect which it produces in presence of water, for dry chlorine does not bleach.

Density.—Chlorine gas is thirty-five and a half times as heavy as hydrogen gas. A wine-bottle would hold about $35\frac{1}{2}$ grains.

SULPHUR, CARBON, IODINE.

The physical properties (color, hardness, weight, etc.) possessed by these elements when they are in the free state are familiar. Their leading chemical characters in the free state will also be understood when a few facts concerning each are made the subject of experiment.

SULPHUR.—Burn a small piece of sulphur; a penetrating odor is produced, due to the formation of a colorless gas. The product is a perfectly definite chemical compound of the oxygen from the air with the sulphur. It is termed sulphurous anhydride or sulphurous acid gas.

CARBON is familiar, in the free form, as soot, coke, charcoal, graphite (or plumbago, popularly termed black lead), and diamond. The presence of combined carbon in wood and in other vegetable and animal matter is at once rendered evident by heat. Place a little tartaric acid on the end of a knife in a flame; the blackening that occurs is due to the separation of carbon. The black matter at the extremity of a piece of half-burned wood is also free carbon.

Carbon, like hydrogen, phosphorus, and sulphur, has a great affinity for oxygen at high temperatures. A striking evidence of

that affinity is the evolution of sufficient heat to make the materials concerned red or even white hot. When ignited in the diluted oxygen of the air, carbon simply burns with a moderate glow, as seen in an ordinary coke or charcoal fire; but when ignited in pure oxygen, the intensity of its combination is greatly exalted. The product of the combination of the two elements, if the oxygen be in excess, is an invisible gaseous body termed carbonic acid gas; if the carbon be in excess, another invisible gas, termed carbonic oxide, results.

IODINE.—A prominent chemical characteristic of free iodine is its great affinity for metals. Place a piece of iodine about the size of a pea in a test-tube with a small quantity of water, and add a few iron filings or small nails. On gently warming this *mechanical* mixture, or even shaking if longer time be allowed, the color and odor of the iodine disappear; it has *chemically* combined with the iron—a *chemical compound* has been produced. If the fluid be filtered, a clear aqueous solution of the compound of the two elements is obtained.

The solution, made as above and mixed with sugar, forms, when of a certain strength, the ordinary syrup of iodide of iron of pharmacy (*Syrupus Ferri Iodidi*, U. S. P., or syrup of ferrous iodide). A strong solution mixed with sugar and liquorice-root constitutes the corresponding pills (*Pilule Ferri Iodidi*, U. S. P., or pills of ferrous iodide). The solid iodide is obtained on removing the water of the above solution by evaporation.

Sulphur and Iron, also, when very strongly heated, *chemically combine* to form a substance which has none of the properties of a *mixture* of sulphur and iron; that is, has none of the characters of sulphur and none of iron, but new properties altogether. The product is termed *Ferrous Sulphide*, U. S. P., or sulphide of iron. Its manufacture and uses will be alluded to in treating of the compounds of iron; it is mentioned here as a simple but striking illustration of the difference between a *chemical compound* and a *mechanical mixture*.

THE ELEMENTS, THEIR SYMBOLS, ETC.

From the foregoing statements a general idea will have been obtained of the nature of several of the more frequently occurring free elements. Some additional facts concerning them may be gathered from the following table, which gives the name in full, the symbol (or shorthand character*) of the name, and the origin of the name.

For the purposes of study by medical and pharmaceutical pupils the elements may be divided into three classes—viz. those frequently, those seldom, and those never used in pharmacy.

* The symbol is also much more than the shorthand character, as will presently be apparent.

Name.	Symbol.	Derivation of Name.
Oxygen	O	From ὄξυς (oxūs), <i>acid</i> , and γένεσις (genesis), <i>generation</i> , i. e. generator of acids. When first discovered it was supposed to enter into the composition of all acids.
Hydrogen . . .	H	From ὑδωρ (hudōr), <i>water</i> , and γένεσις (genesis), <i>generation</i> , in allusion to the product of its combustion in air.
Nitrogen . . .	N	From νίτρον (nitron), and γένεσις (genesis), <i>generator of nitre</i> .
Carbon	C	From carbo, <i>coal</i> , which is chiefly carbon.
Chlorine	Cl	From χλωρὸς (chlōros), <i>green</i> , the color of this element.
Iodine	I	From ἰὼν (ion), <i>a violet</i> , and εἶδος (eidōs), <i>likeness</i> , in reference to the color of its vapor.
Sulphur	S	From sal, <i>a salt</i> , and πῦρ (pur), <i>fire</i> , indicating its combustible qualities. Its common name, <i>brimstone</i> , has the same meaning, being the slightly altered Saxon word, <i>brynstone</i> , i. e. burnstone.
Phosphorus . .	P	Φῶς (phōs), <i>light</i> , and φέρειν (pherein), <i>to bear</i> . The light it emits may be seen on exposing it in a dark room.
Potassium . . . (Kalium)	K	<i>Kalium</i> , from <i>kali</i> , Arabic for <i>ashes</i> (see next paragraph). Manufactories in which compounds of potassium and allied sodium-salts are made are called <i>alkali-works</i> to this day. <i>Potassium</i> , from <i>pot-ash</i> , so called because obtained by evaporating the lixivium of wood-ashes in pots.
Sodium (Natrium)	Na	<i>Natrium</i> , from <i>natron</i> , the old name for certain natural deposits of carbonate of sodium. <i>Sodium</i> , from <i>soda-ash</i> or <i>sodash</i> , the residue of the combustion of masses or <i>sods</i> of marine plants. Sod-ashes were chemically distinguished from pot-ashes by Duhamel in 1736. Previously both were simply <i>kali</i> or ashes from two different sources. Sir Humphry Davy first isolated the two metals in 1807.
Ammonium . .	NH ₄	This body is not an element, but its components exist in all ammoniacal salts, and apparently play the part of such elements as potassium and sodium. Sal ammoniac (chloride of ammonium) was first obtained from near the temple of Jupiter Ammon in Libya; hence the name.
Barium	Ba	From βαρὺς (barūs), <i>heavy</i> , in allusion to the high specific gravity of "heavy spar," the most common of the barium minerals.
Calcium	Ca	<i>Calx</i> , <i>lime</i> , the oxide of calcium.
Magnesium . .	Mg	From <i>Magnesia</i> , the name of the town (in Asia Minor) near which the substance now called "native carbonate of magnesium" was first discovered.
Iron (Ferrum)	Fe	Prehistoric. The spelling may be from the Saxon <i>iren</i> , the pronunciation from the

Name.	Symbol.	Derivation of Name.
Aluminium . .	Al	Gothic " <i>iarn</i> ." The derivation is perhaps Aryan; it probably originally meant <i>metal</i> . The metallic basis of alum was at first confounded with that of sulphate of iron, which was the alum of the Romans, and was so called in allusion to its tonic properties, from <i>alo</i> to <i>nourish</i> .
Zinc	Zn	From Ger. <i>Zinn</i> , <i>tin</i> , with which zinc seems at first to have been confounded.
Arsenium, Arsenicum, or Arsenic	As	<i>Ἀρσενικόν</i> (<i>arsenikon</i>), the Greek name for orpiment, a sulphide of arsenium. Common <i>white</i> " <i>arsenic</i> " is an oxide of arsenium. <i>Arsenium</i> as the name of the <i>element</i> , and <i>arsenic</i> as the old and widely recognized name of the common <i>white oxide</i> , are the <i>distinctive</i> names which science and expediency alike suggest.
Antimony . . . (Stibium)	Sb	<i>Στίβι</i> (<i>stibi</i>), or <i>στίμμι</i> (<i>stimmi</i>), was the Greek name for the native sulphide of antimony. The word <i>antimony</i> is said to be derived from <i>ἀντί</i> (<i>anti</i>), <i>against</i> , and <i>moine</i> , French for <i>monk</i> , from the fact that certain monks were poisoned by it.
Copper (Cuprum)	Cu	From <i>Cyprus</i> , the name of the Mediterranean island where this metal was first worked.
Lead (Plumbum)	Pb	The Latin word is expressive of " <i>something heavy</i> ," and the Saxon <i>læd</i> has a similar signification.
Mercury (Hydrargyrum)	Hg	<i>Hydrargyrum</i> , from <i>ὑδωρ</i> (<i>hudōr</i>), <i>water</i> , and <i>ἀργυρος</i> (<i>arguros</i>), <i>silver</i> , in allusion to its liquid and lustrous characters. <i>Mercury</i> , after the messenger of the gods, on account of its susceptibility of motion. The old name <i>quicksilver</i> also indicates its ready mobility and argentine appearance.
Silver (Argentum)	Ag	<i>Ἀργυρος</i> (<i>arguros</i>), <i>silver</i> , from <i>ἀργός</i> (<i>argos</i>), <i>white</i> . Words resembling the term <i>silver</i> occur in several languages, and indicate a white appearance.

The following are names of some of the less frequently occurring elements, compounds of which, however, are alluded to in the British and United States Pharmacopœias or are met with in pharmacy:

Name.	Symbol.	Derivation of Name.
Bromine	Br	From <i>βρῶμος</i> (<i>brōmos</i>), a <i>stink</i> . It has an intolerable odor.
Fluorine	Fl	From <i>fluo</i> , to flow. Fluoride of calcium, its source, is commonly used as a flux in metallurgic operations.

Name.	Symbol.	Derivation of Name.
Boron	B	From <i>borak</i> or <i>baurak</i> , the Arabic name of <i>borax</i> , the substance from which the element was first obtained.
Silicon	Si	From <i>silex</i> , Latin for flint, which is nearly all silica (an oxide of silicon).
Lithium	L	From <i>λίθιος</i> (<i>litheios</i>), <i>stony</i> , in allusion to its supposed existence in the mineral kingdom only.
Strontium	Sr	This name is commemorative of <i>Strontian</i> , a mining village in Argyleshire, Scotland, in the neighborhood of which the mineral known as <i>strontianite</i> , or carbonate of strontium, was first found.
Cerium	Ce	Discovered in 1803, and named after the planet <i>Ceres</i> , which was discovered on Jan. 1, 1801. The oxalate of cerium is official, but seldom used.
Chromium	Cr	From <i>χρῶμα</i> (<i>chrōma</i>), color, in allusion to the characteristic appearance of its salts.
Manganese	Mn	Probably the slightly altered word <i>magnesia</i> , with whose compounds those of manganese were confounded till 1740.
Cobalt	Co	<i>Cobalus</i> , or <i>Kobold</i> , was the name of a demon supposed to inhabit the mines of Germany. The ores of cobalt were formerly troublesome to the German miners, and hence received the name their metallic radical now bears.
Nickel	Ni	<i>Nickel</i> , from <i>nīl</i> , <i>worthless</i> . Nickel ore was formerly called <i>Kupfernickel</i> , <i>false copper</i> . When a new element was found in the ore, the name nickel was retained for it.
Tin (Stannum)	Sn	Both words are possibly corruptions of the old British word <i>staen</i> , or the Saxon word <i>stan</i> , a stone. Tin was first discovered in Cornwall, and the ore (an oxide) is called <i>tinestone</i> to the present day.
Gold (Aurum)	Au	<i>Aurum</i> (Latin), from a Hebrew word signifying the color of fire. <i>Gold</i> , a similar word is expressive of <i>bright yellow</i> in several old languages.
Platinum	Pt	From <i>platina</i> (Spanish), diminutive of <i>plata</i> , <i>silver</i> . It somewhat resembles silver, but is less white and lustrous.
Bismuth	Bi	Slightly altered from the German <i>Wismuth</i> , derived from <i>Wiesematte</i> , "a beautiful meadow," a name given to it originally by the old miners in allusion to the prettily variegated tints presented by the freshly exposed surface of the crystalline metal.
Cadmium	Cd	<i>Καδμεία</i> (<i>Kadmeia</i>) was the ancient name of calamine (carbonate of zinc), with which carbonate of cadmium was long confounded, the two often occurring together.

Gold, Platinum, Tin and Silicon are classed with the less important elements, because their salts are seldom used in pharmacy.

It will be noticed that the symbol of an element is simply the first letter of its Latin name, which is generally the same as in English. Where two names begin with the same letter, the less important has an additional letter added.

QUESTIONS AND EXERCISES.

Of how many elements is terrestrial matter composed?—In what state do the elements occur in nature?—Distinguish between the *art* and the *science* of chemistry.—What is the difference between an element and a compound?—Enumerate the chief non-metallic elements.—Describe a process for the preparation of oxygen.—How are gases usually stored?—Mention the chief properties of oxygen.—What is the source of animal warmth?—State the proportion of oxygen in air.—Is the proportion constant, and why?—Give a method for the elimination of hydrogen from water.—State the properties of hydrogen.—Why is a mixture of hydrogen and air explosive?—Explain the effects producible by the ignition of large quantities of coal-gas and air.—What is the nature of combustion?—Define a *combustible* and a *supporter of combustion*.—Describe the structure of flame.—State the principle of the Davy safety-lamp.—To what extent is hydrogen lighter than oxygen?—What do you mean by *diffusion* of gases?—State Graham's law concerning diffusion.—Name the source of phosphorus, and give its characters.—Why does phosphorus burn in air?—What remains when ignited phosphorus has removed all the oxygen from a confined portion of air?—Mention the properties of nitrogen.—What office is fulfilled by the nitrogen of the air?—State the proportions of the chief constituents of air.—Mention the minor or occasional constituents of air.—What is the proportion by weight of nitrogen to oxygen in the atmosphere?—Give the specific gravity of nitrogen.—How is chlorine prepared?—Enumerate the properties of chlorine.—Define the terms *deodorizer* and *disinfectant*.—Explain the bleaching effect of chlorine.—What proportion of hydrogen to chlorine is necessary for the formation of hydrochloric acid gas?—State the prominent chemical and physical characters of sulphur.—State those of carbon.—State those of iodine.—Give the derivations of the names of some of the elements.—What are the symbols of oxygen, hydrogen, nitrogen, carbon, chlorine, iodine, sulphur, phosphorus?

THE LEARNER IS RECOMMENDED TO READ THE FOLLOWING PARAGRAPHS ON THE GENERAL PRINCIPLES OF CHEMICAL PHILOSOPHY CAREFULLY ONCE OR TWICE, THEN TO STUDY (EXPERIMENTALLY, IF POSSIBLE) THE SUCCEEDING PAGES, RETURNING TO AND READING OVER THE GENERAL PRINCIPLES FROM TIME TO TIME UNTIL THEY ARE THOROUGHLY COMPREHENDED.

THE GENERAL PRINCIPLES OF CHEMICAL PHILOSOPHY.

DEFINITION OF CHEMICAL ACTION.

The learner may now proceed to study the manner in which substances act chemically on each other. By acting *chemically* it will

be obvious, from the preceding experiments, that what is meant is *so affecting each other that the substances are greatly altered in properties*. A mixture of free oxygen gas and hydrogen gas is still a gas; a *chemical compound* of oxygen and hydrogen is a liquid—namely, water; here is great alteration in properties. Iodine is only slightly soluble in water, and forms a brown-colored solution, and iron is insoluble; but when iodine and iron are *chemically combined*, the product is very soluble in water, forming a light-green solution in which the eye can detect neither iodine nor iron, and which is utterly unlike iron or iodine in any one of their properties. Sand, sugar, and butter, rubbed together, form a *mere mixture*, from which water would extract the sugar, and ether dissolve out the butter, leaving the sand. Tartaric acid, sodium carbonate, and water, mixed together, form a *chemical compound* containing neither an acid nor a carbonate, these bodies having interacted and formed fresh combinations. These illustrations show that *chemical action* is distinguished from all other actions by (a) producing an entire change of properties in bodies. It also is (b) exerted only between definite weights and volumes of matter. This (a and b) cannot be said of any other action—the action of any of the other great forces of nature (gravitation, heat, light, electricity, etc.); hence the statements (a and b) furnish a sharp and precise definition of *chemical action* or the chemical force. Further, (c) chemical action is only exerted when the substances are close together.

ATOMS.

In a chemical compound what has become of its constituents? Let the reader place before him specimens of sulphur, iron, and iron sulphide (*Ferrous Sulphide*, U. S. P.), or iodine, iron, solid iron iodide, and its solution in water or syrup (*Syrupus Ferri Iodidi*, U. S. P.). In the ferrous sulphide what has become of the sulphur and of the iron from which it was made? The *mixture* of sulphur and iron in *combining* to form ferrous sulphide has not lost weight, and, indeed, by certain processes it is possible to recover its sulphur as sulphur and its iron as iron; so that we are compelled to believe (we cannot avoid the conclusion) that ferrous sulphide contains particles of sulphur and of iron. But how small must be these particles! Rub a minute fragment to dust in a mortar and place a trace of the powder under the highest power of the best microscope; no yellow particle is visible, not the minutest portions of lustrous metal, but dull-brown miniature fragments of the original mass. The ultimate particles of sulphur and iron, or of the elements in any other compound (the chlorine and sodium in common salt or the iodine and iron in solution of ferrous iodide), are, in short, too small to be seen. Can they be imagined? Again, no! The mind cannot conceive of an ultimate particle (sulphur, iron, ferrous sulphide, or what not) so small but what the next instant the imagination has divided it. Yet learner and teacher must have some common platform on which to reason and converse. The difficulty is met by speaking of these inconceivably small particles as *atoms*

(ἄτομος, *atomos*, indivisible, from the privative *a* and τέμνω, *temno*, to cut—that which is not cut or divided), an idea first thoroughly worked out by an Englishman, Dalton, at the commencement of the present century.

The Greeks had a vague notion that matter could not be infinitely divisible; that there must be some limit to the divisibility of matter; that all matter must be made up of ultimate particles. Democritus, 400 B. C., held that matter was formed of atoms. But it was Dalton who, chiefly through employing the balance, gave exactitude to this notion, and by his broad mental grasp converted a vague hypothesis into a sound and satisfactory theory that all the world has since accepted and employed in explanation of the facts relating to those changes or alterations in matter which had up to his time proved so inexplicable. Doubtless the theory is only a theory. We may never be able to demonstrate the existence of atoms. But it is a theory supported by all known facts—one of those short reflections of facts dear to healthily constituted minds; moreover, it is the only theory possible to the majority of minds in the present state of knowledge and education.

We cannot speak of iodine and iron uniting lump to lump, as two bricks are cemented together or blocks of wood glued together, for such is not the kind of action. We cannot select minute fragments of each to regard as the combining portions; for the minutest fragment we could obtain is visible, and ferrous iodide contains neither visible iodine nor visible iron. And yet ferrous iodide contains both iodine and iron, or, at least, a given weight of the compound is obtained from the same weight of constituents, and the same weight of constituents is obtainable from an equal weight of the compound. We might say that molecules are concerned in the operation; but a molecule means a little mass of—of what? There is positively no word left with which to carry on conversation and description but *atoms*. Any other mode of treating the matter is too subjective for general employment. Moreover, any difficulty in forming a definite conception of an atom is met by regarding an atom not necessarily as something which *cannot* be divided, but as “a particle of matter which undergoes no further division in *chemical metamorphoses*” (*Kekulé*). Even physicists regard atoms from much the same point of view; indeed, they often speak of still larger portions of matter (molecules) as atoms, meaning thereby “something which is not divided in certain cases that we are considering” (*Clifford*). Dalton's Atomic Theory will again be referred to within the next twenty pages, more especially as explanatory of the curious fixedness of the weights and volumes in which elements and compounds alone combine with each other.

THE CHEMICAL FORCE.

What power binds the atoms of a chemical compound together in such marvellous closeness of union that in the couple or group they lose all individuality? Clearly an attractive force of enormous power—a force remotely resembling, perhaps, that which attracts

a piece of iron to a magnet. Only by such an assumption can we conceive that common salt contains chlorine and a metal (sodium), or that wood contains carbon, hydrogen, and oxygen. Were not this force thus all-powerful, the carbon in wood would show its blackness and other qualities, and the hydrogen and oxygen give indications of their gaseous and other characters. This attractive force is commonly termed *the chemical force*, sometimes *chemical affinity*. The word *chemism* has also been proposed for it, just as the magnetic force is termed magnetism, but the word has not generally been adopted.

Whence comes the chemical force? Whence comes matter? We can neither create nor destroy matter; we can neither create nor destroy force. We can alter matter from one form to another; we can alter force from one form to another. The various forms of compounds are thus co-related; the various forces are co-related. But of the whence and whither, either of matter or of force, we know nothing.

MOLECULES.

A free, uncombined atom probably cannot exist in a state of isolation at common temperatures for any appreciable length of time. For we must regard an atom as the home of an attractive force of great intensity, and the moment such an atom is liberated from a state of combination (say, hydrogen from water or chlorine from salt), it finds itself in proximity to another atom having similar desires for union, so to speak; the result is an impetuous rushing together and formation of either couples, trios, or groups according to the nature of the atoms. It would be as difficult to conceive of separate atoms as to imagine that a strong magnet and a piece of steel could be suspended close to each other without being drawn together. It is doubtless possible to keep some pairs of atoms apart by the aid of heat, just as the magnet and steel may be parted by a superior amount of force; but such a condition of things is abnormal. These pairs and other groups of atoms are conveniently designated by the one word *molecule*, the diminutive of *mole*, a mass—literally, little masses. Dissimilar kinds of atoms seem to have greater attraction for each other than similar kinds; for, first, the masses of matter met with in nature in the great majority of cases contain two or more dissimilar elements; and, secondly, at the moment certain elements are liberated from their combinations they are very specially active in combining with other, different, elements; that is to say, the chances are not equal that the liberated elements will either retain their elementary condition or combine to form compounds, but the cases in which compounds are formed are actually in great majority.

The study of the chemistry of molecules, *quâ* molecules, is of great interest; but the study of the chemistry of the atoms or groups of atoms within molecules is of enormously greater interest. A molecule of nitrogen, for instance, is not very active; an atom of nitrogen has activity which even the most advanced chemist finds difficult of realization.

RECAPITULATION.

It is desirable that the learner should here make some experiment which will serve to bring again under notice in an applied or concrete form what has just been stated respecting the substances termed *chemical compounds*, and concerning the character of that *chemical force* which resides in the *atoms* of *molecules*. The following will usefully serve this purpose; it is the process for detecting a trace of sulphurous acid in common liquid hydrochloric acid:

As already proved, hydrogen gas and chlorine gas, when interunited in a manner presently explained, form hydrochloric acid gas: the latter dissolved in water is the ordinary colorless liquid of the shops termed *hydrochloric acid*, the *Acidum Hydrochloricum* of pharmacopœias. Common yellow acid not infrequently contains as an impurity a trace of sulphurous acid gas, a body also already mentioned and experimentally prepared—a trace too small to be detected by its odor. Obtain a specimen of common liquid hydrochloric acid containing as an impurity a trace of sulphurous acid, or adopt the more simple course of purposely adding a few drops of aqueous solution of sulphurous acid (*Acidum Sulphurosum*,* U. S. P.) to some hydrochloric acid. (If no sulphurous acid is at hand, the object may be accomplished by putting a quarter or half an ounce of liquid hydrochloric acid into a wide-mouthed bottle, then burning a fragment of sulphur on a wire or strip of wood inside the bottle for a few seconds, and shaking the gas and liquid together.) Pour some of the impure liquid hydrochloric acid into a test-tube, add about an equal bulk of water, and then drop in a few fragments of the metal zinc. Effervescence will occur, due to the escape of inodorous hydrogen gas, together with a small quantity of a badly-smelling gas termed sulphuretted hydrogen (*Hydrogen Sulphide* or *Hydrosulphuric Acid*, U. S. P.). Bring the mouth of the tube near the nose; the presence of sulphuretted hydrogen will at once be recognized.

The hydrochloric acid has now been *tested* for sulphurous acid. If the experiment be performed on any commercial specimen of the acid, and a smell of sulphuretted hydrogen be observed, the operator will at once be able to state that the specimen contains sulphurous acid as an impurity.

Now, using Dalton's theory of the atomic constitution of matter, the explanation of what occurs in the successive steps of the foregoing experiment is as follows:

Hydrochloric acid is a *chemical compound* of hydrogen and chlorine. That it is a chemical compound, and not a mere mechanical

* These aqueous solutions of acids are generally, for the sake of brevity, simply termed acids.

mixture of hydrogen and chlorine, is shown by the fact that its properties are altogether different from the properties of its constituents. The attractive power or *chemical force* resident in the *atoms* of the *molecules* of chlorine and of hydrogen has caused the atoms to combine in the closest manner imaginable, and form pairs of atoms or *molecules* of the chemical compound—hydrochloric acid. Zinc being introduced into the acid, and the atoms of zinc and chlorine having even still greater attraction for each other than the hydrogen for the chlorine, the zinc and chlorine atoms combine and form a new molecule (termed zinc chloride), which remains in the liquid, while the hydrogen atoms, having the atoms of no other element to combine with if the acid is pure, unite to form pairs, or molecules, of hydrogen, and in that state escape from the vessel. If the acid be impure from the presence of sulphurous acid (sulphurous acid gas, it will be remembered, is a compound of sulphur and oxygen), some of the hydrogen atoms at the moment of birth, their *nascent* state (from *nascor*, to be born)—the specially active state—finding the atoms of other elements present—namely, the atoms of sulphur and oxygen of the sulphurous acid molecules—combine by preference with these atoms and form new molecules, the sulphur and hydrogen forming sulphuretted hydrogen, and the oxygen and hydrogen producing water; the former escapes with the great bulk of the hydrogen, while the water remains with the water already in the vessel.

Note.—Ordinary hydrogen gas—that is, hydrogen in the molecular, not in the atomic or nascent condition—will not thus attack sulphurous acid. Doubtless the amount or extent of attraction of two atoms of hydrogen for one atom of, say, the sulphur in the sulphurous acid molecule, is a constant amount; but the uncombined, nascent atoms can, it is only fair to suppose, get much nearer to the attacked molecule than they can after they have themselves combined to form a molecule, molecules (but scarcely atoms) having an appreciable amount of space between them, as will be further shown almost immediately. In other words, it is probably distance which prevents an attack which would be inevitable at close quarters. These remarks apply to all similar reactions of other elements. The great activity of nitrogen in what the student will now see is its atomic rather than merely nascent condition, as compared with its slight activity in what now may be termed its molecular condition, has already been alluded to (page 27).

CONDITIONS AND NATURE OF THE MANIFESTATION OF THE CHEMICAL FORCE.

The exertion of chemical affinity is only possible when the masses of the bodies touch. Thus it was necessary to bring the oxygen, hydrogen, phosphorus, chlorine, sulphur, carbon, iodine, and iron into ordinary contact in the respective experiments with those elements before the various reactions occurred. The exact nature of these actions, as indeed of all in which substances act chemically, would seem to be an interchange, most generally a mutual one, of the atoms of which the molecules consist—a change of partners, so to speak. Thus, in the experiment in which hydrogen and chlorine

gases united to form hydrochloric acid gas, a pair of atoms in a hydrogen molecule and a pair of atoms in a chlorine molecule, finding themselves opposite to each other, changed places, the atoms of each of the old molecules unlinking, so to say, and pairing off in fresh couples :

{ Hydrogen } and { Chlorine } become { Hydrogen } and { Hydrogen }
 { Hydrogen } { Chlorine } { Chlorine }


Or, using the symbols of these elements instead of the full names, H H and Cl Cl become H Cl and H Cl. Still further economizing space and trouble, the statement may be made in the following form : H_2 and Cl_2 become $2HCl$. Once more, by using the *plus* sign (+), instead of the words "and" or "added to," and the sign or symbol = or *equal*, instead of the words "become" or "are equal to," we reach the shortest expression : $H_2 + Cl_2 = 2HCl$; and this is the form in which it may be expressed in the student's notebook. It is the shortest and most convenient form, and is instructive and suggestive to the mind.

CHEMICAL NOTATION.

We have thus gradually arrived at a spot in the path of chemical philosophy at which we must halt to more fully discuss the usual method of recording chemical travels. We have arrived at the subject of *chemical notation* (from *noto*, to mark), the art or practice of recording chemical facts by short marks, letters, numbers, or other signs. Already the first capital letter, or the first and one of the following small letters, of the Latin names of the elements have been employed as contractions, or shorthand expressions, or *symbols* of the whole *name*. Thus H has been used for the word "hydrogen," and Cl for "chlorine." A *second function* of such a symbol is that of indicating one *atom*. Thus H stands not only for the word or substance "hydrogen," but for one atom of hydrogen. Large and small figures (2 or ₂) indicate a corresponding number of atoms, the small figure only multiplying the one particular symbol to which it is attached, while a large figure multiplies all the symbols it precedes. Thus H_2 means two atoms of hydrogen, and Cl_2 two atoms of chlorine, while $2HCl$ means two atoms of hydrogen and two atoms of chlorine, or, in one word, two molecules of hydrochloric acid gas. A *third function* of such a symbol as H or Cl is that of indicating one *volume* of the element in the gaseous state. Thus H, Cl, or O stand, first, for the substances named hydrogen, chlorine, and oxygen ; secondly, for single atoms of hydrogen, chlorine, and oxygen ; thirdly, they represent single and equal volumes of chlorine, hydrogen, and oxygen. It will be remembered that one test-tubeful of hydrogen and an equal-sized test-tubeful of chlorine were employed in a previous experiment in forming hydrochloric acid gas, HCl.


The *position* of symbols counts for something. Thus, HCl indicates not only the substances hydrogen and chlorine, single atoms of each of the substances, and equal volumes of each, but also that the two substances are *joined together by the chemical force*.

If the two letters were placed one under the other or at some distance apart, or were separated by a comma or a plus sign (+), they would be understood to mean a mere *mixture* of the elements; but placed as close as the printer's type will conveniently and consistently allow, they must be considered to stand for a *compound* of the elements; that is to say, hydrochloric acid gas (HCl). The collection of symbols representing a molecule is termed a *formula*. H_2 , Cl_2 , and HCl are the *formule* of hydrogen, chlorine, and hydrochloric acid gas.



Such a set of letters, figures, and marks as that on the above line is collectively termed an *equation*, because it indicates the equality of the number and nature of the atoms before and after chemical action. On the left hand of the sign of equality (=) are shown two molecules, and on the right hand two molecules; but of the molecules on the left, one contains two atoms of hydrogen and the other two atoms of chlorine, while of the molecules on the right each contains one atom of hydrogen and one of chlorine. The equation forms a short and convenient plan of recording the facts of experiment.

Instead of an equation, a *diagram* may be employed to exhibit the same facts. Thus:



PHYSICAL AND CHEMICAL CONSTITUTION OF MATTER.

RELATIONS OF GASES, LIQUIDS, AND SOLIDS.

Molecules of gases are not in absolute contact, for a volume of gas may be compressed with very little force to half or one-fourth its bulk—in short, to such an extent that in many cases the molecules sufficiently approximate to form a liquid. In a liquid the molecules are still free to glide about with ease amongst each other; and though in solids they exhibit less mobility, still even solids may be compressed by powerful pressure, so that probably in *no* instance are molecules in absolute contact. (Moreover, from the researches of Cagnard de la Tour and of Andrews there would seem to be no sharp lines of demarcation between the gaseous, liquid, and solid conditions of substances.) One's mental picture of the relative position of the molecules of gaseous, vaporous, liquid, or solid matter must be such a picture as that of the moving particles of dust in the air of a room, or such a relation to each other as that of the planets and stars suspended in space. There is abundant experimental evidence to warrant such a conception. A clear, transparent fluid appears perfectly homogeneous, but is not so. Its particles are

not in contact. Every person who has mixed 5 pints of rectified spirit with 3 pints of water knows that the 100 fluidounces of spirit and 60 fluidounces of water do not when mixed give 160 ounces of "proof" spirit, but only 156 ounces; the molecules of the liquids have gone closer together, having probably a little attraction for each other. Having gone closer together, they were not previously so close together, the necessary conclusion being that even liquids are porous. Why a gas under pressure should immediately return to its original bulk when the pressure is removed, while a liquefied or solidified gas only slowly resumes the gaseous or vaporous state, is a question which requires for discussion a knowledge of the nature of forces other than the chemical. For it must be remembered that the study of the chemical force is mainly the study of the internal constitution of molecules; the study of the properties of entire molecules forming the domain of Physics, sometimes termed Natural Philosophy. (Physics, from *φύσις*, *phusis*, nature; that is, visible and material nature, the study of actions and reactions which do not involve entire and permanent change in the properties of bodies—the study of the action of heat, light, electricity, magnetism, gravitation, etc. on matter.)

It is necessary, however, to state something more about the physical as well as the chemical condition of the molecules of a *gas*, in order that the learner may be prepared for the fact that mixtures of certain gaseous elements, in combining to form gaseous compounds, diminish considerably in volume. Thus, while a pint of hydrogen and a pint of chlorine give a quart of hydrochloric acid gas,

Hydrogen.	Chlorine.
Hydrochloric acid gas.	

two pints of hydrogen and one of oxygen are necessary to produce a quart of gaseous water (steam). It will be remembered that two volumes of hydrogen and one of oxygen were necessary in a previous experiment in which water was formed.

Hydrogen.	Hydrogen.	Oxygen.
Gaseous water (Steam).		

Now, that a pint of hydrogen gas and a pint of chlorine gas should, after chemical reaction or rearrangement of the atoms of the molecules has taken place, form two pints of hydrochloric acid gas is quite what we should expect. For, first, the reader by this time is not astonished that chemical combination is attended by entire change of properties; and, secondly, the experience of years has led him to expect that a pint of one thing added to a pint of another gives two pints of the mixture. But that two pints of hydrogen and one pint of oxygen should, after combination (and under like conditions of temperature and pressure), give, not three, but two pints of product (steam) is perhaps somewhat astonishing and needs explanation. To this end let us *picture* a few of the molecules of hydrogen and as many molecules of chlorine. Draw with a pencil on paper several pairs of crosses (+ +) to represent hydrogen molecules, and circles (O O) for chlorine molecules, or, if colored ink is at hand, red pairs of dots for hydrogen and green for chlorine. Or, at once, for facility in printing, let the following pairs of letters *h h* represent a few (say, nine) molecules of hydrogen, and *c c* molecules (nine) of chlorine, before combination:

<i>h h</i>	<i>h h</i>	<i>h h</i>	<i>c c</i>	<i>c c</i>	<i>c c</i>
<i>h h</i>	<i>h h</i>	<i>h h</i>	<i>c c</i>	<i>c c</i>	<i>c c</i>
<i>h h</i>	<i>h h</i>	<i>h h</i>	<i>c c</i>	<i>c c</i>	<i>c c</i>

Then, after combination, we shall have eighteen molecules of hydrochloric acid gas:

<i>h c</i>	<i>h c</i>	<i>h c</i>	<i>h c</i>	<i>h c</i>	<i>h c</i>
<i>h c</i>	<i>h c</i>	<i>h c</i>	<i>h c</i>	<i>h c</i>	<i>h c</i>
<i>h c</i>	<i>h c</i>	<i>h c</i>	<i>h c</i>	<i>h c</i>	<i>h c</i>

But when two volumes of hydrogen and one of oxygen combine and give two volumes of steam, the mental picture must be, not that of molecules somewhat nearer to each other than before, nor any difference in the size of the molecules, but a picture of molecules each containing three instead of two atoms; thus, still using pairs of letters, just for the moment, to represent a few (the space will allow only twenty-seven) molecules:

<i>h h</i>	<i>h h</i>	<i>h h</i>	<i>h h</i>	<i>h h</i>	<i>h h</i>	<i>o o</i>	<i>o o</i>	<i>o o</i>
<i>h h</i>	<i>h h</i>	<i>h h</i>	<i>h h</i>	<i>h h</i>	<i>h h</i>	<i>o o</i>	<i>o o</i>	<i>o o</i>
<i>h h</i>	<i>h h</i>	<i>h h</i>	<i>h h</i>	<i>h h</i>	<i>h h</i>	<i>o o</i>	<i>o o</i>	<i>o o</i>

The twenty-seven molecules (eighteen hydrogen, nine oxygen) will, after combination, become eighteen molecules of steam:

<i>hoh</i>	<i>hoh</i>	<i>hoh</i>	<i>hoh</i>	<i>hoh</i>	<i>hoh</i>
<i>hoh</i>	<i>hoh</i>	<i>hoh</i>	<i>hoh</i>	<i>hoh</i>	<i>hoh</i>
<i>hoh</i>	<i>hoh</i>	<i>hoh</i>	<i>hoh</i>	<i>hoh</i>	<i>hoh</i>

As already suggested, one's mental picture of a number of molecules may well give them such a relation to each other as that of a number of solar systems in the universe, equally distant from each other, and each occupying a similar space, yet one system containing a sun and one planet, another a sun and two planets, and so on, or even one or more of the planets having one or more moons. Indeed, the atoms in some very complex molecules really appear to have very much the relation to each other of the sun, planets, and moons of a solar system. To indicate such molecules by letters as above would of course require more space than is there given to the assumed pictures of molecules.

Here occurs an opportunity that must not be lost of stating a mode of reasoning by which a molecule of oxygen (or of many other elements) is shown to be a double structure—shown to contain two atoms. Five equal-sized bottles are before us—two filled with hydrogen, one with oxygen, and two with steam. (The bottles are hot enough to prevent the steam condensing to water, and all five are at the same temperature.) Apply heat so that all shall be equally heated, *the three different substances expand equally*. Cool equally, *the contents contract equally*. Apply equal pressure to all five, *each is equally affected*. Diminish pressure equally, *each portion of the three substances equally expands*. Gases (practically steam is gas; it is simply not a permanent gas),—gases thus similarly affected must be, *physically*, similarly constructed or constituted (a law which will again be referred to on page 53); each bottle must contain the same number of particles or molecules, and at any one temperature and pressure the molecules in each must be equally distant from each other. We do not know what actual number or distance, but whatever be the number and distance, it is the same for each bottle. Say that one million is the number; then we shall have a million of molecules in the first hydrogen bottle, a million in the second, a million in the oxygen bottle, and a million in each of the steam bottles. We will cause chemical combination between the two millions of hydrogen molecules and one million of oxygen molecules, producing (as we have seen) two millions of steam molecules having the properties already stated. But a molecule of steam contains an atom of oxygen. Hence two millions of steam molecules contain two millions of oxygen atoms, which two millions of oxygen atoms have been obtained from one million of oxygen molecules. Therefore each molecule of oxygen was a double structure—each molecule of oxygen contained at least two atoms of oxygen. As Clifford says, "You cannot put 50 horses into 100 stables so that there shall be exactly the same amount of horse in each stable; but you can divide 50 *pairs* of horses among 100 stables."

Thus much respecting the construction of gaseous or vaporous matter. Our knowledge of the constitution of liquid and solid matter is still more limited.

With regard to the notation of the subject, it will be sufficient to state here that while a *symbol* usefully represents one volume of any gas, a *formula of any gas or vapor represents two volumes*. By remembering this general rule we may, by looking at a formula,


tell how many volumes of constituents were concerned in the formation of a compound, and therefore what amount of condensation, if any, occurred during the act of formation. By thus reading and interpreting the formula for water, H_2O , we see that two volumes of steam (at any temperature) may be obtained from two volumes of hydrogen and one volume of oxygen (at the same temperature), and thus the extent of condensation when hydrogen and oxygen (at a stated temperature) unite to form gaseous water (at the same temperature) is from three to two. This subject will again be treated of in connection with Chemical Combination and the Specific Gravity of Gases.

FURTHER REMARKS ON GENERAL CHEMICAL NOTATION.

We may now take an experiment already made as an additional example of chemical action, and describe the simplest way of expressing the same by notation. When two volumes of hydrogen and one of oxygen were caused to combine, the production of flame and noise proved that chemical action of some kind had taken place: had the experiment been performed in dry vessels, evidence of the precise action would have been found in the bedewing produced by the condensation of the water on the sides of the tube. Similar evidence was afforded on holding a cool glass surface over the hydrogen flame. The action is expressed in the following equation:

$$2\text{H}_2 + \text{O}_2 = 2\text{H}_2\text{O}.$$


Instead of this equation, the following diagram may be employed:



The foregoing aggregation of symbols or shorthand characters, or *formula*, H_2O , is, then, a convenient picture of the facts that have already come before us—viz. that water is formed of the elements hydrogen, H , and oxygen, O ; moreover, that it is formed of two measures or volumes of hydrogen, H_2 , to one of oxygen, O ; and, thirdly, that the molecule of water (H_2O) is formed of two atoms of hydrogen (H_2) and one of oxygen (O). The formula also fulfils the *fourth function*, of indicating that the two volumes of hydrogen and one of oxygen in combining condensed to two volumes of steam. That the resulting bulk of steam afterward shrunk most considerably in condensing to water is another matter altogether—a physical and not a chemical result, and due to the approximation of the molecules of water *after* formation.

Another experiment already performed, illustrating the character of the manifestations of chemical force (symbolically noted as follows), was that in which the red-hot carbon of wood was plunged into oxygen. The evidence of chemical action in that case was

the sudden inflammation of the carbonaceous extremity of the wood. The particles of carbon and oxygen, having intense attraction or affinity for each other at that temperature, rushed together so impetuously as suddenly to produce a large additional quantity of heat, an amount sufficient to cause the particles to emit an intense white light. The action is expressed on paper in either of the following ways: $C_2 + 2O_2 = 2CO_2$, or,



CO_2 is the formula of the well-known gaseous body commonly termed carbonic acid gas.

The reader should here draw for himself equations or diagrams similar to those on pages 42 and 46, and thus show the formation of the three other bodies he has already produced—namely, phosphoric anhydride (P_2O_5), sulphurous acid gas (SO_2), and ferrous iodide (FeI_2), submitting the same, if possible, to a tutor or other authority to assure himself of their correctness.

Note.—In the foregoing experiments several illustrations occur of the formation of compounds having the gaseous, liquid, and solid conditions, in one of which three forms all matter in the universe apparently exists.

LAWS OF CHEMICAL COMBINATION (BY WEIGHT).

Chemistry as a *science* is little more than a hundred years old, though very many of the facts and operations we now term “chemical” have been known as isolated items of knowledge for centuries. Thus the ancient Egyptians made glass, vitriol, soap, and vinegar, and the Greeks started the idea that matter was composed of a few elements, imagining earth, air, fire, and water to be elements. In short, chemistry as an *art* was already very extensive a hundred or more years ago. But the great general principles which interlace and bind together separate facts, those which from their extensive application and importance are denominated *laws*, have all been brought to light since the year 1770. Scarcely more than a single century ago Lavoisier, by invoking the aid of the balance, converted the *art* into the beginning of a *science* which has since grown by ever-recurring leaps and bounds. (Lavoisier was born in 1743; he was guillotined by Robespierre in 1794. A request for a few days of respite to complete some researches was refused on the ground that “the republic has no need of chemists.”)

First Law relating to Chemical Combination.

Between 1785 and 1800, Bryan Higgins, William Higgins, Wenzel, Richter, and Proust made analyses and researches which led up

to the following generalization: *When compounds unite to form definite chemical substances, they always combine in the same proportions.* The curious character of this fact could but be most striking, and indeed is so now to the mind receiving it for the first time. Thus water (a compound) added to quicklime (a compound) gives slaked lime, a perfectly definite chemical substance. But whereas sand and water, sugar and water, sand and sugar, and such *mechanical* mixtures may be obtained by adding together the ingredients in any proportions whatever—say, 90 of sugar and 10 of sand, or 10 of sugar and 90 of sand—such a *chemical* mixture as slaked lime (say 100 parts) invariably results from the combination of $75\frac{3}{4}$ of quicklime and $24\frac{1}{4}$ of water. If a larger proportion than $75\frac{3}{4}$ per cent. of quicklime be employed, the excess remains as quicklime mixed with the slaked lime; and if more than $24\frac{1}{4}$ per cent. of water be used, an excess of water remains with the slaked lime and evaporates if the mixture be exposed to the air. Dalton discovered that when *elements* unite to form a definite substance, they, like compounds, always combine in the same proportions; and he was the first to set forth the law in a manner which was at once clear and comprehensive enough to include the former generalization. Thus:

A definite compound always contains the same elements in the same proportions.

Take another example: Common salt always contains $39\frac{1}{2}$ per cent. of the metal sodium to $60\frac{3}{8}$ of chlorine, and water always 89 of oxygen to 11 per cent. of hydrogen (more exactly 88.89 to 11.11). As with the quicklime and water, so with the chlorine and sodium and the constituents of many (not *all*) chemical compounds: in such cases, if either be added to the other in any quantity beyond stated proportions, the excess plays no part whatever in the act of combination. (In *some* cases, as will be seen directly, excess of either plays a very simple but very remarkable part.) In short, whether a compound be made directly from its elements or by the combination of other compounds, or indirectly as one of two products of the action of substances chemically on each other, whatever be its origin, if it is a definite compound it *always contains the same elements in the same proportions.* This is the first of the two laws governing chemical combinations.

Second Law relating to Chemical Combination.

Dalton further made such experimental researches as enabled him to lay down a second great law. He found that while many substances only united chemically in one proportion, others combined in two or even more; and he studied several such naturally related bodies. He found that while carbonic oxide (a gas formed when charcoal is burned with a limited supply of air) contains such a proportionate weight of carbon and oxygen as is represented by (to use the simplest figures) 3 and 4, carbonic acid (a gas formed when charcoal is burned with excess of air) contains 3 of carbon to exactly twice 4 of oxygen. He proved that a similar relation existed between two compounds of carbon and hydrogen and between a cluster of compounds of nitrogen and oxygen. The first of the latter, to a

given quantity of nitrogen contains a certain proportion of oxygen ; the next, to the same quantity of nitrogen has exactly twice the proportion of oxygen ; and the others have exactly three, four, and five times as much oxygen as the first, the quantity of nitrogen remaining the same throughout. Dalton thus generalized these facts :

When two elements unite in more than one proportion, the resulting compounds contain, to a constant proportion of one element, simple multiple proportions of the other ; or the weights of the constituent elements bear some similar simple relation to each other.

Thus carbonic oxide gas is a definite compound always containing fixed proportions of carbon and oxygen, and carbonic acid gas is also a definite compound always containing fixed proportions of carbon and oxygen. Both thus obey the first law of combination. But whereas carbonic oxide contains, or may be made from, 30 parts (ounces, grains, or other weights) of carbon and 40 of oxygen, carbonic acid contains, or may be made from, 30 parts of carbon and exactly twice 40 of oxygen.

The second law cannot but be as striking as the first when freshly unveiled to the mind. Sand and sugar, or any substances which do not act *chemically* on each other, may be mixed in the proportions of 30 to 40, 30 to 80, 30 to 60, or any other quantities ; but if an attempt be made to burn 30 parts of carbon in 60 of oxygen, the elements will themselves naturally assert their own special combining powers, and refuse, so to say, to unite in these proportions : the 30 of carbon will first combine with 40 of oxygen and form 70 of carbonic oxide ; and this gas, which, had it the opportunity, would combine with 40 more of oxygen and form carbonic acid gas, finding only half that quantity—namely, 20 of oxygen—present, contents itself by one half (that is 35 of carbonic oxide) accepting the 20 of oxygen and becoming carbonic acid gas, while the other half remains as carbonic oxide. This is a most wonderful fact. Again, if 30 parts of carbon be burnt in more than 80, say 85, of oxygen, only 80 will be used, the other 5 remaining as oxygen merely *mixed* with the resulting carbonic acid gas. If we attempt to burn 30 parts of carbon in less than 40 of oxygen, the oxygen will take up three-fourths its weight of carbon and form carbonic oxide, while the excess of carbon will remain as carbon.

RECAPITULATION.

Nature does not always permit man to mix things in any proportions he pleases. She does sometimes. She does if he only stirs things together, or if he only uses the attractions of adhesion or cohesion in binding the materials together ; but if he employs chemical attraction, she restricts him to special proportions. That is to say, if the things mixed do not attack one another or intimately combine, then admixture may be effected in any proportion ; and the mixture is a mere mixture, having the mean properties of its components. Examples of such mixtures are seen in compound plasters, pill-masses, confections, and plum-puddings. But if the things do unite to form not a mere mixture having the mean properties of its components, but a compound having new and distinct and definite

characters of its own, then Nature does not permit man to combine the things in any proportion he pleases. The proportion is a fixed and constant one; and if he substitutes proportions of his own, the things unite in the proportions fixed by Nature, and the excess he has added either remains in its original uncombined condition, or it combines with the compound already produced to form a second *different* compound. Any one compound—that is, the *same* compound—always contains the same elements in the same proportions, and can only be made from the same elements in the same proportions. An attempt to mix the same elements in other proportions would result in one of two failures—namely, either the extra proportion would remain free and uncombined, or it would combine and convert the first compound or a portion of it into a *different* compound. The fresh compound thus produced, like the first, and indeed like all definite compounds, of course always contains the same elements in the same proportions.

In short (Law 1), any definite compound always contains the same elements in the same proportions, and (Law 2) any two elements uniting in more than one proportion unite in multiples of that proportion, and produce so many different definite compounds. Taking hydrogen as uniting in proportions of 1, oxygen unites in proportions of 16; that is, 16, twice 16, thrice 16, and so on, never in intermediate proportions. Carbon unites in proportions of 12, sulphur of 32, chlorine 35½. Every element (see the table in front of the Index) has its combining proportion fixed by Nature.

The student of chemistry is recommended to accept these two great natural facts, great enough to be dignified by the name of *laws*, in all their inherent solidity and simplicity. Of course he will wonder *why* substances should combine, chemically, only in fixed proportions when forming a definite body, and *why*, when a substance combines in more than one proportion to form different definite bodies, the proportions should only be multiple proportions, and will gladly hail the extremely ingenious and useful explanation of these truths suggested by Dalton. (See foregoing and following paragraphs on the theory that matter is built up of atoms.) But man has not yet succeeded in so questioning Nature as to gain from her a *satisfactory* answer to such questions; and until he does succeed, any hypothesis, even Dalton's, should be held intelligently, but not too tightly. The facts themselves, however, should be grasped with the student's utmost tenacity.

Reciprocal Proportions.—Careful consideration of the foregoing two great laws relating to chemical combination leads to an important truth—namely: *The proportions in which two elements unite with a third are the proportions (or simple multiples or submultiples of the proportions) in which they unite with each other.* Thus oxygen in proportions of 16 unites with hydrogen, and carbon in proportions of 12 unites with hydrogen; therefore 16 and 12 are the proportions in which oxygen and carbon will unite with each other.*

* See Axiom 1 in Hawtrey's *Introduction to the Elements of Euclid*,

THE ATOMIC THEORY.

The laws which Dalton so largely aided to unveil—two grand and wonderful truths—he explained and correlated by a simple and beautiful hypothesis (1803 to 1808). *Why* should any given compound always contain the same elements in such absolutely *fixed* proportions? *Why*, when an element combines in more than one proportion, forming more than one given compound, should it combine in exactly *multiple* proportions? The first answer must be that no one knows; that is to say, that we are unable to refer to any demonstrable *fact* for the answer. But a century ago the same Englishman devised an *explanation* that has satisfied the world. Dalton's explanation was *that matter was not infinitely divisible, but composed of minute particles or ATOMS having an invariable character*. In the words of Wurtz, "*To an old and vague notion he attached an exact meaning by supposing that the atoms of each kind of matter possess a constant weight, and that combination between two kinds of matter takes place not by penetration of their substance, but by juxtaposition of their atoms.*" Dalton raised the old Grecian hypothesis (see page 37) to the dignity and importance of theory, and gave to it a quantitative foundation.

Under this "atomic theory" carbonic oxide is a definite compound always containing the same elements in the same proportions, *because* each particle of it is composed of an atom of carbon and an atom of oxygen chemically united, the weights of the atoms being in the proportion of 3 and 4; that is, having a constant weight of 12 and 16, as we now believe. Carbonic acid gas is also a definite compound always containing the same elements in the same proportions, and the proportion of oxygen is just double that in carbonic oxide, *because* each particle of it is composed of an atom of carbon (weighing 12) and two atoms of oxygen (each weighing 16).

The healthily constituted student-mind asks, "*Why* an element should unite in exactly multiple proportions if it forms several compounds?" Because those compounds are "made" of atoms, which, being indivisible, must, if they unite at all, unite 1 to 1, 2 to 1, 3 to 1, and so on. With such an explanation, although only one explanation, the healthily constituted mind gains desired satisfaction.



Imaginary Pictures of Molecules of Carbonic Oxide Gas and Carbonic Acid Gas.*

Longmans & Co.—a book recommended to any student who is not familiar with the mode of reasoning termed geometrical.

*The size of atoms, their shape, their absolute weight—whether or not they are in actual contact—whether or not they are fixed in relation to each other, free to move about each other or in a constant state of motion—and whether or not the chemical force actuates them as the force of gravitation influences our earth and moon and solar systems, are matters of which, at present, we know almost nothing. The two pictures are not intended to convey any impression that the following formulæ do not give: CO or OC, OCO or CO₂.

Again, the facts that with 12 of carbon oxygen unites in the proportion of 16 or a multiple of 16; that with 12 of carbon sulphur unites in the proportion of 32 or a multiple of 32 (the liquid known as carbon disulphide is a chemical compound of 12 of carbon to twice 32 of sulphur); and, thirdly, that oxygen and sulphur unite in proportions of 16 and 32,—are at once explained on the assumption that these elements exist in atoms which have the respective weights mentioned. Existing in indivisible particles (atoms), which weigh 16, 12, and 32, oxygen, carbon, and sulphur *must* unite in indivisible weights of 16, 12, and 32.

ATOMIC WEIGHTS.

What has just been stated respecting two or three elements is true of all the elements. It is a fact that when elements unite with one another in the peculiar and intimate manner termed chemical, they do not combine in the haphazard proportions of a mere mixture, but in one fixed and constant proportion. Such proportions or weights represent, according to Dalton, the weights of their atoms. Oxygen unites with other elements in proportions of 16; therefore 16 is the weight of the atom of oxygen. Chlorine unites with other elements in proportions of $35\frac{1}{2}$, therefore $35\frac{1}{2}$ is the atomic weight of chlorine. And for a similar reason the atomic weight of hydrogen will be 1, carbon 12, sulphur 32, nitrogen 14, and iodine 127. Of course it will be understood that these are the *relative* weights of atoms, for we cannot know the absolute weights. All that is known is that the chlorine atom, for instance, is 35.5 times as heavy as the hydrogen atom, whatever the absolute weight of the latter may be, and the iodine atom 127 times as heavy. The quantity of metal which with 35.5 of chlorine will form a chloride, and with twice 35.5 a second chloride (dichloride or bichloride), will require 127 of iodine to form an iodide, and twice 127 of iodine to form a second iodide (a diniodide or biniodide).^{*} In other words, the atomic weight of an element is the *ratio* of the weight, quantity of matter, or mass of an atom to the weight, quantity of matter, or mass of an atom of hydrogen.

Notes on Notation.—A *fourth function* of a symbol is to represent atomic weight. Thus the symbols H, Cl, O, etc. not only perform the offices of representing (a) names, (b) single volumes, and (c) single atoms, but (d) definite weights of the respective elements: H = 1, Cl = 35.5, O = 16, I = 127, N = 14, K = 39, etc.

LAWS OF CHEMICAL COMBINATION (BY VOLUME).

In 1809, Gay-Lussac showed it to be a fact that when gaseous elements unite with one another in the intimate manner termed chemical, they do not combine in the haphazard proportions (that is, proportions by measure or volume) of a mere mixture, but in

^{*} Only the atomic weights of the above and a few of the chief metallic elements need be committed to memory; others can be sought out as occasion may require. A table of *combining proportions* of elements, or atomic weights, is given at the end of the volume, in front of the Index.

constant proportions in the case of any single definite compound, and in simple multiple proportions in cases where two elements form more than one definite compound. He thus proved that the laws respecting the constancy of weight with which elements combine hold good with reference to volume, at all events in those cases in which elements exist in or can be made to assume the gaseous condition. A volume of hydrogen gas and an equal one of chlorine gas give hydrochloric acid gas. Two volumes of hydrogen and one of oxygen give water vapor or steam. Such volumes or simple multiples are alone the proportions by bulk in which elements combine. If any excess of either gas be mixed and combination attempted, only the stated proportions really combine, the excess remaining unaltered. Further, following Gay-Lussac, on weighing these similar and equal volumes of hydrogen, chlorine, and oxygen we find that chlorine is 35.5 times as heavy as hydrogen, and oxygen sixteen times as heavy as hydrogen.

Avogadro's "Law."—Avogadro in 1811 and Ampère in 1814, reasoning on the fact that all gases are similarly affected by variations of pressure (Boyle, 1662, verified by Mariotte) and temperature (Charles), (see also page 45), concluded that all gases must be similarly constituted (similarity in properties always indicating similarity in character or nature—a mode of reasoning or deducing or inferring that even children soon naturally adopt in dealing with everything that appeals to their senses); in other words, that if *equal volumes of gases be taken under like conditions*, each will contain the *same number of molecules*, similar in size and equally distant apart. The deduction is obvious. *The weights of molecules of gaseous elements (that is, of pairs of atoms, and therefore of atoms themselves) must differ to the extent that the weights of equal volumes of those elements differ.* Equal volumes of hydrogen, chlorine, and oxygen, weighing respectively 1, 35.5, and 16, and each of these volumes containing an equal number of molecules, each formed of two atoms, it follows that the relative weights of the atoms will be 1, 35.5, and 16.

It will thus be seen that the weight of the *volume* in which an element combines, and the *actual weight* in which it combines, irrespective of volume, are identical. For instance, we should find by experiment that, as a simple matter of fact, oxygen unites with other elements in proportions of 16 by weight, while hydrogen combines in proportions of 1. Turning, then, to experiments on the volumes in which hydrogen or oxygen combine, and having ascertained those volumes, and then having weighed them, we should find that the oxygen volume weighs 16, while the hydrogen weighs 1. In compounds in which proportions of 1 grain of hydrogen were found oxygen would be found in proportions of 16 grains. In gaseous compounds in which hydrogen existed in proportions of, say, 27 ounces by measure, oxygen would be found in proportions of 27 ounces by measure: the 27 ounces of hydrogen would be found to weigh 1 grain, and the 27 ounces of oxygen to weigh 16 grains.

Thus the two great facts or laws respecting chemical compounds which Dalton laid down, by ascertaining the exact weights in which

bodies combine, Gay-Lussac confirmed by experiments on the exact volumes in which elements combine. Further, Gay-Lussac's experiments and Avogadro's reasoning strongly supported Dalton's theory of atoms.

RECAPITULATION.

What are atomic weights or combining weights? First, they are represented by the smallest portion (relative to 1 part of hydrogen) in which an element migrates from compound to compound. Thus, 1 part by weight of hydrogen can be eliminated from 18 similar parts of water by action of certain metals, leaving 1 of hydrogen and 16 of oxygen combined with the metal. From the latter compound 1 more of hydrogen is eliminated by a second experiment with more metal, leaving 16 of oxygen combined with the metal. In these and other well-known reactions 16 parts of oxygen take part in the various operations; 16, therefore, is the probable atomic weight of oxygen; and so with other elements and radicals. Secondly, the weight of the atoms, or the atomic weights, of the gaseous elements already studied must differ from each other to the extent that equal volumes of those elements differ in weight. For equal volumes contain an equal number of molecules equal in size (Avogadro and Ampère), and each molecule of an element is composed of two atoms; so that equal volumes of the gaseous elements contain an equal number of atoms. Now, bulk for bulk, chlorine is thirty-five and a half (35.5) times as heavy as hydrogen; so that the molecule of chlorine must be 35.5 times the weight of the molecule of hydrogen. And as the molecules of chlorine and hydrogen contain two atoms each, the atom of chlorine must be 35.5 times as heavy as that of hydrogen. The actual weight of atoms can never be ascertained; but that is of little consequence if we can only determine, with exactitude, their comparative weights. Comparing, then, all atomic weights (sometimes obscurely termed *equivalents*) with each other, and selecting hydrogen as the standard of comparison (because it is the lightest body known, and therefore, probably, will have the smallest atomic weight), and assigning to it the number 1, we see that the atomic weight of chlorine will be represented by the number 35.5. By parity of reasoning the atomic weight of oxygen is 16, for oxygen is found, by experiment, to be sixteen times as heavy as hydrogen. Similarly, the atomic weight of nitrogen is found to be 14. The atomic weight of carbon is 12—not because its vapor has been proved to be twelve times as heavy as hydrogen, for it has never yet been converted into the gaseous state, but because no gaseous compound of carbon which has been analyzed has been found to contain in 2 volumes (1 of which, if hydrogen, would weigh 1 part) less than 12 parts of carbon. (For an explanation of this reference to *two* volumes see next page.)

By thus weighing equal volumes of gaseous elements or equal volumes of gaseous compounds of non-volatile elements, and ascertaining by analysis the proportion of the non-volatile element, whose atomic weight is being sought, to the volatile element, whose atomic weight is known, the atomic weights of a large number of the

elements have been determined. Some of the elements, however, do not form volatile compounds of any kind: the stated atomic weights of these elements, therefore, are at present simply the proportions by weight in which they combine with or displace elements whose atomic weights have been determined, the proportions being in most cases checked by isomorphic considerations and the relation of the element to other forces, especially heat.* (*Vide infra.*)

MOLECULAR WEIGHT AND MOLECULAR VOLUME.

The weight of a molecule is simply the sum of the weights of its atoms; thus,

$$\text{H}_2 = 2, \text{O}_2 = 32, \text{Cl}_2 = 71, \text{H}_2\text{O} = 18, \text{HCl} = 36.5.$$

The foregoing formulæ are *molecular formulæ* or *two-volume formulæ*. It will be remembered that one volume of hydrogen and one of chlorine gave *two* of hydrochloric acid, and that two of hydrogen and one of oxygen gave *two* of steam, etc.

Molecular Volume.—If the quantities just mentioned be weighed out (in grains or other weights), or if the molecular weights of any gases or liquids be taken and exposed to similar (high) temperatures and pressures, *they will be found to occupy the same volume*. Conversely, if equal volumes of gases or vapors be measured out, and then the whole weighed, the resulting figures (all referred to 2 of hydrogen as a starting-point or standard) are the molecular weights of the respective substances. For equal volumes contain equal numbers of molecules (Avogadro). Why? Because equal volumes have equal physical properties (Boyle, Charles); and various things which have similar properties are by that fact shown to be similar things—the mode of reasoning which from childhood onward teaches us that two separate things (*e.g.* two pennies) are similar things. In the cases now being considered the things differ *chemically*, for they have entirely different chemical properties; they are not similar chemically: the point is that, having similar *physical* properties, they are similar *physically*; whatever the number of molecules in a volume of one of the gases or vapors, there must be a similar number in similar volumes of the others; therefore *the differences in the weights of tangible volumes are the differences in the weights of the intangible molecules*. This subject will again be referred to in connection with quantitative analysis (*vide* Index, “Molecular Weight”); at present the following illustration will suffice:

* *Isomorphous* bodies (ἴσος, *isos*, equal, and μορφή, *morphē*, form) are those which are similar in the shape of their crystals. This identity in crystalline form is so commonly associated with similarity of constitution that non-crystalline substances resembling each other in structure are often regarded as isomorphous. When one element unites with another in more than one proportion, and its atomic weight is so far uncertain, the isomorphism of either of its compounds with some other compound of known constitution is usually accepted as evidence of some value as to which proportion is atomic, especially if the compounds are so closely isomorphous that a crystal of either will “grow” in a solution of the other. The specific heat of elements will be treated of subsequently.

A volume of hydrogen (about 54 fluidounces) which at a temperature of, say, 300° F. or 400° F., and common atmospheric pressure would weigh 2 grains, would in the case of vapor of water (steam) weigh 18 grains. Hence we are justified in considering—indeed, compelled to consider—the molecule of water to contain 2 atoms of hydrogen ($=2$) and one of oxygen ($=16$), and its formula to be H_2O ($=18$), and not H_4O_2 , in which case its vapor would be twice as heavy as it really is found to be.

Construction of Formulæ.—The composition of hydrochloric acid (HCl), water (H_2O), ammonia gas (NH_3), carbonic acid gas (CO_2), or any other compound, as well as the weight of an element that may be concerned in its formation, cannot be ascertained by actual experiment until the student is far advanced in practical chemistry—until he is able to *analyze* not only *qualitatively*, but, by help of a balance, *quantitatively*. The percentage composition of a chemical substance having been determined by quantitative analysis, its formula is constructed by the aid of the foregoing and other theoretical considerations. The correctness of such formulæ can be verified by expert analysis, but must be taken for granted by learners. This subject will again be referred to in the latter part of this Manual.

QUANTIVALENCE OF ATOMS, OR VALENCY.

Turning from the *weights* of atoms, their chemical *value* may now be considered, or their *quantivalence*. The exchangeable chemical value of atoms in relation to each other may be compared to the exchangeable commercial value of coins. As compared with a penny, a groat is four-valued; as compared with hydrogen, carbon is quadrivalent. Here, again, hydrogen is conventionally adopted as the standard of comparison. An oxygen atom in its relations to an atom of hydrogen is bivalent (biv'a-lent, of double worth, from *bis*, twice, and *valens*, being worth); an atom of it will displace two atoms of hydrogen or combine with the same number; nitrogen is usually trivalent (triv'a-lent, from *tres*, three, and *valens*), while carbon is quadrivalent (quad-riv'a-lent, from *quatuor*, four, and *valens*). Chlorine, iodine, and bromine, as well as potassium, sodium, and silver among the metals, are, like hydrogen, univalent (u-niv'a-lent, from *unus*, one, and *valens*). Barium, strontium, calcium, magnesium, zinc, cadmium, mercury, and copper, like oxygen, are bivalent. Phosphorus, arsenium, antimony, and bismuth, like nitrogen, usually exhibit trivalent properties, but the composition of certain compounds of these five elements shows that the several atoms are sometimes quinquivalent (quin-quiv'a-lent, *quinquies*, five times, and *valens*). Gold and boron are trivalent. The atoms of silicon (the characteristic element of flint and sand), tin, platinum, and lead resemble carbon in being quadrivalent. Sulphur, chromium, manganese, iron, cobalt, and nickel are sexivalent (sex-iv'a-lent, from *sex*, six, or *sexies*, six times, and *valens*), but frequently exert only bivalent, trivalent, or quadrivalent activity. This *quantivalence* (quant-iv'a-lence, from *quantitas*, quantity, and *valens*), also termed *atomicity* (maximum quantivalence), *dynamicity*, *equivalence*, and,

simply, *valency*, of atoms, may be ascertained at any time on referring to the Table of the Elements at the end of this volume, where Roman numerals, I, II, III, IV, V, VI, are attached to the symbols of each element to indicate atomic univalence, bivalence, trivalence, quadrivalence, quinquivalence, or sexivalence. Dashes (H' , O'' , N''') similar to those used in accentuating words are often used instead of figures in expressing quantivalence. The quantivalence of elements, as they one after another come under notice, should be carefully committed to memory, for the composition of compounds can often be thereby predicated with accuracy and remembered with ease. For instance, the hydrogen compounds of chlorine, Cl' , oxygen, O'' , nitrogen, N''' , and carbon, C'''' , will be respectively $\text{H}'\text{Cl}'$, $\text{H}'_2\text{O}''$, $\text{H}'_3\text{N}'''$, and $\text{H}'_4\text{C}''''$ —one univalent atom, H' , balancing or saturating one univalent atom, Cl' ; two univalent atoms, H'_2 , and one bivalent atom, O'' , saturating each other; three univalent atoms, H'_3 , and one atom having trivalent activity, N''' , saturating each other; and four univalent atoms, H'_4 , and one quadrivalent atom, C'''' , saturating each other. Carbonic acid gas, $\text{C}''\text{O}''_2$, again, is a saturated molecule containing, in one molecule, one quadrivalent and two bivalent atoms.

The subject of quantivalence will be further explained after the first six metals have been studied, when abundant illustrations of quantivalence will have occurred.

DEFINITIONS.

Chemistry is the study of the force by which matter becomes permanently altered in properties.

The Chemical Force, like other forces, cannot itself be described, for, like them, it is only known by its effects. It is distinguished from other forces by the facts that (*a*) it produces an entire change of properties in the bodies on which it is exerted, and that (*b*) it is exerted only between definite weights and volumes of matter. Like *the force of cohesion*, which is the name given to the attraction which molecules have for each other, and which is great in solids, small in liquids, and apparently absent in gases, and like *the force of adhesion*, which is the name given to the attraction which a mass of molecules has for another mass, *the chemical force* acts only within immeasurable distances; indeed, inasmuch as the chemical force appears to reside in atoms—that is to say, is exerted inside a molecule, while all other forces affect entire molecules—the chemical force may be said to be distinguished (*c*) by being exerted within a smaller distance than that at which any other force is exerted.

An Element is a substance which cannot by any known means be resolved into any simpler form of matter.

An Atom of any element is a particle so small that it undergoes no further subdivision in chemical transformations.

A Molecule is the smallest particle of matter that can exist in a free state.

A Mere Mixture of substances is one in which each ingredient retains its properties.

A *Chemical Compound* is one in which definite weights of constituents have undergone an entire change of properties. A "compound" in *pharmacy* is an intimate mixture of substances, but still only a mixture: it is not a *chemical* compound; the ingredients have not entered into *chemical* union or combination.

Combustion is a variety of chemical combination, a variety in which the chemical union is sufficiently intense to produce heat and, generally, light.

The *Law of Diffusion* is one under which gases mix with each other at a rate which is in inverse proportion to the square roots of their relative weights; that is, irrespective of, and even in spite of, their comparative lightness or heaviness.

A *Chemical Symbol* is a capital letter or a capital and one small letter. It has four functions—namely:

1. It is shorthand for the *name* of the element.
2. It represents one *atom* of the element.
3. It stands for a constant *weight* of the element—the atomic weight or combining weight.
4. Symbols represent equal and *single volumes* of gaseous elements.

A *Chemical Formula* represents (1) a *molecule* either of an element or of a compound. It has four other functions:

2. It indicates at a glance the *names* of the elements in the molecule.
3. Its symbol, or symbols, together with a small figure attached to the foot of any symbol, show the *number of atoms* in the molecule.
4. It stands for a constant weight of a compound—the molecular weight—the sum of the combining *weights* or of the weights of the atoms in the molecule.
5. It represents *two volumes* of the substance (if volatilizable) in the state of gas or vapor, and the number of volumes of gaseous elements from which two volumes of any gaseous compound were obtained.

A *Chemical Equation* or a *Chemical Diagram* is a collection of formulæ and symbols so placed on paper as to form a picture or illustration of the state of things before and after that *metathesis* (interchange) of atoms of molecules which results in the formation of molecules of new substances.

A *Solid* is a substance the molecules of which are more or less immobile, though probably not in absolute contact.

A *Liquid* is a substance the molecules of which so freely move about each other that it readily assumes and retains the form of any vessel in which it is placed.

A *Gas* is a substance the molecules of which are so far apart that they seem to have lost all attraction for each other, and, indeed, to have acquired the property of repulsion to such an extent that they are only prevented from receding to a still greater extent by the pressure of surrounding matter. Motion is especially characteristic of the molecules of gaseous fluids.

*The Two Laws regulating Chemical Combinations
(either by weight or volume).*

First. The Law of Definite Proportions.—A definite compound always contains the same elements and the same proportions of those elements—by weight or by volume.

Second. The Law of Multiple Proportions.—When two elements unite in more than one proportion by weight or by volume, they do so in simple multiple of that proportion, forming *different* compounds, each of which, as regards definiteness of composition, of course obeys the first law.

Reciprocal Proportions.—The proportions in which two elements unite with a third are the proportions in which they unite with each other.

Atomic Weights are, first, the proportions in which elements are found to combine with each other by weight. (The figures showing these proportions are purely relative, but all chemists agree to make this relation fixed by giving the number 1 to hydrogen.) Secondly, they are the weights of equal volumes of gaseous elements (relative to 1 of hydrogen).

Molecular Weights.—These are the weights of equal volumes of gases or vapors under equal circumstances of temperature and pressure, and relative not to 1, but to 2 of hydrogen. In the case of non-volatile bodies molecular weight is deduced from the observed analogies of the bodies with those whose molecular weight admits of proof.

Quantivalence of Atoms.—The chemical value for work of an atom relative to one of hydrogen. (*Caution.*—Quantivalence gives no clue to that varying intensity of union of atoms which results in varying stability of compounds.)

THE LEARNER IS RECOMMENDED TO READ THE FOREGOING PARAGRAPHS ON THE GENERAL PRINCIPLES OF CHEMICAL PHILOSOPHY CAREFULLY ONCE OR TWICE, THEN TO STUDY (EXPERIMENTALLY, IF POSSIBLE) THE FOLLOWING PAGES, RETURNING TO AND READING OVER THE GENERAL PRINCIPLES FROM TIME TO TIME UNTIL THEY ARE THOROUGHLY COMPREHENDED.

Minor principles of Chemical Philosophy will be found scattered throughout the following pages. (Also *vide* Index, under the word "Principles.")

Students of pure Chemistry, especially when fairly well acquainted with chemical facts, will also find the principles of Chemistry, including the probable constitution as distinguished from the mere composition of chemical substances, fully set forth in the larger text-books.

QUESTIONS AND EXERCISES.

What do you understand by *chemical action*? Give examples.—How is the chemical force distinguished from other forces?—Adduce evidence

that elements exist in compounds—that ferrous sulphide, for instance, still contains particles of sulphur and iron, though it possesses properties so different from those elements.—Define the term “atom.”—What condition is essential for the manifestation of chemical force?—Can an atom exist in an uncombined state? and when are atoms most potent to enter into chemical combination?—What is a molecule?—How may the results of chemical reactions be expressed on paper?—Enumerate the functions of a symbol.—Give the additional functions of a chemical formula.—Describe by a diagram or an equation the reaction which ensues when red-hot charcoal is plunged into oxygen gas.—Draw diagrams representing the formation of P_2O_5 , SO_2 , and FeI_2 , respectively.—Enumerate the differences in the physical conditions of the molecules in a solid, a liquid, and a gas.—State the law of constant proportions.—State the law of multiple proportions.—State the principle of reciprocal proportions.—Give illustrations of the above laws.—Describe the origin and use of the atomic theory.—What do you understand by the atomic weight and the molecular weight of an element?—Representing the weight of an atom of hydrogen as 1, what will be the atomic weights of carbon, sulphur, nitrogen, and iodine? Give reasons for considering the stated weights to be correct.—In what proportion, by volume, do elements in the gaseous state chemically combine?—What relation exists between the combining volumes of elements in the gaseous state and their atomic weights? Give the explanation for this. Is there any difference between the molecular volumes of simple and of compound gases?—Define isomorphism.—Explain the value of isomorphism as evidence of atomic weight. What is to be understood by the quantivalence of atoms? Give examples of univalent, bivalent, trivalent, and quadrivalent atoms.—How may the quantivalence of an element be expressed in its atomic symbol?—Give formulæ in which the quantivalence of one atom is saturated by the combined quantivalence of others.

The reader is also recommended to question himself, or be questioned, on the “definitions” given on pp. 57–59.

THE ELEMENTS AND THEIR COMPOUNDS.

HAVING thus obtained a general idea of the nature of such elements as have special interest for the medical and pharmaceutical student, and which, indeed, are all with which any student of Chemistry should at present occupy his attention, we may pass on to consider in detail the relation of the elements to each other. The elements themselves, in the free condition, are seldom used in medicine, being nearly always associated—bound together by the chemical force; in this combined condition, therefore, they must be studied—inorganic combinations first, organic afterward. Most compounds met with in the mineral kingdom may be regarded as containing two parts or roots, two radicals—the one usually metallic, or, to speak more generally, basylous; the other commonly a non-metallic, simple or complex, acidulous radical. In the following pages the basylous radicals, or metals, will be considered first, the acidulous radicals afterward. (They will follow the chemistry of compounds, many of which have not so simple a constitution as that just indicated.) Each radical will be studied from two points of view, the synthetical and the analytical; that is to say, the properties of an element on which the preparation of its compounds

depends will be illustrated by descriptions of actual experiments, and thus the principles of chemistry and their applications to medicine and pharmacy be simultaneously learnt; then the reactions by which the element is detected, though combined with other substances, will be performed, and so the student be instructed in qualitative analysis. Synthetical and analytical reactions are, in truth, frequently identical, the object with which they are performed giving them synthetical interest on the one hand or analytical interest on the other.

A good knowledge of Chemistry may be acquired synthetically by preparing considerable quantities of the salts of the different metals, or analytically by going through a course of pure qualitative analysis. But the former plan demands a larger expenditure of time than most students have to spare, while under the latter system pupils generally lose sight of the synthetical interest which attaches to analytical reactions. Hence the more useful system, now offered, of studying each metal, etc. from both points of view, time being economized by the operator preparing only small specimens of compounds.

Chemical synthesis and analysis, thoughtfully and conscientiously followed, without hurry and mere superficial consideration, but, of course, without undue expenditure of time, will insensibly carry the principles of Chemistry into the mind, and fix them there indelibly.

Note.—As a general rule, throughout this Manual paragraphs describing experiments to be performed are distinguished from paragraphs containing matter merely to be read by being printed in somewhat larger type.

Elements and their Atomic Weights, etc.—For an alphabetical list see the two pages immediately preceding the Index at the end of this volume.

THE BASYLOUS RADICALS.

POTASSIUM.

Symbol, K. Atomic weight, 39.

Formula, K_2 . Probable molecular weight, 78.

Memoranda.—The chief sources of potassium salts* are the chloride found at Stassfurt, in Prussia, as the mineral *Carnallite* (KCl ,

* The ill-defined term *salt* includes most solid definite chemical substances, but more especially those which assume a crystalline form.

$\text{MgCl}_2, 6\text{H}_2\text{O}$); *Kainite*, a double potassium and magnesium sulphate, with magnesium chloride, also occurring among the Stassfurt minerals; the nitrate, found in soils, especially in warm countries; and the compounds of potassium existing in plants. The vegetable salts of potassium are converted into carbonate (other salts are present) when the wood or other parts are burned to ashes. If the ashes be lixiviated with water and the solution evaporated to dryness, the residue when fused constitutes *crude potashes*. The residue, calcined on the hearth of a reverberatory furnace till white, gives the product termed *pearlash*, or impure potassium carbonate. Large quantities of carbonate are thus produced in North America and Russia, and, latterly, from the sugar beet-root marc in France. From the native chloride, and from the carbonate purified by treating the pearlash with its own weight of distilled water, filtering, and evaporating the solution so formed until it thickens, and stirring constantly, "so as to form a granular salt" (*Potassii Carbonas*, B. P., " K_2CO_3 ", with about 16 per cent. of water of crystallization"), nearly all other compounds of potassium are made. Exceptions occur in cream of tartar (*Potassii Bitartras*, U. S. P.), which is the more or less purified natural potassium salt of the grapevine, and in potassium nitrate. Potassium is a constituent of between forty and fifty chemical or galenical preparations of the pharmacopœias.

Potassium carbonate (*Potassii Carbonas*, U. S. P.) is a white crystalline or granular powder, insoluble in alcohol, very soluble in water, rapidly liquefying in the air through absorption of moisture, alkaline and caustic to the taste. It loses all water at a red heat. The official salt should contain 95 per cent. of real potassium carbonate (K_2CO_3).

Preparation.—Potassium itself is isolated with some difficulty by distilling a mixture of its carbonate and charcoal, or by Castner's method (see Sodium). It rapidly oxidizes in the air, hence is always kept below the surface of mineral naphtha, a liquid containing no oxygen. It crystallizes in octahedra.

Quantivalence.—The atom of potassium is univalent, K' .

REACTIONS HAVING (a) SYNTHETICAL AND (b) ANALYTICAL INTEREST.

(a) *Synthetical Reactions.*

These are utilized in making preparations of potassium. The word *synthesis* is from σύνθεσις (*synthēsis*), a putting together; *analysis*, from ἀναλύω (*analuō*), to resolve.

Potassium Hydrate.

Synonyms.—Caustic Potash; Potassium Hydroxide; Potassa; Hydrate of Potassium.

First Synthetical Reaction.—Boil together, for a few minutes, in a basin, 5 or 6 grains of potassium carbonate (K_2CO_3) and a like quantity of slaked lime (Ca_2HO) with a small quantity of water. Set the mixture aside till all solid matter has subsided.

This liquid is a solution of caustic potash, or potassium hydrate (KHO). Made of a prescribed strength, about 5 per cent., it forms the *Liquor Potassæ*, U. S. P.

The mixture is known to be boiled long enough when a little of the clear liquid, poured into a test-tube and warmed, gives no effervescence on the addition of an acid (sulphuric, hydrochloric, or acetic)—a test whose mode of action will be explained hereafter.

In the United States Pharmacopœia the potassium carbonate for this operation is directed to be obtained by boiling a solution of the bicarbonate until effervescence ceases; that is, until the bicarbonate is almost entirely converted into carbonate.

Best Method of Expressing Decomposition.—This will be easy of comprehension if what has already been stated concerning symbols and formulæ on pp. 40 to 42 and 46 and 47 has been carefully and thoughtfully considered. The best means of showing on paper the action which occurs when chemical substances attack each other is by the employment either of equations or diagrams setting forth the formulæ of the molecules concerned in the reaction. In an *equation* the formulæ of the salts used are written on one line, the sign of addition (+) intervening; the sign of equality (=) follows, and then the formulæ of the salts produced, also separated by a plus sign (+). Thus:



In this reaction (the operation just performed) the metals of (the molecules of) the two salts change places: from K_2CO_3 and Ca_2HO there are produced $CaCO_3$ and KHO (two molecules $2KHO$); from potassium carbonate and calcium hydrate there result calcium carbonate (the insoluble portion) and potassium hydrate (in solution).*

In constructing a *diagram* or pictorial illustration of a chemical reaction (the reaction, for instance, just described), *first* the formulæ of the salts used are written under each other on the left side of the leaf of a note-book, thus:




Such formulæ are, in this Manual, always given with the description of the reaction. *Secondly*, on the right is then written the formula of the chief substance produced, thus:


* If the student is already accustomed to the use of ordinary equations, he may pass on to Note 1 on p. 65; if not, the author would strongly recommend the temporary employment of diagrams for expressing chemical changes. Indeed, the occasional use of graphic equations or diagrams is of advantage to all students. For while an equation or a diagram equally well records the formulæ of the salts concerned in the whole reaction, the diagram alone suggests the *mode* in which its writer believes the respective atoms to change their positions. In the paragraphs succeeding the above, detailed explanations are given respecting the use and construction of diagrams.



Thirdly, the formation of this chief body under consideration—that is to say, both the origin of its elements and their destination—is traced out by the help of brackets and letters (which show the source of the elements) and converging lines (which suggest the approach and final union of those elements), thus :




For the next stage (at other stages, perhaps, in other reactions) the reader's own intelligent power of thought and reflection *must* come into exercise. He *must* reason somewhat as follows: "I am converting, and entirely converting, a quantity of potassium carbonate into potassium hydrate. A molecule, the smallest quantity I can picture on paper, of the potassium carbonate (K_2CO_3) contains, I am told, two atoms of potassium (K_2), and a molecule of the hydrate (KHO) one atom (K). Therefore—*therefore*—each molecule of the carbonate (K_2CO_3) will furnish *two* molecules of the hydrate (2KHO). Moreover, I notice that in the formula of a molecule of the calcium hydrate (slaked lime) I employ there are 2 of the HO (that is, 2HO); and this fact confirms me in the deduction that one molecule of the carbonate affords *two* molecules of the hydrate." The pupil will then amend his diagram thus :



Fourthly, the question as to what becomes of the other elements must be cleared up. Indeed, when the reader remembers that he is studying this reaction for the aid it affords him in learning Chemistry, and not because he is desirous of manufacturing caustic potash, he will see that this latter part of the reaction is quite as important as the former. To complete the diagram, then, he must first know what other compound is produced, and its formula. The context of his Manual will afford this information. In this case calcium carbonate is produced (CaCO_3). (This product is, in fact, pre-


precipitated chalk, together with any excess of slaked lime and any natural impurities in the slaked lime. *Pure* "precipitated chalk" is made by an analogous reaction, described subsequently.) The source of the elements of the calcium carbonate, and, finally, their union, must be indicated just as the source and mode of formation of the potash were indicated; that is to say, after the formula of this second substance produced (CaCO_3) is written on the right hand of the diagram, thus—



the source of its elements is shown by writing the symbols for those elements on the right of the bracket attached to the formula containing the symbols of the elements, thus :



Lines converging from the symbols of these elements also, and uniting at the formula of the substance (CaCO_3), are then drawn to suggest approach of the atoms of the elements and their union to form a molecule of the compound. The diagram will now be complete, and will have been built up in the student's note-book thus :



The formation of a third product or a fourth product would be indicated in a similar manner.

Note 1.—It will be seen that the chief data required in making either equationary or diagrammatic notes of decompositions are the symbolic formulæ of the various compounds employed and produced. These formulæ are, in this Manual, given whenever necessary. Chemists obtain them in the first instance by help of quantitative analysis. By the latter means also is obtained a check on the probabilities respecting the relative number of molecules concerned in a reaction.

Note 2.—While an equation or a diagram is an attempt to picture the reaction which ensues when molecules of different substances

act upon one another, it necessarily only represents two or a minimum number of the molecules. The student will of course understand that what is true of these two or three molecules is true of the thousands or millions of molecules forming the mass or whole quantity of material on which he experiments.

Note on Nomenclature.—*Hydrates* are bodies indirectly or directly derived from water by one-half of its hydrogen becoming displaced by an equivalent quantity of another radical. Thus, a piece of potassium thrown on to water (HHO) instantly liberates hydrogen, potassium hydrate (KHO) being formed. The temperature produced at the same time is sufficiently high to cause ignition of the hydrogen, which burns with a purple flame (owing to the presence of a little vapor of potassium), while the potassium hydrate remains dissolved in the bulk of the water. This radical, or root, or group of elements (HO), common to all hydrates, is called *hydroxyl*. Water might be termed hydrogen hydrate, or hydrogen hydroxylide, or hydrogen hydroxide.

Explanation.—With regard to the group of atoms represented by the symbols CO_3 and HO only a few words need be said here. The former (CO_3) is the grouping (root or radical) found in all the molecules of all carbonates; it is termed the carbonic radical, and is as characteristic of the molecules of carbonates as potassium (K) is of the molecules of potassium salts. HO (hydroxyl) is characteristic of all molecules of all hydrates. CO_3 is a bivalent root, HO is univalent; hence the group of atoms represented by CO_3 is found united with two univalent atoms, as in potassium carbonate, K_2CO_3 , or with one bivalent atom, as in calcium carbonate, CaCO_3 ; and HO is found united in single proportion with univalent atoms, as in a molecule of potassium hydrate, KHO, or in double proportion with bivalent atoms, as in a molecule of calcium hydrate, Ca_2HO . The quantivalence of a metal has only to be learnt, and the formulæ of its carbonate and hydrate are ascertained without seeing the formula of either; and this principle applies to formulæ of all other metallic salts. But, beyond committing to memory the formulæ and quantivalence of the various groupings characteristic of carbonates, hydrates, nitrates, sulphates, acetates, etc. (see the following table), special attention should not at present be devoted to the subject of the constitution of salts, but restricted to what may be called the metallic or basylous side of salts. The formulæ and quantivalence of the chief acidulous groupings referred to, and the symbols and quantivalence of allied elementary bodies, are included in the following table:

Formulæ and Quantivalence of Acidulous Radicals.

All chlorides contain	Cl	Univalent radicals.
“ bromides “	Br	
“ iodides “	I	
“ cyanides “	CN	
“ hydrates “	HO	
“ nitrates “	NO ₃	
“ chlorates “	ClO ₃	
“ acetates “	C ₂ H ₃ O ₂	Bivalent radicals.
“ oxides “	O	
“ sulphides “	S	
“ sulphites “	SO ₃	
“ sulphates “	SO ₄	
“ carbonates “	CO ₃	
“ oxalates “	C ₂ O ₄	Trivalent radicals.
“ tartrates “	C ₄ H ₄ O ₆	
“ citrates “	C ₆ H ₅ O ₇	
“ phosphates “	PO ₄	
“ borates “	BO ₃	

Radicals.—The above elements and compounds are termed *radicals*, each being the common *root* (*radix*) in a series of salts. Why compound radicals (as NO₃, SO₄, PO₄, etc.) differ in quantivalence need not be fully explained at present. Their constituent atoms doubtless always exert the same amount of attractive force, nearly but not quite all this force being exerted in retaining the atoms in one group, and the remainder probably determining the quantivalence. Compound radicals are more or less stable groups of atoms, capable of migration without change, but not necessarily capable of existing in the free state.*

Liquor Potassæ.

Liquor Potassæ, B. P., is officially directed to be made as follows:†

* Some modern chemical authors term these roots *radicles*, a word more usefully expressive of little roots or rootlets. The word *radicle* is indeed thus used as a diminutive in botany.

† The official processes will not usually be reprinted in this Manual, belonging, as they do, more to the manufacturing than to the educational side of practical chemistry. But pupils should take care that they *understand* every word of the processes, whether as regards chemical or mechanical details: the medical student in order that he may learn chemical principles—the principles on which the successful practice of modern medicine depends; the pharmaceutical student in order that he too may know the principles on which his avocation largely depends, and

Dissolve 1 pound of carbonate of potassium in 1 gallon of water; heat the solution to the boiling-point in a clean iron vessel, gradually mix with it 12 ounces of washed slaked lime (obtained from about 13 ounces of slaked lime washed with distilled water until a little of the washings, acidified with nitric acid, gives no cloudiness with nitrate of silver), and continue the ebullition for ten minutes with constant stirring. Then remove the vessel from the fire, and when, by the subsidence of the insoluble matter, the supernatant liquor has become perfectly clear, transfer it by means of a siphon or by decantation to a green-glass bottle furnished with an air-tight stopper, and add distilled water, if necessary, to make it correspond with the tests of *specific gravity* and *neutralizing power*. (These tests and the modes of applying them will be explained in subsequent sections.)

Liquor Potassæ, U. S. P., is prepared in a very similar way, starting with the bicarbonate, which by ebullition resolves itself into potassium carbonate, water, and carbonic acid gas:




Potassa cum Calce, U. S. P., is a grayish-white powder, made by rubbing together equal weights of solid potash and quicklime.

Solid Potash.—Solution of potash evaporated to dryness in a silver or clean iron vessel, and the residue fused and poured into moulds, constitutes *Potassa*, U. S. P. It often contains chlorides, detected by silver nitrate, and sulphates, detected by a barium salt, as described subsequently in connection with hydrochloric and sulphuric acids.

Sulphurated Potash.

Synonyms.—Sulphurated Potassa; Liver of Sulphur; Impure Potassium Sulphide.

Second Synthetical Reaction.—Into a test-tube put a few grains of a mixture of *previously dried* potassium carbonate with half its weight of sulphur. Heat the mixture gradually until it ceases to effervesce. The resulting fused mass, poured on a slab and quickly bottled, is the *Potassa Sulphurata*, sulphurated potash, of the United States Pharmacopœia.




in order that he may apply those principles in his pursuit of that avocation. With such views before him each student should carefully read through the above process, and in future, after making an officially interesting synthetical experiment, should turn to the corresponding monograph in his pharmacopœia and assure himself that he *understands* all that is there stated.

As met with in pharmacy, this salt is not a single definite chemical compound, but a mixture of several; in short, its chemical character is well indicated by its vague name. When fresh, and if carefully prepared with the official proportions of dry ingredients, it is of the color of liver (whence the old name "liver of sulphur"), and consists, as shown by J. Watts, of the salts mentioned in the foregoing equation, together with a little undecomposed potassium carbonate, with perhaps higher sulphides of potassium (K_2S_4 and K_2S_5); but, rapidly absorbing oxygen from the air, it soon becomes green, and yellow potassium sulphite (K_2SO_3) and sulphate ($K_2S_4O_4$) are successively formed, and ultimately a useless mass of a dirty-white color results, consisting of sulphate and hyposulphite, with generally some potassium carbonate and free sulphur. Moreover, if overheated in manufacture, the hyposulphite $4(K_2S_2O_3)$ is decomposed into potassium sulphate $3(K_2S_4O_4)$ and sulphide (K_2S_5). Recently made, about 50 per cent. should be soluble in rectified spirit. It is occasionally employed in the form of ointment (*Unguentum Potassæ Sulphuratæ*, B. P.).

"On triturating 1 grm. of sulphurated potash with 1 grm. of crystallized copper sulphate and 10 cc. of water, and filtering, the filtrate should remain unaffected by hydrogen sulphide, corresponding to at least 12.85 per cent. of sulphur combined with potassium to form sulphide."—*U. S. P.*

The extremely—indeed, most unusually—complicated nature of the decomposition will probably cause failure in any attempt by a student to draw out an equation or a diagram of the reaction without the aid of the printed equation given above. He may therefore content himself in this case by introducing into his note-book a diagram founded directly on the equation and on the numbers of molecules there stated. With this printed equation, and the details of construction of diagrams given in connection with the first syn-



Crucibles of Various Forms.

thetical reaction, he will be able to give a diagram of this second synthetical reaction without unduly troubling his reasoning powers, while at the same time he will be familiarizing himself with the more mechanical portions of the diagram.

In preparing large quantities of sulphurated potash the test-tube is replaced by an earthenware vessel termed a *crucible*


(possibly from *crux*, a cross, for originally a cross was impressed upon the melting-pot as used by alchemists and goldsmiths: others derive the word from *crux*, an instrument of torture, the sense here being symbolical).

Heating Crucibles.—Crucibles of a few ounces' capacity may be heated in an ordinary grate-fire. Larger ones require a stove with a good draught—that is, a *furnace*. Even the smaller ones are more conveniently and quickly heated in a furnace. Half-ounce or one-ounce experimental porcelain crucibles may be heated in a spirit or gas flame, the air-gas flame already described being generally the most suitable.

Potassium Acetate.

Synonyms.—Acetate of Potassium; Acetate of Potash.

Third Synthetical Reaction.—Place 10, 20, or more grains of potassium carbonate in a small dish, and *saturate* (*satur*, full) with acetic acid; that is, add acetic acid so long as effervescence is thereby produced; the resulting liquid is a strong, slightly acid solution of potassium acetate. Evaporate most of the water in an open dish (see Figs. 15 and 16, p. 71), stirring with a glass rod* to promote evolution of vapor; a white salt remains, which fuses on the further careful application of heat: this is the official potassium acetate (*Potassii Acetas*, U. S. P.), formerly termed acetate of potash. If fused in the open vessel, the acetate is liable to become slightly charred and discolored; this is prevented by transferring the solid residue to a test-tube or Florence flask before finally fusing. It forms a white deliquescent, foliaceous, satiny mass, neutral to test-paper and wholly soluble in spirit. A 10 per cent. solution in water forms the "Solution of Acetate of Potassium," B. P.



Explanation of Formulæ.—The formula for one molecule of acetic acid (hydrogen acetate) is $\text{HC}_2\text{H}_3\text{O}_2$, and one of potassium acetate $\text{KC}_2\text{H}_3\text{O}_2$. The grouping, $\text{C}_2\text{H}_3\text{O}_2$, is characteristic of all acetates; it is univalent.

Explanation of Process.—When two molecules of acetic acid ($2\text{HC}_2\text{H}_3\text{O}_2$) and one of potassium carbonate (K_2CO_3) react, two molecules of potassium acetate ($2\text{KC}_2\text{H}_3\text{O}_2$) and one of carbonic acid (H_2CO_3) are produced, the latter at once splitting up into water (H_2O) and carbonic acid gas (CO_2), as already shown in the equation.

* Glass rod is usually purchased in the form of long sticks. The pieces may be cut to convenient lengths of from six to twelve inches (*vide* p. 16), sharp ends being rounded off by holding the extremity in a flame for a few minutes.

Diagram of the Reaction.—The nature of the above operation is indicated by an *equation*; it (and succeeding reactions) should be expressed in the student's note-book as a *diagram*, and, if possible, without the aid of the above equation.

Note.—The foregoing reaction has a general as well as a special synthetical interest. It represents one of the commonest methods of forming salts—namely, the saturation of a carbonate by an acid, or *vice versâ*; carbonates added to acetic acid yield acetates, to nitric acid nitrates, to sulphuric acid sulphates. Many illustrations of this general process occur in pharmacy.

Evaporation of water from a liquid is best conducted in wide shallow vessels rather than in narrow deep ones, as the steam can thus quickly diffuse into the air and be conveyed rapidly away; hence a small round-bottomed basin, heated as shown in Fig. 15, is far more suitable than a test-tube for such operations. On the manufacturing scale iron or iron lined with enamel or semi-porcelain, copper, tinned copper, or solid tinned pans are used. Up to twelve or eighteen inches diameter, pans, basins, or dishes made of Wedgwood ware or porcelain composition (Fig. 16) may be employed. Small dishes may be supported by retort stands (Fig. 15), larger by cylinders (Fig. 16), to which the dish is, if less in diameter than the cylinder, adapted by such flat rings or diaphragms as are shown in the figure.




Fig. 15.





Fig. 16.

Evaporation from Small and Large Basins.

Potassium Bicarbonate.

Synonyms.—Bicarbonate of Potassium; Bicarbonate of Potash.

Fourth Synthetical Reaction.—Make a *strong* solution of potassium carbonate by heating in a test-tube a mixture of several grains of the salt with rather less than an equal weight of water. Through the cool solution pass carbonic acid gas slowly but continuously; after a time a white crystalline precipitate of potassium acid carbonate or bicarbonate (KHCO_3) of the Pharmacopœia (*Potassii Bicarbonas*, U. S. P.) will be formed.




The carbonic acid gas for this operation is economically and conveniently prepared from small lumps of marble, though it might be obtained from any carbonate; thus the previous synthetical reaction could be made available for this purpose, the carbonic acid gas evolved on the addition of the acetic acid to the potassium carbonate being conducted into a strong solution of more potassium carbonate by a glass tube bent and fitted as described when treating of oxygen gas. Economy also causes hydrochloric acid to be used in preference to acetic or any other acid.

Generate the carbonic acid gas by adding common hydrochloric acid, diluted with twice its bulk of water, to a few fragments of marble contained in a test-tube or small flask, and conduct the gas into the solution of potassium carbonate by a glass tube bent to a convenient angle or angles and fitted to the test-tube by a cork in the usual way (see Fig. 10, p. 28, though no heat is necessary). The tube may be replenished with marble or acid, or both, when the evolution of gas is becoming slow. In working on any larger quantity than a few grains of the carbonate a wide delivery-tube should be employed, or the end of the narrow tube occasionally be cleared from any bicarbonate that may have been deposited in it. A more economical arrangement of the apparatus employed in this process will be described under the corresponding sodium salt (p. 88).

Deposition of the Bicarbonate explained.—Potassium bicarbonate is to a certain extent soluble in water, but as it is less so than the potassium carbonate, and as a *saturated* solution of the latter has been used, a precipitation of a part of the bicarbonate inevitably occurs. In other words, the quantity of water present is sufficient to keep the carbonate, but insufficient to retain the equivalent quantity of bicarbonate, in solution.

Properties.—Prepared on the large scale, potassium bicarbonate occurs in colorless, non-deliquescent right rhombic prisms; it has a saline, feebly alkaline, non-corrosive taste. Heated to redness, it loses 31 per cent. of its weight, and is converted into carbonate (K_2CO_3), water (H_2O), and carbonic acid gas (CO_2).



The foregoing equation and accompanying molecular weights (see page 55) show how potassium bicarbonate, the molecular weight of which happens to be 100, *must* lose 31 per cent. ($9 + 22$) when completely decomposed by heat.

Effervescing Solution of Potash.—A solution of 30 grains of this bicarbonate in 1 pint of water, charged with five times its bulk of carbonic acid gas by pressure, is medicinal “potash-water,” *Liquor Potassæ Effervescens*, B. P.


Notes on Nomenclature.—The prefix *bi-* in the name “potassium bicarbonate” serves to recall the fact that to a given amount of potassium this salt contains *twice* as much carbonic radical as the carbonate. The salt is really “potassium and hydrogen carbonate” (KHCO_3); it is intermediate between potassium carbonate (K_2CO_3) and hydrogen carbonate or true carbonic acid (H_2CO_3); it is “acid potassium carbonate” or “hydrogen potassium carbonate.” Hence in constitution it is an acid salt, although not acid to the taste.

Salts whose specific names end in the syllable “*ate*” (carbonate, sulphate, etc.) are in general conventionally so termed when they contain the radical, or the characteristic elements, of an acid whose name ends in “*ic*,” and from which acid they have been or may be formed. Thus the syllable “*ate*” in the words sulphate, nitrate, acetate, carbonate, etc. indicates that the respective salts contain the radical of an acid whose name ends in *ic*, the previous syllables, sulph-, nitr-, acet-, carbon-, indicating what that acid is—sulphuric, nitric, acetic, or carbonic. Occasionally a letter or syllable is dropped from or added to a word to render the name more euphonious; thus the sulphuric radical forms *sulphates*, not sulphurates, and the tartaric radical yields *tartrates*, not tartarates.

Potassium Citrate.

Synonyms—Citrate of Potassium; Citrate of Potash.

Fifth Synthetical Reaction.—Dissolve a few grains or more of potassium bicarbonate in water, and add citric acid ($\text{H}_3\text{C}_6\text{H}_5\text{O}_7$) until it no longer causes effervescence. The resulting liquid is a solution of potassium citrate ($\text{K}_3\text{C}_6\text{H}_5\text{O}_7$). Evaporate to dryness in an open dish cautiously, so as to avoid charring; a pulverulent or granular residue is obtained, which is the official *Potassii Citras* (U. S. P.), a white deliquescent powder.



Citrates.—The citric radical or group of elements, which with three atoms of hydrogen forms a molecule of citric acid, and with three of potassium, potassium citrate, is a trivalent grouping; hence the three atoms of potassium in a molecule of the citrate. The full chemistry of citric acid and other citrates will be described subsequently.

Potassium Nitrate.

Synonyms.—Nitrate of Potassium; Nitrate of Potash; Nitre.

Potassium Nitrate (KNO_3) (*Potassii Nitras*, U. S. P.), the old nitrate of potash, and

Potassium Sulphate.

Synonyms.—Sulphate of Potassium; Sulphate of Potash.


Potassium Sulphate (K_2SO_4), (*Potassii Sulphas*, U. S. P.), could

obviously also be made by saturating nitric acid (HNO_3) and sulphuric acid (H_2SO_4), respectively, with potassium carbonate. Practically, they are not made in that way—the nitrate occurring, as already stated, in nature, and the sulphate as a by-product in many operations. Both salts will be alluded to hereafter in connection with nitric acid.

Potassium Tartrate.

Synonyms.—Tartrate of Potassium; Tartrate of Potash.

Sixth Synthetical Reaction.—Place a few grains of potassium carbonate in a test-tube with a little water, heat to the boiling-point, and then add acid potassium tartrate ($\text{KHC}_4\text{H}_4\text{O}_6$) till there is no more effervescence; a solution of neutral potassium tartrate ($\text{K}_2\text{C}_4\text{H}_4\text{O}_6$) results, the *Potassii Tartras* of the United States Pharmacopœia, the old tartrate of potash or “soluble tartar.” Crystals (four- or six-sided prisms) may be obtained on concentrating the solution by evaporation and setting the hot liquid aside. Larger quantities are made in the same way, 20 of acid tartrate and 9 of carbonate (with 50 of water) being about the proportions necessary for neutrality.




Tartrates.— $\text{C}_4\text{H}_4\text{O}_6$ are the elements characteristic of all tartrates: they form a bivalent grouping; hence the formula of the hydrogen tartrate, or tartaric acid, is $\text{H}_2\text{C}_4\text{H}_4\text{O}_6$; that of the potassium tartrate, $\text{K}_2\text{C}_4\text{H}_4\text{O}_6$; of the intermediate salt, the acid potassium tartrate (cream of tartar), $\text{KHC}_4\text{H}_4\text{O}_6$. If the acid tartrate of one metal and the carbonate of another react, a neutral dimetallic tartrate results, as seen in Rochelle salt ($\text{KNaC}_4\text{H}_4\text{O}_6$), the *Soda Tartarata* of the British Pharmacopœia (*Potassii et Sodii Tartras*, U. S. P.).

Acid Salts (e. g. $\text{KHC}_4\text{H}_4\text{O}_6$)—that is, salts intermediate in composition between a neutral salt (e. g. $\text{K}_2\text{C}_4\text{H}_4\text{O}_6$) and an acid (e. g. $\text{H}_2\text{C}_4\text{H}_4\text{O}_6$)—will frequently be met with. All acidulous radicals, except those which are univalent, may be concerned in the formation of such acid salts.


Potassium Iodide, or Iodide of Potassium.

Seventh Synthetical Reaction.—Into solution of potash, heated in a test-tube, flask, or evaporating basin according to quantity, stir a little solid iodine. The deep color of the iodine disappears entirely. This is due to the formation of the colorless salts, potassium iodide (KI) and potassium iodate (KIO_3), which remain dissolved in the liquid. Continue the addition of iodine so long as its color, after a few minutes' warming and stirring, disappears. When the whole of the potash in the solution of potash has been converted into the salts mentioned,

the slight excess of iodine remaining in the liquid will color it, and thus show that this stage of the operation is completed.



Separation of the Iodide from the Iodate.—Evaporate the solution to dryness. If each salt were required, the resulting mixture might be digested in spirit of wine, which dissolves the iodide, but not the iodate. But the iodide only is needed. Intimately mix the residue, therefore (reserving a grain or two for a subsequent experiment), with excess (about a third of its weight) of charcoal, and gently heat in a test-tube or crucible until slight deflagration ensues.* The crucible may be held in a spirit or air-gas flame or other fire by tongs. (Scissors-shaped and other “crucible-tongs” are sold by all makers of apparatus.) The iodide remains unaffected, but the iodate loses all its oxygen, and is thus also reduced to the state of iodide.



Treat the mass with a little water, and filter to separate excess of charcoal; a solution of pure potassium iodide results (*Potassii Iodidum*, U. S. P.). The latter may be used as a reagent or it may be evaporated to a small bulk and set aside to crystallize.

Properties.—Potassium iodide crystallizes in small cubical crystals, very soluble in water, less so in spirit. One part in twenty of water forms “Solution of Iodide of Potassium,” U. S. P. Exposed to air and sunlight, pure potassium iodide becomes slightly brown, owing to the liberation of iodine. Under these circumstances a little potassium carbonate is produced by action of the atmospheric carbonic acid, hydriodic acid (HI) is set free, and the latter, attacked by oxygen, yields a trace of water and of free iodine.

The addition of charcoal in the above process is simply to facilitate the removal of the oxygen of the potassium iodate. Potassium iodate (KIO_3) is analogous in constitution and in composition, so far as the atoms of oxygen are concerned, to potassium chlorate


* If, in the operation of heating potassium iodate with charcoal, excess of the latter be employed, slight incandescence rather than deflagration occurs; if the charcoal be largely in excess, the reduction of the potassium iodate to iodide is effected without visible deflagration or even incandescence.

Deflagration means violent burning, from *flagratus*, burnt (*flagro*, to burn), and *de*, a prefix augmenting the sense of the word to which it may be attached. Paper thrown into a fire simply burns, nitre deflagrates. *Detonate* (*detono*) is a precisely similar word, meaning to explode with violent noise.

(KClO_3), which has already been stated to be more useful than any other salt for the actual preparation of oxygen gas itself. Hence the removal of the oxygen of the iodate might be accomplished by heating the residue without charcoal. In that case the liberated oxygen would be detected on inserting the incandescent extremity of a strip of wood into the mouth of the test-tube in which the mixture of iodide and iodate had been heated. The charcoal, however, burns out the oxygen more quickly, and thus economizes both heat and time.

Note.—The formula of potassium iodide (KI) shows that the salt contains potassium and iodine in atomic proportions. A reference to the table of Atomic Weights at the end of the volume and a rule-of-three sum would therefore show what weight of salt is producible from any given weight of iodine.

Detection of Potassium Iodate in Potassium Iodide.—Potassium iodate remaining as an impurity in potassium iodide may be detected by adding to a solution of the latter salt some weak acid (say, tartaric), shaking, and then adding starch mucilage; blue "starch iodide" is formed if a trace of iodate be present, but not otherwise. By the reaction of the added acid and the potassium iodate iodic acid (HIO_3) is produced; and by reaction of the added acid and the potassium iodide hydriodic acid (HI) is produced; neither of these alone attacks starch, but by reaction on each other they give free iodine, which then forms the blue color. This experiment should be tried on a grain or two of pure iodide and on the impure iodide reserved from the previous experiment. Potassium iodide containing iodate would obviously yield free iodine, which is excessively corrosive, on the salts coming into contact with the acids of the stomach.




Note on Nomenclature.—The syllable *ide* attached to the syllable *iod* in the name "potassium iodide" indicates that the *element* iodine is combined with the potassium. An *iodate*, as already explained, is a salt containing the characteristic elements of iodic acid and of all iodic compounds. Inorganic salts, one of whose names ends in *ide*, are those which are, or may be, formed from *elements*. The names of salts which are, or may be, formed from compounds include other syllables, *ate* being one (see page 73). The only other syllable is *ite*, which is included in the names of salts which are, or may be, formed from the acids and radicals whose names end in *ous*: thus sodium sulphite, etc. To recapitulate: An inorganic salt whose name ends in *ate* contains a compound acidulous radical whose name ends in *ic*; a salt whose name ends in *ite* contains a compound acidulous radical whose name ends in *ous*; an inorganic salt whose name ends in *ide* contains an element for its acidulous radical. Thus sulphide relates to sulphur, sulphite to the

sulphurous radical, sulphate to the sulphuric radical, and so on with other inorganic "ides," "ites," or "ates."


Potassium Bromide, or Bromide of Potassium.

Potassium Bromide (*Potassii Bromidum*, U. S. P.).—This salt is identical in constitution with potassium iodide, and is made in exactly the same way, bromine being substituted for iodine. The formula of bromic acid is HBrO_3 . It will be noticed that the following equations are similar in character to those showing the preparation of potassium iodide:




Potassium Manganates, or Manganates of Potassium.

Eighth Synthetical Reaction.—Place a fragment of solid caustic potash (KHO), with about the same quantity of potassium chlorate (KClO_3) and of black manganese oxide (MnO_2), on a piece of platinum-foil.* Hold the foil by a small pair of forceps or tongs in the flame of a blowpipe for a few minutes until the fused mixture has become dark green, apparently black. This color is that of *potassium manganate* (K_2MnO_4).



Ninth Synthetical Reaction.—*Potassium Permanganate* ($\text{K}_2\text{Mn}_2\text{O}_8$) (*Potassii Permanganas*, U. S. P.), the old permanganate of potash, which is purple, is obtained, or rather a solution of it, on placing the foil and its adherent mass in water and boiling for a short time.



On the large scale the potash set free in the reaction is neutralized by sulphuric or, better, carbonic acid, and the solution evaporated to the crystallizing point. Further details will be given in connection with Manganese.

* The foil may be one inch broad by two inches long. No ordinary flame will melt the platinum, fused caustic alkalies only slowly corrode it, and very few other chemical substances affect it at all; hence the same piece may be used in experiments over and over again. Most metals form a fusible alloy with platinum, and phosphorus rapidly attacks it; hence such substances, as well as mixtures likely to yield them, should be heated in a small porcelain crucible.

Solutions of (green) potassium and sodium manganates (or purple) permanganates so readily yield their oxygen to organic matter that they are used on the large scale as disinfectants.

Synthetical Reactions, bringing under consideration the remaining official compounds (namely, potassium bichromate, arsenite, chlorate, cyanide, ferrocyanide, and ferricyanide) are deferred at present.

Crystallization.—This operation will have been performed several times in the course of the foregoing synthetical experiments. Obviously it offers a mode of separating soluble crystallizable substances from soluble *amorphous* (*a*, without; *μορφή*, *morphe*, shape) substances; also of separating from each other substances of varying degrees of solubility or which crystallize with varying degrees of readiness—*fractional crystallization*.

(b) Reactions having Analytical Interest (Tests).

Note.—These are reactions utilized in searching for small quantities of a substance (in the present instance potassium) in a solution. They are best performed in test-tubes or other small vessels. Each reaction should be expressed in the form of an equation or diagram in the student's note book. *All previous or future equations given in this volume should be transferred to the note-book in the form of diagrams, constructed as described on pages 63 to 66, unless the student can with ease construct equations without the aid of the Manual.*

First Analytical Reaction.*—To a solution of any potassium salt (chloride,† for example) add a few drops of hydrochloric acid and of a solution of platinum perchloride‡ (PtCl_4), and

* As already indicated, chemical reactions are scarcely analytical or synthetical in themselves, but rather performed with an analytical or synthetical object. Indeed, not infrequently one and the same reaction is both a synthetical and an analytical reaction. Thus this first so-called "analytical reaction" is a synthetical reaction if performed with the object of preparing a specimen of the double platinum and potassium chloride. It is an analytical reaction, or rather has analytical interest, if performed with the object of demonstrating the presence of potassium. Chemical reactions in themselves are operations not so much of analysis (resolution) or synthesis (combination), or of analysis and synthesis conjoined, as of what has sometimes been termed *metathesis* (transposition). Molecules are not torn to atoms in an operation performed with an analytical object, nor are the atoms put together or set together in an operation (perhaps the same operation) performed with a synthetical object; but in both operations the atoms of the molecules undergo metathesis; that is, exchange places or are transposed. In short, chemists use the words "analytical" and "synthetical" in a conventional rather than a strictly etymological sense.

† A few fragments of potassium carbonate, two or three drops of hydrochloric acid, and a small quantity of water give a solution of potassium chloride at once, $\text{K}_2\text{CO}_3 + 2\text{HCl} = 2\text{KCl} + \text{H}_2\text{O} + \text{CO}_2$.

‡ Experiments with such expensive reagents as platinum perchloride are economically performed in watch-glasses, drops of the liquids being operated on.

stir the mixture with a glass rod; a yellow granular or slightly crystalline precipitate* slowly forms. (This precipitate is the double platinum and potassium chloride, and its composition is expressed by the formula $\text{PtCl}_4 \cdot 2\text{KCl}$.)

Memoranda.—When the precipitate is long in forming, it is sometimes of an orange-yellow tint. If potassium iodide happens to be in the potassium salt under examination, some platinum iodide (PtI_4) will also be formed, giving a red color to the solution, and a larger quantity of the *precipitant* (that is, the precipitating agent) will be required.

Precaution.—Only potassium chloride forms this characteristic compound; hence if the potassium salt in the solution is known not to be a chloride, or if its composition is unknown, a few drops of hydrochloric acid must be added, otherwise some of the platinum perchloride will be utilized for its chlorine only, the platinum being wasted. Thus, if potassium nitrate (KNO_3) be the salt present, a little hydrochloric acid enables the potassium to assume the form of chloride when the platinum perchloride is added, nitric acid (HNO_3) being set free.

Explanations.—The precipitate is, practically, insoluble in water. It is for this reason that a very small quantity of any soluble potassium salt (or, rather, of the potassium in that salt) is thrown out of solution by platinum perchloride.

Note on Nomenclature.—When distinct molecules of salts unite and form a single crystalline compound, the product is often termed a *double salt*. The double potassium and platinum chloride is such a body.

Educational Note.—The thoughtful student will not confuse *the test with the chemistry of the test*. The test itself appeals to the senses—commonly to the eye, sometimes to the nose, occasionally to the ear. A person may be able to apply a test, and yet never know anything of the chemistry of a test.

Acid Potassium Tartrate.

Synonyms.—Acid Tartrate of Potassium; Acid Tartrate of Potash; Cream of Tartar.

Second Analytical Reaction.—To a solution of any salt of potassium add excess of strong solution of tartaric acid ($\text{H}_2\text{C}_4\text{H}_4\text{O}_6$), and shake or well stir the mixture; a white granular precipitate (acid potassium tartrate, $\text{KHC}_4\text{H}_4\text{O}_6$) will be formed.

Note.—By “excess” of any test-liquid (such as the “solution of tartaric acid” just mentioned) is meant such a quantity as is *probably* rather more than sufficient to convert the whole weight of the

* By *precipitation* (from *præcipitare*, to throw down suddenly) is simply meant the formation of particles of solid in a liquid, no matter whether the solid, the *precipitate*, subsides or floats, and no matter whether the operation be entire and complete or partial and fractional.

compound attacked into the compound produced. Thus in the present case enough acid must be added to convert the whole of the potassium salt operated on into acid potassium tartrate. What the weight of salt operated on was must be mentally estimated, roughly, by the operator. It is not necessary in analyzing to know the exact weights of salts employed. The analyst must use his judgment, founded on his knowledge of the reaction (as shown by an equation) and of the molecular weights of the substances employed in the reaction, as well as by the rough estimate of the amount of material on which he is experimenting.

Limits of the Tests.—Acid potassium tartrate is soluble in about 180 parts of cold and in 6 parts of boiling water. Hence, in applying the tartaric test for potassium the solutions must not be hot. Even if cold, no precipitate will be obtained if the solutions are very dilute. This test, therefore, is of far less value than that first mentioned. The acid potassium tartrate is less soluble in diluted alcohol than in water, so that the addition of spirit of wine renders the reaction somewhat more delicate.

Cream of Tartar.—The precipitate is the *Potassii Bitartras*, U. S. P., the old acid tartrate or bitartrate of potash or cream of tartar, though the official preparation is not formed in the above manner; on the contrary, the acid is derived from the salt, which, often mixed with some calcium tartrate, occurs naturally in the juice of many plants.

Third Analytical Reaction.—The *flame-test*. Dip the looped end of a platinum wire into a solution containing a potassium salt, and introduce the loop into the lower part of a spirit flame, the flame of a mixture of gas and air, a blowpipe flame, or other slightly colored flame. A light violet or lavender tint will be communicated to the flame, an effect highly characteristic of potassium salts.

Fourth Analytical Fact.—Potassium salts are not readily volatile. Place a fragment of carbonate, nitrate, or other potassium salt on a piece of platinum-foil, and heat the latter in the flame of a lamp; the salt may fuse to a transparent liquid and flow over the foil; water also, if present, will escape as steam, and black carbon be set free if the salt happen to be a tartrate, citrate, etc.; but the potassium compound itself will not be vaporized. This is a valuable negative property, as will be evident when the analytical reactions of ammonium come under notice.

Other Tests.—Sodium cobaltic nitrite is an official test for potassium.

QUESTIONS AND EXERCISES.

Name the sources of potassium.—Give the source, formula, and characters of potassium carbonate.—Distinguish between synthetical and

analytical reactions.—How is the official *Liquor Potassæ* prepared?—What is the systematic name of caustic potash?—State the chemical formula of caustic potash.—Construct an equation or diagram expressive of the reaction between potassium carbonate and slaked lime.—Define a *hydrate*.—What group of atoms is characteristic of all carbonates?—Define the term *radical*.—How is “sulphurated potash” made, and of what salts is it a mixture?—What is the formula of the radical of all acetates?—Draw a diagram showing the formation of potassium acetate.—Give a process for the conversion of carbonates into other salts.—What is the difference between potassium carbonate and bicarbonate?—How is the latter prepared?—What is the relation between salts whose specific names end in the syllable “ate” and acids ending in “ic”?—Draw out diagrams or equations descriptive of the formation of potassium tartrate from the acid tartrate, and citrate from potassium carbonate.—Distinguish between a neutral and an acid salt.—How is potassium iodide made?—Illustrate the process either by diagrams or equations.—Work out a sum showing how much potassium iodide is producible from 1000 grains of iodine. *Ans.* 1307 grains.—Give a method for the detection of iodate in potassium iodide.—Explain the reaction.—Has the syllable “ide” any general signification in chemical nomenclature?—State the relations between sulphides, sulphites, and sulphates.—Mention the chemical relation of potassium bromide to iodide.—Describe the formation of potassium permanganate, giving equations or diagrams.—How do manganates and permanganates act as disinfectants?—Enumerate the tests for potassium, explaining by diagrams or equations the various reactions which occur.

SODIUM.

Symbol, Na. Atomic weight, 23.

Formula, Na₂. Probable molecular weight, 46.

Memoranda.—Most of the sodium salts met with in pharmacy are obtained directly from sodium carbonate, which is now manufactured on an enormous scale from sodium chloride (common salt, sea-salt, bay-salt, or rock-salt), the natural source of the sodium salts. When pure, salt (*Sodii Chloridum*, U. S. P.) occurs “in small white crystalline grains or transparent cubic crystals, free from moisture;” the best varieties commonly contain a little magnesium chloride and sometimes other impurities. Besides the direct and indirect use of sodium carbonate—or carbonate of soda, as it is commonly called—in medicine, it is largely used for household cleansing purposes under the name of “soda” and in the manufacture of soap. Sodium nitrate also occurs in nature, but is valuable for its nitric constituents rather than its sodium. Sodium is a constituent of about forty chemical or galenical preparations of the pharmacopœias.

Sodium (*Sodium*, B. P.) is prepared by a process similar to that for potassium, but with less difficulty. Castner obtains it comparatively cheaply by distillation from a mixture of soda, carbon, and iron contained in steel vessels. It has a bright metallic lustre when freshly cut, but rapidly absorbs oxygen and carbonic acid gas from the air, and thus becomes coated with sodium carbonate. It displaces hydrogen from water, yielding solution of sodium hydrate; but unless the sodium is confined to one spot by placing it on a small floating piece of filter-paper, the action is not sufficiently intense to cause ignition of the escaping hydrogen. When the

latter does ignite, it burns with a yellow flame, due to the presence of a little vapor of sodium.



It similarly attacks alcohol, yielding "sodium ethylate" (see Index). It may be kept beneath the surface of a liquid containing neither moisture nor oxygen (mineral naphtha). It crystallizes in octahedra. Its atom is univalent, Na'.

REACTIONS HAVING (a) SYNTHETICAL AND (b) ANALYTICAL INTEREST.

(a) Reactions having *Synthetical Interest.*

Sodium Hydrate.

Synonyms.—Sodium Hydroxide; Caustic Soda; Hydrate of Sodium; Hydrate of Soda; Soda.

First Synthetical Reaction.—The formation of solution of sodium hydrate or caustic soda, NaHO (*Liquor Sodæ*, U. S. P.), the old hydrate of soda. This operation resembles that of making solution of potash, already described.



The practical student should apply to this solution the remarks made concerning solution of potash.


The official *Liquor Sodæ* is made from 170 grms. of crystals of sodium carbonate, 50 of washed slaked lime, and 780 cc. of water, under precisely similar circumstances to those detailed for *Liquor Potassæ* (p. 63). If the solution be evaporated to dryness and the residue fused and poured into moulds, solid sodium hydrate (*Soda*, U. S. P.) is obtained.

Commercial and cheap caustic soda is largely employed in various manufactures. This variety is a by-product in the manufacture of sodium carbonate, but, though highly useful as a chemical agent, is too impure for medicinal use. The United States Pharmacopœia recognizes liquor sodæ made from solid caustic soda about 56 parts and distilled water about 944, or from caustic soda of any other strength if only an equivalent amount be used and the product complies with the official tests.

Second Synthetical Reaction.—The reaction of sulphur and sodium carbonate at a high temperature resembles that of sulphur and potassium carbonate, but as the product is not used in medicine, the experiment may be omitted. It is mentioned here to draw attention to the close resemblance of the potassium salts to those of sodium.

Sodium Acetate, or Acetate of Sodium.


Third Synthetical Reaction.—Add sodium carbonate (in powder, or, better, in fragments) to some strong acetic acid in an evaporating basin as long as effervescence occurs, and then boil off some of the water.* When the solution is cold, crystals of Acetate of Sodium, B. P., *Sodii Acetas*, U. S. P. ($\text{NaC}_2\text{H}_3\text{O}_2, 3\text{H}_2\text{O}$), the old acetate of soda, will be deposited. A 10 per cent. solution in distilled water forms the "Solution of Acetate of Sodium," B. P.



Sodium acetate effloresces in dry air, and loses all its waters of crystallization when gently heated. It supports a temperature of 270° or 280° F. without decomposition, but above 300° soon chars.

Sodium Bicarbonate.

Fourth Synthetical Reaction.—The action of carbonic acid (H_2CO_3), or carbonic acid gas (CO_2) and water (H_2O), on sodium carbonate (Na_2CO_3) resembles that on potassium carbonate, but is applied in a different manner. The result is sodium bicarbonate (NaHCO_3), (*Sodii Bicarbonas*, *Bicarbonate of Sodium*, U. S. P.), the old bicarbonate of soda.



Process.—Heat crystals of sodium carbonate in a porcelain crucible until no more steam escapes. Rub the product in a mortar with two-thirds its weight of more of the crystals, and place the powder in a test-tube or small bottle, into which carbonic acid gas may be conveyed by a tube passing through a cork and terminating at the bottom of the vessel. To generate the carbonic acid gas, fill a test-tube having a small hole in the bottom (or a similar piece of glass tubing of which one end is plugged by a grooved cork) with fragments of marble, insert a cork and a delivery-




Fig. 17.
Preparation of Sodium Bicarbonate.

*The "water" alluded to occurs in the acid, which, though commonly termed "acetic acid," is really a solution of that acid in water.

tube, and connect the latter with the similar tube of the vessel containing the sodium carbonate by a piece of india-rubber tubing. Now plunge the tube of marble into a test-glass or other vessel containing a mixture of 1 part hydrochloric acid and 2 parts water, and loosen the cork of the sodium carbonate tube until carbonic acid gas, generated in the marble tube, may be considered to fill the whole arrangement; then replace the cork tightly and set the apparatus aside. As the gas is absorbed by the sodium carbonate, hydrochloric acid rises into the marble tube and generates fresh gas, which, in its turn, drives back the acid liquid, and thus prevents the production of any more gas until further absorption has occurred. When the salt is wholly converted into bicarbonate (NaHCO_3) it will be found to have become damp through the liberation of water from the crystallized carbonate ($\text{Na}_2\text{CO}_3 \cdot 10\text{H}_2\text{O}$). (It would be inconveniently moist, even semi-fluid, if a part of the carbonate had not previously been rendered anhydrous.) On the large scale the resulting bicarbonate may be freed from any carbonate or traces of other salts by adding half its bulk of cold distilled water, setting aside for about half an hour, shaking occasionally, draining the undissolved portion, and drying it by exposure on filtering paper.

This arrangement of apparatus for *Sodii Bicarbonas*, U. S. P., may be adopted for *Potassii Bicarbonas*, U. S. P., 1 part of carbonate dissolved in $2\frac{1}{2}$ parts of water being subjected to the action of the gas, and not the solid carbonate, as in the case of the sodium salt.

The pure may be prepared from impure bicarbonate by washing out any carbonate or traces of other salts, after introducing it into a percolator and passing water through it till the washings cease to precipitate a solution of magnesium sulphate, when the sodium bicarbonate is removed from the percolator and dried on bibulous paper in a warm place.

The sodium carbonate may be placed, not in a test-tube or bottle, but in a vertical tube, the bottom of which is loosely closed by a grooved cork. Any water of crystallization that is set free then runs off (into a basin or cup beneath), and takes with it impurities (chlorides or sulphates, etc.) that may have been present in the original salt.

Sodium Carbonates by "The Ammonia Process."

Sodium bicarbonate is prepared by bringing to the elements of ammonium bicarbonate a strong solution of common salt; the sodium bicarbonate is precipitated.



The resulting ammonium chloride is reconverted into carbonate (p. 93), the latter more fully carbonated, and again used for producing sodium bicarbonate. Sodium carbonate (*Sodii Carbonas*, B. P.) is made by heating the bicarbonate thus obtained, the carbonic acid then liberated serving for the conversion of some ammonium carbonate into ammonium bicarbonate.



A crystal of sodium carbonate is sodium carbonate plus water; on heating it more or less of the water is evolved, and *anhydrous* sodium carbonate is partially or wholly produced (*Sodii Carbonas Exsiccatus*, U. S. P.).



According to the United States Pharmacopœia, dried sodium carbonate is to be prepared by exposing crystals of sodium carbonate to warm air for several days to effloresce, and then to a temperature of about 45° C. until half the original weight is obtained. 286 parts would thus become 143, and the latter would thus still retain 37 parts of water. In other words, the dried carbonate contains 72.6 per cent. of anhydrous carbonate and 27.4 per cent. of water. The crystals contain, obviously, a little more than 37 per cent. of anhydrous carbonate and nearly 63 per cent. of water. *The student should verify all these figures.*

Note on Nomenclature.—*Anhydrous* bodies (from *a* and *ὕδωρ*, *hudōr*, i. e. without water) are compounds from which water has been taken, but whose essential chemical properties are unaltered. Salts containing water are *hydrous* bodies; of these the larger portion are crystalline, and their water is then termed *water of crystallization*. Non-crystalline *hydrous* compounds were formerly spoken of as *hydrated* substances: hydrates are, however, a distinct class of bodies, salts derived from water by one-half of its hydrogen becoming displaced by an equivalent quantity of another radical. *Anhydrides* form still another distinct class of chemical substances; they are derived from acids: in short, they are acids from which not exactly water as water, but the elements of water, have been removed, the essential chemical (acid) properties being thereby greatly altered. (For illustrations, see Index, "Anhydrides.")

Water of Crystallization.—The water in crystallized sodium carbonate is in the solid condition, and, like ice and other fusible substances, requires heat for its liquefaction. Many salts (freezing mixtures) when dissolved in water give a very cold solution. This is because they and their solid water, if they have any, are then, absorbing some heat from surrounding media, converted into liquids. Take away from water some of its heat, the result is ice. Give to

ice (at 32° F.) more heat than it contains already, the result is water (still at 32° F.). (Heat thus taken into a substance without increasing its temperature is said to become *latent*—from *latens*, hiding; it is no longer discoverable by the sense of touch or the thermometer. The term *latent* gives a somewhat incorrect idea, however, of the conditions; for our knowledge of the extent and readiness with which one form of force is convertible into another renders highly probable the assumption that heat is in these cases converted into motion, the latter enabling the molecules of a solid to take up the new positions demanded by their liquid condition.) The only apparent difference between ice and the water in such crystals as sodium carbonate is that ice is solid water in the free, and water of crystallization solid water in the combined state. The former can only exist at and below 32° F.; the latter may exist at ordinary temperatures. Salts which unite with little or even no water of crystallization at common temperatures, but take up much at very low temperatures, are termed *cryohydrates* (*κρυος*, *kruos*, icy cold, frost). All water of crystallization is dispelled at high temperatures. In chemical formulæ the symbols representing water are usually separated by a comma from those representing salts. The crystals of sodium acetate (of the third reaction) contain water in this loose state of combination—water of crystallization ($\text{NaC}_2\text{H}_3\text{O}_2, 3\text{H}_2\text{O}$). It is possible, however, that this so-called water of crystallization is in a more intimate state of combination than is indicated by such a formula as that just given.

“*Soda-water*.”—A solution of sodium bicarbonate in water charged with carbonic acid gas under pressure constitutes the official *Liquor Sodæ Effervescens*, B. P., and like the official “potash-water” is a true *medicine*, an antacid. The ordinary *beverage* “soda-water” is commonly a simple solution of carbonic acid gas in water, and would be more appropriately termed “aërated water:” any medicinal effect it may possess is due to the sedative influence of its carbonic acid gas on the coats of the stomach. Originally it contained a little “soda” and was a mild antacid, but the public took to it as a mere beverage, refusing to drink it if it contained any “soda,” yet persisted, and still persist, in calling it “soda-water.” At common temperatures water dissolves about its own volume of carbonic acid gas, both being under the same pressure. 1 pint of the official soda-water contains 30 grains of sodium bicarbonate and a pint of carbonic acid gas; but the solution is under a pressure of four atmospheres, in addition to the ordinary pressure of our atmosphere, five atmospheres altogether, so that 5 pints of the gas at ordinary atmospheric pressure are required for the quantity mentioned.

Solubility of Gases in Water.—Whatever the weight and volume of a gas dissolved by a liquid at ordinary atmospheric pressure, that weight is doubled by double pressure, the two original volumes of gas thereby being reduced to one; trebled at treble pressure, the three original volumes of gas being reduced to one; quadrupled at quadruple pressure, the four original volumes of gas being reduced to one; and so on. This is a general law (Henry and Dalton) regarding the solubility of gases in liquids under given temperatures.

An average bottle of "soda-water" contains about five times the weight of carbonic acid gas which can exist in it without artificial pressure, so that on removing its cork four times its bulk escapes, its own bulk remaining dissolved.


Note.—Sodium bicarbonate may also be medicinally administered in the form of lozenge (*Trochisci Sodii Bicarbonatis*, B. P.).

Sodium and potassium carbonate is officially distinguished from the bicarbonate by the former giving a reddish color with phenolphthalein, and the latter no color at all.

Potassium and Sodium Tartrate.

Synonyms.—Tartrate of Potassium and Sodium; Tartrate of Potash and Soda; Rochelle Salt.

Fifth Synthetical Reaction.—To some hot strong solution of sodium carbonate (about 3 parts), in a test-tube or larger vessel, add acid potassium tartrate (about 4 parts) till no more effervescence occurs; when the solution is cold crystals of the potassium and sodium tartrate (*Soda Tartarata*, B. P., *Potassii et Sodii Tartras*, U. S. P.) will be deposited ($\text{KNaC}_4\text{H}_4\text{O}_6, 4\text{H}_2\text{O}$). The crystals are usually halves of right rhombic prisms.



Very close analogy will be noticed in the above formulæ, indicative of the analogy in the constitution of the molecules of these salts. When the other tartrates come under notice, it will be found that they also have a similar constitution.

Sodium Hypochlorite.

Synonyms.—Hypochlorite of Sodium; Hypochlorite of Soda; Chlorinated Soda.

Sixth Synthetical Reaction.—Triturate in a mortar 75 grms. of chlorinated lime with 200 cc. of water; make up to 400 cc., and filter. Wash with 100 cc. of water. Dissolve 150 grms. of sodium carbonate in 300 cc. of hot water, and add to the clear solution of chlorinated lime; thoroughly mix and filter and wash with enough water to make the product weigh 1000 grms. and *Liquor Sodæ Chloratæ*, U. S. P. ("Labarraque's solution") results. It should have a sp. gr. of 1.052.



This solution is an old and very useful disinfectant, formerly known as *Labarraque's Liquor* and *Eau de Javelle*. It contains at least 2.6 per cent. of available chlorine.

Sodium Iodide and Bromide.

Synonyms.—Bromide and Iodide of Sodium.

These salts (NaI and NaBr), *Sodii Iodidum*, U. S. P., and *Sodii Bromidum*, U. S. P., are similar to potassium iodide and bromide in constitution, and are prepared with the same manipulations, soda being used in place of potash. The sodium bromide, however, must be crystallized from warm solutions, or rhombic prisms containing water ($\text{NaBr} \cdot 2\text{H}_2\text{O}$) will be deposited.

Other Sodium Compounds.

Synthetical Reactions portraying the chemistry of the remaining official compounds (namely, sodium nitrate, sulphate, hyposulphite, borate, arseniate, valerianate, and ethylate) are deferred until the several acidulous radicals of these salts have been described.

Sodium Phosphate.—The preparation and composition of this salt will be most usefully studied after bone-ash, the source of it and other phosphates, has been described. Bone-ash is calcium phosphate. (See page 113.)

The official *Citro-Tartrate of Sodium* (*Sodii Citro-tartras Effervescens*, B. P.), the old citro-tartrate of soda, is a mixture of sodium bicarbonate (17 parts), citric acid (6), tartaric acid (9), and sugar (5), mixed and heated (to 200° or 220° F.) until the particles aggregate to a granular condition. When required for medicinal use a dose of the mixture is placed in water; escape of carbonic acid gas at once occurs, and an effervescing liquid results. This substance may be regarded as the official representative of the popular "effervescing citrate of magnesia," so called, which will be further noticed in connection with the salts of magnesium (p. 121).

"*Lemon and Kali*" is a pulverulent mixture of sodium bicarbonate, tartaric acid, sugar, and essence of lemon. It was invented and so named by one Charles Gomond Cooke, who long retained a trade monopoly in the article by thus hiding the "soda" under its ancient name of "kali" (see p. 32), heightening the mystery by the prefix "*Lemon and*".

Sodii Phosphas Effervescens, B. P., and *Sodii Sulphas Effervescens*, B. P., are the respective *anhydrous* salts mixed with sodium bicarbonate and tartaric and citric acids.

Soda Powders (*Pulveres Effervescentes*, U. S. P. 1870).—Formed of 30 grains of sodium bicarbonate and 25 of tartaric acid, wrapped separately in papers of different colors. When mixed with water carbonic acid gas escapes, and sodium tartrate ($\text{Na}_2\text{C}_4\text{H}_4\text{O}_6$) results, a little bicarbonate also remaining.

In the manufacture of sodium carbonate, the old carbonate of soda, the source of the sodium is sodium chloride, and of the carbonic radical calcium carbonate in the form of limestone. By "the Le Blanc process" the chloride is converted into sulphate, the sul-

phate then roasted with coal and limestone, and the resulting *black ash* lixiviated (*lixivia*, from *lix*, lye—water impregnated with alkaline salts; hence *lixiviation*, the operation of washing a mixture with a view of dissolving out salts). If relatively small quantities of solvents be employed, the solution by lixiviation will be more or less *fractional*, salts of varying solubility being thus more or less separated from each other. The lye, evaporated to dryness, yields crude sodium carbonate (soda-ash). By “the ammonia process” the sodium carbonate is obtained by heating sodium bicarbonate, the latter by mixing strong solutions of sodium chloride and ammonium bicarbonate. The last-named salt results from the action of carbonic acid gas (liberated on heating the sodium bicarbonate) on ammonium carbonate, and this again from ammonium chloride and limestone. By either process common salt and limestone are the ordinary prime sources respectively of the sodium and the carbonic radical in sodium carbonate. The processes will be further described in connection with Carbonates.

Deliquescence and Efflorescence.—The sodium and potassium carbonates, chemically closely allied, differ physically. Potassium carbonate quickly absorbs moisture from the air and becomes damp, wet, and finally fluid; it is *deliquescent* (*deliquescent*, melting away). Sodium carbonate, on the other hand yields water of crystallization to the air, the crystals becoming white, opaque, and pulverulent; it is *efflorescent* (*efflorescent*, blossoming forth, in allusion to the appearance of the product).

Analogy of Sodium Salts to Potassium Salts.—Other synthetical reactions might be described similar to those given under Potassium, and thus sodium citrate, iodate, bromate, chlorate, manganate and permanganate, and many other salts be formed. But enough has been stated to show how chemically analogous sodium is to potassium. Such analogies will constantly present themselves. In few departments of knowledge are order and method more perceptible; in few is there as much natural law, as much *science*, as in Chemistry.

Substitution of Potassium and Sodium Salts for Each Other.—Sodium salts being cheaper than potassium salts, the former may sometimes be economically substituted. That one is employed rather than the other is often merely a result due to accident or fashion. But it must be borne in mind that in some cases a potassium salt will crystallize more readily than its sodium analogue, or that a sodium salt is stable when the corresponding potassium salt has a tendency to absorb moisture, or one may be more soluble than the other, or the two may have different medicinal effects. For these or similar reasons a potassium salt has come to be used in medicine or trade instead of the corresponding sodium salt, and *vice versa*. Whenever the acidulous portion only is to be utilized, the least expensive salt of the class would nearly always be selected.

(b) Reactions having Analytical Interest.

1. *The chief analytical reaction for sodium is the flame-test.* When brought into contact with a flame in the manner

described under Potassium (page 80), an intensely yellow color is communicated to the flame by any sodium salt. This is highly characteristic—indeed, almost too delicate a test; for if the point of the wire be touched by the fingers, enough salt (which is contained in the moisture of the hand) adheres to the wire to communicate a very distinct sodium reaction. These statements should experimentally be verified, the chloride, sulphate, or any other sodium salt being employed.

2. Sodium salts, like those of potassium, are not volatile. Prove this fact by the means described when treating of the effect of heat on potassium salts (p. 80).

QUESTIONS AND EXERCISES.

How is the official solution of soda prepared? Give a diagram or equation.—Explain the action of sodium or potassium on water. What colors do these elements respectively communicate to flame?—How much sodium bicarbonate can be obtained from 2240 pounds of crystallized carbonate? *Ans.* 1316 lbs., nearly.—Sodium acetate: give formula, process, and equation.—Give a diagram showing the formation of sodium bicarbonate.—Why is a mixture of dried and undried sodium carbonate employed in the preparation of the bicarbonate?—State the difference between anhydrous and crystallized sodium carbonate.—Define the terms *anhydrous*, *hydrous*, *hydrate*, *anhydride*.—What do you understand by *water of crystallization*?—What is the nature of the official "*Liquor Sodæ Effervesces*"?—How many volumes of gas (reckoned as at ordinary atmospheric pressure) are contained in any given volume of the official "soda-water"?—What is the general law regarding the solubility of gases in liquids under pressure?—What is the systematic name of Rochelle salt? and how is the salt prepared?—What is the relation of Rochelle salt to cream of tartar and tartaric acid?—Give the mode of preparation of the official solution of chlorinated soda, expressing the process by a diagram.—How is the effervescing sodium citro-tartrate prepared?—Define *deliquescence*, *efflorescence*, and *lixiviation*.—State the relations of potassium salts to those of sodium.—How are sodium salts distinguished from those of potassium?

AMMONIUM.

Formula, NH_4 . Combining weight, 18.

Memoranda.—The elements nitrogen and hydrogen, in the proportion of 1 atom to 4 (NH_4), are those characteristic of all the compounds about to be studied, just as potassium (K) and sodium (Na) are the characteristic elements of the potassium and sodium compounds. Ammonium is a univalent nucleus, root, or radical, like potassium or sodium, and the ammonium compounds closely resemble those of potassium or sodium. In short, if, for an instant, potassium or sodium be imagined to be compounds, the analogy between these three series of salts is complete. Ammonium is said to have been isolated, by Weyl, as an unstable dark-blue liquid, possessing a metallic lustre.

Source.—The source of nearly all the ammoniacal salts met with

in commerce is the ammonia gas (NH_3) obtained in distilling all kinds of coal in the manufacture of ordinary illuminating gas and of coke; being derived, doubtless, from the nitrogen of the plants from which the coal has been produced. It is also a by-product in distilling paraffin oil from shale, and may be washed out of the furnace gases of iron-works. It is possible, however, to produce ammonia from its elements. Thus, coal-dust, air, and vapor of water, all at a red heat, yield, according to Rickman and Thompson, gaseous ammonia. Salt added to the mixture prevents the further combustion of the formed ammonia, and ammonium chloride sublimes. Nitrogen and hydrogen passed over spongy platinum yield traces of ammonia.

Ammonia.—When this gas (NH_3) comes into contact with the condensed steam (H_2O) in the process of cooling the coal-gas, the resulting “ammoniacal liquor” is believed to contain ammonium *hydrate* (NH_4HO), the analogue of potassium or sodium hydrate, KOH or NaOH . The grounds for this belief are the observed analogy of the well-known ammoniacal salts to those of potassium and sodium, the similarity of action of solutions of potash, soda, and ammonia on salts of most metals, and the asserted existence of crystals of an analogous sulphur salt (NH_4HS).

Ammonium Chloride.—The ammonia of the “ammoniacal liquor” of the gas-works, liberated by heat and the concurrent action of lime on sulphhydrate, carbonate, and other salts present, and passed into hydrochloric acid, yields crude *ammonium chloride* (sal-ammoniac), $\text{NH}_3 + \text{HCl} = \text{NH}_4\text{Cl}$; and from this salt, purified, the others used in pharmacy are directly or indirectly made. Ammonium chloride (*Ammonii Chloridum*, U. S. P.) occurs in inodorous colorless minute crystals, or in translucent fibrous masses, tough and difficult to powder, soluble in water (1 in 10 is the “Solution of Chloride of Ammonium,” U. S. P.), almost insoluble in alcohol.

Ammonium chloride generally contains slight traces of iron oxychloride, tarry matter, and possibly chlorides of compound ammoniums. (*Vide* “Artificial Alkaloids” in Index.)

Ammonium Sulphate ($\text{NH}_4)_2\text{SO}_4$, the old sulphate of ammonia, results when the ammonia from the “ammoniacal liquor” is neutralized by sulphuric acid. It is largely used as a constituent of artificial manure, and when purified by recrystallization is employed in pharmacy for producing double sulphate of ammonium and iron, etc.

Volcanic Ammonia.—A very pure form of ammonia is that met with in volcanic districts and obtained as a by-product in the manufacture of borax; the crude boracic acid as imported contains 5 to 10 per cent. of ammonium salts, chiefly sulphate, and double sulphates of ammonium with magnesium, sodium, and manganese (*Howard*).

REACTIONS HAVING (a) GENERAL, (b) SYNTHETICAL, AND (c) ANALYTICAL INTEREST.

Ammonium Amalgam (?).

(a) *General Reaction*.—To 40 or 50 grains of dry mercury in a dry test-tube add one or two small pieces of sodium (freed


from adhering naphtha by gentle pressure with a piece of filter-paper), and amalgamate by gently warming the tube. To this amalgam, when cold, add some fragments of ammonium chloride and a strong solution of the same salt. The sodium amalgam soon begins to swell and rapidly increase in bulk, probably overflowing the tube. The light spongy mass produced is the so-called ammonium amalgam, and the reaction is usually adduced as evidence of the existence of ammonium. The sodium of the amalgam unites with the chlorine of the ammonium chloride, while the ammonium is supposed to form an amalgam with the mercury. As soon as formed the amalgam gives off hydrogen and ammonia gases; this decomposition is nearly complete after some minutes, and mercury remains.

(b) *Reactions having Synthetical Interest.*


Ammonium Hydrate.

Synonyms.—Hydrate of Ammonium; Ammonia.

First Synthetical Reaction.—Heat a few grains of sal-ammoniac with about an equal weight of calcium hydrate (slaked lime) damped with a little water in a test-tube; ammonia gas is given off, and may be recognized by its well-known odor. It is very soluble in water. Pass a delivery-tube fitted to the charged test-tube, as described for the preparation of oxygen and hydrogen, into a second test-tube, at the bottom of which is a little water; again heat, the end of the delivery-tube being only just beneath the surface of the water (or, possibly, all the water might rush back into the generating-tube, water absorbing ammonia gas with great avidity); solution of ammonia (*Liquor Ammoniaë*, B. P., or *Liquor Ammoniaë Fortior*, B. P.) will thus be formed.



A molecule of ammonia gas is composed of one atom of nitrogen with three atoms of hydrogen; its formula is NH_3 . Two volumes of the gas contain one volume of nitrogen combined with three similar volumes of hydrogen. Its constituents have, therefore, in combining suffered condensation to one-half their normal bulk. Its conversion into ammonium hydrate may thus be shown:



Solutions of Ammonia, prepared by this process on a large scale and in suitable apparatus, are met with in pharmacy—the one (sp. gr. 0.891) containing 32.5 per cent., the other (sp. gr. 0.959) 10 per cent. by weight of ammonia gas, NH_3 , or 66.9 and 20.6 of ammonia,


NH_4HO (*Aqua Ammoniac Fortior* and *Aqua Ammoniac*, U. S. P. : 1 part, by measure, of the former and 2 of water form the latter). On the large scale, bottles are so arranged in a series as to condense all the ammonia evolved during the operation. *Spiritus Ammoniac*, U. S. P., is an alcoholic solution of ammonia containing 10 per cent., by weight, of the gas (NH_3).

Ammonium Acetate.

Synonyms.—Acetate of Ammonium ; Acetate of Ammonia.

Second Synthetical Reaction.—To acetic acid and water in a test-tube add powdered commercial ammonium carbonate (acid carbonate and carbamate) until effervescence ceases. A solution containing 7 per cent of ammonium acetate forms the official *Liquor Ammonii Acetatis*, U. S. P., the old “spirit of Minde-
rerus.”

On evaporating and cooling, ammonium acetate (the old acetate of ammonia) may be obtained in crystals.



Solution of ammonium acetate can, of course, be just as easily made by reaction of acetic acid and solution of ammonia (NH_4HO) ; but the liquid, owing to absence of dissolved carbonic acid, is too vapid for use in pharmacy.

Ammonium Carbonates.

Synonyms.—Carbonates of Ammonium ; Carbonates of Ammonia.

Commercial Ammonium Carbonate is made by heating a mixture of chalk and sal-ammoniac ; calcium chloride (CaCl_2) is produced, ammonia gas (NH_3) and water (H_2O) escape, and the ammoniacal carbonate distils, or, rather, sublimes* in cakes (*Ammonii Carbonas*, U. S. P., the old carbonate of ammonia). The best form of apparatus to employ is a retort with a short wide neck and a cool receiver. On the large scale the retort is usually iron and the receiver earthenware or glass ; on the small scale glass vessels are employed. The salt is purified by resublimation at a low temperature— 150° F. is said to be sufficient.


The salt, the empirical formula of which is $\text{N}_3\text{H}_{11}\text{C}_2\text{O}_5$, is probably a mixture of one molecule (sometimes two) of acid ammonium carbonate (NH_4HCO_3) and one of a salt termed ammonium carbamate ($\text{NH}_4\text{NH}_2\text{CO}_2$). The latter belongs to an important class of salts known as *carbamates*, but is the only one of direct interest to the

* *Sublimation* (from *sublimis*, high), vaporization of a solid substance by heat, and its condensation on an upper and cooler part of the vessel or apparatus in which the operation is performed. Substances sublime at different temperatures, hence a mixture of volatile solids may sometimes be separated or fractionated by sublimation.

pharmacist. Cold water extracts it from the commercial ammonium carbonate, leaving the acid carbonate or bicarbonate undissolved if the amount of liquid used be very small. Alcohol extracts the carbamate, leaving the acid carbonate undissolved. In water the carbamate soon changes into the neutral ammonium carbonate,



so that an aqueous solution of commercial ammonium carbonate contains both acid and neutral ammonium carbonate. If to such a solution some ordinary solution of ammonia be added, a solution of the *neutral ammonium carbonate* only is obtained: and this is the common reagent always found on the shelves of the analytical laboratory. Thus, "Solution of Carbonate of Ammonium," B. P., is formed by dissolving $\frac{1}{2}$ an ounce of the salt in 10 ounces of water, to which $\frac{3}{4}$ of an ounce of solution of ammonia has been added.



Neutral ammonium carbonate is the salt formed on adding strong solution of ammonia to the commercial carbonate in preparing a pungent mixture for toilet smelling-bottles; but it is unstable, and on continued exposure to air is reduced to a mass of crystals of bicarbonate.


If ammonium carbonate contain more than traces of empyreumatic matters (from the gas-liquors), an aqueous solution, with excess of sulphuric acid added, will decolorize a dilute solution of potassium permanganate at once.

Sal Volatile (*Spiritus Ammoniae Aromaticus*, U. S. P.) is a spirituous solution of about $1\frac{1}{4}$ per cent. of ammonia gas (NH_3), nearly $3\frac{1}{2}$ per cent. of neutral ammonium carbonate, $(\text{NH}_4)_2\text{CO}_3$, and the oils of nutmeg, lemon, and lavender. Commercial samples contain salts equivalent to from 1 to nearly 3 per cent. of ammonia gas, the official spirit yielding a total of $2\frac{1}{2}$ per cent. of the gas. Fetid spirit of ammonia (*Spiritus Ammoniae Fetidus*, B. P.) is an alcoholic solution of the volatile oil of asafœtida mixed with solution of ammonia.

Ammonium Nitrate.

Synonyms.—Nitrate of Ammonium; Nitrate of Ammonia.

Third Synthetical Reaction.—To some diluted nitric acid add ammonium carbonate, until, after well stirring, a slightly ammoniacal odor remains. The solution contains ammonium nitrate (*Ammonii Nitras*, U. S. P., the old nitrate of ammonia).



From a strong hot solution of ammonium nitrate crystals may be obtained containing much water ($\text{NH}_4\text{NO}_3, 12\text{H}_2\text{O}$). On heating these in a dish to about 320°F . the water escapes. The anhydrous salt remaining (NH_4NO_3) may be poured on to an iron plate. On further heating the crushed nitrate at 350° to 450°F ., it is resolved

into *nitrous oxide* gas ("laughing gas") and water, $\text{NH}_4\text{NO}_3 = \text{N}_2\text{O} + 2\text{H}_2\text{O}$.

Nitrous Oxide is thus prepared for use as an anæsthetic. When required for inhalation, it is washed from any trace of acid or nitric oxide by being passed through solution of potash and solution of ferrous sulphate, the former absorbing acid vapors, the latter nitric oxide. It is slightly soluble in warm water, more so in cold. It supports combustion almost as well as oxygen. By pressure it may be liquefied to a colorless fluid, and by simultaneous cooling solidified.

Ammonium Citrate, Phosphate, and Benzoate.

Synonyms.—Citrate, etc. of Ammonium; Citrate, etc. of Ammonia.

Fourth Synthetical Reaction.—To solution of citric acid ($\text{H}_3\text{C}_6\text{H}_5\text{O}_7$) add solution of ammonia (NH_4HO) until the well-stirred liquid smells faintly of ammonia. Solutions having the specific gravities, respectively, of 1.062 and (four times the strength) 1.209 form the official solutions of ammonium citrate, $(\text{NH}_4)_3\text{C}_6\text{H}_5\text{O}_7$ (*Liquor Ammonii Citratis*, B. P., and *Liquor Ammonii Citratis Fortior*, B. P.).

Ammonium Phosphate, $(\text{NH}_4)_2\text{HPO}_4$ (*Ammonii Phosphas*, B. P.), and *Ammonium Benzoate* ($\text{NH}_4\text{C}_7\text{H}_5\text{O}_2$) (*Ammonii Benzoas*, U. S. P.), are also made by adding solution of ammonia to phosphoric acid (H_3PO_4) and benzoic acid ($\text{HC}_7\text{H}_5\text{O}_2$), respectively, evaporating (keeping the ammonia in slight excess by adding more of its solution), and setting aside for crystals to form.



Ammonium phosphate occurs in transparent colorless prisms soluble in water, insoluble in spirit; benzoate in crystalline plates soluble in water and in spirit.


Ammonium Bromide (U. S. P.) will be noticed in connection with Hydrobromic Acid and other bromides, and the *Ammonii Iodidum* (U. S. P.) with the other iodides.

Ammonium Oxalate.

Synonyms.—Oxalate of Ammonium; Oxalate of Ammonia.

Fifth Synthetical Reaction.—To a nearly boiling solution of 1 part of oxalic acid in about 8 of water add ammonium carbonate until the liquid is neutral to *test-paper* (see following paragraphs), filter while hot, and set aside for crystals to form. The product is Ammonium Oxalate, U. S. P. (Oxalate of Ammonium, B. P.), $(\text{NH}_4)_2\text{C}_2\text{O}_4 \cdot \text{H}_2\text{O}$, the old oxalate of ammonia. The mother-

liquor is useful as a reagent in analysis; one of the pure salt in forty of water forms "Solution of Oxalate of Ammonium," B. P.



Neutralization.—Thus far, the methods by which the student has avoided excess of either acid matter on the one hand or alkaline on the other have been the rough aid of taste, cessation of effervescence, presence or absence of odor, etc. More delicate aid is afforded by *test-papers*.

Test-papers.—*Litmus* (B. P.) is a blue vegetable pigment prepared from various species of *Roccella* lichen, exceedingly sensitive to the action of acids, which turn it red. When thus reddened, alkalies (potash, soda, and ammonia) and other soluble hydrates readily turn it blue. The student should here test for himself the delicacy of this action by experiments with paper soaked in solution of litmus and dipped into very dilute solutions of acids, acid salts ($\text{KHC}_4\text{H}_4\text{O}_6$, *e. g.*), alkalies, and such neutral salts as potassium nitrate, sodium sulphate, or ammonium chloride.

Solution of Litmus (U. S. P.).—This is prepared from purified litmus. Gently boil litmus with three times its bulk of spirit of wine for an hour. Pour away the fluid and repeat the operation twice. Digest the residual litmus in distilled water and filter.

Blue Litmus-paper (U. S. P.) is unsized white paper colored by a solution of litmus and dried by exposure to the air.

Red Litmus-paper (U. S. P.) is unsized white paper colored with solution of litmus previously reddened by the smallest requisite quantity of hydrochloric acid, and dried by exposure to the air.

Turmeric-paper (U. S. P.), similarly prepared from tincture of turmeric (one of washed turmeric-root or rhizome to six of rectified spirit, macerated for several days), is occasionally useful as a test for alkalies, which turn its yellow color to brown; acids do not affect it.

Other "indicators" of alkalinity or acidity are used.

Ammonium Sulphydrate.

Synonyms.—Sulphydrate of Ammonium; Sulphydrate of Ammonia.


Sixth Synthetical Reaction.—Pass sulphuretted hydrogen gas (H_2S) through a small quantity of solution of ammonia in a test-tube, until a portion of the liquid no longer causes a white precipitate in solution of magnesium sulphate; the product is a solution of ammonium sulphydrate (NH_4HS), erroneously termed ammonium sulphide, a valuable chemical reagent, as will presently be apparent:



"Ammonium Sulphide Test-solution," U. S. P., of official strength is made by passing the gas, prepared as described on the next page,


into 300 cc. of ammonia solution (*Aqua Ammoniacæ*) so long as the gas continues to be absorbed, then adding 200 cc. more of ammonia solution, and preserving in a well-stoppered bottle.

Sulphuretted Hydrogen (*Hydrogen Sulphide*, *Sulphuric Acid*, or *Hydrosulphuric Acid*) is a compound of noxious odor; hence the above operation, and many others described farther on, in which this gas is indispensable, can only be performed in the open air or in a *fume-cupboard*—a chamber so contrived that deleterious gases and vapors shall escape into a chimney in connection with the external air. In the above experiment the small quantity of gas required can be made in a test-tube, after the manner of hydrogen itself. To some fragments of ferrous sulphide (FeS) add water, and then sulphuric acid; the gas is at once evolved, and may be conducted by a tube into the solution of ammonia. Ferrous sulphate remains dissolved:



Sulphuretted-hydrogen Apparatus.—As no heat is necessary in making sulphuretted hydrogen (U. S. P.), the test-tube of the foregoing operation may be advantageously replaced by a bottle, especially when larger quantities of the gas are required. In analytical operations the gas should be purified by passing it through water contained in a second bottle.

Fig. 18.



Sulphuretted-hydrogen Apparatus.


The most convenient arrangement for experimental use is prepared as follows: Two common wide-mouth bottles are selected, the one having a capacity of about half a pint, the other a quarter pint; the former may be called the *generating-bottle*, the latter the *wash-bottle*. Fit two corks to the bottles. Through each cork bore two holes, with a round file or other instrument, of such size that glass tubing of about the diam-

eter of a quill pen shall fit them tightly. Through one of the holes in the cork of the generating-bottle pass a funnel-tube, so that its extremity may nearly reach the bottom of the bottle. To the other hole adapt a piece of tubing six inches long and bent in the middle to a right angle. A similar "elbow-tube" is fitted to one of the holes in the cork of the wash-bottle, and another elbow-tube, one arm of which is long enough to reach to near the bottom of the wash-bottle, fitted to the other hole. Removing the corks, two or three ounces of water are now poured into each bottle, an ounce or two of ferrous sulphide put into the generating-bottle, and the corks replaced. The elbow-tube of the generating-bottle is now attached by a short piece of india-rubber tubing to the long-armed elbow-tube of the wash-bottle, so that gas coming from the generator may pass through the water into the wash-bottle. The delivery-tube of the wash-bottle is then lengthened by attaching to it, by india-rubber tubing, another piece of glass tubing several inches in length. The apparatus is now ready for use. Strong sulphuric acid is poured down the funnel-tube in small quantities at a time, until brisk effervescence is established, and more added from time to time as the evolution of gas becomes slow. The gas passes through the tubes into the wash-bottle, where, as it bubbles up through the water, any trace of sulphuric acid or other matter mechanically carried over is arrested, and thence the gas flows out at the delivery-tube into any vessel or liquid that may be placed there to receive it. The generator must be occasionally dismounted and the ferrous sulphate washed out.

Luting (lutum, mud).—If the corks of the above apparatus are sound and the tube-holes well made, no escape of gas will occur. If rough corks have been employed or the holes are not cylindrical, linseed-meal lute may be rubbed over the defective parts. The lute is prepared by mixing linseed meal with water to the consistence of dough. A neat appearance may be given to the lute by gently rubbing a well-wetted finger over its surface.

(c) *Reactions having Analytical Interest (Tests).*

First Analytical Reaction.—To a solution of an ammonium salt (chloride, for example) in a test-tube add solution of soda (or potash, or slaked lime), and well shake or warm; a characteristic odor (ammonia, NH_3) results:



Though ammonium itself cannot be obtained in the free state, its compounds are stable. Ammonia is easily expelled from those compounds by action of the stronger alkalies, caustic potash, soda,

or lime. As a matter of exercise the student should here draw out equations in which ammonium acetate ($\text{NH}_4\text{C}_2\text{H}_3\text{O}_2$), sulphate ($(\text{NH}_4)_2\text{SO}_4$), nitrate (NH_4NO_3), or any other ammoniacal salt, not already having the odor of ammonia, is supposed to be under examination; also equations representing the use of the other hydrates (KHO or Ca_2HO).

The *odor* of ammonia is the best means of recognizing its presence; but the following tests are occasionally useful: Into the upper part of the test-tube insert a glass rod moistened with strong hydrochloric acid (that is, with the solution of hydrochloric acid gas conventionally termed hydrochloric acid, the *Acidum Hydrochloricum* of the pharmacopœias); white fumes of ammonium chloride will be produced: $\text{NH}_3 + \text{HCl} = \text{NH}_4\text{Cl}$. Hold a piece of moist red litmus-paper in a tube in which is ammonia gas; the red color will be changed to blue.

Second Analytical Reaction.—To a few drops of a solution of an ammonium salt add a drop or two of hydrochloric acid and a like small quantity of solution of platinum perchloride (PtCl_4); a yellow crystalline precipitate (the double platinum and ammonium chloride, $\text{PtCl}_4 \cdot 2\text{NH}_4\text{Cl}$) will be produced, similar in appearance to the corresponding potassium salt, the remarks concerning which (p. 79) are equally applicable to the precipitate under notice.

Third Analytical Reaction.—To a moderately strong solution of an ammonium salt add a strong solution of tartaric acid, and shake or well stir the mixture; a white granular precipitate (acid ammonium tartrate) will be formed.

For data from which to draw out an equation representing this action see the remarks and formulæ under the analogous potassium salt (p. 79).

Fourth Analytical Fact.—Evaporate a few drops of a solution of an ammonium salt to dryness, or place a fragment of a salt in the solid state on a piece of platinum-foil, and heat in a flame; the salt is readily *volatilized*. As already noticed, the salts of potassium and sodium are *fixed* under these circumstances, a point of difference of which advantage will frequently be taken in analysis. A porcelain crucible may often be advantageously substituted for platinum-foil in experiments on volatilization.

Salts of ammonium with the more complex acidulous radicals seldom volatilize unchanged when heated. The oxalate, when warmed, loses its water of crystallization, and at a higher temperature decomposes, yielding carbonic oxide, carbonic acid gas, ammonia gas, water (the three latter sometimes in combination),

and several organic substances. The phosphate yields more or less phosphoric acid as a residue.

A *wire triangle* may be used in supporting crucibles (Fig. 19). It is made by twisting together each pair of ends of three (five- or six-inch) crossed pieces of wire (Fig. 20). A piece of tobacco-pipe stem (about two inches) is sometimes placed in the centre of each wire before twisting, the transference of any metallic matter to the sides of the crucible being thus prevented (Fig. 21).

Fig. 19.




Fig. 20.



Fig. 21.



Triangular Supports for Crucibles.

Practical Analysis.

With regard to those of the preceding experiments which are useful rather as means of detecting the presence of potassium, sodium, and ammonium than as illustrating the preparation of salts, the so-called "tests," the student should proceed to apply them to certain solutions of any of the salts of potassium, sodium, and ammonium, with the view of ascertaining which metal is present; that is, proceed to *practical analysis*.* A little thought will enable him to apply these reactions in the most suitable order and to the best advantage for the contemplated purpose; but the following arrangements are perhaps as good as can be devised:

* Such solutions are prepared in educational laboratories by a tutor. They should, under other circumstances, be mixed by a friend, as it is not desirable for the student to know previously what is contained in the substance he is about to analyze.

The analysis of solutions containing only one salt serves to impress the memory with the characteristic tests for the various metals and other radicals, and familiarize the mind with chemical principles. Medical students seldom have time to go farther than this. More thorough analytical and general chemical knowledge is only acquired by working on such mixtures of bodies as are met with in actual practice, beginning with solutions which may contain any or all of the members of a group. Hence in this Manual two tables of short directions for analyzing are given under each group. Pharmaceutical students should follow the second.

DIRECTIONS FOR APPLYING THE FOREGOING ANALYTICAL REACTIONS TO THE ANALYSIS OF AN AQUEOUS SOLUTION OF A SALT OF **ONE** OF THE METALS, POTASSIUM, SODIUM, AMMONIUM.

To a small portion of the solution to be examined in a test-tube add caustic soda, and warm the mixture; the odor of ammonia gas reveals the presence of an ammonium salt.

If ammonium be not present, apply the platinum perchloride test to another portion of the liquid; a yellow precipitate proves the presence of potassium.

(It will be observed that potassium can only be detected in the absence of ammonium, salts of the latter radical giving similar precipitates.)

The flame test is sufficient for the recognition of sodium.

DIRECTIONS FOR APPLYING THE FOREGOING ANALYTICAL REACTIONS TO THE ANALYSIS OF AN AQUEOUS SOLUTION OF SALTS OF **ONE, TWO, OR ALL THREE** OF THE ALKALI METALS.

Commence by testing a small portion of the solution for an ammonium salt. If it be present, make a memorandum to that effect, and then proceed to get rid of the ammoniacal compound to make way for the detection of potassium; advantage is here taken of the volatility of ammonium salts and the fixity of those of potassium and sodium. Evaporate the original solution to dryness in a small basin, transfer the solid residue to a porcelain crucible, and heat the latter to low redness, or until dense white fumes (of ammoniacal salts) cease to escape. (See Fig. 19.) This operation should be conducted in a fume-cup-board, to avoid contamination of the air of the laboratory. When the crucible is cold, dissolve out the solid residue with a small quantity of hot water, and test the solution for potassium by the platinum perchloride test, and for sodium by the flame test.

When the starting test has shown absence of ammonium, the original solution may, of course, at once be tested for potassium and sodium.

Flame Test.—The violet tint imparted to flame by potassium salts may be seen when masked by the intense yellow color due to sodium if the flame be observed through a piece of dark-blue glass, a medium which absorbs the yellow rays of light.

Note on Nomenclature.—The operations of *evaporation* and heating to redness, or *ignition*, are frequently necessary in analysis, and are usually conducted in the above manner. If vegetable or animal matter be also present, carbon is set free, and ignition is accompanied by *carbonization*; the material is said to *char*. When all carbonaceous matter is burnt off, the crucible being slightly inclined and its cover removed to facilitate combustion, and mineral matter, or *ash*, alone remains, the operation of *incineration* has been effected.

Note on the Classification of the Elements.—The compounds of potassium, sodium, and ammonium have many analogies. Their carbonates, phosphates, and other common salts are soluble in water. The atoms of the three radicals are univalent; that is, each displaces or is displaced by one atom of hydrogen. In fact, these radicals constitute by their similarity in properties a distinct group or family. All the elements thus naturally fall into classes—a fact that should constantly be borne in mind, and evidence of which should always be sought. It would be impossible for the memory to retain the details of Chemistry without a system of classification and leading principles. Classification is also an important feature in the art as well as in the science of Chemistry, for without it practical analysis could not be undertaken. The classification adopted in this volume is founded on the quantivalence of the elements and on their analytical relations.

QUESTIONS AND EXERCISES.

Why are ammoniacal salts classed with those of potassium and sodium?—Mention the sources of the ammonium salts.—Describe the characters of ammonium chloride.—Give the formula of ammonium sulphate.—Adduce evidence of the existence of ammonium.—How is the official solution of ammonia prepared? Give a diagram.—How is the official solution of ammonium acetate prepared?—What is the composition of commercial ammonium carbonate?—Define *sublimation*.—What ammoniacal salts are contained in *Spiritus Ammonizæ Aromaticus*, U. S. P.?—Give diagrams or equations illustrating the formation of ammonium citrate, phosphate, and benzoate.—Give the formula of ammonium oxalate.—How is ammonium hydrate converted into sulphhydrate?—Describe the preparation of sulphuretted hydrogen gas.—Enumerate and explain the tests for ammonium.—How is potassium detected in a solution in which ammonium has been found?—Give equations illustrating the action of sodium hydrate on ammonium acetate; potassium hydrate on ammonium sulphate; and calcium hydrate on ammonium nitrate.—What are the effects of acids and alkalis on litmus and turmeric?—Describe the analysis of an aqueous liquid containing potassium, sodium, and ammonium salts.—What meanings are commonly assigned to the terms *evaporation*, *ignition*, *carbonization*, and *incineration*?—Write a short article descriptive of the analogies of potassium, sodium, and ammonium, and their compounds.

BARIUM, CALCIUM, MAGNESIUM.

These three elements have many analogies. Their atoms are bivalent.

BARIUM.

Symbol, Ba. Atomic weight, 137.

The analytical reactions of this metal are those which are of chief interest to the general student of pharmacy. The barium nitrate, the old nitrate and baryta or nitrate of barytes (Ba_2NO_3), and the chloride ($\text{BaCl}_2 \cdot 2\text{H}_2\text{O}$), are the soluble salts in common use in analysis (Chloride of Barium, B. P., and test-solution of barium chloride, 1 in 10 of water, U. S. P.); and these and others are made by dissolving the native carbonate (BaCO_3), the mineral *withelite*, in acids, or by heating the other common natural barium compound, the sulphate, *heavy white* or *heavy spar* (BaSO_4), with coal, which yields barium sulphide (BaS), ($\text{BaSO}_4 + \text{C} = 4\text{CO} + \text{BaS}$) and dissolving the sulphide in appropriate acids. When the nitrate is strongly heated, it is decomposed, barium oxide or *baryta* (BaO) remaining. Baryta, on being moistened, assimilates the elements of water with great avidity, and yields barium hydrate (Ba_2HO). The latter is tolerably soluble, giving *baryta-water*; and from this solution crystals of barium hydrate ($\text{Ba}_2\text{HO} \cdot 8\text{H}_2\text{O}$) are obtained on evaporation.

The operations above described may all be performed in test-tubes and small porcelain crucibles heated by the gas flame. Quantities of 1 oz. to 1 lb. require a coke furnace.

Barium Peroxide.

Synonyms.—Peroxide of Barium; Barium Dioxide.

Barium Peroxide (BaO_2) is formed on passing air over baryta heated to about 600°F . On raising the temperature oxygen is evolved and baryta remains. This is Boussingault's old process, but the baryta loses its absorbing power after a time. If the air be freed from carbonic acid gas, and the peroxide be not exposed to a much higher temperature than 800°F . (by heating in a vacuum), the baryta can be used over and over again. This improvement is by Messrs. Brin, who sell the oxygen compressed within strong metal cylinders. Barium peroxide is now official (*Barii Dioxidum*, U. S. P.). It is a heavy white amorphous powder without odor or taste; exposed to air it gradually absorbs moisture and carbon dioxide, and is thus slowly decomposed. It is almost insoluble in, but forms a definite hydrate with, cold water.

Quantivalence.—The atom of barium is bivalent, Ba'' .

Hydrogen Peroxide, or Peroxide of Hydrogen.

Hydrogen peroxide (H_2O_2) is prepared by the action of dilute acid on barium peroxide. A solution of the oxide is thus obtained, the old *oxygenated water*. *Aqua Hydrogenii Dioxidii*, U. S. P., is officially prepared by suspending barium peroxide in water, and well shaking every few minutes for half an hour, keeping the temperature below

10° C. Then continue to shake occasionally until all the peroxide has been hydrated, which is easily determined by the fact that hydrated oxide of barium does not settle to the bottom on standing. Dilute phosphoric acid (nine of water to two of acid) is then carefully added until the liquid remains just neutral after being well agitated. The precipitated barium phosphate is then allowed to settle, and filtered. Dilute sulphuric acid is then added to the filtrate, until small portions filtered give no further precipitate on the addition of more acid. Starch may be used to assist filtration. The amount of H_2O_2 is finally ascertained by the official method (see Index) and made up to the proper strength.

The solution of hydrogen peroxide made as above should contain about 3 per cent. by weight of the pure peroxide. It is a liquid without color or odor, but having a slightly acid taste and a specific gravity about 1.006–1.012. When exposed to the air the solution loses water, and if heated is liable to decompose suddenly. In the official solution a little free acid is allowed to remain to preserve it; on evaporation should not leave more than 0.5 per cent. of solid matter.

Tests.—If some water be mixed in a test-tube with a drop of potassium chromate, a drop or two of dilute sulphuric acid, and a few cc. of ether run on the top, then one or two drops of a solution of hydrogen peroxide added, on well shaking the solution the ether becomes colored blue.

On neutralizing a solution of hydrogen peroxide with ammonia and adding potassium permanganate, oxygen gas is given off, its volume indicating the oxygen "volume strength" of the original solution.

REACTIONS HAVING ANALYTICAL INTEREST (TESTS).


First Analytical Reaction.—To the aqueous solution of any soluble barium salt (nitrate or chloride, for example) add dilute sulphuric acid; a white precipitate is obtained. Set the test-tube aside for two or three minutes, and when some of the precipitate has fallen to the bottom pour away the supernatant liquid, wash the precipitate by adding water, shaking, setting aside, and again decanting; and then add strong nitric acid, and boil; the precipitate is insoluble.

The production of a white precipitate by sulphuric acid, insoluble even in hot nitric acid, is highly characteristic of barium. The name of this precipitate is barium sulphate; its formula is $BaSO_4$.

Antidotes.—In cases of poisoning by soluble barium salts obvious antidotes would be solution of alum or of any sulphates, such as those of magnesium and sodium (Epsom salt, Glauber's salt).

Second Analytical Reaction.—To a barium solution add solution of the yellow potassium chromate (K_2CrO_4); a pale yellow precipitate (barium chromate, $BaCrO_4$) falls. Add acetic acid to a portion, it is insoluble. Add hydrochloric or nitric acid to another portion, it is soluble.


"*Neutral Chromate.*"—The red potassium chromate (or bichromate) (K_2CrO_4, CrO_3) must not be used in this reaction, or the barium will be only imperfectly precipitated; for the red salt gives rise to the formation of free acid, in which barium chromate is to some extent soluble:



The yellow chromate is obtained on adding potassium bicarbonate, 200 grains, in small quantities at a time, to a hot solution of the red chromate, about 295 grains, until effervescence ceases. The product, diluted to 10 fluidounces, is "*Solution of Yellow Chromate of Potassium,*" B. P.



For analytical purposes solution of a *neutral chromate* is still more readily prepared by simply adding solution of ammonia to solution of red potassium chromate, until the liquid turns yellow, and, after stirring, smells of ammonia.



Other Analytical Reactions.—To a barium solution add a soluble carbonate (ammonium carbonate— $(NH_4)_2CO_3$ —will generally be rather more useful than the others); a white precipitate (barium carbonate, $BaCO_3$) results. To more of the solution add an alkaline phosphate or arseniate (sodium phosphate— Na_2HPO_4 —is the most common of these chemically analogous salts, but ammonium phosphate— $(NH_4)_2HPO_4$ —or arseniate— $(NH_4)_2HAsO_4$ —will subsequently have the preference); white barium phosphate ($BaHPO_4$), insoluble in pure water, but slightly soluble in aqueous solutions of some salts, or barium arseniate ($BaHAsO_4$), both soluble even in acetic and other weak acids, will be precipitated. To another portion add ammonium oxalate $(NH_4)_2C_2O_4$; white barium oxalate (BaC_2O_4) is precipitated, soluble in the diluted mineral acids and sparingly so in acetic acid. Barium salts moistened with hydrochloric acid impart a greenish color to flame.

Mem.—Good practice will be found in writing out equations descriptive of each of the foregoing reactions.

QUESTIONS AND EXERCISES.

What is the quantivalence of barium?—Write down the formulæ of barium oxide, hydrate, chloride, nitrate, and sulphate, and state how they are prepared.—Describe the preparation of hydrogen peroxide.—Which of the tests for barium are most characteristic? Give equations of the reactions.—Name the antidote in cases of poisoning by soluble barium salts, and explain its action.—How is hydrogen peroxide officially made?

CALCIUM.


Symbol, Ca. Atomic weight, 40.

Calcium compounds form a large proportion of the crust of our earth. Calcium carbonate is met with as chalk, marble, limestone, calc-spar, etc.; the sulphate as gypsum and plaster of Paris (*Calcii Sulphas Exsiccatus*, U. S. P., "native sulphate of calcium— $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ —rendered nearly anhydrous by heat") and alabaster; the silicate in many minerals; calcium fluoride as fluor-spar. The phosphate is also a common mineral. The element itself is only isolated with great difficulty. The atom of calcium is bivalent, Ca'' .

REACTIONS HAVING SYNTHETICAL INTEREST.

Calcium Chloride, or Chloride of Calcium.

First Synthetical Reaction.—To some hydrochloric acid add calcium carbonate (chalk, or the purer form, white marble, *Marmor Album*, B. P.) (CaCO_3) until effervescence ceases; filter; solution of calcium chloride (CaCl_2), the most common soluble salt of calcium, is formed.



This solution contains carbonic acid, and will give a precipitate of calcium carbonate on the addition of lime-water. It may be obtained quite neutral by well boiling before filtering off the excess of marble. It is a serviceable test-liquid in analytical operations.

Solution of calcium chloride evaporated to a syrupy consistence yields crystals ($\text{CaCl}_2 \cdot 6\text{H}_2\text{O}$). These are extremely deliquescent. The solution, evaporated to dryness, and the white residue heated to about 200°C ., gives solid calcium chloride ($\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$) in a porous form. The resulting agglutinated lumps (*Calcii Chloridum*, U. S. P.) are used for drying gases and for freeing certain liquids from water. The salt is also soluble in alcohol. 1 part of this dried chloride with 10 of water forms the "Calcium Chloride Test-solution," U. S. P.


Mem.—The practical student has already met with solution of calcium chloride as a by-product or secondary product in the preparation of carbonic acid gas.

Marble often contains ferrous carbonate (FeCO_3), which in the above process becomes converted into ferrous chloride, rendering the calcium chloride impure:



If absolutely pure calcium chloride be required, a few drops of the solution should be poured into a test-tube or test-glass, diluted with water, and examined for iron (by adding ammo-

mium sulphhydrate, which gives a black precipitate with salts of iron), and, if the latter is present, calcium hypochlorite (in the form of chlorinated lime) and slaked lime be added to the remaining bulk of the liquid, and the whole boiled for a few minutes. Iron (as ferric hydrate) is thus precipitated; on filtering a pure solution of calcium chloride is obtained:




This is the official process, and may be imitated on the small scale after adding a minute piece of iron to a fragment of the marble before dissolving in acid.

The names, formulæ, and reactions of the compounds of iron will be considered later.

Calcium Oxide.

Synonyms.—Oxide of Calcium; Quick Lime; Caustic Lime; Lime.

Second Synthetical Reaction.—Place a small piece of chalk in a strong grate fire or furnace, and heat until a trial fragment, chipped off and cooled, no longer effervesces on the addition of acid; caustic lime, CaO (*Calx*, U. S. P.), remains.



Note.—Etymologically considered, this action is analytical (*ἀναλύνω*, *analuō*, I resolve), and not synthetical (*σύνθεσις*, *synthesis*, a putting together); but conventionally it is synthetical, and not analytical; for in this, the usual sense, synthesis is the application of chemical action with the view of producing something; analysis, the application of chemical action with the view of finding out the composition of a substance. In the etymological view of the matter there is scarcely an operation performed, either by the analyst or by the manufacturer, but includes both analysis and synthesis; that is, includes interchange, or *metathesis*.

Lime-kilns.—On a large scale the above operation is carried on in what are termed *lime-kilns* (*kiln*, Saxon *cyln*, from *cylene*, a furnace).

Calcium Hydrate.

Synonyms.—Hydrate of Calcium; Slaked Lime; Hydrate of Lime.

Slaked Lime.—When cold, add to the lime about half its weight of water, and notice the evolution of steam and other evidence of strong action; the product is *slaked* lime, hydrate of calcium (Ca_2HO) (*Calcii Hydras*, B. P.), the old hydrate of lime, with whatever slight natural impurities the lime may

contain. The slaking of hard or "stony" lime may be accelerated by using hot water.



Lime-water.—Place the calcium hydrate (washed with a little water to remove traces of soluble salts) in about a hundred times its weight of water; in a short time a saturated solution, known as *lime-water* (*Liquor Calcis*, U. S. P.), results. It contains about 0.15 per cent. of slaked lime, or about 16 grains of calcium hydrate (Ca_2HO), equivalent to about 11 or 12 grains of lime (CaO) in 1 (Imperial) pint at 60°F .; at higher temperatures less is dissolved. Sp. gr. 1.0015.


Strong Solution of Lime.—Slaked lime is more soluble in aqueous solution of glycerin and much more soluble in aqueous solution of sugar than in pure water. The *Syrupus Calcis*, U. S. P., is such a solution, containing 6.5 parts of lime and 40 of sugar in 100 parts, by weight, of fluid. It is a more efficient precipitant of hydrates, carbonates, and phosphates than lime-water. The official (B. P.) process is as follows: Mix 1 ounce of lime and 2 of sugar by trituration in a mortar. Transfer the mixture to a bottle containing 1 pint of water, and, having closed this with a cork, shake it occasionally for a few hours. Finally, separate the clear solution with a siphon, avoiding unnecessary exposure to air, and keep it in a well-stoppered bottle.

Solutions of calcium hydrate absorb carbonic acid gas on exposure to air, a semi-crystalline precipitate of carbonate being deposited. When the saccharated solution is heated there is precipitated a compound consisting of three molecules of lime with one of sugar. When it is freely exposed to air, oxygen is absorbed and the solution becomes colored.

Calcium Carbonate.


Synonyms.—Carbonate of Calcium; Carbonate of Lime.

Third Synthetical Reaction.—To a solution of calcium chloride add excess of sodium carbonate, or about 5 parts of dry chloride to 13 of carbonate; a white precipitate of calcium carbonate (*Calcii Carbonas Præcipitatus*, U. S. P., the old precipitated carbonate of lime) (CaCO_3) results. If the solutions of the salts be made hot before admixture and the whole be set aside for a short time, the particles aggregate to a greater extent than when cold water is used, and the product is finally granular or slightly crystalline. The official variety is thus prepared:



Collect and purify this *precipitated chalk* by pouring the mixture into a paper cone supported by a funnel, and, when the liquid has passed through the filter, pour water over the precipitate three or four times, until the whole of the sodium chloride is washed away. This operation is termed *washing a precipitate*. When dried by aid of a *water-bath* (p. 112) or other means the precipitate is fit for use.

Fig. 22.



Construction of Paper Filters.

Filtering Paper, or *Bibulous Paper* (from *bibo*, I drink), is simply good unsized paper made from the best white rags—white blotting-paper, in fact, of unusually good quality. Students' or analysts' filters, on which to collect precipitates, are *circular* pieces (*a*, Fig. 22) of this paper, from three to six inches in diameter, twice folded (*b*, *c*), and then opened out so as to form a hollow cone (*d*). Square pieces of filter-paper are rounded by scissors *after* twice folding. The cone is supported by a glass or earthenware funnel.

Filters should *always* be cut round, so as to form a cone. If the square piece of paper is folded and used without being so cut or trimmed, an ugly angular filter results, from which it is difficult to wash all "mother-liquor" (the solution of sodium chloride is the "mother-liquor" in the previous reaction). Moreover, if a spirituous or other volatile fluid is being passed through such an angular filter, much of the liquor will also be wasted by evaporation from the unnecessarily large surface exposed.

Paper filters of large size are apt to break at the point of the cone. This may be prevented and the rate of filtration be much accelerated by supporting the paper cone in a cone of muslin.

Washing-bottle.—Precipitates are best washed by a fine jet of water directed on to the different parts of the filter. A common narrow-necked bottle, of about half-pint capacity (Fig. 23), is fitted with a cork; two holes are bored through the cork, the one for a glass tube reaching to the bottom of the bottle within, and externally bent to a slightly acute angle, the other for a tube bent to a slightly obtuse angle, the inner arm terminating just within the bottle. The outer arms may be about three inches in length. The extremity of the outer arm continuous with the longer tube should be previously drawn out to a fine capillary opening by holding the original tube (before bending) in a flame, and, when soft, slowly pulling the halves away from each other until the heated portion is reduced to the thinness of a knitting-needle. The tube is now cut at the thin part by a file, and the sharp edges rounded off by placing

in a flame for a second or two. The outer extremity of the shorter tube should also be made smooth in the flame. The apparatus being

Fig. 23.



Washing-bottle.

Fig. 24.




Washing-flask.

put together and the bottle nearly filled with water, air blown through the short tube by the lungs forces water out in a fine stream at the capillary orifice.

For a hot-water washing-flask (Fig. 24) the tubes and cork are fitted to a flask which may be heated. A strip of thin leather tied round the neck will protect the fingers.

Decantation.—Precipitates may also be washed by allowing them to settle, pouring off the supernatant liquid (Fig. 25), agitating

Fig. 25.




Decantation.

Fig. 26.



Decantation.

Fig. 27.



Siphon in Action.

with water, again allowing to settle, and so on. This is washing by *decantation* (*de*, from; *canthus*, an edge). If a stream of liquid flowing from a basin or other vessel exhibits any tendency to run down the outer side of the vessel, it should be guided by a glass rod placed against the point whence the stream emerges (Fig. 26).

If the vessel be too large to handle with convenience, the wash-water may be drawn off by a *siphon*, as shown in miniature in Fig. 27. A siphon is a tube of glass, metal, gutta-percha, or india-rubber bent into the form of a V or U, filled with water and inverted.

One end is immersed in the wash-water, and the other allowed to hang over the side of the vessel. So long as the outer orifice of the instrument is below the level of any liquid in the vessel, so long will that liquid flow from within outward.*


Prepared calcium carbonate (*Creta Præparata*, U. S. P.) is merely washed chalk (*Creta*, B. P.) or *whiting*, only that in pharmacy fashion demands that the chalk be in little conical lumps, about the size of thimbles, instead of in the larger rolls characteristic of "whiting." Wet whiting pushed, portion by portion, through a funnel, and each separately dried, gives the conventional *Creta Præparata*. Its powder is amorphous.

If either the precipitated or the prepared calcium carbonate contains alumina, magnesian salts, iron oxides, or phosphates, its solution in acid, evaporated and redissolved in water, will yield a precipitate of hydrates or phosphates on the addition of saccharated solution of lime.


Calcium Phosphate.

Synonyms.—Phosphate of Calcium; Phosphate of Lime; Bone Phosphate.

Fourth Synthetical Reaction.—Digest bone-ash (bones burnt in an open crucible with free access of air until all animal and carbonaceous matter has been removed—impure calcium phosphate—*Os Ustum*, B. P.) with twice its weight of hydrochloric acid (diluted with four times its bulk of water) in a test-tube or larger vessel; the phosphate is dissolved.



Dilute with water, boil, filter, and, when cold, add excess of solution of ammonia; the calcium phosphate, now practically pure (*Calcii Phosphas Præcipitatus*, U. S. P.), the old phosphate of lime, is reprecipitated as a light white amorphous powder. After well washing the precipitate should be dried over a *water-bath* (see next page) or at a temperature not exceeding 212° F., to prevent undue aggregation of the particles.



* *The nature of the action of a siphon* is simple. The column of water in the outer limb is longer, and therefore heavier, than the column of similar area in the inner limb. (The length of the inner limb must be reckoned from the surface of the liquid, the portion below the surface playing no part in the operation.) Being heavier, it naturally falls by gravitation, the liquid in the shorter limb instantly following because pressed upward by the air. The air, be it observed, exerts a similar amount of pressure on the liquid in the outer limb; in short, atmospheric pressure causes the retention of liquid in the instrument, while gravitation determines the direction of the flow.

Bone-ash or bone-earth contains small quantities of calcium carbonate and sulphide. These are decomposed in the above process by the acid, calcium chloride being formed; on boiling the mixture, carbonic acid gas and sulphuretted hydrogen gas are evolved. Any carbonaceous or siliceous matter, etc. is removed by filtration. In bones the calcium phosphate is always accompanied by a small quantity of an allied substance, magnesium phosphate: a trace of calcium fluoride (CaF_2) is also present.

A *water-bath* for the evaporation of liquids or for drying moist solids at temperatures below 212° F. is an iron, tin, or earthenware pan, the mouth of which can be narrowed by iron or tin diaphragms of various sizes, and having orifices adapted to the diameters of basins or plates. (See Fig. 16, p. 71.) In the British Pharmacopœia, "when a *water-bath* is directed to be used it is to be understood that this term refers to an apparatus by means of which water or its vapor, at a temperature not exceeding 212° , is applied to the outer surface of a vessel containing the substance to be heated, which substance may thus be subjected to a heat near to, but necessarily below, that of 212° . In the *steam-bath* the vapor of water at a temperature above 212° , but not exceeding 230° , is similarly applied." Evaporation *in vacuo* is performed by simply placing the vessel of liquid over or by the side of a small reservoir of strong sulphuric acid or other absorbent of moisture, on the plate of an air-pump, covering with a capacious glass hood or "receiver," and exhausting.

Bone-black, or *Animal Charcoal* (*Carbo Animalis*, U. S. P.), is the residue obtained on subjecting dried bones to a red heat without access of air. It is a mixture of about 9 parts of mineral matter with 1 of carbonaceous matter. The operation may be imitated by heating a few fragments of bone in a covered porcelain crucible in a fume-chamber until smoke and vapor cease to be evolved. Purified animal charcoal (*Carbo Animalis Purificatus*, U. S. P.) may be obtained as follows: Boil powdered animal charcoal with a mixture of twice its weight of hydrochloric acid and twice its weight of water; filter; again boil the drained residue with half the amount of such diluted acid as was previously employed; again filter; wash the residual charcoal with distilled water until the washings give little or no turbidity with solution of silver nitrate; dry the product in a warm place. It should not yield more than 10 per cent. of moisture when dried at a high temperature, nor more than 15 per cent. of ash when thoroughly incinerated. Thirty grains well shaken with fifteen ounces of distilled water containing 0.005 per cent. of ordinary commercial caramel

should remove at least four-fifths of the color from the fluid (Hodgkin).


Wood Charcoal (*Carbo Ligni*, U. S. P.) is wood similarly ignited without access of air. On incineration it yields about 2 per cent. of ash.

Decolorizing Power of Animal Charcoal.—Animal charcoal, in small fragments, is the material employed in decolorizing solutions of common brown sugar with the view of producing white lump sugar. Its power, and the nearly equal power of an equivalent quantity of the purified variety, may be demonstrated on solution of litmus or logwood.

Syrup of calcium lacto-phosphate (*Syrupus Calcii Lactophosphatis*, U. S. P.) is a flavored solution of precipitated calcium phosphate in lactic acid.

Sodium Phosphate.—Calcium phosphate is converted into sodium phosphate (*Sodii Phosphas*, U. S. P., the old phosphate of soda) ($\text{Na}_2\text{HPO}_4, 12\text{H}_2\text{O}$), as follows: Mix, in a mortar, 300 grms. of ground bone-earth with 100 cc. of sulphuric acid; set aside for twenty-four hours to promote reaction; mix in about 300 cc. of water, and put in a warm place for two days, a little water being added to make up for that lost by evaporation; stir in another 300 cc. of water, warm the whole for a short time, filter, and wash the residual calcium sulphate on the filter to remove adhering acid calcium phosphate; concentrate the *filtrate* (the liquid portion), which is a solution of acid calcium phosphate, to about 300 cc.; filter again if necessary, add solution of (about 450 grms. of crystals of) sodium carbonate to the hot filtrate until a precipitate (a calcium phosphate, CaHPO_4) ceases to form, and the liquid is faintly alkaline; filter, evaporate, and set aside to crystallize.

Sodium phosphate occurs "in transparent, colorless, rhombic prisms, terminated by four converging planes; efflorescent, tasting like common salt." 1 part in 10 of water constitutes "Solution of Phosphate of Sodium," B. P. The following equations show the two decompositions which occur during the operations:




Ordinary sodium phosphate ($\text{Na}_2\text{HPO}_4, 12\text{H}_2\text{O}$) effloresces rapidly in the air until nearly half its water has escaped, when it has a permanent composition represented by the formula

$\text{Na}_2\text{HPO}_4, 7\text{H}_2\text{O}$. Sodium phosphate has an alkaline reaction. Neutralization by acid results in the removal of half its sodium and formation of the salt $\text{NaH}_2\text{PO}_4, \text{H}_2\text{O}$.


Calcium Hypochlorite, or Hypochlorite of Calcium.

Fifth Synthetical Reaction.—Pass chlorine, generated as already described, into damped slaked lime contained in a piece of wide tubing, open at the opposite end to that in which the delivery-tube is fixed. (A test-tube, the bottom of which has been accidentally broken, is very convenient for such operations.) The product is ordinary *bleaching-powder*, a compound of calcium hypochlorite and chloride, commonly called *chloride of lime* or *chlorinated lime*, the *Calx Chlorata* of the United States Pharmacopœia.



Chlorinated lime exposed to air and moisture, as in disinfecting the atmosphere of sick-rooms, slowly yields hypochlorous acid (HClO). Free hypochlorous acid soon breaks up into water, chloric acid (HClO_3), and free chlorine. Chloric acid is also unstable, decomposing into oxygen, chlorine, water, and perchloric acid (HClO_4). The small quantity of hypochlorous acid diffused through an apartment when bleaching-powder is exposed thus yields fourteen-fifteenths of its chlorine in the form of chlorine gas—one of the most efficient of known disinfectants.

Constitution of Bleaching-powder.—Treated with alcohol, bleaching-powder does not yield its calcium chloride to the solvent; hence the powder is not a mere mixture of calcium chloride and hypochlorite; water, also, does not dissolve out first one salt and then the other, but both together, in the molecular proportions of the above formula. On the other hand, when the aqueous solution is cooled, or evaporated *in vacuo*, crystals are obtained which Kingzett has shown to be nearly pure calcium hypochlorite, the solution containing calcium chloride. While the former fact indicates that the powder is a compound, and not a mere mixture, the latter indicates that it is a feeble compound—an adhesion of molecules of hypochlorite and chloride, as shown in the equation, rather than any more intimate or closer combinations of atoms. If it be regarded as a single rather than a double salt, then the following formula may be employed:




Bleaching-liquor.—Digest chlorinated lime in water, in which the bleaching compound is soluble, filter from undissolved lime, etc., and test the bleaching powers of the clear liquid by adding a few drops to a decoction of logwood slightly acidulated. 1 pound of this bleaching-powder, shaken several times during three hours with 1 gallon of water, forms the official solution of chlorinated lime (*Liquor Calcis Chlorinatæ*, B. P.).

Sixth Synthetical Reaction.—Heat to a bright redness a mixture of 17 parts of powdered dried calcium sulphate and 9 of charcoal in a crucible having a luted cover until the contents are no longer black. Some of the sulphate is reduced to calcium sulphide, with production of carbon oxides. The product, when cold, rubbed to powder, constitutes sulphurated lime (*Calx Sulphurata*, U. S. P.). It should contain not less than 60 per cent. of pure calcium sulphide (CaS); the remainder is sulphate, with, probably, calcium sulphite and hyposulphite.

Official Tests of Strength.—"If 1 grm. of sulphurated lime be gradually added to a boiling solution of 2.08 grms. of cupric sulphate in 50 cc. of water, the mixture digested on a water-bath for fifteen minutes, and filtered when cold, no color should be imparted to the filtrate by one drop of potassium ferrocyanide."—U. S. P.

The B. P. test is similar—eight grains of the substance and fourteen of copper sulphate with a little acid, and warmed as above.

The explanation of the mode of action of the B. P. test is as follows: Copper sulphate and calcium sulphide, in the presence of the acid, react on each other, giving insoluble copper sulphide and calcium sulphate, thus:



On adding up the atomic weights of the constituent elements of crystallized copper sulphate, 249.5 will be found to be the molecular weight, while CaS will similarly represent 72 parts. As 249.5 are to 72, so are 14 to 4. But only half of the sulphurated lime is calcium sulphide; 8 grains of such sulphurated lime will react with 14 of copper sulphate. If the 8 grains are below the stated strength, then they will not attack 14 grains of copper salt, and in that case potassium ferrocyanide will reveal copper in the filtered liquid.

Calcium Sulphate.

Synonyms.—Plaster of Paris; Gypsum (CaSO_4); Sulphate of Lime.

It is found native, and the dried substance is now official, *Calcei Sulphas Exsiccatus*, U. S. P. It should contain about 95 per cent.

of anhydrous calcium sulphate, and then it occurs in a fine white amorphous powder, void of odor and taste, very slightly soluble in water, soluble in dilute nitric and hydrochloric acids, and in saturated solutions of potassium nitrate, sodium hyposulphite, and other ammonium salts; insoluble in alcohol. Calcium sulphate, when mixed with half its weight of water and made into a paste, rapidly hardens.

Solution of Calcium Sulphate.—A $\frac{1}{4}$ ounce of that (dried) form of sulphate of calcium known as plaster of Paris (CaSO_4) digested in 1 pint of water for a short time, with occasional shaking, and the mixture filtered, yields the official test-liquid termed “Solution of Sulphate of Calcium,” B. P. About 400 parts of the solution contain 1 of calcium sulphate.

Calcium Gummate.

Calcium Gummate is the only official calcium salt that remains to be noticed. This compound is, in short, *arabin*, the ordinary gum-acacia or gum-arabic (*Acaciæ*, U. S. P.), a substance too well known to need description. A solution of gum-arabic in water (*Mucilago Acaciæ*, U. S. P.) yields a white precipitate of calcium oxalate on the addition of solution of ammonium oxalate. Or a piece of gum burnt to an ash in a porcelain crucible yields a calcareous residue, which, dissolved in dilute acids, affords characteristic reactions with any of the following analytical reagents for calcium. In some specimens of gum-arabic a portion of the calcium is displaced by an equivalent quantity of potassium or magnesium. The gummie or arabic radical may be precipitated as opaque gelatinous lead gummate by the addition of solution of lead oxyacetate (*Liquor Plumbi Subacetatis*, U. S. P.) to an aqueous solution of gum. These statements should be experimentally verified.

Tragacanth (*Tragacantha*, U. S. P.) is a mixture of soluble arabinoid gum and a variety of calcium gum insoluble in water, termed *bassorin*. With water a gelatinous mucilage is formed (*Mucilago Tragacanthæ*, U. S. P.) containing 6 parts of tragacanth, 18 of glycerin, and 76 of water.

REACTIONS HAVING ANALYTICAL INTEREST (TESTS).

First Analytical Reaction.—Add sulphuric acid, *very* highly diluted, to a calcium solution contained in a test-tube or small test-glass; calcium sulphate ($\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$) is formed, but is *not* precipitated, it being, unlike barium sulphate, slightly soluble in water.

Second Analytical Reaction.—Add yellow potassium chromate (K_2CrO_4) or other neutral chromate (KNH_4CrO_4) to a calcium

solution slightly acidified with acetic acid; calcium chromate (CaCrO_4) may be formed, but is *not* precipitated. Barium is precipitated by the chromate.

These two negative reactions are most valuable in analysis, as every precipitant of calcium is also a precipitant of barium, but the above two reagents are precipitants of barium only. Hence, calcium, which when alone can be readily detected by the following reactions, cannot by any reaction be detected in the presence of barium. But by the sulphuric or chromic test barium is easily removed, and then either of the following reagents will throw down the calcium.

Other Analytical Reactions.—Add ammonium carbonate, sodium phosphate, ammonium arseniate, and ammonium oxalate to calcium solutions, as described under the analytical reactions of barium, and write out descriptive equations. The precipitates correspond in appearance to those of barium; their constitution is also similar, hence their correct formulæ can easily be deduced. Of these precipitants, ammonium oxalate is that most commonly used as a reagent for calcium salts, barium being absent. The calcium oxalate is insoluble in acetic, but soluble in hydrochloric or nitric, acid. Calcium compounds impart a reddish color to flame.

QUESTIONS AND EXERCISES.

Enumerate some of the common natural compounds of calcium.—Explain by an equation the action of hydrochloric acid on marble. What official compound results?—Why is calcium chloride used as a desiccator for gases?—How would you purify calcium chloride which has been made from ferruginous marble? Give diagrams.—Write a few lines on the chemistry of the lime-kiln.—In what sense is the conversion of chalk into lime an analytical action?—What occurs when lime is “slaked”?—To what extent is lime soluble in water? to what in syrup?—Describe the preparation of the official precipitated calcium carbonate; in what does it differ from prepared chalk?—In what does filtering-paper differ from other kinds of paper?—Explain the construction of a “washing-bottle” for cleansing precipitates by water.—Define *decantation*.—Describe the construction and manner of employment of a siphon.—Explain the mode of action of a siphon.—State the difference between *Os Ustum* and *Calci Phosphas*.—How is “bone-earth” purified for use in medicine?—Explain the action of hydrochloric acid on animal charcoal in the processes of purification.—What is the chemical difference between *Carbo Animalis*, U. S. P., and *Carbo Ligni*, U. S. P.?—Give equations showing the conversion of calcium phosphate into sodium phosphate.—Write a short article on the manufacture, composition, and uses of “bleaching-powder.”—How may calcium be detected in gum-arabic?—State the chemical nature of tragacanth.—To what extent is calcium sulphate soluble in water?—Can calcium be detected in a solution containing barium?—Barium being absent, what reagents may be used for the detection of calcium? Which is the chief test?

MAGNESIUM.

Symbol, Mg. Atomic weight, 24.

Source.—Magnesium is abundant in nature as magnesian or mountain limestone, termed *dolomite* (after Dolomieu, a geologist), a double magnesium and calcium carbonate in very common use as a building-stone (*e. g.* the Houses of Parliament and School of Mines in London), and *magnesite*, a tolerably pure magnesium carbonate, though too “stony” for direct use in medicine, even if very finely powdered. Magnesium chloride and magnesium sulphate (Epsom salt) also occur in sea-water and the water of many springs. A monohydrous sulphate ($\text{MgSO}_4 \cdot \text{H}_2\text{O}$), termed *kieserite*, occurs near Stassfurt in Prussia. Metallic magnesium may be obtained from the chloride by the action of sodium. It burns readily in the air, emitting a dazzling light, due to the white heat to which the resulting particles of magnesia (MgO) are exposed. The chloride employed as a source of the metal is obtained by dissolving the carbonate in hydrochloric acid, adding some ammonium chloride, evaporating to dryness, heating the residue in a deep vessel (on the small scale, a large test-tube or flask) until the ammonium chloride is all volatilized, and the magnesium chloride remains as a clear fused liquid. The latter is poured on to a clean earthenware slab. The ammonium chloride prevents reaction between magnesium chloride and water in the last stages of the operation, and consequent formation of magnesium oxide (or oxychloride) and hydrochloric acid gas.


Quantivalence.—The atom of magnesium is bivalent, Mg^{++} .

REACTIONS HAVING SYNTHETICAL INTEREST.**Magnesium Sulphate.**

Synonyms.—Sulphate of Magnesium; Sulphate of Magnesia; Epsom Salt.

First Synthetical Reaction.—To a few drops of sulphuric acid and a little water in a test-tube, made hot (or to larger quantities in larger vessels), add powdered native magnesium carbonate, *magnesite*, MgCO_3 , until effervescence ceases, subsequently boiling to aid in the expulsion of the carbonic acid gas. The filtered liquid is a solution of magnesium sulphate (MgSO_4), crystals of which, *Epsom salt* ($\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$) (*Magnesiæ Sulphas*, U. S. P., the old sulphate of magnesia), may be obtained on evaporating most of the water and setting the concentrated solution aside to cool. This is an ordinary manufacturing process. Instead of *magnesite* *dolomite*, the common magnesian limestone (magnesium and calcium carbonate— $\text{CaCO}_3, \text{MgCO}_3$), may be employed, any iron being removed by evaporating the solution (filtered from the calcium sulphate produced) to dryness, gently igniting to decompose the ferrous sulphate, dissolving in water, filtering from iron oxide, and

crystallizing. (If neither mineral be at hand, the practical student may use a little of the ordinary manufactured carbonate of pharmacy, for the chemical action is almost identical, and it is the chemistry, and not, just now, the commercial economy of the matter, that he is studying. The manufacturer must, of course, commence with one of the above mineral carbonates furnished by Nature, from that make his sulphate, and from the latter, as will be seen directly, make the pure pulverulent carbonate of pharmacy.)



Magnesium sulphate readily crystallizes in large, colorless, transparent, rhombic prisms, but from concentrated solutions the crystals are deposited in short, thin needles, a form more convenient for manipulation, solution, and general use in medicine.

Iron may be detected in magnesium sulphate by adding the common alkaline solution of chlorinated lime or chlorinated soda to some aqueous solution of the salt; brown ferric hydrate ($\text{Fe}_2\text{O}_3 \cdot \text{H}_2\text{O}$) is then precipitated. Ammonium sulphhydrate will also give a black precipitate if iron be present.

Effervescent Sulphate of Magnesium (Magnesii Sulphas Effervescentis, B. P.) is magnesium sulphate out of which nearly half its water of crystallization has been dried, and then mixed with citric and tartaric acids and sugar.

Magnesium Carbonates.

Synonyms.—Carbonates of Magnesium; Carbonates of Magnesia.

Second Synthetical Reaction.—To solution of magnesium sulphate add solution of sodium carbonate, and boil; the resulting precipitate is *light* magnesium carbonate (*Magnesii Carbonas Levis, B. P., Magnesii Carbonas, U. S. P.*), the old light carbonate of magnesia, a white, partly amorphous, partly minutely crystalline mixture of magnesium carbonate and hydrate ($3\text{MgCO}_3, \text{Mg}(\text{OH})_2, 4\text{H}_2\text{O}$). A denser, slightly granular precipitate of similar chemical composition (*Magnesii Carbonas Ponderosa, B. P.*), the old heavy carbonate of magnesia, is obtained on mixing strong solutions of the above salts, evaporating to dryness, then removing the sodium sulphate by digesting the residue in hot water, filtering, washing, and drying the precipitate.



The official (B. P.) proportions for the light carbonate are 10 of magnesium sulphate and 12 of crystals of sodium carbonate, each

dissolved in 80 of cold water, the solutions mixed, boiled for fifteen minutes, the precipitate collected on a filter, well washed, drained, and dried over a water-bath. The heavier carbonate is made with the same proportions of salts, each dissolved in 20 instead of 80 of water, the mixture evaporated quite to dryness, and the residue washed by decantation or filtration until all sodium sulphate is removed (shown by a white precipitate—barium sulphate—ceasing to form on the addition of solution of barium chloride or nitrate to a little of the filtrate).

Another (Pattinson's) Process.—Considerable quantities of magnesium carbonate are now prepared by treating dolomite (p. 118) with carbonic acid gas under pressure. Of the two carbonates, that of magnesium is dissolved first, and is precipitated from the clear liquid by the heat of a current of steam. (See next reaction.)

Third Synthetical Reaction.—Pass carbonic acid gas, generated as described on page 72, into a mixture of water and magnesium carbonate contained in a test-tube. After some time separate any undissolved carbonate by filtration; the filtrate contains *normal* magnesium carbonate ($\text{MgCO}_3 \cdot 3\text{H}_2\text{O}$) dissolved in carbonic acid. When of a strength of about 10 grains of *official* carbonate in 1 ounce, such a solution constitutes "*Fluid Magnesia*" (*Liquor Magnesii Carbonatis*, B. P.). It is possible to obtain a strength of about 3 per cent. at about 55°F. , which is reduced to $2\frac{1}{2}$ per cent. at 70° and to about 2 per cent. at 80° .

Officially, 1 pint is directed (B. P.) to be made from freshly-prepared carbonate. The latter is obtained by adding a hot solution of 2 ounces of magnesium sulphate in $\frac{1}{2}$ pint of water to one of $2\frac{1}{2}$ ounces of crystals of sodium carbonate in another $\frac{1}{2}$ pint of water, boiling the mixture for a short time (to complete decomposition), filtering, thoroughly washing the precipitate, placing the latter in 1 pint of distilled water, and transmitting carbonic acid gas through the liquid (say, at the rate of three or four bubbles per second) for an hour or two, then leaving the solution in contact with the gas under pressure of about three atmospheres for twenty-four hours, and, finally, filtering from undissolved carbonate; then, after passing in a little more gas, keeping in a well-corked bottle. Slight pressure is best produced by placing the carbonate and water in a bottle fitted with a cork and tubes as for a wash-bottle (p. 97), conveying the gas by the tube which reaches to the bottom, and allowing excess of gas to flow out by the upper tube, the external end of which is continued to the bottom of a common phial containing about an inch of mercury. The phial should be loosely plugged with cotton wool, to prevent loss of metal by spurting during the flow of the gas through it. (Each inch in depth of mercury through which the gas escapes corresponds to about $\frac{1}{2}$ pound pressure on every square inch of surface within the apparatus.)

Heat a portion of the solution: true magnesium carbonate con-

taining combined water ($\text{MgCO}_3, 3\text{H}_2\text{O}$) is precipitated. The water in this compound is probably in the state of water of crystallization, for a salt having the same composition is deposited in crystals by the spontaneous evaporation of the solution of magnesium carbonate. The official "carbonate" ($3\text{MgCO}_3, \text{Mg}2\text{HO}, 4\text{H}_2\text{O}$) is another of these very common *hydrous* compounds.


Exposed to cold, the solution of "fluid magnesia" sometimes affords large thick crystals ($\text{MgCO}_3, 5\text{H}_2\text{O}$), which, in contact with the air, lose water, become opaque, and then have the composition of those deposited by evaporation ($\text{MgCO}_3, 3\text{H}_2\text{O}$).

It is difficult to understand the *constitution* of the official carbonate. It is probably a single molecule, of the constitution of which we are at present ignorant.

Magnesium Oxide.

Synonyms.—Oxide of Magnesium ; Magnesia ; Calcined Magnesia.

Fourth Synthetical Reaction.—Heat light dry magnesium carbonate in a porcelain crucible over a lamp (or in a larger earthen crucible in a furnace) till it ceases to effervesce on adding water and acid to a small portion ; the residue is light magnesia (MgO) (*Magnesia Levis*, B. P., *Magnesia*, U. S. P.). The same operation on the heavy carbonate yields heavy magnesia (MgO) (*Magnesia Ponderosa*, U. S. P.). Both are sometimes spoken of as "calcined magnesia." A given weight of the official light magnesia occupies three and a half times the bulk of the same weight of heavy magnesia.



A trace only of magnesia is dissolved by water. Moisten a grain or two of magnesia with water, and place the paste on a piece of red litmus-paper ; the wet spot after a time becomes blue, showing that the magnesia is slightly soluble.

"*Effervescing Citrate of Magnesia*," so called, is generally a mixture of sodium bicarbonate, citric acid, tartaric acid, sugar, either magnesium carbonate or sulphate, or both, and flavoring essences. True magnesium citrate is easily made by combining together calcined magnesia and citric acid ; it is frequently prescribed in France in doses of 2 ounces.

The "Granulated Citrate of Magnesium" (*Magnesi Citras Effervescens*, U. S. P., *Effervescent Magnesium Citrate*) is made as follows: Mix 10 parts of magnesium carbonate intimately with 30 of citric acid, and enough distilled water to make a thick paste ; dry this at a temperature not exceeding 30°C . (86°F .), and reduce it to a fine powder. Then mix it intimately with 8 of sugar (No. 60 powder), 34 of sodium bicarbonate, and 16 of citric acid previously reduced to a very fine powder. Dampen the mass with a sufficient

quantity of alcohol, and rub it through a No. 20 tinned-iron sieve to form a coarse, granular powder. Lastly dry it in a moderately warm place.

The official effervescing "Solution of Magnesium Citrate" (*Liquor Magnesii Citratis*, U. S. P.) is made by dissolving magnesium carbonate in slight excess of solution of citric acid, adding lemon syrup, placing the diluted liquid in an aerated-water bottle, dropping in crystals of potassium bicarbonate, corking at once, "wiring," and shaking till the crystals are dissolved.

The formula of magnesium citrate deposited from solution is $\text{Mg}_3\text{C}_6\text{H}_5\text{O}_{17}\cdot 14\text{H}_2\text{O}$.

REACTIONS HAVING ANALYTICAL INTEREST (TESTS).

First Analytical Reaction.—Add solution of ammonium hydrate or carbonate to a magnesium solution (sulphate, for example), and warm the mixture in a test-tube; the precipitation of part only of the magnesium as hydrate ($\text{Mg}2\text{HO}$) or carbonate (MgCO_3) occurs. Add now to a small portion of the mixture of precipitate and liquid a considerable excess of solution of ammonium chloride; the precipitate is dissolved.

This is an important reaction, especially as regards magnesium carbonate, the presence of ammonium chloride enabling the analyst to throw out from a solution barium and calcium by an alkaline carbonate, magnesium being retained. The cause of this retention is found in the tendency of magnesium to form soluble double salts with potassium, sodium, or ammonium. In analysis the ammonium chloride should be added before the carbonate, as it is easier to prevent precipitation than to redissolve a precipitate once formed.

Second Analytical Reaction.—To some of the solution resulting from the last reaction add solution of sodium or ammonium phosphate; a white granular precipitate (magnesium and ammonium phosphate, MgNH_4PO_4) results.—*Third.*—To another portion add ammonium arseniate; a precipitate similar in appearance falls (magnesium and ammonium arseniate, $\text{MgNH}_4\text{AsO}_4$).

Note.—Barium and calcium are also precipitated by alkaline phosphates and arseniates. The other precipitants of magnesium are also precipitants of barium and calcium. In other words, there is no *direct* test for magnesium. Hence the analyst always removes any barium or calcium by an alkaline carbonate, as above indicated; the sodium phosphate (or ammonium arseniate or phosphate) then becomes a very delicate test of the presence of magnesium. In speaking of magnesium tests, the absence of barium and calcium salts is to be understood.

QUESTIONS AND EXERCISES.

Name the natural sources of the various salts of magnesium.—Give a process for the preparation of Epsom salt.—Draw diagrams illustrative

of the formation of magnesium sulphate from *magnezite* and from *dolomite*. Show by an equation the process for the preparation of the official magnesium carbonate.—What circumstances determine the two different states of aggregation of the official magnesium carbonates (*Magnesii Carbonas Ponderosa*, B. P., and *Magnesii Carbonas Levis*, U. S. P.)?—What are the relations of *Magnesia Ponderosa*, B. P., and *Magnesia Levis*, B. P., to the official magnesium carbonates?—How much denser is the one than the other?—Is magnesia soluble in water?—How is “Fluid Magnesia” prepared?—Mention the effects of heat and cold on “Fluid Magnesia.”—Ascertain how much magnesia (MgO) can be obtained from 100 grains of Epsom salt.—Calculate the amount of official magnesium carbonate which will yield 100 grains of magnesia.—Can magnesium be detected in presence of barium and calcium?—Describe the analysis of an aqueous liquid containing salts of barium, calcium, and magnesium.—How may magnesium be precipitated from solutions containing ammoniacal salts?

Quantivalence, or Valency.

On reviewing the foregoing statements regarding compounds of the three univalent radicals—potassium, sodium, and ammonium—and the three bivalent elements—barium, calcium, and magnesium—the doctrine of quantivalence, or valency, will be more clearly understood and its usefulness be more apparent. Quantivalence, or the value of atoms, is, in short, in chemistry, closely allied to value in commercial barter. A number of articles, differing much in weight, appearance, and general characters, may be of equal money value; and if these be regarded, for convenience, as having a sort of unit of value, others worth double as much might be termed bivalent, three times as much trivalent, and so on. In like manner, chemical radicals, no matter whether elementary, like potassium (K), iodine (I), or sulphur (S), or compound, like those of nitrates (NO_3), sulphates (SO_4), or acetates ($\text{C}_2\text{H}_3\text{O}_2$), have a given chemical value in relation to each other, and are exchangeable for, or will unite with, each other to an extent determined by that value. Two such radicals may be considered to be present in a molecule of most ordinary salts, a basylous and an acidulous radical, one quantitatively balancing the other. The formulæ of the chief of these radicals and their quantivalence are given in the following table. Examples of formulæ of salts containing univalent, bivalent, and trivalent radicals are given in the succeeding table:

QUANTIVALENCE OF COMMON RADICALS.

Univalent Radicals, or <i>Monads</i> .		Bivalent Radicals, or <i>Dyads</i> .		Trivalent Radicals, or <i>Triads</i> .	
Acidulous.	Basylous.	Acidulous.	Basylous.	Acidulous.	Basylous.
H	H	O	Ca	PO_4	As
Cl	K	SO_4	Mg	BO_3	Sb
I	Na	CO_3	Zn	$\text{C}_6\text{H}_5\text{O}_7$	Bi
HO	NH_4	C_2O_4	Cu	AsO_3	{ $\text{Fe}^{\text{iii}}(\text{ic})$ or $\text{Fe}^{\text{vi}}_2(\text{ic})$
NO_3	Ag	$\text{C}_4\text{H}_4\text{O}_6$	Hg(ic)	AsO_4	
$\text{C}_2\text{H}_3\text{O}_2$	Hg(ous)	S	Fe(ous)	$\text{C}_4\text{H}_3\text{O}_5$	

Note 1.—Hydrogen (H) as the basylous part of salts has entirely different functions to hydrogen (H) as the acidulous part. Acidulous hydrogen gives compounds commonly termed *hydrides* (e.g. AsH_3); basylous hydrogen is the basylous radical of acids (e.g. HCl , H_2SO_4). On the other hand, in compound radicals, e.g. $\text{C}_2\text{H}_3\text{O}_2$ or NH_4 , these properties of hydrogen are no longer apparent; the chemical force resident with the atoms of such radicals seems to be mainly exerted in binding those atoms together.

Note 2.—The name, symbol, quantivalence, and atomic weight of all the important elements are given in a table immediately preceding the Index.

Examples of Formulæ of Salts containing Univalent, Bivalent, and Trivalent Radicals.

The reader will find instructive practice in writing twenty or thirty imaginary formulæ of salts by placing in juxtaposition acidulous and basylous radicals, as in the following table of examples. Just as in a pair of scales a 2-lb. weight must be balanced by two 1-lb. weights, or a 4-lb. weight by two 2-lb. weights, or by one 3-lb. and one 1-lb. weight, so a bivalent radical unites with a bivalent radical or with two univalent radicals, a quadrivalent radical with two bivalent radicals, or with one trivalent and one univalent radical, and so on.

(R = any basylous radical. R = any acidulous radical.)

General Formulæ.	Examples.
$R' R'$. . .	KI , NaCl , $\text{NH}_4\text{C}_2\text{H}_3\text{O}_2$, AgNO_3 .
$R'' R'_2$. . .	CaCl_2 , $\text{Zn}_2\text{C}_2\text{H}_3\text{O}_2$, Pb_2NO_3 ($\text{BaNO}_3\text{C}_2\text{H}_3\text{O}_2$).
$R''' R'_3$. . .	Bi_3NO_3 , AsH_3 , SbCl_3 .
$R'_2 R''$. . .	{ K_2CO_3 , Na_2SO_4 , $\text{H}_2\text{C}_4\text{H}_4\text{O}_6$.
$R' R' R''$. . .	
$R'_3 R''$. . .	{ KHCO_3 , NaHSO_4 , $\text{KNaC}_4\text{H}_4\text{O}_6$.
$R'_2 R' R''$. . .	
$R'_3 R'''$. . .	{ $(\text{NH}_4)_3\text{PO}_4$, $\text{K}_3\text{C}_6\text{H}_5\text{O}_7$, H_3AsO_3 .
$R'_2 R' R'''$. . .	
$R'_2 R''$. . .	Na_2HPO_4 , Na_2HAsO_4 .
$R'' R''$. . .	CaCO_3 , MgO , CuSO_4 , HgO , FeSO_4 .
$R''_3 R'''$. . .	Ca_3PO_4 , $\text{Ca}_3\text{C}_6\text{H}_5\text{O}_7$.
$R'' R' R'''$. . .	MgNH_4PO_4 , CuHAsO_3 .
$R''' R' R'$. . .	BiONO_3 .
$R'''_2 R'' R''$. . .	$\text{Bi}_2\text{O}_2\text{CO}_3$.
$R'''_2 R'''$. . .	As_2O_3 , Sb_2O_3 , Fe_2O_3 , Fe_2SO_4 .
$R''' R'''$. . .	$\text{BiC}_6\text{H}_5\text{O}_7$.
$R'''_2 R'_6$. . .	Fe_2Cl_6 , Fe_2NO_3 , $\text{Fe}_2\text{C}_2\text{H}_3\text{O}_2$.

Quadrivalent Radicals or *Tetrads*, Quinivalent Radicals or *Pentads*, and Sexivalent Radicals or *Hexads*, are known.

Cautions.—1. The student will not mistake valency for strength. Bivalent atoms, for example, will vary as much in *power* of chemical attachment as several two-armed boys will vary from each other in *power* of grasp. The greater the quantivalence of an atom, the more compounds it will form; the stability of those compounds is another affair altogether. Indeed, it often happens that the greater the complexity, the less the stability of a molecule. 2. A "radical"

is a single or whole substance in a broad and general sense only, the great majority of radicals themselves admitting of subdivision. A molecule, like a crystal, is only broadly a whole or single thing, open to attack or cleavage from without, and probably then will split in more than one or two directions.

Exercise.—Write an exposition of the doctrine of quantivalence within the limits of a sheet of note-paper.

DIRECTIONS FOR APPLYING THE FOREGOING ANALYTICAL REACTIONS TO THE ANALYSIS OF AN AQUEOUS SOLUTION* OF A SALT OF **ONE** OF THE METALS, BARIUM, CALCIUM, MAGNESIUM.

Add yellow potassium chromate to a portion of the solution to be examined; a precipitate indicates barium.

If no barium is present, add ammonium chloride and carbonate, and boil; a precipitate indicates calcium.

If barium and calcium are proved to be absent, add ammonium chloride, ammonia, and then either sodium phosphate or ammonium arseniate; a white granular precipitate indicates magnesium.

Ammonia is here added to yield the necessary elements to ammonio-magnesian phosphate or ammonio-magnesian arseniate, both of which are highly characteristic precipitates; and ammonium chloride is added to prevent a mere partial precipitation of the magnesium by the ammonia.

DIRECTIONS FOR APPLYING THE FOREGOING ANALYTICAL REACTIONS TO THE ANALYSIS OF AN AQUEOUS SOLUTION OF SALTS OF **ONE, TWO, OR ALL THREE** OF THE METALS, BARIUM, CALCIUM, MAGNESIUM.

Add potassium chromate to the solution; barium, if present, is precipitated. Filter, if necessary, and add to the *filtrate* (that is, the liquid which has run through the filter) ammonium chloride, hydrate, and carbonate, and boil; calcium, if present, is precipitated. Filter, if requisite, and add sodium phosphate; magnesium, if present, is precipitated.

Note.—Red potassium chromate must not be used in these operations, or a portion of the barium will remain in the liquid and be thrown down with, or in the place of, the calcium carbonate. (*Vide* p. 105). The yellow chromate must not contain potassium carbonate,

* In preparing such solutions for analysis, salts should be selected which do not decompose each other. Chlorides will serve in most cases, but nitrates and acetates are still more convenient.

or calcium will be precipitated with, or in the place of, barium. The absence of carbonate is proved by the non-occurrence of effervescence on the addition of hydrochloric acid to a little of the solution of the chromate, previously made hot in a test-tube. If the yellow chromate has been prepared by adding excess of ammonia to solution of red potassium chromate, its addition to the liquid to be analyzed must be preceded by that of solution of ammonium chloride; the precipitation of a portion of the magnesium (by the free ammonia in the yellow chromate) is thus prevented; for solution of ammonium chloride is a good solvent of magnesium hydrate (and carbonate), as already stated on page 122.

TABLE OF SHORT DIRECTIONS FOR APPLYING THE FOREGOING ANALYTICAL REACTIONS TO THE ANALYSIS OF AN AQUEOUS SOLUTION OF SALTS CONTAINING **ANY** OR **ALL** OF THE METALLIC ELEMENTS HITHERTO CONSIDERED.

To the solution add NH_4Cl , NH_4HO , $(\text{NH}_4)_2\text{CO}_3$; boil and filter.

Precipitate Ba Ca.		Filtrate Mg NH_4 Na K.	
Wash, dissolve in $\text{HC}_2\text{H}_3\text{O}_2$, add K_2CrO_4 , and filter.		Add $(\text{NH}_4)_2\text{HPO}_4$, shake, filter.	
Precipitate Ba.*	Filtrate Ca. Test by $(\text{NH}_4)_2\text{C}_2\text{O}_4$.	Precipitate Mg.	Filtrate NH_4 Na K. Evap. to dryness, ignite, dissolve residue in water. Test for K by PtCl_4 . “ “ Na by flame. Test orig. sol. for NH_4 .

Note 1.—The analysis of solutions containing the foregoing metals is commenced by the addition of ammonium chloride (NH_4Cl) and ammonia (NH_4HO), simply as a precautionary measure, the

* It is perhaps scarcely necessary to state that this precipitate is not barium (Ba) itself, but barium chromate (BaCrO_4), as any reader who has carefully gone through the “foregoing analytical reactions” will know. Chemical symbols and formulæ are often used for mere shorthand purposes, but no intelligent student will thereby be misled. The occurrence of barium chromate here, however, and under the circumstances described, is abundant evidence of the presence of barium (Ba, in some form or other) in the liquid analyzed; which was a part of the problem to be solved by the operator. Similar remarks apply to the Ca, which is finally precipitated as oxalate (CaC_2O_4), to Mg, which is thrown out as ammonio-phosphate (MgNH_4PO_4), to NH_4 , Na, and K, and to the elements similarly alluded to in the other subsequent tables of “short” directions for analysis.

former compound preventing partial precipitation of magnesium, the latter neutralizing acids. The ammonium carbonate, $(\text{NH}_4)_2\text{CO}_3$, is the important group-reagent—the precipitant of barium and calcium.

Note 2.—In the above and in subsequent charts of analytical processes the leading precipitants will be found to be ammonium salts. These being volatile, can be got rid of toward the end of the operations, and thus the detection of potassium and sodium be in no way prevented—an advantage which could not be had if such salts as potassium chromate or sodium phosphate were the group precipitants employed.

Note 3.—Acetic, and not hydrochloric or nitric, acid is used in dissolving the barium and calcium carbonates, because barium chromate—on the precipitation of which the detection of barium depends—is soluble in the stronger acids, and therefore could not be thrown down in their presence.

Note on Classification.—The compounds of barium, calcium, and magnesium, like those of the alkali metals, have many analogies; the carbonate, phosphate, and arseniate of each are insoluble in water, which sufficiently distinguishes them from the members of the class first studied. They possess, however, well-marked differences, so that their separation from each other is easy. The solubility of their hydrates in water marks their connection with the *alkali metals*; the slightness of that solubility, diminishing as we advance farther and farther from the alkalies, baryta being most and magnesia least soluble in water, points to their connection with the next class of metals, the hydrates of which are insoluble in water. These considerations must not, however, be overvalued. Though the solubility of their hydrates places barium nearest and magnesium farthest from the alkali metals, the solubility of their sulphates gives them the opposite order, magnesium sulphate being most soluble, calcium sulphate next, strontium sulphate third (strontium is a rarer element, mentioned subsequently), while barium sulphate is insoluble in water. These elements are sometimes spoken of as *the metals of the alkaline earths*.


Note.—In connection with the bivalence of the atoms of barium, calcium, and magnesium, it is interesting to note that just as bivalent acidulous radicals give salts containing two atoms of univalent basylous radicals ($\text{K}_2\text{SO}_4, \text{H}_2\text{CO}_3, \text{KNaC}_4\text{H}_4\text{O}_6$), so bivalent basylous radicals yield salts containing two atoms of univalent acidulous radicals, as seen in acetonitrate of barium, $\text{BaC}_2\text{H}_3\text{O}_2\text{NO}_3$, a salt which is a definite compound, and not a mixture of acetate with nitrate. A large number of such salts is known.

Distillation.

The water with which, in analysis, solution of a salt or dilution of a liquid is effected should be pure. Well and river-waters are unfit for the purpose, because they contain alkaline and earthy salts (some 20 to 60 grains per gallon), derived from the soil through which the water percolates, and rain-water is not infrequently contaminated with the dust and *débris* which fall on the roofs whence it is usually collected. Such water is purified by *distillation*, an

operation in which the water is, by ebullition, converted into steam, and the steam condensed again to water in a separate vessel, the fixed earthy and other salts remaining in the vessel in which the water is boiled. On the large scale ebullition is effected in metal boilers having a hood or head in which is a wide lateral channel

Fig. 28.



Distillation, on a Small Scale.

through which passes the steam; on the small scale either a common glass flask is employed, into the neck of which, by a cork, is inserted a glass tube bent to an acute angle, or a *retort* is used (*a*, Fig. 28), a sort of long-necked Florence flask, dextrously bent near the body by the glass-worker to an appropriate angle (hence the name *retort*, from *retorqueo*, I bend back). *Condensation* is effected by surrounding the lateral steam-tube with cold water. In large stills the steam-tube, or *condensing worm*, is usually a metal (tin) pipe, twisted into a spiral form for the sake of compactness, and so fixed in a tub that a few inches of one end of the pipe may pass through and closely fit a hole bored near the bottom of the tub. Cold water is kept in contact with the exterior of the pipe, provision being made for a continuous supply to the bottom, while the lighter water, heated by the condensing steam, runs off from the top of the column. The condenser for a flask or retort may be a simple glass tube of any size placed within a much wider tube (a common long, narrow lamp-glass answers very well for experimental operations), the inner tube being connected at the extremities of the wider by bored corks; a stream of water passes into one end of the enclosed space (the end farthest from the retort), through a small glass tube inserted in the cork, and out at the other end through a similar tube. The common (Liebig's) form of laboratory condenser is a glass tube three-fourths of an inch wide and a yard long (*b*, Fig. 28), surrounded by a shorter tin or zinc tube (*c*, Fig. 28) two inches in diameter, and having at each extremity a neck, through which the glass tube passes. The ends of the necks of the tin tube and small portions of the glass tube near them are connected by means of a strip of sheet caoutchouc carefully bound round, or by short wide india-rubber tubes (*d* and *e*, Fig. 28). An aperture (*f*, Fig. 28) near the lower part of the tin tube provides for the admission of a

current of cold water, by glass tubing or india-rubber tubing, from the house supply or from a vessel placed above the apparatus; and a similar aperture near the top (*g*, Fig. 28) allows the escape of heated water into a vessel or sink. The inner tube may thus constantly be surrounded by cold water, and heated vapors passing through it be perfectly cooled and condensed, and collected in any receiver (*h*, Fig. 28).

In distilling several gallons of water for analytical or medicinal purposes (*Aqua Destillata*, U. S. P.) the first two or three pints should be rejected, because they are likely to contain ammoniacal and other volatile impurities.

Water (*Aqua*, B. P.) is defined as natural water, the purest that can be obtained, cleared, if necessary, by filtration; free from odor, unusual taste, and visible impurity. The official "water" (*Aqua*, U. S. P.) is not to contain more than 1 part in 2000 of soluble salts, and to be so free from organic matter that when tinted rose-red with potassium permanganate the color should not be destroyed after boiling the fluid for five minutes, or, in the case of distilled water, after setting the vessel aside, well covered, for ten hours. In dispensing prescriptions *aqua* should be understood to mean distilled water.

Rectification is the process of redistilling a distilled liquid. *Rectified spirit* is spirit of wine which has thus been treated.

Dry or destructive distillation is distillation in which the condensed products are directly formed by the decomposing influence of the heat applied to the dry or non-volatile substances in the retort or still.

Exercise.—Write from memory a short description of *distillation*.

RECAPITULATORY.

The subject just alluded to (distillation) naturally causes wonder respecting the cause of the physical differences between solid, liquid, and gaseous water. Common observation will have suggested to the student that the force of heat has much to do with the differences; and if he will turn to the chapter on *latent heat* in any book on Physics, he will find that, as already indicated (p. 86), when ice liquefies by heat a very large amount of heat must be given before the slightest rise of temperature occurs. Afterward the addition of heat makes the water hotter and hotter, until one other point is reached (the boiling-point), when here again a great amount of heat is absorbed without causing the slightest rise of temperature. Afterward more heat makes the gaseous water hotter and hotter, until, like a bar of iron, the steam, under special conditions, is made red hot or even white hot. Different bodies absorb different amounts of heat in changing their physical condition from solid to liquid or liquid to gas (or vapor). The amount is constant for any one body; hence definite comparative numbers may be used for expressing the *latent heats* of substances.

The absorption of heat at particular (liquefying and vaporizing) points must not be confounded with an analogous physical action—namely, the absorption of heat which goes on when a body is rising in temperature. The amount of this absorption, also, differs with different substances. That is to say, if equal weights of several substances, all at the same temperature, be heated to a stated higher temperature, very different amounts of fuel will be required. The particular or specific amount in each case is always the same; hence the *specific heats* of substances may be expressed by numbers. See the chapter on “Specific Heat” in any manual of Physics.

But after reading what has been stated respecting the constitution of matter (pp. 42 to 46), the chemical student will, in connection with the subject of distillation, be led once more to think over the subject of molecular constitution of solid, liquid, and gaseous water, and of the molecular condition of bodies generally. As previously stated, little can be told him respecting the molecular condition of solids and liquids, for temperature and pressure affect them unequally; whence we conclude that though the relation to each other of the molecules of any one substance is constant, this relation is different in different bodies. Different *gases*, however, are not differently affected, but similarly affected, by temperature and pressure; whence we conclude that their molecular constitution—the relation of their molecules to one another—is similar.

Another gas, ammonia, has been brought before the reader since the molecular constitution of gases was considered.


A small quantity of ammonia gas enclosed in the upper part of a roughly graduated test-tube over *mercury* (water would dissolve it), and exposed to the continuous action of the electric spark by means of wires of platinum fused in the sides of the tube, is decomposed into its elements, nitrogen and hydrogen, the bulk of gas operated on being exactly doubled. This expansion is not due to the gaseous molecules receding from each other, but to every two molecules becoming four similar-sized molecules:



Here each space (rectangular chiefly for convenience in printing) represents a molecule and each letter an atom. Each space, if regarded as the side of a double cube, may also, for the moment, represent two volumes, such two volumes yielding, in the decomposition, one volume of nitrogen and three volumes of hydrogen, or the four such double-cube volumes of ammonia shown in the diagram yielding two volumes of nitrogen and six volumes of hydrogen.

Remembering that a symbol (of a gas) represents one volume, and that a *formula* (of a gas) always represents two volumes, the pupil will now see how full of meaning is such an equation as the following, including, as it does, names of the elements, number of atoms, nature of the molecules, number of the molecules, weights

of atoms of the molecules, and therefore weights of bulks of the bodies, or extent of expansion in the disunion of the elements, and therefore their extent of contraction in the act of union :



AT THIS STAGE THE LEARNER IS AGAIN RECOMMENDED TO READ THE PARAGRAPHS ON THE GENERAL PRINCIPLES OF CHEMICAL PHILOSOPHY (PAGES 35-57), AND TO RETURN TO THEM FROM TIME TO TIME UNTIL THEY ARE THOROUGHLY COMPREHENDED.

ZINC, ALUMINIUM, IRON.

These three elements are classed together for analytical convenience rather than for more general analogies.

ZINC.

Symbol, Zn. Atomic weight, 64.9.

Source.—Zinc is tolerably abundant in Nature as sulphide (ZnS) or *blende*, and carbonate (ZnCO_3) or *calamine* (from *calamus*, a reed, in allusion to the appearance of the mineral). The ores are roasted to expel sulphur, carbonic acid gas, and some impurities, and the resulting oxide heated with charcoal, when the metal vaporizes and readily condenses. Zinc is a brittle metal, but at a temperature somewhat below 300°F . is malleable and may be rolled into thin sheets. Above 400° it is again brittle, and may then be pulverized. At 773°F . it melts, and at a bright red heat is volatile. Zinc in exceptionally fine powder ignites spontaneously, especially if damp or if stored in a warm place.

Uses.—Its use as a metal is familiar: alloyed with nickel, it yields German silver; with twice its weight of copper it forms common brass, and as a coating on iron (the so-called *galvanized iron*) greatly retards the formation of rust. Most of the salts of zinc are prepared, directly or indirectly, from the metal (*Zincum*, U. S. P.).

Quantivalence.—The atom of zinc is bivalent, Zn'' .


Molecular Weights.—Some remarks on this point will be made under Mercury.

REACTIONS HAVING SYNTHETICAL INTEREST.

Zinc Sulphate, or Sulphate of Zinc.

First Synthetical Reaction.—Heat zinc (4 parts) with water (20 parts) and sulphuric acid (3 fluid parts) in a test-tube (or larger vessel) until gas ceases to be evolved; solution of zinc sulphate results. Filter (to separate the particles of lead, carbon, etc. present in common zinc) and concentrate the solu-

tion in an evaporating-dish; on cooling, colorless, prismatic crystals of zinc sulphate ($\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$) are deposited (*Zinci Sulphas*, U. S. P.).



Ordinary zinc does not displace hydrogen from pure sulphuric acid alone nor from pure water alone, yet it does from the mixture. The possible explanation is that as sulphuric acid combines with several different quantities of water to form definite hydrous compounds ($\text{H}_2\text{SO}_4 \cdot \text{H}_2\text{O}$; $\text{H}_2\text{SO}_4 \cdot 2\text{H}_2\text{O}$; etc.), it is one or more of these that is decomposed with elimination of hydrogen. At present we can only say that an unknown (x) amount of water is required in the reaction.


Note.—This reaction affords hydrogen and zinc sulphate; it also develops electricity. Of several methods of evolving hydrogen it is the most convenient; of the two or three means of preparing zinc sulphate it is that most commonly employed; and of the many reactions which may be utilized in the development of voltaic electricity it is one of the most convenient. The apparatus in which the reaction is effected differs according to the requirements of the operator: if the zinc sulphate alone is wanted, an open dish is all that is necessary, the action being, perhaps, accelerated by heat; if hydrogen, a closed vessel and delivery-tube; if electricity, certain vessels called cells and various complementary materials, forming altogether what is termed a battery. In each operation for one of the three the other two are commonly wasted. It would not be difficult for the operator, as a matter of amusement, to construct an apparatus from which all three should be collected.

Purification.—Impure zinc sulphate may be purified in the same manner as impure chloride. (See next Reaction.)

Zinc sulphate is isomorphous with magnesium sulphate, and, like that salt, loses six-sevenths of its water of crystallization at 100°C . An old name for it is *white vitriol* (p. 144).

Zinc Chloride, or Chloride of Zinc.


Second Synthetical Reaction.—Digest zinc in hydrochloric acid mixed with half its bulk of water; the resulting solution contains zinc chloride. Evaporate the liquid till no more steam escapes; zinc chloride (ZnCl_2) in a state of fusion remains, and, on cooling, is obtained as an opaque white solid (*Zinci Chloridum*, U. S. P.). It is soluble in water, alcohol, or ether.




This reaction is analogous to that previously described. Burnett's deodorizing or disinfecting liquid is solution of zinc chloride.

Purification of Zinc Chloride or Sulphate.—Zinc sometimes contains traces of iron or lead. and these, like zinc, are dissolved by most acids, with formation of soluble salts; they may be recognized

in the liquids by applying the test described hereafter (p. 136) to a little of the solution in a test-tube. Should either be present in the above solution, a little chlorine-water is added to the liquid till the odor of chlorine is permanent, and then the whole well shaken with some zinc hydrate or the common official zinc "carbonate" (really hydrato-carbonate; see next page). In this way iron is precipitated as ferric hydrate, and lead as peroxide:



In the British Pharmacopœia the possible presence of impurities in the zinc is recognized, and the process of purification just described incorporated with the process of preparation of *Zinci Chloridum*, *Liquor Zinci Chloridi*, and *Zinci Sulphas*. In the purification of the zinc sulphate the action of chlorine on any ferrous sulphate will result in the formation of ferric sulphate as well as ferric chloride:



Zinc carbonate will then give chloride as well as zinc sulphate, and thus the whole quantity of zinc sulphate be slightly contaminated by chloride. On evaporating and crystallizing, however, the zinc chloride will be retained in the mother-liquor. These processes of purification admit of *general* application.

For *Liquor Zinci Chloridi*, B. P., 1 pound of zinc is placed in a mixture of 44 fluidounces of hydrochloric acid and 20 of water, the mixture ultimately warmed until no more gas escapes, filtered into a bottle, chlorine-water added until the liquid, after shaking, smells fairly of chlorine, about $\frac{1}{2}$ an ounce or somewhat more of zinc carbonate shaken up with the solution until a brown precipitate of ferric hydrate or lead peroxide, or both, appears, the whole filtered, and the filtrate evaporated to 40 fluidounces. 1 fluidounce contains 366 grains of zinc chloride. If, on testing a little of the solution first produced with ammonia and ammonium sulphhydrate, the precipitate is quite white, neither iron nor lead was present in the zinc, and the treatment with chlorine-water and zinc carbonate is to be omitted.

The solution of zinc chloride (*Liquor Zinci Chloridi*, U. S. P.) is prepared by a somewhat similar process; nitric acid, however, is used instead of chlorine-water; the solution contains "about 50 per cent. of the salt (ZnCl_2)," sp. gr. about 1.535. It is miscible with alcohol in all proportions, indicating absence of basic chloride of zinc.

* It will be noticed that the atom of iron is represented, in these equations, as exerting both bivalent and trivalent activity; this will be alluded to when iron comes under consideration.

Zinc Bromide, ZnBr_2 (*Zinci Bromidum*, U. S. P.), may be made by the action of zinc on hydrobromic acid and evaporation to dryness. It is a white powder, but may be sublimed in needles.

Zinc Iodide, ZnI_2 (*Zinci Iodidum*, U. S. P.), may be made from its elements. It is a white powder, but when volatilized condenses in acicular prisms.

Carbonate of Zinc, or Zinc Carbonate.

Third Synthetical Reaction.—To solution of any given quantity of zinc sulphate in twice its weight of water (in a test-tube, evaporating-basin, or other large or small vessel) add about an equal quantity of sodium carbonate, also dissolved in twice its weight of water, and boil; the resulting white precipitate is so-called zinc carbonate (*Zinci Carbonas*, B. P., *Zinci Carbonas Præcipitatus*, U. S. P.), or precipitated zinc carbonate, a mixture of carbonate (ZnCO_3) and hydrate (Zn_2HO) in the proportion of one molecule of the former and two of the latter, together with a molecule of water (H_2O); these proportions, however, vary considerably. It may be washed, drained, and dried in the usual manner. It is used in the arts under the name of *zinc-white*.




Calamina Præparata, B. P., is a smooth, pale pinkish-brown powder, obtained by calcining and powdering native zinc carbonate or *calamine*, and freeing the product from gritty particles by elutriation. *Prepared calamine* is chiefly zinc carbonate with some iron oxide, etc.

Elutriation (Lat. *elutriatus*; *elutreo*; *eluo*; I wash out).—This fractional operation consists in straining off water (or other liquids—light like ether, or heavy like chloroform) containing lighter particles in suspension from heavier and coarser particles which have become deposited. The decanted fluid yields a sediment of the fine particles on standing. By allowing a varying number of seconds to elapse between the shaking and the decantation, and by the use of fluids of different specific gravities and different degrees of limpidity or viscosity, substances of different specific gravities or particles of different degrees of fineness of any one substance may be separated from each other.

Zinc Acetate, or Acetate of Zinc.


Fourth Synthetical Reaction.—Collect on a filter the precipitate obtained in the last reaction, wash with distilled water, and dissolve a portion in strong acetic acid; the resulting solution contains zinc acetate, and on evaporating and setting aside

yields lamellar pearly crystals ($\text{Zn}_2\text{C}_2\text{H}_3\text{O}_2 \cdot 2\text{H}_2\text{O}$), *Zinci Acetas*, U. S. P.



Zinc Oxide, or Oxide of Zinc.

Fifth Synthetical Reaction.—Dry the remainder of the precipitated carbonate (by placing the open filter on a plate over a dish of water kept boiling), and then heat it in a small crucible till trial samples taken out of the crucible from time to time cease to effervesce on the addition of water and acid; the product is zinc oxide (*Zinci Oxidum*, U. S. P), much used in the form of ointment (*Unguentum Zinci Oxidi*, U. S. P.).



Note.—This oxide is yellow while hot, and of a very pale yellow or slight buff tint when cold—not actually white, like the oxide prepared by the combustion of zinc in air. The preparation of the latter variety, which also occurs in commerce, can only be practically accomplished on the large scale; but the chief features of the action may be observed by heating a piece of zinc on charcoal in the blow-

Fig. 29.





Fig. 30.



The Blowpipe.

pipe flame (Fig. 29) till it burns; flocks escape, float about in the air, and slowly fall. These are the old *Flores Zinci*, *Lana Philosophica*, or *Nihilum Album*.

A clear blowpipe flame consists more or less of two portions (see Fig. 30)—an inner cone, at the apex of which are hot hydrocarbon gases greedy of oxygen, and an outer cone, at the apex of which is excess of hot oxygen. At the latter point oxidizable metals, etc. are readily oxidized, as in the foregoing experiment, and that part of the flame is therefore termed *the oxidizing flame*; in the inner flame oxides and other compounds (a grain of lead acetate may be

employed for illustration) are reduced to the metallic state; hence that part is termed *the reducing flame*. A blowpipe flame is much altered in character by slight variations in the position of the nozzle of the blowpipe, by the form of the nozzle, by the force with which air is expelled from the blowpipe, and by the character of the jet of gas.

Zinc oxide slowly absorbs carbonic acid from moist air, and is partly reconverted into the hydrato-carbonate.

Zinc Valerianate, or Valerianate of Zinc.

Sixth Synthetical Reaction.—The zinc valerianate ($\text{Zn}_2\text{C}_5\text{H}_9\text{O}_2, \text{H}_2\text{O}$), *Zinci Valerianas*, U. S. P., is prepared by mixing strong solutions of zinc sulphide and sodium valerianate, cooling, separating the white pearly crystalline substance, evaporating at 200°F. to a low bulk, cooling, again separating the lamellar crystals, washing the whole product with a small quantity of cold distilled water, draining, and drying by exposure to air at ordinary temperatures. Zinc valerianate is soluble in ether, alcohol, or hot water.



Other Zinc Compounds.

Oleatum Zinci, U. S. P., is a zinc soap or zinc oleate dissolved in a considerable excess of oleic acid. It is prepared by dissolving 5 parts of zinc oxide in 95 of oleic acid.

Zinc Sulphide and *Hydrate* are mentioned in the following analytical paragraphs. The formula of zinc sulphite is $\text{ZnSO}_3, 3\text{H}_2\text{O}$.

REACTIONS HAVING ANALYTICAL INTEREST (TESTS).

First Analytical Reaction.—To solution of a zinc salt (sulphate, *e. g.*) in a test-tube add solution of ammonium sulphhydrate (NH_4HS); a white precipitate (zinc sulphide, ZnS) falls, insoluble in acetic acid, soluble in the stronger acids.

Note.—This is the only white sulphite that will be met with. Its formation, on the addition of the ammonium sulphhydrate, is therefore highly characteristic of zinc. If the zinc salt contains iron or lead as impurities, the precipitate will have a dark appearance, the sulphides of those metals being black. Aluminium hydrate, which is also white and precipitated by ammonium sulphhydrate, is the only substance for which zinc sulphide is likely to be mistaken, and *vice versa*; but, as will be seen immediately, there are good means of distinguishing these from each other.

Second Analytical Reaction.—To solution of a zinc salt add ammonia; a white precipitate (zinc hydrate, Zn_2HO) appears.

Add excess of ammonia; the precipitate is redissolved. This reaction at once distinguishes a zinc salt from an aluminium salt, aluminium hydrate being insoluble in dilute ammonia.

Other Analytical Reactions.—The fixed alkali-hydrates afford a similar reaction to that just mentioned, zinc hydrate redissolving if the alkali is free from carbonate. Ammonium carbonate yields a white precipitate of carbonate and hydrate, soluble in excess. The fixed alkaline carbonates give a similar precipitate, which is not redissolved if the mixed solution and precipitate be well boiled. Potassium ferrocyanide precipitates white zinc ferrocyanide (Zn_2FeCy_6).

Magnesium sulphate, which is isomorphous with, and sometimes indistinguishable in appearance from, zinc sulphate, is not precipitated from its solutions either by potassium ferrocyanide or ammonium sulphhydrate.

Antidotes.—There are no efficient chemical means of counteracting the poisonous effects of zinc. Large doses, fortunately, act as powerful emetics. If vomiting has not occurred or apparently to an insufficient extent, solution of sodium carbonate (common washing soda), immediately followed by white of egg and demulcents, may be administered, and then the stomach be cleared.

QUESTIONS AND EXERCISES.

Give the sources and uses of metallic zinc.—Give a diagram of the action of zinc on diluted sulphuric acid.—How may solutions of zinc chloride or sulphate be purified from salts of iron? Give equations for the reactions.—State the formula of the official zinc carbonate, and illustrate by a diagram the reaction which takes place in its production.—Give an equation for the synthesis of zinc acetate.—In what respect does zinc oxide, resulting from the ignition of the carbonate, differ from that produced during the combustion of the metal?—How is zinc valerianate prepared?—What are the properties of zinc valerianate?—Name the more important tests for zinc.—How would you distinguish, chemically, between solutions of zinc sulphate and alum?—Describe the treatment in cases of poisoning by zinc salts.—Give reactions distinguishing zinc sulphate from magnesium sulphate.


ALUMINIUM, OR ALUMINUM.

Symbol, Al. Atomic weight, 27.

Note on Nomenclature.—The author prefers the spelling *aluminium*, as giving a more euphonious word than *aluminum*. In the United States Pharmacopœia the second *i* is omitted both in the English and the Latin names.

Note on Quantivalence.—In the formulæ of aluminium salts it will be observed that to one atom of metal there are three atoms of other

univalent radicals; hence, apparently, the atom of aluminium is trivalent, Al''' . But possibly it is quadrivalent, for one molecule of aluminium compounds includes two atoms of the metal, three-fourths only of whose power may be supposed to be exerted in retaining the other constituents of the molecule, the remaining fourth enabling the aluminium atoms themselves to keep together. This is graphically shown in the following formula of aluminium chloride (Al_2Cl_6), which represents each aluminium atom as a body having four arms or bonds, three of which are engaged in grasping the arms of univalent chlorine atoms, while the fourth grasps the



corresponding arm of its brother aluminium atom. Such *structural formulae*—or *graphic formulae*, as they are called—are useful in facilitating the acquirement of hypotheses regarding the constitution of chemical substances, especially if the error be avoided of supposing that they are pictures either of the position or absolute power of atoms in a molecule, or, indeed, the true representation of a molecule at all; for on this

point man knows little or nothing. $AlCl_3$ may be the formula at very high temperatures.

Source.—Aluminium is abundant in nature, chiefly as silicate in clays, slate, marl, granite, basalt, and a large number of minerals. *Mica* or *talc* consists chiefly of aluminium, iron, magnesium, and potassium silicates. The sapphire and ruby are almost pure aluminium oxide. *Rottenstone* is a soft, friable aluminium silicate containing a little organic matter.

The metal aluminium is obtained from the double aluminium and sodium chloride by the action of metallic sodium, the source of the chloride being the mineral *bauxite*, a more or less ferruginous aluminium hydrate; also by electrolysis. It is readily attacked by inorganic and organic acids.

Aluminium readily alloys with other metals. 1 part fused with 9 of copper gives *aluminium bronze*. *Aluminium steel* is a hard and tenacious alloy of a little aluminium with the iron.

Alum (*Alumen*, U. S. P.), a double aluminium and ammonium sulphate ($Al_2(SO_4)_3 \cdot Am_2SO_4 \cdot 24H_2O$) or aluminium and potassium ($Al_2(SO_4)_3 \cdot K_2SO_4 \cdot 24H_2O$), may be obtained from aluminous schist (from $\sigma\chi\iota\sigma\tau\omicron\varsigma$, *schistos*, divided), a sort of pyritous slate or shale, by exposure to air; oxidation and chemical change produce aluminium sulphate, iron sulphate, and silica from the aluminium silicate and iron bisulphide (iron pyrites) originally present in the shale. The aluminium sulphate and iron sulphate are dissolved out of the mass by water, and ammonium or potassium sulphate or chloride added; on concentrating the liquid, alum crystallizes out, while the more soluble iron salt remains in the mother-liquor.

It is more frequently prepared by directly decomposing the aluminium silicate in the calcined shale of the coal-measures by hot sulphuric acid, ammonium or potassium salts being added from time to time until a solution strong enough to crystallize is obtained.

The liquid, well agitated during cooling, deposits alum in minute crystals termed *alum-flour*, which is afterward recrystallized.

Alums.—There are several *alums*, iron or chromium taking the place of aluminium, and potassium or sodium that of ammonium, all crystallizing in an eight-sided form, the octahedron—a sort of double pyramid. They are apparently alike in chemical constitution, and their general formula (M = either metal) is M'''_2SO_4 , $M'_2SO_4 \cdot 24H_2O$. The alum of the manufacturer commonly occurs in colorless, transparent, octahedral crystals, massed in lumps, which are roughly broken up for trade purposes, but still exhibit the faces of octahedra. It contains ammonium sulphate or potassium sulphate, according as one or other is the cheaper. At the present time potassium alum is the variety met with in trade, and this is the official recognized variety.

Note on Constitution or Structure.—In presence of the fact that a great change in the properties of aluminium sulphate, potassium sulphate, and water ensues when those three substances unite to form alum, it would seem to be wrong to picture alum by a formula still reflecting those three substances; thus, $Al_2SO_4 \cdot K_2SO_4 \cdot 24H_2O$. Many chemists admit this, and find reasonable excuse in sheer inability to offer any more probable formula in the present stage of chemical knowledge. Many chemists, on the other hand, defend the formula, and explain that while no doubt the particles in each of the original separate substances are united by *atomic* or ordinary chemical attraction or affinity, those substances are, in alum, united by *molecular* chemical attraction. But this view involves the assumption of the existence of either a new force or of two forms of chemical force, in which case the old position that permanent and entire alterations in properties are due to the action of a single force, the chemical force, is no longer tenable. It is to be feared that both learner and teacher must be content to remain for the present in somewhat of a dilemma. The discovery that resolves this dilemma will probably lay bare the cause of the properties and phenomena attendant upon the formation of all salts containing what is termed “water of crystallization,” as well as of “double salts,” “solutions,” “alloys,” “amalgams,” and perhaps what is conveniently spoken of as “the variations in the valency of atoms.”


Preparation of Alum.—Prepare alum by heating a small quantity of powdered pipeclay (aluminium silicate) with about twice its weight of sulphuric acid for some time, dissolving out the resulting aluminium sulphate and excess of sulphuric acid by water, and adding potassium carbonate to the clear filtered solution until, after well stirring, the excess of acid is neutralized. (If too much carbonate be added, aluminium hydrate precipitated when the carbonate is first poured in will not be redissolved on well mixing the whole. Perhaps the readiest indication of neutrality in this and similar cases is the presence of a *little* precipitate after stirring and warming the mixture.) On evaporating the clear solution, crystals of alum are obtained.

Aluminium Sulphate, or “*Alum-cake*” ($Al_2SO_4 \cdot 9H_2O$), prepared from natural silicates in the manner just described, is a common

article of trade, serving most of the manufacturing purposes for which alum was formerly employed. It is a pure variety, official in the United States Pharmacopœia (*Alumini Sulphas*). It may be made by dissolving aluminium hydrate in diluted sulphuric acid, with subsequent removal of water by evaporation.



Aluminium hydrate (*Alumini Hydras*, U. S. P.) is to be prepared by the addition of solution of alum to solution of sodium carbonate, the precipitated hydrate being collected on a filter and well washed.




Dried Alum (*Alumen Exsiccatum*, U. S. P.) is alum from which the water of crystallization has been expelled by heat, the temperature not exceeding 205° C. or 400° F. By calculation from the molecular weight of alum, it will be found that the salt contains between 47 and 48 per cent. of water. At temperatures above 400° F. ammonium alum is decomposed, ammonium sulphate and sulphuric anhydride escaping, and pure aluminium oxide (Al_2O_3) remaining. Dried alum rapidly reabsorbs water from the atmosphere, and is slowly but completely soluble in water.

Roche Alum, or *Rock Alum* (*roche*, French, rock), is the name of an impure native variety of alum containing iron. The article sold under this name is generally an artificial mixture of common alum with ferric oxide.

The *Ammonio-ferric Alum* or *Ammonio-ferric Sulphate* or *Ferric Ammonium Sulphate* of American pharmacy (*Ferri et Ammonii Sulphas*, U. S. P.) may be made by adding ammonium sulphate to a hot solution of iron persulphate, and setting the liquid aside to crystallize. It forms pale violet octahedral crystals, expressed by the formula $\text{Fe}_2\text{SO}_4, (\text{NH}_4)_2\text{SO}_4, 24\text{H}_2\text{O}$.

REACTIONS HAVING ANALYTICAL INTEREST (TESTS).

First Analytical Reaction.—To a solution of an aluminium salt (alum, for example, which contains aluminium sulphate) add ammonium sulphhydrate (NH_4HS); a gelatinous white precipitate (aluminium hydrate) falls.



Second Analytical Reaction.—To solution of alum add ammonia, NH_4HO ; aluminium hydrate falls: add excess of ammonia; the precipitate is, practically, insoluble.

Principle of Dyeing by Help of Mordants.—The precipitated aluminium hydrate, alumina, has great affinity for vegetable coloring-matters and also for the fibre of cloth. Once more perform the above experiment, but before adding the ammonia introduce some decoction of logwood, solution of cochineal, or

other similar colored liquid into the test-tube. Add now the ammonia, and set the tube aside for the alumina to fall; the latter takes down with it all the coloring principle. In dye-works the fabrics are passed through liquids holding the alumina but weakly in solution, and then through the coloring solutions: from the first bath the fibres abstract alumina, and from the second the alumina abstracts coloring-matter. Some other metallic hydrates, notably those of tin and iron, resemble alumina in this property; they are termed *mordants* (from *mordens*, biting); the substances they form with coloring-matters have the name of *lakes*.

Third Analytical Reaction.—To the solution of alum add solution of potash; again aluminium hydrate falls. Add excess of potash and agitate; the precipitate dissolves.

Aluminium hydrate may be precipitated from this solution by neutralizing the potash with hydrochloric acid and adding ammonia, until, after shaking, the mixture has an ammoniacal smell, or by adding solution of ammonium chloride to the potash liquid. But the former way is the better, for it is difficult to know when a sufficiency of ammonium chloride has been poured in; whereas reaction with blue and red litmus-paper at once enables the operator to know when excess of hydrochloric acid or of ammonia has been added.

Alkaline carbonates, phosphates, arseniates, and salts of other acidulous radicals also decompose solutions of aluminium salts, and produce insoluble compounds of that metal with the several acidulous radicals (except the carbonic), but the resulting precipitates are of no special interest.

QUESTIONS AND EXERCISES.

What is there remarkable about the quantivalence of aluminium?—Practically, what is the quantivalence of the atom of aluminium?—Enumerate the chief natural compounds of aluminium.—Write down a formula which will represent either of the alums.—Which alum is official, and commonly employed in the arts?—State the source and explain the formation of alum.—What is the crystalline form of alum?—Work a sum showing how much dried alum is theoretically producible from 100 pounds of alum. *Ans.* 52 lb. 6 oz.—Show by figures how ordinary ammonium alum is capable of yielding 11.356 per cent. of aluminium oxide.—Why are aluminium compounds used in dyeing?—How are salts of aluminium analytically distinguished from those of zinc?

IRON.

Symbol, Fe. Atomic weight, 56.

Sources.—Compounds of iron are abundant in nature. *Magnetic Iron Ore*, or *Lodestone* (*Lodestone* or *Leadstone*, from the Saxon *lædan*, to lead, in allusion to its use, or rather to the use of magnets made from it, in navigation), is the chief ore from which Swedish iron is made: it is a compound of ferrous and ferric oxide ($\text{FeO}, \text{Fe}_2\text{O}_3$). Much of the Russian iron is made from *Specular Iron Ore* (from *speculum*, a mirror, in allusion to the lustrous nature of the crystals of this mineral); this and *Red Hæmatite* (from *αἷμα*, *haima*, blood, so named from the color of its streak), an ore raised in Lancashire, are composed of ferric oxide only (Fe_2O_3). *Brown Hæmatite*, an oxyhydrate, is the source of much of the French iron. *Spathic Iron Ore* (from *spatha*, a slice, in allusion to the lamellar structure of the ore) is a ferrous carbonate (FeCO_3). An impure ferrous carbonate forms the *Clay Ironstone* whence most of the English iron is derived. The chief Scotch ore is also an impure carbonate, containing much bituminous matter: it is known as *Black Band*. *Iron Pyrites* (from *πῦρ*, *pur*, fire, in allusion to the production of sparks when sharply struck), (FeS_2), is a yellow lustrous mineral of use only for its sulphur. As met with in coal it is commonly termed *coal brasses*. Ferrous carbonate (FeCO_3), chloride ($\text{FeCl}_2, 4\text{H}_2\text{O}$), and sulphate ($\text{FeSO}_4, 7\text{H}_2\text{O}$) sometimes occur in springs, the water of which is hence termed *chalybeate* (*chalybs*, steel).

Process.—Iron is obtained from its ores by processes of roasting, and reduction of the resulting impure oxide with coal or charcoal in the presence of chalk, the latter uniting with the sand, clay, etc. to form a fusible slag. The *cast iron* thus produced may be converted into *wrought iron* by burning out the 4 or 5 per cent. of carbon, silicon, and other impurities present by oxidation in a furnace—the old operation termed *puddling*. *Steel* is iron containing from 1 to 2 per cent. of carbon, and is made by the now celebrated Bessemer process of burning out from cast iron the variable amount of carbon it contains, and then adding melted iron containing a known proportion of carbon. The official varieties of the metal are “metallic iron, in the form of fine, bright, and non-elastic wire” (*Ferrum*, U. S. P.); and “annealed iron wire,” having a diameter about 0.005 of an inch (about No. 35 wire gauge), or wrought-iron nails free from oxide (*Ferrum*, B. P.), these being the forms in which iron is conveniently employed for conversion into its compounds. In the form of a fine powder (see the Seventeenth Reaction) metallic iron is employed as a medicine.

Properties.—The specific gravity of pure iron is 7.844; of the best bar iron, 7.7. Its color is bluish-white or gray. Bar iron requires the highest heat of a wind-furnace for fusion, but below that temperature assumes a pasty consistence, and in that state two pieces may be joined or *welded* (Germ. *wellen*, to join) by the pressure of blows from a hammer. A little sand thrown upon the hot metal facilitates this operation by forming with the superficial iron oxide a fusible slag which is dispersed by the blows: the purely

metallic surfaces are thus better enabled to come into thorough contact and enter into perfect union. Iron is highly ductile, and of all common metals possesses the greatest amount of tenacity. At a high temperature it burns in the air, forming red iron oxide. Ordinary *iron rust* is chiefly ferric oxide, with a little ferrous oxide and carbonate; it is produced by action of the moist carbonic acid of the air and subsequent oxidation. Steam passed over scrap iron heated to redness gives hydrogen gas and black iron oxide.

Quantivalence.—Iron combines with other elements and radicals in two proportions: those salts in which the atom of iron appears to possess inferior affinities (in which the other radicals are in the less amount) are termed *ferrous*, the higher being *ferric* salts. In the former the atom exerts bivalent (Fe'' or Fe_2''), in the latter trivalent activity (Fe''' or Fe_2''') as seen in the formulæ of the chlorides, FeCl_2 (possibly Fe_2Cl_4) and Fe_2Cl_6 (perhaps FeCl_3 at very high temperatures).

The atom of iron is also sometimes considered to be sexivalent, on account of the analogy of its compounds with those of chromium, which is sexivalent if the formula of its fluoride (CrF_6) be correct, and because the composition of *potassium ferrate* (K_2FeO_4), a deep-purple salt (obtained on warming a mixture, in a test-tube, of a few fragments of solid potassium hydrate and of iron perchloride with a few drops of bromine), is best explained on the assumption of the sexivalence of its iron.

Why the quantivalence of such atoms as that of iron should vary is not at present known.

The Nomenclature of Iron Salts.—For educational and descriptive purposes, also, the two classes of compounds are very conveniently spoken of as *ferrous* and *ferric*, the syllable "*ferr*," common to all, indicating their allied ferruginous character, the syllables *ous* and *ic* indicating the lower and higher class respectively—functions fulfilled by these two syllables in other similar cases (sulphurous and sulphuric, mercurous and mercuric). In the British Pharmacopœia the iron salts are known by other names; thus, *Sulphate of Iron* (*Ferri Sulphas*) and *Phosphate of Iron* (*Ferri Phosphas*)—names which are chemically inexplicit, for there are two sulphates and two phosphates, and the terms do not define which salt is intended. Consistency and uniformity would demand that the names ferrous sulphate, ferrous phosphate, or similar terms should be employed, as is done in the new United States Pharmacopœia. Practically, however, the old names cause no confusion, inasmuch as only one sulphate, phosphate, etc. is used in medicine; moreover, the higher salts usually have the prefix *per* attached (as persulphate, perchloride). These names are already well known, can easily be rendered in Latin, and then admit of simple abbreviations and adaptations such as are employed in prescriptions—advantages not possessed by the more rational terms. While, therefore, the comprehension of the chemistry of iron is rendered simple and intelligible by the use of the terms ferrous and ferric, the employment of older and less definite names may be continued in pharmacy as being more convenient under certain circumstances.

REACTIONS HAVING SYNTHETICAL INTEREST.

FERROUS SALTS.

Ferrous Sulphate.

Synonyms.—Green Sulphate of Iron ; Green Vitriol.

First Synthetical Reaction.—Place iron (small tacks) in sulphuric acid diluted with eight times its bulk of water (in a test-tube, basin, or other vessel of any required size), accelerating the action by heat until effervescence ceases.



The solution contains what is generally known as sulphate of iron—that is, ferrous sulphate—the lower of the two sulphates, and will yield crystals of that substance ($\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$), (*Ferri Sulphas*, U. S. P.), on cooling or on further evaporation ; or if the hot concentrated solution be poured into alcohol, the mixture being well stirred, the sulphate is at once thrown down in minute crystals (*Ferri Sulphas Granulatus*, U. S. P.), the precipitated sulphate of iron of U. S. P. 1880. At a temperature of 300° F. ferrous sulphate loses six-sevenths of its water, and becomes the *dried Ferrous Sulphate* (*Ferri Sulphas Exsiccatus*, U. S. P.), a salt used in the preparation of *Pilulæ Aloes et Ferri*, U. S. P. (For the nature of the chemical action with iron and diluted sulphuric acid see the analogous zinc reaction on p. 132.)

Other Sources of Ferrous Sulphate.—In the laboratory ferrous sulphate is often obtained as a by-product in making sulphuretted hydrogen.



In manufactories it occurs as a by-product in the decomposition of aluminous shale, as already noticed (p. 138).

10 grains of granulated ferrous sulphate dissolved in 1 ounce of water constitutes “Solution of Sulphate of Iron,” B. P. “The solution should be recently prepared,” because of its liability to absorb oxygen and become spoilt through formation of ferric oxy-sulphate (see below).

Notes.—Ferrous sulphate is sometimes termed *green vitriol*. Vitriol (from *vitrum*, glass) was originally the name of any transparent crystalline substance : it was afterward restricted to the sulphates of zinc, iron, and copper, which were, and still are, occasionally known as white, green, and blue vitriol. *Copperas* (probably originally *copper-rust*, a term applied to verdigris and other green incrustations of copper) is another name for this ferrous iron sulphate, sometimes distinguished as *green copperas*, copper sulphate being blue copperas. Ferrous sulphate forms a light-green double salt with ammonium sulphate (see p. 91).

Ferrous sulphate, when exposed to the air, gradually turns brown through absorption of oxygen, ferric oxysulphate ($\text{Fe}_2\text{O}_2\text{SO}_4$) being formed. The latter is not completely dissolved by water, owing to the formation of a still lower insoluble oxysalt ($\text{Fe}_4\text{O}_5\text{SO}_4$) and the soluble ferric sulphate: $5(\text{Fe}_2\text{O}_2\text{SO}_4) = \text{Fe}_4\text{O}_5\text{SO}_4 + 3(\text{Fe}_2\text{SO}_4)$.

Iron heated with undiluted sulphuric acid gives sulphurous acid gas and ferrous sulphate:



Ferrous Carbonate, or Carbonate of Iron.

Second Synthetical Reaction.—To solution of ferrous sulphate, boiling, in a test-tube, add a solution of ammonium carbonate ($(\text{NH}_4)_2\text{CO}_3$) in recently boiled hot water; a white precipitate of ferrous carbonate (FeCO_3) is thrown down, rapidly becoming light-green, bluish-green, and, after a long time, red, through absorption of oxygen, evolution of carbonic acid gas, and formation of ferric oxyhydrate.



Saccharated Ferrous Carbonate.—The above precipitate, rapidly washed with hot, well-boiled distilled water, and the moist powder mixed with sugar and quickly dried—in short, all possible precautions taken to avoid exposure to air—forms the saccharated ferrous carbonate (*Ferri Carbonas Saccharatus*, U. S. P.).

The official proportions are 10 of the sulphate dissolved in 40 of hot water, and 7 of the bicarbonate dissolved in 100 of warm water, and each filtered. The former is then added to the latter in a flask, the mixture shaken, the precipitate washed by decantation until the washings give only a very slight turbidity with barium chloride, drained, and while still somewhat moist mixed with 16 parts of sugar, and finally dried over a water-bath.

Carbonate of iron, mixed with honey and sugar, forms the *Massa Ferri Carbonatis*, U. S. P., mass of carbonate of iron or *mass of ferrous carbonate*. Saccharated ferrous carbonate, mixed with a fourth of its weight of confection of roses, forms the *Pilula Ferri Carbonatis*, B. P.

Notes.—The red powder formerly termed carbonate or subcarbonate of iron (*Ferri Carbonas* or *Ferri Subcarbonas*) was ferrous carbonate washed and dried with free exposure to air, the product thus, by the absorption of oxygen and the elements of water and elimination of carbonic acid gas, becoming ferric oxyhydrate, a compound which will come under notice subsequently (pp. 151–153). Ferrous carbonate is said to be more easily dissolved in the stomach than any other iron preparation. It is so unstable and prone to oxidation that it must be washed in water containing no dissolved air and mixed with the sugar (which protects it from oxidation) as quickly as possible. In making the official compound mixture of

iron (*Mistura Ferri Composita*, U. S. P.), "Griffith's mixture," the prepared ingredients, including the potassium carbonate, should be placed in a bottle of the required size, space being left for the crystals or solution of ferrous sulphate, which should be added last, the bottle immediately filled up with rose-water and securely corked; the minimum of oxidation is thus ensured. More than two molecular weights of the potassium carbonate to one of the ferrous sulphate are ordered in the official mixture; hence, as the ferrous carbonate decomposes, the carbonic acid produced does not necessarily escape, but converts potassium carbonate into bicarbonate. *Pilule Ferri Compositæ* (U. S. P. 1880) are made from myrrh, sodium carbonate, ferrous sulphate, and syrup: ferrous carbonate is gradually formed. *Iron Pill*, or "Blaud's pill," is also prepared with ferrous sulphate and potassium carbonate.




Ferrous Arseniate, or Arseniate of Iron.

Third Synthetical Reaction, by which the lower arseniate of iron, ferrous arseniate (*Ferri Arsenias*, B. P.) (Fe_2AsO_4), partially oxidized, is formed. This will be noticed again under Arsenium.

Ferrous Phosphate, or Phosphate of Iron.

Fourth Synthetical Reaction.—To a hot solution of ferrous sulphate in a test-tube add a hot solution of sodium phosphate and a little of a solution of sodium bicarbonate; the lower phosphate of iron, or ferrous phosphate (Fe_2PO_4), is precipitated (*Ferri Phosphas Solubilis*, U. S. P.).



According to the British Pharmacopœia, solutions of 3 ounces of sulphate of iron in 30 of hot water, and $2\frac{3}{4}$ of phosphate in 30 of hot water, together with $\frac{3}{4}$ of an ounce of bicarbonate of sodium dissolved in a little water, are well mixed, filtered, the precipitate well washed with hot water, and, to prevent oxidation as much as possible, dried at a temperature not exceeding 120° F. These proportions will be found to fairly accord with the molecular weights of the crystalline salts, multiplied as indicated in the above equation. $3(\text{FeSO}_4, 7\text{H}_2\text{O}) = 834$; $2(\text{Na}_2\text{HPO}_4, 12\text{H}_2\text{O}) = 716$; $2(\text{NaHCO}_3) = 168$. This reaction occurs in making *Syrupus Ferri Phosphatis*, B. P.

In the United States Pharmacopœia the ferric phosphate is directed to be made from the citrate instead of from the sulphate. 50 grms. of ferric citrate are dissolved in water, and 55 grms. of sodium phosphate added; the mixture is then evaporated on the water-bath to

the consistency of treacle, and spread on glass plates; it is thus obtained in the form of scales.

The use of the sodium bicarbonate is to ensure the absence of free sulphuric acid in the solution. Sulphuric acid is a powerful solvent of ferrous phosphate. It is impossible to prevent the separation of sulphuric acid if only the 834 parts of ferrous sulphate and 716 parts of sodium phosphate be employed. Ferrous phosphate is white, but soon oxidizes and becomes slate-blue. The official salt should contain at least 47 per cent. of hydrous ferrous phosphate, $\text{Fe}_3(\text{PO}_4)_2 \cdot 8\text{H}_2\text{O}$, with ferric phosphate and some oxide.

Ferrous Sulphide, or Sulphide of Iron.

Fifth Synthetical Reaction.—In a gas flame or spirit flame strongly heat sulphur with about twice its weight of iron filings in a test-tube (or in an earthen crucible in a furnace); ferrous sulphide (FeS) is formed. When cold, add water to a small portion, and then a few drops of sulphuric acid; sulphuretted hydrogen gas (H_2S), known by its odor, is evolved.



Sticks of sulphur pressed against a white-hot bar of cast iron give a pure form of ferrous sulphide. The liquid sulphide thus formed is allowed to drop into a vessel of water (Sulphide of Iron, B. P.). Or melted sulphur may be poured into a crucible full of red-hot iron nails, when a quantity of fluid ferrous sulphide is at once formed, and may be poured out on a slab.

Ferrous Iodide, or Green Iodide of Iron.

Sixth Synthetical Reaction.—Place a piece of iodine, about the size of a pea, in a test-tube, with a small quantity of water, and add a few iron filings, small nails, or iron wire. On gently warming, or merely shaking if longer time be allowed, the iodine disappears, and, on filtering, a clear light-green solution of ferrous iodide (FeI_2) is obtained. On evaporation the solid iodide remains.

Solid ferrous iodide contains about 18 per cent. of water of crystallization and a little iron oxide. It is deliquescent, and liable to absorb oxygen from the air with formation of insoluble ferric oxy-iodide or hydrato-iodide. Ferrous iodide thus spoilt may be purified by re-solution in water, addition of a little more iodine and some iron, warming, filtering, and evaporating as before. Syrup of ferrous iodide which has become brown may usually be restored by immersing the bottle in a water-bath and slowly warming.

Ferrous Bromide (FeBr_2) occasionally used in medicine, could be made, as might be expected, in the same way as the iodide. *Syrupus Ferri Bromidi*, U. S. P. 1880, contained 10 per cent. of ferrous bromide.

Ferri Iodidum Saccharatum, U. S. P., or *Saccharated Ferrous*

Iodide, is made by mixing 6 parts of iron, 17 of iodine, and 20 of water, shaking until reaction ceases, filtering into 40 parts of sugar of milk, evaporating to dryness with frequent stirring and mixing the product in a hot mortar with 20 additional parts of sugar of milk and one of reduced iron. It is a grayish or yellowish-white hygroscopic powder.


Syrupus Ferri Iodidi, or *Syrup of Ferrous Iodide*, U. S. P., contains 10 per cent. of the iodide.

FERRIC SALTS, ETC.

Ferric Chloride, or Anhydrous Perchloride of Iron.

Seventh Synthetical Reaction.—Pass chlorine (generated from black manganese oxide and hydrochloric acid in a flask) through strong sulphuric acid contained in a small bottle, and thence by the ordinary narrow glass tubing quite to the bottom of a test-tube containing twenty or thirty small iron tacks (or a Florence flask containing 2 or 3 ounces; see Fig. 31), the latter kept hot

Fig. 31.



Preparation of Anhydrous Ferric Chloride.

by a gas flame; the higher chloride of iron, ferric chloride, or the perchloride* of iron (Fe_2Cl_6) is formed and condenses in the upper part of the tube or flask as a mass of small, dark, iridescent crystals. When a tolerably thick crust of the salt is formed, break off the part of the glass containing it, being careful that the remaining corroded tacks are excluded, and place it in ten or twenty times its weight of water; the result-

* The prefix *per* (and *hyper*) used here and elsewhere is from *ὑπέρ*, *hyper*, over or above, and simply means the "highest" of several; thus perchloride, the highest chloride.

ing solution, poured off from any pieces of glass, is a pure neutral solution of ferric chloride, and will be serviceable for analytical reactions.

Precaution.—The above experiment must be conducted in the open air or in a cupboard having a draught outward.

Anhydrous Ferrous Chloride.—In breaking up the vessel scaly crystals of this substance (FeCl_3), of a light buff color, will be observed adhering to the nails.

Note.—Solution of ferric chloride evolves some hydrochloric acid on boiling, while a darker-colored solution of ferric oxychloride remains.

Formula of Ferric Chloride.—Qualitative analysis shows that ferric chloride contains iron and chlorine. Quantitative analysis shows that to 56 parts of iron there are 106.5 parts of chlorine; total, 162.5. And as 56 parts of iron are indicated by the symbol Fe, and 106.5 of chlorine by the symbol and figure Cl_3 , the formula for ferric chloride will, so far, be FeCl_3 . But equal volumes of gases and vapors contain equal numbers of molecules (see p. 531). A volume which, if water vapor, weighs 18 grains, will if ammonia weigh 17 grains, or if carbonic acid gas, 44 grains, and if iron perchloride vapor, 325 grains. And as these respective volumes contain equal numbers of molecules, one molecule of each will be represented by the same figures respectively. That is to say, the equal volumes differing in weight as the figures 18, 17, 44 and 325 differ, and the volumes containing equal numbers of molecules, the respective molecules themselves will differ in weight as the figures 18, 17, 44 and 325 differ. The formula H_2O represents the 18 parts of water, or one molecule; NH_3 , the 17 of ammonia, or one molecule; CO_2 , the 44 of carbonic acid gas, or one molecule; and the 325 parts by weight of ferric chloride can only be represented by the formula Fe_2Cl_6 , for FeCl_3 would only represent half the number thus obtained by actual experiment. Hence Fe_2Cl_6 is the formula for a molecule of ferric chloride, and not FeCl_3 —at all events, at the temperatures of the experiments (320° to 440°C.).

Hydrous Ferrous Chloride, or Green Chloride of Iron.

Solution of Hydrous Ferric Chloride.

Eighth Analytical Reaction.—Dissolve iron tacks or wire, in a test-tube, in hydrochloric acid; hydrogen escapes, and the solution on cooling, or on evaporation and cooling, deposits *crystallized ferrous chloride*, containing four molecules of water of crystallization ($\text{FeCl}_2 \cdot 4\text{H}_2\text{O}$). A syrup of ferrous chloride is official (*Syrupus Ferri Subchloridi*, B. P.).


Through a portion of the solution of ferrous chloride pass chlorine gas; the ferrous chloride becomes *ferric chloride*.

The excess of chlorine dissolved by the liquid in this experiment may be removed by ebullition, but the ferric chloride is liable to be

slightly decomposed. The free chlorine is better carried off by passing a current of air through the liquid for some time.

Hydrous Ferric Chloride (*the Official Process*).

Ninth Synthetical Reaction.—To another portion of the solution of ferrous chloride, in a test-tube, add a little hydrochloric acid; heat the liquid, and continue to drop in nitric acid until the black color it first produces disappears; the resulting reddish-brown liquid is solution of ferric chloride.



The black substance is a compound of nitrous oxide gas (NO) with a portion of the ferrous salt; it is decomposed by heat.

This is the process for producing the solution of chloride of iron, *Liquor Ferri Chloridi*, U. S. P., or *Solution of Ferric Chloride*, definite weights of materials being employed and the solution of ferrous chloride being poured slowly into the nitric acid. The sp. gr. of the liquor is about 1.387. It contains some free hydrochloric acid. Percentage of anhydrous chloride, 37.8. Practically, it is impossible so to apportion the acids that a solution shall result containing neither excess of acid nor of metal, nor contain ferric nitrate. For most medicinal purposes, however, solution of ferric chloride containing hydrochloric acid is said to be unobjectionable. On the large scale some time may be saved by adding the acid solution of ferrous chloride to the nitric acid.

If 25 cc. of the above solution be diluted with alcohol up to 100 cc., the tincture of chloride of iron will be formed (*Tinctura Ferri Chloridi*, U. S. P.), or *Tincture of Ferric Chloride*.


Note.—The spirit in the latter preparation is unnecessary, useless, and deleterious, for it acts neither as a special solvent nor as a preservative—the offices usually performed by alcohol (*Tincturæ et Succii*, U. S. P. and B. P.)—but, unless the liquid contain excess of acid, decomposes the ferric chloride and causes the formation of an insoluble oxychloride of iron. Even if the tincture be acid, it slowly loses color, ferrous chloride and chlorinated ethereal bodies being formed.

Solution of ferric chloride, evaporated, yields a mass of yellow crystals composed of $\text{Fe}_2\text{Cl}_6 \cdot 12\text{H}_2\text{O}$, or, rarely, red crystals having the formula $\text{Fe}_2\text{Cl}_6 \cdot 5\text{H}_2\text{O}$.

Ferric Sulphate, or Persulphate of Iron.

Tenth Synthetical Reaction.—Dissolve about $\frac{3}{4}$ of an ounce of ferrous sulphate and a sixth of its weight of sulphuric acid in $1\frac{1}{2}$ ounces of water in an evaporating-dish, heating the mixture and dropping in nitric acid until the black color it first produces disappears. The resulting liquid, when made of a certain prescribed strength, is the solution of ferric sulphate,

or higher sulphate, "Solution of Tersulphate of Iron" of the Pharmacopœia, a heavy, dark-red liquid, sp. gr. 1.320 (*Liquor Ferri Tersulphatis*, or *Solution of Ferric Tersulphate*, U. S. P.). Solution of subsulphate of iron, *Liquor Ferri Subsulphatis* or *Solution of Ferric Subsulphate*, U. S. P. (Monsel's solution), is a similar fluid, made with less acids, containing, therefore, ferric oxysulphate, $\text{Fe}_4\text{O}_5\text{SO}_4$ (sp. gr. about 1.320 at 15°C .).



The black color, as in the previous reaction, is due to a compound of ferrous salt with nitric oxide ($2\text{FeSO}_4 + \text{NO}$).

The definite official solution of ferric sulphate (*Liquor Ferri Persulphatis*, B. P.) is made by adding 6 fluidrachms of sulphuric acid to $\frac{1}{2}$ pint of water, warming, dissolving 8 ounces of crystals of ferrous sulphate in the liquid, then pouring it into nitric acid (6 fluidrachms or rather more) slightly diluted, until the mixture turns to a reddish color and ruddy nitrous vapors cease to be produced. The whole should measure 11 fluidounces, being diluted or further evaporated, as the case may be, to this bulk.

Note.—In reactions in which iron passes from ferrous to ferric condition it assumes different properties, the atoms of the metal, as we believe, passing from bivalent to trivalent activity.



Ferric Oxyhydrate, or Hydrated Oxide of Iron.

Eleventh Synthetical Reaction.—Pour a portion of the solution of ferric sulphate into excess of solution of ammonia; moist *ferric hydrate*, $\text{Fe}_2\text{6HO}$, is precipitated. Wash the precipitate by decantation or on a filter, and dry it on a plate over boiling water; *ferric oxyhydrate*, $\text{Fe}_2\text{O}_2(\text{HO})_2$ (*Ferri Peroxidum Hydratum*, B. P.), remains. When rubbed to a fine powder it is fit for medicinal use.

Ferric Hydrate.


Synonyms.—Ferric Hydroxide; Hydrated Oxide of Iron; Ferric Oxyhydrate.

Ferric hydrate, suspended in a certain quantity of water, forms the *Ferri Oxidum Hydratum*, U. S. P.




Either of the other alkalies (potash or soda) will produce a similar reaction, but ammonia is most convenient.

Ferric hydrate is an antidote to arsenic if administered directly after the poison has been taken. It converts the soluble arsenic (As_2O_3) into insoluble ferrous arseniate:




Dried ferric hydrate (then become an oxyhydrate, $\text{Fe}_2\text{O}_22\text{HO}$) has less action on arsenic. Even the moist, recently prepared hydrate (Fe_26HO) loses much of this power as soon as it has become converted into an oxyhydrate ($\text{Fe}_4\text{O}_36\text{HO}$)—a change which occurs though the hydrate be kept under water (W. Procter, Jr.). According to T. and H. Smith, this decomposition occurs gradually, but in an increasing ratio; so that after four months the power of the moist mass is reduced to one-half, and after five months to one-fourth. Now the mere loss of water is not usually followed by any alteration of the essential chemical properties of a compound. It would seem, therefore, that ferric hydrate (two molecules) (Fe_412HO) probably suffers, on standing, actual decomposition into oxyhydrate ($\text{Fe}_4\text{O}_36\text{HO}$) and water ($3\text{H}_2\text{O}$), and does not merely lose water already existing in it as water. Ferric hydrate is also far more readily soluble in hydrochloric acid, tartaric acid, citric acid, and acid potassium tartrate than ferric oxyhydrate. Any formulæ exhibiting ferric hydrate (Fe_26HO) as a combination of ferric oxide and water ($\text{Fe}_2\text{O}_3, 3\text{H}_2\text{O}$), or the oxyhydrate ($\text{Fe}_2\text{O}_22\text{HO}$) as a similar combination ($\text{Fe}_2\text{O}_3, \text{H}_2\text{O}$), are apparently, for these and other reasons, scarcely correct.

Ferri Oxidum Hydratum cum Magnesia (Hydrated Oxide of Iron with Magnesia, or *Ferric Hydrate with Magnesia*).—As a more trustworthy arsenic antidote a mixture of solution of ferric sulphate and magnesia is recommended in the United States Pharmacopœia. Bottles containing (a) 50 cc. of the official solution of ferric sulphate mixed with twice its weight of water, and (b) 10 parts of magnesia well mixed and diluted with 700 of water, are to be kept on hand ready for immediate use. Their contents are simply mixed, shaken together, and administered to the patient.



Ferric Oxide, or Peroxide of Iron.

The above oxyhydrate, $\text{Fe}_2\text{O}_22\text{HO}$, sometimes shortly termed *Peroxide of Iron* (B. P.), further decomposes when heated to low redness, ferric oxide or true iron peroxide (Fe_2O_3) remaining.



The six univalent atoms of the HO (hydroxyl), the characteristic elements of all hydrates, are thus, by two successive steps, split up into water and oxygen. But between the hydrate and oxide there obviously may be another oxyhydrate in which only 2HO is displaced by O' ; and such a compound is well known; it is a variety of brown iron ore. The other oxyhydrate, $\text{Fe}_2\text{O}_22\text{HO}$, is also native (needle iron ore), as well as being the *Ferri Peroxidum Hydratum*, B. P.

"Ferri Peroxidum Humidum"	$\text{Fe}'''_2\text{O}'''_2$	6HO
A variety of brown iron ore	$\text{Fe}'''_2\text{O}'''_2$	4HO
"Ferri Peroxidum Hydratum" (needle ore)	$\text{Fe}'''_2\text{O}'''_2$	2HO
Ferric oxide	$\text{Fe}'''_2\text{O}'''_3$	

The moist ferric hydrate, as already stated, when kept for some months, even under water, loses the elements of water, and is converted into an oxyhydrate, having the formula $\text{Fe}_4\text{H}_6\text{O}_9$ (limonite or brown hæmatite), which is either a compound of the above oxyhydrates ($\text{Fe}_2\text{O}_4\text{HO} + \text{Fe}_2\text{O}_2\text{HO}$) or is a definite intermediate oxyhydrate ($\text{Fe}_4\text{O}_3\text{HO}$).

By ebullition with water for seven or eight hours ferric hydrate is decomposed into water and an oxyhydrate having the formula $\text{Fe}_4\text{H}_2\text{O}_7$ (Saint-Gilles), which is either a mixture of the official oxyhydrate ($\text{Fe}_2\text{O}_2\text{HO}$) with ferric oxide (Fe_2O_3) or a definite intermediate body ($\text{Fe}_4\text{O}_3\text{HO}$). The relation of these bodies to each other will be apparent from the following table, in which, for convenience, the formulæ of ferric hydrate and oxide are doubled:

Ferric hydrate (and as stalactite)	Fe_4	12HO
Kilbride mineral (?)	$\text{Fe}_4\text{O}10\text{HO}$	
Brown iron ore (Huttenrod and Raschau)	$\text{Fe}_4\text{O}_3\text{HO}$	
Old, or frozen, ferric hydrate (and limonite)	$\text{Fe}_4\text{O}_3\text{HO}$	
Ferric oxyhydrate (B. P.) (and gothite)	$\text{Fe}_4\text{O}_4\text{HO}$	
Boiled ferric hydrate (and turgite)	$\text{Fe}_4\text{O}_5\text{HO}$	
Ferric oxide (red hæmatite, etc.)	Fe_4O_6	

A ferric oxycarbhydrate ($\text{Fe}_4\text{OCO}_3\text{HO}$) has been obtained.

The official ferric oxyhydrate ($\text{Fe}_2\text{O}_2\text{HO}$) was formerly made by mixing solutions of ferrous sulphate and sodium carbonate, and exposing the resulting ferrous carbonate to the air until it was nearly all converted into ferric oxyhydrate; hence its old names, still sometimes seen on shop-bottles, of *Ferri Carbonas* and *Ferri Subcarbonas*. Possibly it sometimes contained a little of the oxycarbhydrate just mentioned.


Ferric Oxide (another process).

Twelfth Synthetical Reaction.—Roast a crystal or two of ferrous sulphate (mixed with a little sulphur to aid the reduction) in a small crucible until fumes cease to be evolved; the residue is a variety of ferric oxide (Fe_2O_3), or iron peroxide, known in trade as *red oxide of iron*, *colcothar*, *crocus*, *rouge* (mineral), *Venetian red*, etc. It has sometimes been used in pharmacy in mistake for the official oxyhydrates (*vide* Eleventh Synthetical Reaction); from which it differs not only in composition, but in the important respect of being almost insoluble in acids.

Ferric Acetate, or Acetate of Iron.

Thirteenth Synthetical Reaction.—Digest recently washed

and drained ferric hydrate in glacial acetic acid; ferric acetate, $\text{Fe}_26\text{C}_2\text{H}_3\text{O}_2$, is produced.



The solution of acetate of iron (*Liquor Ferri Acetatis*, U. S. P.) is an aqueous *Solution of Ferric Acetate*, containing 31 per cent. of $\text{Fe}_26\text{C}_2\text{H}_3\text{O}_2$. It is made by dissolving the ferric hydrate prepared from a known quantity of ferric sulphate in a definite weight of glacial acetic acid. Sp. gr. 1.160. 50 parts of this solution, 30 of alcohol, and 20 of acetic ether form the *Tinctura Ferri Acetatis*, U. S. P. 1880.

The "Scale" Compounds of Iron.

Fourteenth Synthetical Reaction.—Repeat the Eleventh Reaction, introducing a little solution of citric or tartaric acid or acid potassium tartrate, before adding to the alkali (soda, potash, or ammonia), and notice that now no precipitation of ferric hydrate occurs. This experiment serves to illustrate not the manufacture of a scale compound, but the chemistry of the manufacture. The effect is due to the formation of double compounds, termed ammonio-citrate, potassio-citrate, ammonio-tartrate, potassio-tartrate, and similar sodium compounds of iron, which remain in solution along with the secondary product—sulphate of the alkali metal. Such ferric compounds, made with certain prescribed proportions of recently prepared ferric hydrate (from which all alkaline sulphate has been washed), and the respective acids (tartaric or citric) or acid salts (acid potassium tartrate), etc., and the solutions evaporated to a syrupy consistence and spread on smooth plates to dry, form the scale preparations known as *Ferri et Ammonii Citras*, U. S. P., *Ferri Citras*, U. S. P. (also *Liquor Ferri Citratis*, U. S. P.), and *Ferri Potassio-tartras*, or, rather, *Ferrum Tartaratum*, B. P., *Ferri et Potassii Tartras*, U. S. P.

1 part of strychnine and 1 of citric acid with a solution containing 98 parts of ferric and ammonium citrate, the mixture being evaporated at a temperature not exceeding 60°C . or 140°F . to a syrupy consistence, and then sealing in the usual way, yield *Ferri et Strychninæ Citras*, U. S. P.

Specimens of these substances may be prepared by attending to the following details: It is essential, first, that the ferric hydrate be thoroughly washed, or an insoluble oxysulphate will be formed; secondly, that the ferric hydrate be rapidly washed, or an insoluble ferric oxyhydrate will be produced; thirdly, that the whole operation be conducted quickly, or reduction to green ferrous salt will occur; fourthly, that the

solutions of the salts be not evaporated at a higher temperature than that stated, or decomposition will take place; and, fifthly, that excess of ferric hydrate be employed.

In the pharmacopœial process for the scale compounds the ferric hydrate is in each case freshly made from solution of ferric sulphate by precipitation with solution of ammonia:



the solution of ferric sulphate being made of a definite strength from a known weight of ferrous sulphate. The reason for adopting this course is that ferric hydrate is unstable and cannot be weighed, because it cannot be dried without decomposing and becoming insoluble, as explained under the Eleventh Reaction. This definite solution of ferric sulphate (*Liquor Ferri Persulphatis*, B. P.) is made as already described (see p. 151).

Ferric Citrate, or Citrate of Iron (*Ferri Citras*, U. S. P.), and *Iron and Ammonium Citrate*, or Citrate of Iron and Ammonium (*Ferri et Ammonii Citras*, U. S. P.).—Ferric hydrate is dissolved in solution of citric acid, and the whole evaporated to dryness with or without ammonia.

To prepare the ferric hydrate, dilute 105 parts of official solution of ferric sulphate with water; pour this into water containing excess of "solution of ammonia." (If the opposite course were adopted—the alkaline liquid poured into the ferric solution—the precipitate would contain ferric oxysulphate or hydrato-sulphate, which interferes with the brilliancy of the scales.) Thoroughly stir the mixture (it will smell strongly of ammonia if enough of the latter has been used), allow the precipitate to subside, pour away the supernatant liquid, add more water, and repeat the washing until a little of the liquid tested for by-product (ammonium sulphate) by solution of barium chloride or nitrate ceases to give a white precipitate (barium sulphate). Collect the ferric hydrate on a filter, drain, and place it, while still moist, with 30 parts of citric acid in an evaporating-basin over a water-bath; stir frequently until the hydrate has dissolved. Filter, and either evaporate until the liquid weighs 100 parts to form the solution of citrate of iron (*Liquor Ferri Citratis*, or *Solution of Ferric Citrate*, U. S. P.), (sp. gr. about 1.250; strength 35.5 per cent. of $\text{Fe}_2\text{2C}_6\text{H}_5\text{O}_7$), or evaporate to a syrup at 60° C. and spread on glass plates to dry (*Ferri Citras*, U. S. P., $\text{Fe}_2\text{2C}_6\text{H}_5\text{O}_7, 6\text{H}_2\text{O}$), or to 3 parts of the liquor add 1 of ammonia-water and evaporate to form scales (*Ferri et Ammonii Citras*, U. S. P.).

Ferri et Quininæ Citras Solubilis, U. S. P., and *Ferri et Quininæ Citras*, B. P., are prepared similarly. Ferric hydrate and pure quinine are dissolved in solution of citric acid, ammonia added, and the whole evaporated to dryness. The product contains iron citrate, quinine citrate, and ammonium citrate.

The ferric hydrate is obtained from 4½ fluidounces of the solution

of ferric sulphate, with all the precautions described in the previous paragraph, a proportionate quantity of ammonia being employed.

While the ferric hydrate is being washed, prepare the quinine by dissolving 1 ounce of the ordinary quinine sulphate in 8 ounces of distilled water, acidified with sufficient sulphuric acid to dissolve the sulphate (about 12 fluidrachms of the official "diluted sulphuric acid"), and to the clear liquid add solution of ammonia, well mixing the product by stirring, until the whole of the quinine is precipitated (that is, until the mixture, after thorough agitation, smells of ammonia). Collect the precipitate on a filter, let it drain, and wash away adhering solution of ammonium sulphate by passing through it about $1\frac{1}{2}$ pints of distilled water.

(It will be observed that the principle involved in the preparation of quinine from its sulphate is identical with that which obtains in the precipitation of alumina, ferric hydrate, or zinc hydrate, etc. Such a soluble sulphate—or, indeed, any similar soluble salt—has its acidulous constituent removed by the superior affinity of the basylous radical in ammonia or other alkali, an insoluble precipitate and a new soluble sulphate being formed. The latter is washed away, leaving the former pure. In such manipulations, when economy has to be practised, soda is the alkali generally employed. Ammonia, however, has the advantage of showing the moment when its work of removing an acidulous radical is completed, for the salts which ammonium forms with such acidulous radicals as those represented by the formulæ SO_4 , Cl , NO_3 , and $\text{C}_2\text{H}_3\text{O}_2$ are inodorous, while ammonia has a powerful odor: so long, therefore, as the salt to be decomposed is not wholly attacked, the addition of ammonia does not give an ammoniacal odor to the mixture, the ammonia, as such, being, in fact, destroyed; but when the work is accomplished the quantity of ammonia last added remains as ammonia, and communicates its natural smell to the liquid.)

The ferric hydrate and quinine being now washed and drained, dissolve the former, and afterward the latter, in a solution of 3 ounces of citric acid in 5 of distilled water, the acid liquid being warmed over a water-bath, and portions of the precipitates stirred in as fast as solution is effected. "Let the solution cool, then add, in small quantities at a time, 12 fluidrachms of solution of ammonia diluted with 2 fluidounces of distilled water, stirring the solution briskly, and allowing the quinine which separates with each addition of ammonia to dissolve (in the acid) before the next addition is made (excess of ammonia must be avoided, or the quinine will be precipitated). Filter the solution, evaporate to the consistence of a thin syrup, and then dry in thin layers on flat porcelain or glass plates, at a temperature of 100°F .—*B. P.*

When dry the scales are deliquescent and of a golden-yellow color, with a bitter taste, soluble in water, but only partially so in alcohol.

The *Ferri et Quininae Citras*, U. S. P., is prepared in the same way as the soluble salt, with the exception that the ammonia is omitted. They are scales of a reddish-brown color.

Iron and Potassium Tartrate, or Potassio-ferric tartrate (*Ferri et Potassii Tartaras*, U. S. P.; *Ferrum Tartaratum*, B. P.).—Ferric hydrate is dissolved in solution of acid potassium tartrate, and the whole evaporated to dryness.

The ferric hydrate obtainable from 6 fluidounces of the official solution of ferric sulphate by the action of ammonia, in the manner detailed in the previous paragraphs, is mixed (in a mortar), while still moist, but well drained, with 2 ounces of acid potassium tartrate. The whole is set aside for about twenty-four hours, occasionally triturated to promote contact and reaction of the molecules (otherwise somewhat sluggish in attacking each other), and then heated in a dish over a water-bath to a temperature not exceeding 140°F. ; a pint of distilled water is then added, and the mixture kept warm until nothing more will dissolve; filtered, evaporated at a temperature not exceeding 140°F. (greater heat causes decomposition), and, when the mixture has the consistence of syrup, spread on panes of glass and allowed to dry (in any warm and open place shown by a thermometer to be not much hotter than 100°F.). The dry salt is thus obtained in flakes. It should be kept in well-closed bottles.

Ferri et Ammonii Tartras, Iron and Ammonium Tartrate, U. S. P., is made by saturating solution of acid ammonium tartrate with ferric hydrate, evaporating, and scaling. The acid ammonium tartrate is prepared by exactly neutralizing half of any quantity of tartaric acid by ammonia-water, and then adding the other half.


The foregoing are the only official scale preparations of iron. Many others of similar character might be formed. The citrate dissolves slowly in cold, but readily in warm, water. Few crystallize or give other indications of definite chemical composition. Their properties are only constant so long as they are made with unvarying proportions of constituents. Want of chemical compactness, the loose state in which the iron is combined, precludes their recognition as well-defined chemical compounds, yet possibly enables them to be more readily assimilated as medicines than some of the more definite ferrous and ferric salts. A definite *ferrous tartrate* ($\text{FeC}_4\text{H}_4\text{O}_6$) and *ferrous citrate* ($\text{FeHC}_6\text{H}_5\text{O}_7, \text{H}_2\text{O}$) have been obtained by reaction of iron and acid in hot water. They occur as white, gritty masses of microscopic crystals. A *sodio-ferrous citrate* ($\text{FeNaC}_6\text{H}_5\text{O}_7$) and *hydrato-citrate* ($\text{FeNa}_2\text{HOC}_6\text{H}_5\text{O}_7$) may be obtained in scales. *Ferric phosphate* (Fe_2PO_4), when freshly precipitated, is soluble in solutions of citrates of the alkali-metals. The official *Ferri Phosphas Solubilis*, U. S. P., is to be made by adding 55 parts of sodium phosphate to an aqueous solution of 50 parts of ferric citrate, and the mixture evaporated on glass plates to yield scales. It is a mixture of ferric phosphate and sodium citrate.

Wine of Iron, or "steel" wine (*Vinum Ferri*, B. P.), made by digesting iron wire in sherry wine, probably contains a little potassium tartrate and iron and other iron salts, formed by action of the metal on the small quantities of acid potassium tartrate and tartaric, citric, malic and acetic acids present in the wine. *Vinum Ferri Citratis*, B. P., is a solution of iron ammonio-citrate in orange wine.


Ferric Nitrate, or Pernitrate of Iron.

Fifteenth Synthetical Reaction.—Place a few iron tacks in

dilute nitric acid and set aside; solution of ferric nitrate, or iron perntrate, is formed (Fe_26NO_3).



Precipitate ferric hydrate from solution of ferric sulphate, wash, and dissolve it in nitric acid.




The latter is the official method for preparing *Liquor Ferri Nitratis*, solution of nitrate of iron or *Solution of Ferric Nitrate*, U. S. P., definite quantities of solution of ferric sulphate and of nitric acid being employed. Sp. gr. 1.050. Strength, about 6.2 per cent. of anhydrous nitrate.

Ferric nitrate and ferric acetate unite to form various aceto-nitrates, amongst which is one having the formula $\text{Fe}_2(\text{C}_2\text{H}_3\text{O}_2)_4\text{NO}_3\text{HO}, 4\text{H}_2\text{O}$, crystallizing in hard, shining, brownish-red prisms.

Reduced Iron.

Sixteenth Synthetical Reaction.—Pass hydrogen gas (dried by passing over pieces of calcium chloride contained in a tube or through sulphuric acid in a wash-bottle) over a small quantity of ferric oxyhydrate contained in a tube arranged horizontally (a test-tube the bottom of which has been accidentally broken serves very well), the oxyhydrate being kept hot by a gas flame; oxygen is removed by the hydrogen, steam escapes at the open end of the tube, and after a short time, when moisture ceases to be evolved, metallic iron, in a minute state of division, remains. (See Fig. 32.)




While still hot throw the iron out into the air; it takes fire and falls to the ground as oxide.

If the ferric oxide is reduced in an iron tube heated by a strong furnace, the particles of iron aggregate to some extent, and, when cold, are only slowly oxidized in dry air. This latter form of reduced iron is *Fer réduit*, or, *Quevenne's Iron*, the *Ferri Pulvis* or *Ferrum Reductum*, U. S. P.—a fine grayish-black powder, strongly attracted by the magnet, and exhibiting metallic streaks when rubbed with pressure in a mortar. It is often administered in the form of lozenges (*Trochisci Ferri Redacti*, B. P.), gum and sugar protecting the iron from oxidation as well as forming a vehicle for its administration.

Note 1.—The spontaneous ignition of the iron in the above experiment is an illustration of the influence of minute division on chemical affinity. The action is the same as occurs whenever iron

rusts, and the heat evolved and the amount of oxide formed are not greater from a given quantity of iron; but the surface exposed to the action of the oxygen of the air is, in the case of this variety of reduced iron, so enormous compared with the weight of the iron, that heat cannot be conducted away sufficiently fast to prevent

Fig. 32.




Preparation of Reduced Iron.

elevation of temperature to a point at which the whole becomes incandescent. In the slow rusting of iron escape of heat occurs, but is not observed because spread over a length of time; in the spontaneous ignition of reduced iron the whole is evolved at one moment. The mixture of lead and carbon (lead pyrophorus) resulting when lead tartrate is heated in a test-tube until fumes cease to be evolved, spontaneously ignites when thrown into the air, and for the same reason. Many substances, solid and liquid, if sufficiently finely divided and liable to oxidation, and especially if exposed in a warm place, become hot and even occasionally burst into flame spontaneously. Oil on cotton waste, powdered charcoal, coal (especially if pyritic, porous, or powdered), resins in powder, and even flour and hay, are familiar illustrations of materials liable to "heat," char, or even burn spontaneously.

Note 2.—The student having time and opportunity for the experiment is advised to convert this sixteenth reaction into a roughly quantitative one, by way of realizing what has been stated (see again the General Principles of Chemical Philosophy, pp. 35–60) respecting the action of chemical force on definite weights only of matter. Three tubes, similar to the oxide-tube shown in the engraving, should be prepared, the second being connected to the first, and the third to the second, by india-rubber tubing in the usual manner. The first tube should contain pieces of calcium chloride to absorb any traces of moisture not retained by the sulphuric acid. The second tube (the ends of the small tube being temporarily closed by small corks) should be weighed in any ordinary scales which will turn with a quarter or half a grain, and, the weight being noted, 160 grains of dry ferric oxide should be neatly placed in the middle of the tube. (The oxide before being weighed must be heated gently

in a small crucible over a lamp to remove all traces of moisture.) The third tube should contain pieces of calcium chloride to absorb the water produced in the reaction, and just before being connected should be weighed. The operation is now carried out. At its close, and when the middle tube is cold, the latter tube and the third tube are again weighed. The oxide-tube should weigh 48 grains less than before, and the terminal tube 54 grains more than before.



The operation is more quickly and easily performed if one-half or one-quarter of the weight of oxide be taken; in that case one-half or one-quarter of the weights of iron and of water will be obtained. Indeed, any weight of oxide may be employed; the amount of iron and water resulting will be *always exactly proportionate* to the weights just mentioned. Thus 16 parts of oxide yield 11.2 of iron and 5.4 of water. Iron, hydrogen, and oxygen always combine in proportions of 56, of 1, and of 16 respectively. Such facts justify us in agreeing that the symbol Fe shall stand for 56 parts by weight of iron, H for 1 part by weight of hydrogen, and O for 16 parts by weight of oxygen.

Ferric Pyrophosphate.

Seventeenth Synthetical Reaction.—To solution of sodium pyrophosphate add solution of ferric sulphate; a yellowish-white precipitate of ferric pyrophosphate ($\text{Fe}_4\text{P}_2\text{O}_7, 9\text{H}_2\text{O}$) separates.

The official *Ferri Pyrophosphas Solubilis*, or *Soluble Ferric Pyrophosphate*, the Pyrophosphate of Iron, U. S. P. 1880, is to be made by adding 10 parts of sodium pyrophosphate to an aqueous solution of 10 parts of ferric citrate, evaporating and scaling. The apple-green product is a mixture of ferric pyrophosphate and sodium citrate.

Dialyzed Iron.

“*Liquor Ferri Dialysatus*, B. P.—This solution of dialyzed iron, so called, is a solution of about 5 per cent. of highly basic ferric oxychloride or chloroxide, from which most of the acidulous matter has been removed by dialysis.” It is “a clear dark, reddish-brown liquid, free from any marked ferruginous taste.” Its specific gravity is about 1.047. (*Vide* “Dialysis” in Index.)

REACTIONS HAVING ANALYTICAL INTEREST (TESTS).

(The iron occurring as a ferrous salt.)

First Analytical Reaction.—Pass sulphuretted hydrogen

(H_2S) through a solution of a ferrous salt (*e. g.* ferrous sulphate) slightly acidulated by hydrochloric acid; no precipitate occurs. This is a valuable negative fact, as will be evident presently.

Second Analytical Reaction.—Add ammonium sulphhydrate (NH_4HS) to solution of a ferrous salt; a black precipitate (ferrous sulphide, FeS) falls.



Third Analytical Reaction.—Add solution of potassium ferrocyanide (yellow prussiate of potash), $\text{K}_4\text{Fe}'''\text{Cy}_6$, or $\text{K}_4\text{Fcy}'''$ to solution of a ferrous salt; a precipitate ($\text{K}_2\text{Fe}'''\text{Fe}'''\text{Cy}_6$, or $\text{K}_2\text{Fe}'''\text{Fcy}$) falls, at first white or bluish gray, but rapidly becoming blue, owing to absorption of oxygen.

Fourth Analytical Reaction.—To a ferrous salt add potassium ferricyanide (red prussiate of potash, $\text{K}_6\text{Fe}'''\text{Cy}_{12}$, or K_6Fdcy); a precipitate falls ($\text{Fe}'''\text{Fe}'''\text{Cy}_{12}$, or $\text{Fe}'''\text{Fdcy}$) (Turnbull's blue) resembling prussian blue in color.

Other Analytical Reactions.—The precipitates produced from ferrous solutions on the addition of alkaline carbonates, phosphates, and arseniates, as already described in the synthetical reactions of ferrous salts, are characteristic, and hence have a certain amount of analytical interest, but are inferior in this respect to the four reactions above mentioned.

Note.—Alkalies (potash, soda, or ammonia) are incomplete precipitants of ferrous salts, hence are almost useless as tests. To solution of ferrous salt add ammonia (NH_4HO); on filtering off the whitish ferrous hydrate and testing the solution with ammonium sulphhydrate, iron will still be found. To another portion of the ferrous solution add a few drops of nitric acid or excess of chlorine-water, and boil; this converts the ferrous into ferric salt, and now alkalies will wholly remove the iron, as already twice seen during the performance of the synthetical experiments.

In actual analysis the separation of iron as ferric hydrate is an operation of frequent performance. This is always accomplished by the addition of alkali after (if the iron occurs as a ferrous salt) previous ebullition with a little nitric acid. Potassium ferrocyanide and ferricyanide are the reagents used in distinguishing ferrous from ferric salts.

(The iron occurring as a ferric salt.)

Fifth Analytical Reaction.—Through a ferric solution (ferric chloride, *e. g.*) pass sulphuretted hydrogen; a white precipitate of the sulphur of the sulphuretted hydrogen falls. The ferric

is simultaneously reduced to a ferrous salt, the latter remaining in solution. This reaction is of frequent occurrence in practical analysis: $2\text{Fe}_2\text{Cl}_6 + 2\text{H}_2\text{S} = 4\text{FeCl}_2 + 4\text{HCl} + \text{S}_2$.

Sixth Analytical Reaction.—Add ammonium sulphhydrate to a ferric solution; the latter is reduced to the ferrous state, and a black substance (ferrous sulphide, FeS) is precipitated as in the second analytical reaction, sulphur being set free.

Seventh Analytical Reaction.—To a ferric solution add potassium ferrocyanide (K_4FeCy_6 or $\text{K}_4\text{Fcy}'''$); a precipitate of prussian blue (the common pigment) occurs ($\text{Fe}'''_4\text{Fe}''\text{Cy}_6$ or $\text{Fe}''_4\text{Fcy}_3'''$).

Eighth Analytical Reaction.—To a ferric solution add solution of potassium ferricyanide; no precipitate occurs, but the liquid is darkened to a brownish red, or to a greenish or olive hue if the salts are not quite pure,

Ninth Analytical Reaction.—The production of a red precipitate (ferric hydrate) on adding alkalies to ferric salts. It is identical with the eleventh synthetical reaction.

Note.—This reaction illustrates the conventional character of the terms synthesis and analysis. It is of equal importance to the manufacturer and the analyst, and is synthetical or analytical according to the intention with which it is performed.

Other ferric reactions have occasional analytical interest. In neutral ferric solutions the tannic acid in aqueous infusion of galls occasions a bluish-black inky precipitate, the basis of most black writing-inks. (The *Mistura Ferri Aromatica* of the British Pharmacopœia, made by digesting metallic iron in an infusion of various vegetable substances, contains iron tannate, or rather tannates; it is commonly known in Ireland by the name of Heberden's ink, after the physician by whom it was first used. It contains about 1 grain of iron in 1 pint.)

Potassium sulphocyanate (KCyS) causes the formation of ferric sulphocyanate, which is of a deep blood-red color. There is no normal ferric carbonate; alkaline carbonates cause the precipitation of ferric hydrate, whilst carbonic acid gas escapes.

Note.—Cyanogen (CN , or Cy'), ferro-cyanogen (FeC_6N_6 , or FeCy_6 , or simply Fcy'''), and ferri-cyanogen ($\text{Fe}_2\text{Cy}_{12}$, or Fdcy'') are radicals which play the part of non-metallic elements, just as ammonium in its chemical relations resembles the metallic elements. They will be referred to again.

Memorandum.—The reader must on no account omit to write out equations or diagrams expressive of each of the reactions of iron, analytical as well as synthetical. It is assumed that this has already been done immediately after each reaction has been performed.

DIRECTIONS FOR APPLYING THE FOREGOING ANALYTICAL REACTIONS TO THE ANALYSIS OF AN AQUEOUS SOLUTION OF SALTS CONTAINING **ONE** OF THE METALS, ZINC, ALUMINIUM, IRON.

Add solution of ammonia gradually :

A dirty-green precipitate indicates iron in the state of a ferrous salt.

A red precipitate indicates iron in the state of a ferric salt.

A white precipitate, insoluble in excess, indicates the presence of an aluminium salt.

A white precipitate, soluble in excess, indicates zinc.

These results may be confirmed by the application of some of the other tests to fresh portions of the solution.

TABLE OF SHORT DIRECTIONS FOR APPLYING THE FOREGOING ANALYTICAL REACTIONS TO THE ANALYSIS OF AN AQUEOUS SOLUTION OF SALTS OF **ONE, TWO, OR ALL THREE** OF THE METALS, ZINC, ALUMINIUM, IRON.

Boil about half a test-tubeful of the solution with a few drops of nitric acid. This ensures the conversion of ferrous into ferric salts, and enables the next reagent (ammonia) *completely* to precipitate the iron. Add excess of ammonia, and shake the mixture. Filter.

Precipitate AlFe.*		Filtrate Zn.
Dissolve in HCl, add excess of KHO, stir, filter.		Test by NH_4HS (white ppt.).
Ppt. Fe (red ppt.).	Filtrate Al.	
	Make slightly acid by HCl, and add excess of $\text{NH}_4\text{HO}^\dagger$ (white ppt.).	

* The aluminium precipitate ($\text{Al}_2\text{6HO}$) is white, the iron ($\text{Fe}_2\text{6HO}$), red. If the precipitate is red, iron must be and aluminium may be present; if white, iron is absent, and further operations on the ppt. are unnecessary.

This precipitate ($\text{Al}_2\text{6HO}$ and $\text{Fe}_2\text{6HO}$) may also, if sufficient is at disposal, be analyzed by simply well shaking a washed portion in a tube with solution of potash or soda; the ferric hydrate is not thereby affected, while the aluminium hydrate is dissolved, and may be detected in the clear decanted fluid by neutralizing all alkali by a little excess of acid, and then adding excess of ammonia.

† Alumina, when in small quantity, is sometimes prevented from

Note 1.—If iron is present, portions of the original solution must be tested by potassium ferricyanide for ferrous, and by ferrocyanide for ferric salts; dark-blue precipitates with both indicate both ferrous and ferric salts.

Note 2.—If no ferrous salt is present, ebullition with nitric acid is unnecessary. It is perhaps therefore advisable always to determine this point *previously* by testing a little of the original solution with ferricyanide; if no blue precipitate occurs, the nitric-acid treatment may be omitted.

CHART FOR ALL METALS HITHERTO CONSIDERED.

The following table (*vide* p. 165) is perhaps the best, but not the only, adaptation of the ordinary reactions to systematic analysis. In it the analytical scheme for the third group is added to that of the first two groups. As before, analysis is commenced by the addition of ammonium chloride to prevent partial precipitation of magnesium, and by ammonium to neutralize any acid, for acid destroys the group-precipitant, ammonium sulphhydrate, preventing its useful action, and causing a precipitation of any free sulphur it may contain. Any precipitate by the ammonia may be disregarded, for the sulphhydrate attacks both solid and liquid.

Note.—When a test gives no reaction, absence of the body sought may fairly be inferred. If a group-test (that is, a test which precipitates a group of substances) gives no reaction, the analyst is saved the trouble of looking for any of the members of that group.

QUESTIONS AND EXERCISES.

Name the chief ores of iron.—How is the metal obtained from the ores?—What is the chemical difference between cast iron, wrought iron, and steel?—Explain the process of “welding.”—What is the nature of chalybeate waters?—Illustrate by formulæ the difference between ferrous and ferric salts.—Under what different circumstances may the atom of iron be considered to exert bivalent, trivalent, and sexivalent activity? Write a paragraph on the nomenclature of iron salts.—Give a diagram of the process for the preparation of ferrous sulphate.—In what respects do the official ferrous sulphate, granulated ferrous sulphate, and dried ferrous sulphate differ?—How is ferrous sulphate obtained on the large scale?—Give the chemical names of white, green, and blue vitriol.—Why does ferrous sulphate become brown on exposure to air?—Show the formation of ferrous carbonate by a diagram.—Describe the action of atmospheric oxygen on ferrous carbonate; can the effect be prevented?—In what order would you mix the ingredients of *Mistura Ferri Composita*, and why?—Write out an equation illustrative of the formation of the official phosphate of iron.—Why is sodium bicarbonate used in the preparation of ferrous phosphate?—Name four compounds of iron which

being precipitated by ammonia through the presence of organic matter derived from the filter-paper by action of the potash. In case of doubt, therefore, before adding ammonia boil the liquid with a little nitric acid, which destroys any organic matter. Avoid *great* excess of ammonia.

TABLE OF SHORT DIRECTIONS FOR THE ANALYSIS OF AN AQUEOUS SOLUTION OF SALTS OF ANY OR ALL OF THE METALLIC ELEMENTS HITHERTO CONSIDERED.

Add NH_4Cl ; NH_4HO ; NH_4HS ; warm, filter.

Precipitate Fe Al Zn. Wash, dissolve in HCl .* boil (to remove H_2S), filter (to remove S), add excess of KHO ,† str, filter.		Filtrate Ba Ca Mg NH_4 Na K. Add $(\text{NH}_4)_2\text{CO}_3$, boil, filter.			
Ppt. Fe (test orig. sol. by K_4FeCy and K_6Fcy).	Filtrate Al Zn. Neutralize with HCl . Add NH_4HO , str, filter.	Ppt. Ba (yellow).	Filtrate Ca Add $(\text{NH}_4)_2\text{C}_2\text{O}_4$, (white ppt.).	Ppt. Mg (white).	Filtrate NH_4 Na K. Evap., ignite, dissolve. Na by flame; K by PtCl_4 ; orig. sol. for NH_4 .
	Ppt. Al (white).				
	Filtrate Zn. Add NH_4HS (white ppt.).				

* Add, also, a few drops of HNO_3 if Fe be present—i. e., if the ppt. be black. (Tide notes on pp. 161 and 164.)

† Or add excess of ammonia, filter, and test filtrate for zinc. Iron and aluminium hydrates may then be separated by an alkali, as described in the foot-note to the previous table, p. 163.

may be formed by the direct union of their elements.—Give the official method for the preparation of solution of ferric chloride.—Of what use is the spirit in tincture of ferric chloride?—How may ferrous be converted into ferric sulphate?—What is the formula of ferric acetate? and how is it prepared for use in pharmacy?—Give the formula for *Ferri Per-oxidum Hydratum*, B. P.—How does ferric hydrate act as an antidote to arsenic?—What are the properties of anhydrous ferrous oxide?—What are the general characters and mode of production of the medicinal scale preparations of iron?—In what state is the iron in *Vinum Ferri*, B. P.?—What other form of wine of iron is official in B. P.?—Give a diagram showing the formation of ferric nitrate.—Work out a sum showing how much ferric oxide will yield, theoretically, one hundredweight of iron. *Ans.* 160 lbs.—Explain the action of the following reagents for iron, distinguishing between ferrous and ferric reactions, and illustrating each by an equation or a diagram: *a.* Ammonium sulphhydrate. *b.* Potassium ferrocyanide. *c.* Potassium ferricyanide. *d.* Caustic alkalies. *e.* Potassium sulphocyanate.—Describe the action of ammonia on salts of iron, aluminium, and zinc respectively.—What precautions must be used in testing for calcium a solution containing iron?—How is magnesium detected in the presence of zinc?—How is aluminium detected in presence of magnesium?—Draw up a scheme for the analysis of an aqueous liquid containing salts of iron, barium, and potassium.—How may zinc, magnesium, and ammonium be consecutively removed from aqueous solution?

ARSENUM, AND STIBIUM OR ANTIMONY.

Nomenclature.—The word *arsenic* is, throughout the British Empire and the United States, understood, and always has been understood, to mean common white arsenic, sometimes termed arsenous acid. Any attempt to restrict the word “arsenic” to the element itself seems hopeless. Under these circumstances the author retains the word *arsenum* (the analogue of the name of the sister element *stibium*) for the element itself.

These elements resemble metals in appearance and in the character of some of their compounds, but they are still more closely allied to the non-metals phosphorus and nitrogen. Their atoms are quinivalent (As^v , Sb^v), as seen in arsenic anhydride (As_2O_5) and antimony pentachloride (SbCl_5), but usually exert trivalent activity only (As^{III} , Sb^{III}), as seen in the hydrogen and other compounds (AsH_3 , AsCl_3 , AsBr_3 , AsI_3). The hydrogen compounds of the four members of this group have the formulæ NH_3 , PH_3 , AsH_3 , SbH_3 . A few preparations of these two elements are used in medicine, but all are more or less powerful poisons, and hence have considerable toxicological interest.

Arsenum is an exception to the rule that the atomic weights (taken in grains, grammes, or other weights) of elements, under similar circumstances of temperature and pressure, give equal volumes of vapor, the equivalent weight (75) of arsenum only occupying half such a volume. Hence while the molecular weights (that is, double the atomic weights) of oxygen ($\text{O}_2 = 32$), hydrogen ($\text{H}_2 = 2$), nitrogen ($\text{N}_2 = 28$), etc., give a similar bulk of vapor at any given temperature and pressure, the double atomic weight of arsenum ($\text{As}_2 = 150$), at the same temperature and pressure, only

affords half this bulk. It would appear, therefore, that the molecule of arsenum contains four atoms, and that its formula is As_4 . As in the case of sulphur, however, arsenum, in the state ordinarily known to us, may be abnormal, and conditions yet be found in which the molecular weight is double (instead of quadruple) the atomic weight.

From observed analogy between the two metals the molecular constitution of stibium is probably similar to that of arsenum.

ARSENUM.

Symbol, As. Atomic weight, 75.

Sources.—Arsenical ores are frequently met with in Nature, the commonest being the iron arsenio-sulphide ($FeSAs$). This "mispickel" is roasted in a current of air, the oxygen of which, combining with the arsenum, forms common *white arsenic*, arsenous oxide, sometimes called *anhydrous arsenous acid*, or, better, *arsenous anhydride* (As_2O_3 , or, perhaps, As_4O_6) or *arsenous acid* (*Acidum Arsenosum*, U. S. P.), which is condensed in chambers or long flues. It commonly "occurs as a heavy white powder or in sublimed masses, which usually present a stratified appearance, caused by the existence of separate layers differing from each other in degrees of opacity." When very much diluted the vapor of arsenic has a garlic odor, but it is very poisonous. The vitreous or amorphous arsenic is far more soluble than the crystalline variety, and in other respects they differ in properties. Such differences between the crystalline and amorphous varieties of an element or compound are not frequent: they have not yet been satisfactorily explained. *Realgar* (red algar) is the red native arsenum sulphide (As_2S_2), and orpiment (*auripigmentum*, the golden pigment), the yellow native sulphide (As_2S_3). The arsenum iodide or tri-iodide (AsI_3), *Arseni Iodidum*, U. S. P., may be made from its elements or by dissolving white arsenic in aqueous hydriodic acid and evaporating. It occurs in small orange-colored crystals, readily and almost entirely soluble in water and in rectified spirit. Its aqueous solution has a neutral reaction, and gives a yellow precipitate with sulphuretted hydrogen. Heated in a test-tube, it almost entirely volatilizes, violet vapors of iodine being set free. An aqueous solution of about 1 per cent. of this arsenous iodide and 1 per cent. of mercuric iodide forms the *Liquor Arseni et Hydrargyri Iodidi*, U. S. P., or *Donovan's Solution*.

REACTIONS HAVING SYNTHETICAL INTEREST.

Alkaline Solution of Arsenic.

First Synthetical Reaction.—Boil a grain or two of powdered arsenic (As_2O_3) in water containing a little potassium carbonate, and, if necessary, filter. The solution, colored with compound

tincture of lavender, and containing 1 per cent. of arsenic, forms the *Liquor Potassii Arsenitis*, Solution of Potassium Arsenite, U. S. P. (*Fowler's Solution*).

Note.—This official solution does not generally contain potassium arsenite, for the arsenic does not decompose the potassium carbonate, or only after long boiling. From concentrated solutions carbonic acid gas is more quickly eliminated.

Arsenous Acid and other Arsenites.

Arsenic, or *arsenous anhydride* (the so-called arsenous acid), when dissolved in water is said to yield true arsenous acid (H_3AsO_3), hydrogen arsenite.



When arsenic is dissolved in excess of solutions of potash or soda, arsenites are formed having the formulæ KH_2AsO_3 and NaH_2AsO_3 . Boiled with excess of arsenic, one molecule of these salts combines with one of the arsenic. The usual character of such compounds is that of oily alkaline liquids.

Arsenic fused with alkaline carbonates yields pyroarsenates ($\text{Na}_4\text{As}_2\text{O}_7$ or $\text{K}_4\text{As}_2\text{O}_7$) and metallic arsenum. Arsenites have the general formula $\text{R}'_3\text{AsO}_3$.

Acid Solution of Arsenic.

Second Synthetical Reaction.—Boil white arsenic with diluted hydrochloric acid. Such a solution, made with prescribed proportions of acid (2 per cent.) and water, and containing 1 per cent. of arsenic (As_2O_3), forms the *Solution of Arsenous Acid* (*Liquor Acidi Arsenosi*, U. S. P.). *De Valangin's Solution* contains $1\frac{1}{2}$ grains per ounce.

Note.—No decomposition occurs in this experiment. The liquid is simply a solution of arsenic in the acid. The two solutions may be preserved for analytical operations.

Mem.—The practical student should boil arsenic in water only, and thus have an acid, alkaline, and aqueous solution for analytical comparison.

Arsenum.

Third Synthetical Reaction.—Place a grain or less of white arsenic at the bottom of a narrow test-tube, cover it with about half an inch or an inch of small fragments of dry charcoal, and hold the tube, nearly horizontally, in a flame, the mouth being loosely covered by the thumb. At first let the bottom of the tube project slightly beyond the flame, so that the charcoal may become nearly red-hot; then heat the bottom of the tube. The arsenic will sublime, become deoxidized by the charcoal, carbonic oxide being formed, and the element *arsenum*, sometimes termed *arsenicum*, and also, unfortunately, *arsenic*, be

deposited in the cooler part of the tube as a dark mirror-like metallic incrustation.

There is a characteristic odor, resembling garlic, emitted during this operation, probably due to a partially oxidized trace of arsenum which escapes from the tube, for white arsenic does not give this odor; moreover, arsenum being a freely oxidizable element, its vaporous particles could scarcely exist in the air in an entirely unoxidized state.

Metallic arsenum may be obtained in large quantities by the above process if the operation be conducted in vessels of commensurate size. But performed with great care, in narrow tubes, using not charcoal alone, but *black flux* (a mixture of charcoal and potassium carbonate obtained by heating acid potassium tartrate in a test-tube or other closed vessel till no more fumes are evolved), the reaction has considerable analytical interest, the garlic odor and the formation of the mirror-like ring being highly characteristic of arsenum. Compounds of mercury and antimony, however, give sublimate which may be mistaken for arsenum.

Arsenic Acid and other Arsenates.

Fourth Synthetical Reaction.—Boil a grain or two of arsenic with a few drops of nitric acid until red fumes cease to be evolved; evaporate the solution in a small dish to dryness, to remove excess of nitric acid; dissolve the residue in water: the product is arsenic acid (H_3AsO_4).

Arsenic acid, when strongly heated, loses the elements of water, and arsenic anhydride remains (As_2O_5).


Arsenic anhydride readily absorbs water and becomes arsenic acid (H_3AsO_4). Arsenic acid is reduced to arsenous by the action of sulphurous acid ($\text{H}_3\text{AsO}_4 + \text{H}_2\text{SO}_3 = \text{H}_3\text{AsO}_3 + \text{H}_2\text{SO}_4$).

Salts analogous to arsenic acid, the hydrogen arsenate, are termed *arsenates*, and have the general formula $\text{R}'_3\text{AsO}_4$. The ammonium arsenate, $(\text{NH}_4)_2\text{HAsO}_4$, may be made by neutralizing arsenic acid with ammonia. Its solution in water forms a useful reagent. Arsenic acid is used as an oxidizing agent in the manufacture of the well-known dye, magenta.


Sodium arsenite and arsenate are used in the cleansing operations of the calico-printer.

Sodium Pyroarsenate and Arsenate.

Fifth Synthetical Reaction.—Fuse two or three grains of common white arsenic (As_2O_3) with sodium nitrate (NaNO_3) and dried sodium carbonate (Na_2CO_3) in a porcelain crucible, and dissolve the mass in water; solution of sodium arsenate (Na_2HAsO_4) results.



The official proportions (B. P.) are 10 of arsenic to $8\frac{1}{2}$ of sodium nitrate and $5\frac{1}{2}$ of dried carbonate, each powdered, the whole well mixed, fused in a crucible at a red heat till effervescence ceases, and the liquid poured out on a slab. The product is sodium pyroarsenate ($\text{Na}_4\text{As}_2\text{O}_7$). Dissolved in water, crystallized, and dried, the salt has the formula $\text{Na}_2\text{HAsO}_4, 7\text{H}_2\text{O}$ (*Sodii Arsenas*, U. S. P.), the old arseniate of soda.




Heated to 300°F. , the crystals lose all water. A solution of 1 per cent. of the anhydrous salt (Na_2HAsO_4) in water forms the *Liquor Sodii Arsenatis*, *Solution of Sodium Arsenate*, U. S. P. It has about half the arsenical strength of *Liquor Arsenicalis*, B. P. The anhydrous salt is used because the crystallized is of somewhat uncertain composition. The fresh crystals are represented by the formula $\text{Na}_2\text{HAsO}_4, 12\text{H}_2\text{O}$ ($= 53.7$ per cent. of water); these soon effloresce and yield a stable salt having the formula $\text{Na}_2\text{HAsO}_4, 7\text{H}_2\text{O}$ ($= 40.4$ per cent. of water). To avoid the possible employment of a mixture of these bodies, the invariable anhydrous salt is officially used, constancy in the strength of a powerful preparation being thereby secured.

The student will find useful practice in verifying, by calculation, the above numbers representing the centesimal proportion of water in the two sodium arsenates. This will be easy if what has already been stated respecting a symbol representing a number as well as a name, and the remarks concerning molecular weight, be remembered.

The shape of each of the two varieties of sodium arsenate ($\text{Na}_2\text{HAsO}_4, 12\text{H}_2\text{O}$ and $\text{Na}_2\text{HAsO}_4, 7\text{H}_2\text{O}$) is identical with that of the corresponding sodium phosphate ($\text{Na}_2\text{HPO}_4, 12\text{H}_2\text{O}$ and $\text{Na}_2\text{HPO}_4, 7\text{H}_2\text{O}$); the structure of the molecule of the 12-arsenate is the same as that of the 12-phosphate, and the 7-arsenate as that of the 7-phosphate; the two former are isomorphous, the two latter are isomorphous. This is only one instance of the strong analogy of arsenium and its compounds with phosphorus and its corresponding compounds. The preparation and characters of the next substance, ferrous iron arsenate, will remind the learner of ferrous iron phosphate.

Ferrous Arsenate, or Arsenate of Iron.

Sixth Synthetical Reaction.—To a hot solution of sodium arsenate add a hot solution of ferrous sulphate and a little solution of sodium bicarbonate; a precipitate of ferrous arsenate occurs ($\text{Fe}_3, 2\text{AsO}_4$) (*Ferri Arsenias*, B. P.). On a larger scale, $15\frac{3}{4}$ parts of dried arsenate dissolved in 100 of hot water, and $20\frac{3}{4}$ of sulphate in 120 of hot water, with $4\frac{1}{2}$ of bicarbonate, may be employed. The precipitate should be collected on a calico filter, washed, squeezed, and dried at a low temperature (100°F.) over a water-bath, to avoid excessive oxidation.



The use of the sodium bicarbonate is to ensure the absence of free sulphuric acid in the solution. This acid is a solvent of ferrous arsenate. It is impossible to prevent its separation if only the ferrous sulphate (three molecular weights) and sodium arsenate (two molecular weights) be employed.

At the instant of precipitation ferrous arsenate is white, but rapidly becomes of a green or greenish-blue color, owing to absorption of oxygen and formation of a ferrosoferrie arsenate. When dry, it is a tasteless, amorphous, much-oxidized powder, soluble in acids.

The *Arsenum Hydrides* and *Sulphides* and the *Copper* and *Silver Arsenites* and *Arsenates* are mentioned in the following analytical paragraphs.

REACTIONS HAVING ANALYTICAL INTEREST (TESTS).

First Analytical Reaction.—Repeat the third synthetical reaction, operating on not very much more white arsenic than the bulk of a small pin's head, and using not charcoal alone, but the *black flux* already mentioned (p. 169) or a well-made and perfectly dry mixture of charcoal and potassium carbonate, the latter best obtained by heating potassium bicarbonate. The tube employed should be a narrow test-tube, or, better, a tube (easily made from glass tubing) having the form (Berzelius's) shown in Fig. 33.

Fig. 33.



The white arsenic and black flux are placed in the bulb of the tube, which is then heated in a flame; the arsenum condenses on the constricted portion of the tube. If now the bulb be carefully fused off in a flame, the arsenum may be chased up and down the narrower part of the tube until the air in the tube has reoxidized it to arsenous anhydride.


If the operation has been performed in a less delicate manner in an ordinary test-tube, cut or break off portions of the tube containing the sublimate of arsenum, put them into a test-tube and heat the bottom of the latter, holding it near horizontally, and partially covering the mouth with the finger or

thumb: the arsenum will absorb oxygen from the air in the tube, and the resulting arsenous anhydride (As_2O_3) be deposited on the cool part of the tube in brilliant, generally imperfect, octahedral crystals.

Microscopic Test.—Prove that the crystals are identical in form with those of common white arsenic by heating a grain or less of the latter in another test-tube, examining the two sublimate by a good lens or compound microscope.


The appearance of a sublimate of white arsenic is peculiar and quite characteristic. The primary form of each crystal is

Fig. 34.



A Sublimate of White Arsenic (magnified).

Fig. 34a.



A Perfect Octahedron.

an octahedron (*ὀκτῶν, okto*, eight; *ἑδρα, hedra*, side) (Fig. 34a), or, rarely, a tetrahedron, and in a sublimate a few perfect octahedra are generally present. Usually, however, the crystals are modifications of octahedra, such as are shown in Fig. 34, which is drawn from actual sublimate.

Second Analytical Reaction.—Place a thin piece of copper, about a quarter inch wide and half inch long, in a solution of white arsenic, acidified by hydrochloric acid, and boil (nitric acid must not be present, or the piece of metal will be dissolved); arsenum is deposited on the copper in a metallic condition. (*Memorandum.*—An equivalent proportion of copper goes into solution. The experiment forms an illustration of a class of chemical changes appropriately termed changes by *substitution*.) Pour off the supernatant liquid from the copper, wash the latter with water, dry the piece of metal by holding it in the clean fingers and passing through a flame, and finally place it at the bottom of a clean, dry, narrow test-tube or a Berzelius tube, and sublime as described in the last reaction, again noticing the form of the resulting crystals.

This is commonly known as Reinsch's test for arsenum, it having


been introduced by Reinsch in 1843. The tube may be reserved for subsequent comparison with an antimonial sublimate.

Note.—Copper itself frequently contains arsenum—a fact that may not, perhaps, much trouble an operator so long as he is performing experiments in practical chemistry merely for educational purposes, but when he engages in the analysis of bodies of unknown composition, he must assure himself that neither his apparatus nor materials already contain the element for which he is searching.

The detection of arsenum in metallic copper is best accomplished by distilling a mixture of a few grains of the sample with five or six times its weight of ferric hydrate or chloride (free from arsenum) and excess of hydrochloric acid. The arsenum is thus volatilized in the form of arsenum chloride, and may be condensed in water and detected by sulphuretted hydrogen (Sixth Analytical Reaction) or Reinsch's test. The ferric chloride solution is, if necessary, freed from any trace of arsenum by evaporating once or twice to dryness with excess of hydrochloric acid.

Third Analytical Reaction—The Hydrogen Test or "Marsh's" Test.—Generate hydrogen in the usual way from water by zinc and sulphuric acid, a bottle of about four or six ounces capacity being used, and a funnel-tube and short delivery-tube passing through the cork in the usual manner (see following figure). Dry the escaping hydrogen (except in rough experiments, when it is unnecessary) by adapting to the delivery-tube, by a pierced cork, a short piece of wider tubing filled with fragments of calcium chloride (*a*). To the opposite end of the drying-tube fit a piece of narrow tubing ten or twelve inches long, made of hard German glass, and having its aperture narrowed by drawing out in the flame of the blowpipe. When the hydro-

Fig 35.



The Hydrogen Test for Arsenum.

gen has been escaping for a sufficient number of minutes, and at such a rate as to warrant the operator in concluding that *all*

the air originally existing in the bottle has been expelled, set light to the jet, and then pour eight or ten drops of the aqueous solution of arsenic, or three or four drops of the acid or alkaline solution of white arsenic, previously prepared, into the funnel-tube, washing the liquid into the generating-bottle with a little water. The white arsenic is at once reduced to the state of arsenum, and the latter combines with some of the hydrogen to form arsenum hydride or arseniuretted hydrogen gas (AsH_3). Immediately hold a piece of earthenware or porcelain (the lid of a porcelain crucible (*b*) if at hand) in the hydrogen jet at the extremity of the delivery-tube; a brown spot of arsenum is deposited on the porcelain. Collect several of these spots, and retain them for future comparison with antimonial spots similarly obtained (p. 185).

The separation of arsenum in the flame is due to the decomposition of the arseniuretted hydrogen by the heat. The cool porcelain at once condenses the arsenum, and thus prevents its oxidation to white arsenic, which would otherwise take place at the outer edge of the flame.

Hold a small beaker (*c*) or wide test-tube over the flame for a few minutes; a white film of arsenic (As_2O_3) will be slowly deposited, and may be further examined in contrast with a similar antimonial film (p. 185).

During these experiments the effect produced by the arsenical vapors on the color of the hydrogen flame will have been noticed; they give it a dull, livid, bluish tint. This is characteristic.

Apply the flame of a gas-lamp to the middle of the hard glass delivery-tube (*d*); the arseniuretted hydrogen, as before, is decomposed by the heat, but the liberated arsenum immediately condenses in the cool part of the tube, beyond the flame, as a dark metallic mirror. The tube may be removed and kept for comparison with an antimonial deposit (p. 185).

Note 1.—Zinc, like copper, frequently itself contains arsenum. When a specimen free from arsenum is met with, it should be reserved for analytical experiments, or a quantity of guaranteed purity should be purchased of the dealers in such articles. Sulphuric acid is more easily obtained free from arsenic.


Note 2.—In delicate and important applications of Marsh's test magnesium may be substituted for zinc with safety, as arsenum has not yet been, and is not likely to be, found in magnesium. Magnesium in rods is convenient for this purpose, and may be obtained from most dealers in chemical substances. Both magnesium and zinc, if perfectly pure, react with acids extremely slowly; the addition of a very little platinum perchloride, however, at once promotes an abundant evolution of hydrogen. But platinum has a

tendency to hold back arsenum. According to Dyer, rod zinc has a similar tendency, while granulated zinc at once gives arseniuretted hydrogen.

Note 3.—Sulphuric acid, which is often used for drying gases, decomposes arseniuretted hydrogen. Calcium chloride is the appropriate desiccating agent for this gas.

Note 4.—The original apparatus proposed by Mr. Marsh, a pharmacist of Woolwich, England, in 1836, was a U-shaped tube, one limb of which was short and closed by a stopcock, so that the whole of a small quantity of arseniuretted hydrogen could be collected and be examined at leisure.

Fourth Analytical Reaction—Fleitmann's Test.—Generate hydrogen by heating in the test-tube, to near the boiling-point, a strong solution of caustic soda or potash and some pieces of zinc ($\text{Zn} + 2\text{NaHO} = \text{H}_2 + \text{Na}_2\text{ZnO}_2$, sodium zincate). Add a drop of arsenical solution. Now spread over the mouth of the tube a cap of filter-paper moistened with one drop of solution of silver nitrate. Again heat the tube, taking care that the liquid itself shall not spurt up on to the cap. A plug of cotton wool may even be placed in the mouth of the test-tube to prevent this spurting. The arsenic is reduced to arsenum, the latter uniting with the hydrogen, as in Marsh's test; and the arseniuretted hydrogen, passing up through the cap, reacts on the silver nitrate, and gives rise to an excellent test by causing the production of a purplish-black spot (of silver).



Note 1.—This reaction is particularly valuable, enabling the analyst to quickly distinguish arsenum in the presence of its sister element antimony, which, although it combines with the hydrogen evolved from dilute acid and zinc, does not combine with the hydrogen evolved from solution of alkali and zinc, and therefore does not give the effect just described.

Note 2.—Aluminium answers as well as zinc for Fleitmann's test (Gatchouse), or magnesium may be used, or, instead of zinc and alkali, weak sodium amalgam may be employed (Davy).

Fifth Analytical Reaction—Bettendorff's Test.—To a solution of stannous chloride in strong hydrochloric acid add a very small quantity of any arsenical solution. Arsenum then separates, especially on the application of heat, giving the mixture a yellowish and then brownish hue or grayish-brown turbidity, or even a sediment of gray-brown flocks, according to the amount present. Much water prevents the reaction; its presence, therefore, must be avoided as far as possible; indeed, a liquid saturated by hydrochloric acid gas gives best results. Arsenic in sulphuric or hydrochloric acid or in tartar emetic,

etc. may be detected by this method. Nitrates, such as bismuth subnitrate, must first be heated with sulphuric acid to remove the nitric radical before applying this reduction-test for arsenum. The stannous is converted into stannic salt during the reaction.

Distinction between Arsenous and Arsenic Combinations.—The above tests are those for arsenum, whether existing in the arsenous or arsenic condition, though the element is not so easily attacked when it is in the latter as when it is in the former state. Of the following reactions, that with silver nitrate at once distinguishes arsenous acid and other arsenites from arsenic acid and other arsenates.

Mem.—The exact nature of all these analytical reactions will be more fully evident if the student will perseveringly trace them out by diagram or equations.

Sixth Analytical Reaction.—Through an acidified solution of arsenic pass sulphuretted hydrogen; a yellow precipitate (arsenum sulphide or arsenous sulphide, As_2S_3) quickly falls. Add an alkaline hydrate or sulphhydrate to a portion of the precipitate; it readily dissolves. The precipitate consequently would not be obtained on passing sulphuretted hydrogen through an alkaline solution of arsenic. To another portion of the precipitate, well drained, add *strong* hydrochloric acid; it is insoluble, unlike antimony sulphide. (Neither sulphide is soluble in the *weak* acid.)

Note 1.—Cadmium also affords a yellow sulphide in an acid solution by action of sulphuretted hydrogen, but this sulphide is insoluble in alkaline liquids. Under certain circumstances tin, too, yields a yellow sulphide, but tin is otherwise easily distinguished. (*Vide* "Tin" in Index.)


Note 2.—A trace of arsenum sulphide is sometimes met with in sulphur (distilled from arsenical pyrites). It may be detected by digesting the sulphur in solution of ammonia, filtering, and evaporating to dryness; a yellow residue of arsenum sulphide is obtained if that substance be present.

Seventh Analytical Reaction.—Through an acidified solution of arsenic acid, or any other arsenate, pass a rapid current of sulphuretted hydrogen; a yellow precipitate (arsenic sulphide, As_2S_5) slowly falls. Brauner and Tornieck state that by a slow current the arsenic acid is gradually reduced to the arsenous, and a yellow precipitate of arsenous sulphide and sulphur ($\text{As}_2\text{S}_3 + \text{S}_2$) slowly falls. The precipitate is soluble in alkaline hydrates and sulphhydrates. This reaction is more rapid if the solution be warmed.

Chemical Analogy of Sulphur and Oxygen.—The potassium arsenite and sulph-arsenite, arsenate and sulph-arsenate, have the composition represented by the following formulæ :



and the corresponding ammonium and sodium salts have a similar composition :



Eighth Analytical Reaction.—To an aqueous solution of arsenic add two or three drops of solution of copper sulphate, and then cautiously add diluted solution of ammonia, drop by drop, until a green precipitate is obtained. The production of this precipitate is characteristic of arsenum. To a portion of the mixture add an acid; the precipitate dissolves. To another portion add alkali; the precipitate dissolves. These two experiments show the advantage of testing a suspected arsenical solution by litmus-paper before applying this reaction—if acid, cautiously adding alkali; if alkaline, adding acid till neutrality is obtained. Or a special copper reagent may be used. (See a note to the Eleventh Analytical Reaction.)

The precipitate is copper arsenite ($\text{Cu}''\text{HAsO}_3$) or *Scheele's Green*. More or less pure or mixed with copper acetate or, occasionally, carbonate, it is used as a pigment under many names, such as Brunswick Green and Schweinfurth Green, by painters and others.

Ninth Analytical Reaction.—Apply the test just described to a solution of arsenic acid or other arsenate; a somewhat similar colored precipitate (copper arsenate) is obtained.

Tenth Analytical Reaction.—Repeat the eighth reaction, substituting silver nitrate for copper sulphate: in this case a *yellow* precipitate (silver arsenite, Ag_3AsO_3) falls, also soluble in acids and alkalies.

Eleventh Analytical Reaction.—Apply the silver test to a solution of arsenic acid or other arsenate; a *chocolate-colored* precipitate (silver arsenate, Ag_3AsO_4) falls.

This reaction may be utilized for the detection of arsenic when occurring in ores and other substances, as ordered by the U. S. Pharmacopœia in the case of *Antimonii Sulphidum Purificatum* :

“If 2 grms. of the sulphide be mixed and cautiously ignited, in a porcelain crucible, with 8 grms. of pure sodium nitrate, and, after cooling, the fused mass be boiled with 25 cc. of water, there will remain a residue which should be white or nearly so, and not yellowish nor brownish (absence of other metallic sulphides). On boiling the filtrate separated from the last-mentioned residue with a

slight excess of nitric acid until no more nitrous vapors are evolved, then dissolving it in 0.1 grm. of silver nitrate, filtering again if necessary, and cautiously pouring a few drops of ammonia-water on top, not more than a white cloud, but no red or reddish precipitate, should appear at the line of contact of the two liquids (absence of more than about 0.1 per cent. of arsenic)."

Copper and Silver Reagents for Arsenum.—The last four reactions may be performed with increased delicacy and certainty of result if the copper and silver reagents be previously prepared in the following manner: To solution of pure copper sulphate (about 1 part in 20 of water) add ammonia until the blue precipitate at first formed is nearly, but not quite, redissolved; filter and preserve the liquid as an arsenum reagent, labelling it *Solution of Ammonio-Sulphate of Copper* (B. P.). Treat solution of silver nitrate (about 1 part in 40) in the same way, and label it *Solution of Ammonio-Nitrate of Silver* (B. P.). The composition of these two salts will be referred to subsequently.

Arsenous and Arsenic Compounds.—While many reagents may be used for the detection of arsenum, only silver nitrate, as already stated, will readily indicate in which state the arsenum exists; for the two sulphides and the two copper precipitates, though differing in composition, resemble each other in appearance, whereas the two silver precipitates differ in color as well as in composition.

Soluble arsenates give insoluble arsenates with solutions of salts of barium, calcium, zinc, and other metals.

In a group-testing, arsenum, if existing as arsenic acid or other arsenate, is not readily affected by such tests as sulphuretted hydrogen or even by nascent hydrogen. Hence, if its presence in that state is suspected, the liquid under analysis should be warmed with a little sulphurous acid (*vide* p. 169), and then tested with sulphuretted hydrogen.

Antidote.—In cases of poisoning by arsenic or arsenical preparations the most effective antidote is recently precipitated moist ferric hydrate (*Ferri Oxidum Hydratum*, U. S. P.), administered as soon as possible. It is perhaps best administered in the form of a mixture of, or solution of, ferric sulphate (*Liquor Ferri Tersulphas*, U. S. P.), or ferric chloride (*Liquor* or *Tinctura*) with sodium carbonate—2 to 3 ounces of the former to about 1 ounce of the crystals of the latter. Instead of the sodium carbonate, about $\frac{1}{4}$ ounce of calcined magnesium may be used. (See *Ferri Oxidum Hydratum cum Magnesia*, U. S. P., page 152.) These quantities will render at least 10 grains of arsenic insoluble. Emetics should also be given, and the stomach-pump, or a common india-rubber tube worked as a siphon (p. 116), be applied as quickly as possible.

The above statements regarding the antidote for arsenic may be verified by mixing the various substances together, filtering, and proving the absence of arsenum in the filtrate by applying some of the foregoing tests.

Mode of Action of the Antidote.—The action of the sodium carbonate or the magnesia is to precipitate ferric hydrate ($\text{Fe}_2\text{O}_3 \cdot 3\text{H}_2\text{O}$)—sodium chloride (NaCl) or magnesium chloride (MgCl_2) being formed, which are harmless, if not beneficial, under the circumstances. The reaction between the ferric hydrate and the arsenic results in the formation of insoluble ferrous arsenate. (See also p. 152.)



The so-called solution of dialyzed iron (see Index) is also, as might be expected from its composition, an antidote to arsenic. It should be administered with a little sodium or potassium bicarbonate, or with magnesia, or with any other salt which serves to neutralize any acid that may be present.

QUESTIONS AND EXERCISES.

What is the formula of a molecule of arsenum?—In what form does arsenum occur in nature?—Describe the characters of white arsenic. Name the official preparations of arsenum.—What compound of arsenum is contained in *Liquor Potassii Arsenitis*, U. S. P., and in *Liquor Acidi Arsenosi*, U. S. P.?—By what method may arsenic be reduced to arsenum?—Give the formulæ of arsenous and arsenic acids and anhydrides.—Explain, by diagrams, the reactions which occur in converting arsenic into sodium arsenate by the official process.—Why is anhydrous instead of crystallized sodium arsenate employed in the preparation of *Liquor Sodii Arsenatis*, U. S. P.?—In the preparation of ferrous arsenate from ferrous sulphate and sodium arsenate why is sodium bicarbonate included? Describe the manipulations necessary to obtain arsenic in its characteristic crystalline form.—How is Reinsch's test for arsenum applied? and under what circumstances may its indications be fallacious?—Give the details of Marsh's test for arsenum, and the precautions to be observed.—Explain the reactions by diagrams.—What peculiar value has Fleitmann's test for arsenum?—Describe the conditions under which sulphuretted hydrogen becomes a trustworthy test for arsenum.—How may a trace of arsenum sulphide be detected in sulphur?—How are salts of copper and silver applied as reagents for the detection of arsenum?—How are arsenites distinguished from arsenates?—Mention the best antidote in case of poisoning by arsenic; explain the process by which it may be most quickly prepared, and describe its action.—Do you know of any other antidote to arsenic? If so, describe the mode of administration.

ANTIMONY.

Symbol, Sb (Stibium). Atomic weight, 120.

Source and Uses.—Antimony occurs in nature chiefly as sulphide, Sb_2S_3 . The crude or black antimony of pharmacy is this native sul-

phide freed from impurities by fusion: it has a striated, crystalline, lustrous fracture; subsequently powdered, and if it contains any soluble arsenum salt, the latter removed by digestion in solution of ammonia, it forms the grayish-black crystalline *Antimonii Sulphidum*, Sulphide of Antimony, or *Antimony Sulphide*, U. S. P. The metal is obtained from the sulphide by roasting, the resulting oxide being reduced with charcoal and sodium carbonate. The resulting scoria is known as *crocus of antimony* or *glass of antimony*. Metallic antimony is an important constituent of *type-metal*, *Britannia metal* (tea- and coffee-pots, spoons, etc.), and the best varieties of *pewter*. The old *pocula emetica*, or everlasting emetic cups, were made of antimony; wine kept in them for a day or two was said to have acquired emetic quality. The metal is not used in making official antimonial preparations, the sulphide alone being, directly or indirectly, employed for this purpose.

Antimony has very close chemical analogies with arsenum. Its atom, in the common salts, exerts trivalent activity (e. g. SbCl_3), but sometimes it is quinquivalent (e. g. SbCl_5).

Antimony, like arsenum, unites with iodine to form a tri-iodide (SbI_3). A bromide (SbBr_3) also is known.

REACTIONS HAVING SYNTHETICAL INTEREST.

Antimonious Chloride, or Chloride of Antimony.

First Synthetical Reaction.—Boil $\frac{1}{2}$ an ounce or less of antimony sulphide with four or five times its weight of hydrochloric acid in a dish placed in a fume-chamber or in the open air; sulphuretted hydrogen is evolved and solution of antimony chloride, SbCl_3 , is obtained.



This solution, cleared by subsidence, is what is commonly known as *Butter of Antimony* (*Liquor Antimonii Chloridi*, B. P.). If pure sulphide has been used in its preparation, the liquid is nearly colorless; but much of that met with in veterinary pharmacy is simply a by-product in the generation of sulphuretted hydrogen from native ferruginous antimony sulphide and hydrochloric acid, and is more or less brown from the presence of iron chloride. It not infrequently darkens in color on keeping: this is due to absorption of oxygen from the air and conversion of light-colored ferrous into dark-brown ferric chloride or oxychloride.


True Butter of Antimony (SbCl_3) is obtained on evaporating the above solution to a low bulk and distilling the residue. The butter condenses as a white crystalline semi-transparent mass in the neck of the retort; at the close of the operation it may be easily melted and run down into a bottle, which should be subsequently well stoppered.

Antimony Pentachloride (SbCl_5), or *Antimonic Chloride*, is a fuming liquid, obtained on passing chlorine over the lower chloride.

Antimonious Oxychloride, or Oxychloride of Antimony.

Second Synthetical Reaction.—Boil the solution of antimony chloride produced in the last reaction, and pour it into several ounces of water; a white precipitate of antimony oxychloride ($2\text{SbCl}_3, 5\text{Sb}_2\text{O}_3$) falls, some antimony chloride remaining in the supernatant acid liquid.

This precipitate is the old *pulvis Algarothi*, *pulvis angelicus*, or *mercurius vitæ*. It varies somewhat in composition, according to the amount of water with which the chloride may be mixed, but on standing under water gradually becomes crystalline, and has the composition above given.

**Antimonious Oxide, or Oxide of Antimony.**

Well wash the precipitate with water by decantation (*vide* p. 110), and add solution of sodium carbonate; the chloride remaining with the oxide is thus decomposed, and antimony oxide (Sb_2O_3) alone remains. This is *Antimoni Oxidum*, U. S. P. It is of a light buff or grayish-white color, or quite white if absolutely free from iron, insoluble in water, soluble in hydrochloric acid, fusible at a low red heat. The moist antimony oxide may be well washed and employed for the next reaction. Or it may be dried over a water-bath, for at temperatures above 100°C . oxygen is absorbed and other antimony oxides formed. The presence of the latter is detected on boiling the powder in solution of acid potassium tartrate, in which antimony oxide (Sb_2O_3 ; Sb_4O_6 ?) is soluble, but *antimonic anhydride* (Sb_2O_5) and the double oxide, or so-called *antimonious anhydride* (Sb_4O_8), is insoluble.




The Higher Antimony Oxide (Sb_2O_5), termed *antimonic oxide* or *anhydride*, corresponding with arsenic anhydride, is obtained on decomposing the pentachloride by water or on boiling metallic antimony with nitric acid. The variety obtained from the chloride differs in saturating power from that obtained from the metal, and is termed *metantimonic acid* ($\mu\epsilon\tau\acute{\alpha}$, *meta*, beyond).

Tartar Emetic.

Third Synthetical Reaction.—Mix the moist antimony oxide obtained in the previous reaction with about an equal quantity

of cream of tartar (6 of the latter to 5 of the dry oxide) and sufficient water to form a paste; set aside for a day to facilitate complete combination; boil the product with water and filter; the resulting liquid contains antimony and potassium tartrate ($\text{KSbOC}_4\text{H}_4\text{O}_6$), antimony potassio-tartrate, or tartarated antimony, or tartar emetic (emetic, from $\xi\mu\acute{\epsilon}\omega$, *emcō*, I vomit; tartar, from *Tάρταρος*, *tartaros*; see Index).



On evaporation, the salt is obtained in colorless, transparent triangular-faced crystals containing one molecular proportion of water to every two of anhydrous salt, forming the *Tartrate of Antimony and Potassium*, *Antimonii et Potassii Tartras*, U. S. P., or *Antimony and Potassium Tartrate* ($\text{KSbOC}_4\text{H}_4\text{O}_6$)₂.H₂O.

The formula for tartar emetic is apparently inconsistent with the general formula for tartrates ($\text{R}'\text{R}''\text{C}_4\text{H}_4\text{O}_6$); this will be subsequently fully explained in connection with Tartaric Acid. The salt appears to be an oxytartrate, $\text{K}'\text{Sb}'''\text{O}''(\text{C}_4\text{H}_4\text{O}_6)''$.

Tartar emetic is soluble in water, and slightly so in proof spirit. Dissolved in sherry wine, it forms the official *Vinum Antimonii*, U. S. P. It may be externally applied as an ointment, *Unguentum Antimonii Tartarati*, B. P.




"Sulphurated Antimony" and other Antimony Oxy-sulphides.

Fourth Synthetical Reaction.—Boil a gramme or two of antimony sulphide with solution of soda in a test-tube, and filter (or larger quantities in larger vessels, 1 part of sulphide to 12 of soda and 30 of water, for two hours, frequently stirring, and occasionally replacing water lost by evaporation). Into the filtrate, before cool, stir diluted sulphuric acid until the liquid is slightly acid to test-paper; a reddish-brown precipitate, sulphurated antimony, is formed, *Antimonium Sulphuratum*, U. S. P.; filter, wash, and dry over a water-bath. It is a mixture of antimony sulphide (Sb_2S_3) with a little oxide (Sb_2O_3). The oxide results from the double decomposition of antimony sulphide and caustic soda.





This is one of the many varieties of *mineral kermes*, so called from their similarity in color to the *insect kermes*. *Kermes* is the name, now obsolete, of the *Coccus Ilidis*, a sort of cochineal insect, full of reddish juice, and used for dyeing from the earliest times. The term *mineral kermes* was apparently applied originally to the amorphous or precipitated orange antimony sulphide (Sb_2S_3). It afterward included any oxy-sulphide and pentasulphide. The color of the brownish-red variety is affected by the temperature as well as

state of dilution of the alkaline liquid when the acid is added. When this alkaline liquid is boiled, especially if long exposed to air, oxygen is absorbed by some of the antimony, whose sulphur, uniting with the trisulphide, forms a portion of the lighter yellow pentasulphide. The process of the British Pharmacopoeia includes the use of as much sulphur as black antimony sulphide; the product is of an orange-red color and contains much antimony pentasulphide.

Explanation of Processes.—Antimony sulphides and oxides, like those of arsenium, react with the sulphides, hydrates, and oxides of certain metals to form salts of greater or less degree of solubility. Thus, sodium antimonite (Na_3SbO_3) is formed and remains in solution, and sodium sulphantimonite (Na_3SbS_3) is formed, and is deposited in brilliant yellow tetrahedral crystals when a hot alkaline solution of antimony sulphide is set aside to cool. Sulphur being present, the slightly soluble sodium antimoniate (Na_3SbO_4) and sodium sulphantimoniate (Na_3SbS_4) are produced.



In the hot solutions of these sulphur salts and oxygen salts, antimony sulphides and oxides are soluble, and are reprecipitated in an indefinite state of combination, partially on cooling or wholly on the addition of acid. The acid also decomposes the oxyalts with precipitation of oxides, and the sulphur salts with precipitation of sulphides of antimony. The acid should be added to the liquids before much oxysulphide has deposited (that is, before the solution is cool) if uniformity of product is desired.



The oxides and sulphides indicated in these equations, together with excess of antimony sulphide originally dissolved by the alkaline liquid, are all precipitated when the acid is added, and form

the varieties of *kermes*. *Kermes* may be formed by fusion as well as by aqueous solution of the components. The student is strongly recommended carefully to study the foregoing paragraphs; for, although neither the official nor any other variety of *kermes* is itself of much importance in modern practical pharmacy, a thoughtful consideration of their chemistry will, by revealing chemical actions and analogies that are general, aid in sowing the germs of chemical principles in the mind.

The previous four synthetical reactions illustrate the official process for the respective substances. In pharmacy the solution of antimony chloride is only used in the preparation of oxide; the oxide, besides its use in the preparation of tartar emetic, is mixed with twice its weight of calcium phosphate (purified bone-earth) to form *Pulvis Antimonialis*, U. S. P., or "James's Powder."

Antimony Sulphides and *Hydrides* are incidentally mentioned in the following analytical paragraphs.

REACTIONS HAVING ANALYTICAL INTEREST (TESTS).

First Analytical Reaction.—Through an acidified antimonial solution pass sulphuretted hydrogen; an orange precipitate (amorphous antimony sulphide) falls. It has the same composition as the crystalline black sulphide (Sb_2S_3), into which, indeed, when dried, it is quickly converted by heat. Like arsenum sulphide, it is soluble in alkaline solutions. Collect a portion on a filter, and, when well drained, add *strong* hydrochloric acid; it dissolves—unlike arsenum sulphide.

A higher antimony sulphide (Sb_2S_5), corresponding to the higher arsenum sulphate, exists. It is formed on passing sulphuretted hydrogen through an acidified solution of the higher chloride (SbCl_5), or less pure, on boiling black antimony sulphide and sulphur with an alkali and decomposing the resulting filtered liquid by an acid.

Note.—The arsenous and antimonious compounds are those chiefly employed in medicine; sodium and iron arsenates are, however, sometimes employed. The arsenates, and rarely, an antimoniate, are useful in analysis, and the antimonie chloride in chemical research. The higher compounds of both elements are noticed here chiefly to draw attention to the close analogy existing between arsenum and stibium—an analogy carried out in the numerous other compounds of these elements.

Second Analytical Reaction.—Dilute two or three drops of the solution of antimony chloride with water; a white precipitate (oxychloride) occurs, the formation of which has been

explained under the similar synthetical reaction. The occurrence of a precipitate under the circumstances distinguishes antimony from arsenum, but is a reaction that cannot be fully relied upon in analysis, because requiring the presence of too much material and the observance of too many conditions. Add a sufficient quantity of hydrochloric acid to dissolve the precipitate, and boil a piece of copper in the solution, as directed in the corresponding test for arsenum (*vide* p. 172); antimony is deposited on the copper. Wash, dry, and heat the copper in a test-tube, as before; the antimony, like the arsenum, is volatilized off the copper, and condenses on the side of the tube as white oxide; but the sublimate, from its low degree of volatility, condenses close to the copper; moreover, it is destitute of crystalline character.

Shake out the copper and boil water in the tube for several minutes. Do the same with the arsenical sublimate similarly obtained. The deposit of arsenic slowly dissolves, and may be recognized in the solution by silver ammonio-nitrate; the antimonial sublimate is insoluble.

Third Analytical Reaction.—Perform the experiments described under Marsh's test for arsenum (p. 173), carefully observing all the details there mentioned, but using a few drops of solution of antimony chloride or tartar emetic instead of the arsenical solution. Antimoniuretted hydrogen, or antimony hydride (SbH_3), is formed and decomposed in the same way as arseniuretted hydrogen.

To one of the arsenum spots on the porcelain lid (p. 173) add a drop of solution of "chloride of lime" (bleaching-powder); it quickly dissolves. Do the same with an antimony spot; it is unaffected. Heat more quickly causes the volatilization of an arsenum than an antimony spot; ammonium sulphhydrate more readily dissolves the antimony than the arsenum.

Boil water for several minutes in the beaker or wide test-tube containing the arsenous sublimate (p. 173); it slowly dissolves, and may be recognized in the solution by the yellow precipitate given on the addition of solution of silver ammonio-nitrate. The antimonial sublimate, similarly treated, does not dissolve.

Pass a slow current of sulphuretted hydrogen through the delivery-tube removed from the hydrogen apparatus (p. 173), and when the air may be considered to have been expelled from the tube, gently heat that portion containing the deposit of arsenum: the latter will be converted into a *yellow* sublimate of arsenum sulphide. Remove the tube from the sul-

phuretted-hydrogen apparatus, and repeat the experiment with a similar antimony deposit: it is converted into *orange* antimony sulphide, which, moreover, owing to inferior volatility, condenses nearer to the flame than arsenum sulphide.

Pass dry hydrochloric acid gas through the two delivery-tubes. This is accomplished by adapting first one tube and then the other by a cork to a test-tube containing a few lumps of common salt on which a little sulphuric acid is poured during the momentary removal of the cork. The antimony sulphide dissolves and disappears; the arsenum sulphide is unaffected.

Thorough perception of the chemistry of arsenum and antimony will be obtained on constructing equations or diagrams descriptive of each of the foregoing reactions.

Antidote.—The introduction of poisonous doses of antimonials into the stomach is fortunately quickly followed by vomiting. If vomiting has not occurred or apparently to an insufficient extent, any form of tannic acid may be administered (infusion of tea, nutgalls, cinchona, oak-bark, or other astringent solutions or tinctures), an insoluble antimony tannate being formed, and absorption of the poison consequently somewhat retarded. The stomach-pump or stomach-siphon must be applied as quickly as possible.

Recently precipitated moist ferric hydrate is also, according to T. and H. Smith, a perfect absorbent of antimony from its solutions, the chemical action being probably, they say, similar to that which takes place between ferric hydrate and arsenous anhydride. It may be given in the form of a mixture of ferric chloride with either sodium carbonate or other soluble carbonate or bicarbonate, or with magnesia.

These statements may be verified by mixing together the various substances, filtering, and testing the filtrate for antimony in the usual manner.

DIRECTIONS FOR APPLYING THE FOREGOING REACTIONS TO THE ANALYSIS OF AN AQUEOUS SOLUTION OF A SALT OF ONE OF THE ELEMENTS ARSENUM AND ANTIMONY.

Acidify the liquid with hydrochloric acid, and pass through it sulphuretted hydrogen:

A *yellow* precipitate indicates arsenum;

An *orange* precipitate indicates antimony.

The result may be confirmed by the application of other tests.

DIRECTIONS FOR APPLYING THE FOREGOING REACTIONS TO THE ANALYSIS OF AN AQUEOUS SOLUTION OF SALTS OF BOTH ARSENUM AND ANTIMONY.

Acidify a small portion of the liquid with hydrochloric acid, and pass through it sulphuretted hydrogen.

Note 1.—If the precipitate by sulphuretted hydrogen is unmistakably orange, antimony may be put down as present, and arsenum only further sought by the application of Fleitmann's test to the solution of the sulphides in aqua regia* freed from sulphur by boiling, or, better, to the original solution.

Note 2.—Antimony sulphide is far less readily soluble than arsenum sulphide in solution of ammonium carbonate. But this fact possesses limited analytical value; for the color of the sulphides is already sufficient to distinguish the one from the other when they are unmixed; and when mixed, much antimony sulphide will prevent a little arsenum sulphide from being dissolved by the alkaline carbonate, while much arsenum sulphide will carry a little antimony sulphide into the solution. When the proportions are, apparently, from the color of the precipitate, less wide, solution of ammonium carbonate will be found useful in roughly separating the one sulphide from the other. On filtering and neutralizing the alkaline solution by an acid the yellow arsenum sulphide is reprecipitated. The orange antimony sulphide will remain on the filter.

Note 3.—Solution of potassium bisulphite is said by Wöhler to be a good reagent for separating the sulphides of arsenum and antimony, the former being soluble, the latter insoluble, in the liquid.


Note 4.—Another reagent for separating the sulphides of arsenum and antimony is *strong* hydrochloric acid. As little water as possible must be present. On boiling, antimony sulphide dissolves, while arsenum sulphide remains insoluble. The liquid, slightly diluted, filtered, more water added, and sulphuretted hydrogen again transmitted, gives orange antimony sulphide. The process should previously be tried on the precipitated mixed sulphides. The presence of arsenum may be confirmed by the application of Fleitmann's test to the original solution.

Note 5.—If the precipitate by sulphuretted hydrogen is unmistakably yellow, arsenum may be put down as present, and any antimony be detected by the previous or one of the following two processes. These two processes are rather long, and require much care in their performance, but are useful, because a small quantity of antimony in much arsenum, or *vice versâ*, may be detected by their means.

First Process.—Generate hydrogen, and pass it through a

* *Aqua Regia* is a mixture of 4 parts hydrochloric and 3 parts nitric acid. It was so called from its property of dissolving gold, the "king" of metals. Diluted with rather more than four times its bulk of water, it forms the *Acidum Nitro-hydrochloricum Dilutum*, B. P.

small wash-bottle containing solution of lead acetate, to free the gas from any trace of sulphuretted hydrogen it may possess, and then through a dilute solution of silver nitrate contained in a test-tube. When the apparatus is in good working order, pour into the generating-bottle the solution to be examined, adding it gradually to prevent violent action. After the gas has been passing for five or ten minutes examine the contents of the silver nitrate tube; arsenum, if present, will be found in the solution in the state of arsenous acid—



while antimony, if present, will be found in the black precipitate that has fallen, according to the following equation:



The arsenous radical may be detected in the clear, filtered supernatant liquid, which still contains much silver nitrate, by cautiously neutralizing with a very dilute solution of ammonia, or by adding a few drops of solution of silver ammonio-nitrate, yellow silver arsenite being produced. The antimony may be detected by washing the black precipitate, boiling it in an open dish with solution of tartaric acid, acidulating with hydrochloric acid, filtering and passing sulphuretted hydrogen through the solution, the orange sulphide being precipitated (Hofmann).

Second Process.—Obtain the metallic deposit in the middle of the delivery-tube, as already described under Marsh's test. Act on the deposit by sulphuretted hydrogen gas, and then by hydrochloric acid gas, as detailed in the third analytical reaction of antimony (p. 186). If both arsenum and antimony are present, the deposit, after the action of sulphuretted hydrogen, will be found to be of two colors, the yellow arsenum sulphide being usually farther removed from the heated portion of the tube than the orange antimony sulphide. Moreover, subsequent action of hydrochloric acid gas causes disappearance of the antimonial deposit, which is converted into antimony chloride and carried off in the stream of gas.

The chief objection to this process is the liability of the operator mistaking sulphur, deposited from the sulphuretted hydrogen gas by heat, for arsenum sulphide. But the presence or absence of arsenum is easily confirmed by applying Fleitmann's test to the original solution, while the process is most useful for the detection of a small quantity of a salt of antimony when mixed with much arsenical matter. On the whole, Hofmann's method is to be recommended.

The laboratory student may now proceed to the analysis of aqueous solutions of salts of any of the metallic elements hitherto considered. The method followed may be that for the separation of the previous three groups, sulphuretted hydrogen being first passed through the solution to throw out arsenum and antimony. The whole scheme of analysis is given in the accompanying table, p. 190. Three or four solutions should be examined before proceeding to the next and last group of metals.

Learners who have no opportunity of working at practical analysis will gain much knowledge by endeavoring, not to remember, but to understand, these methods of separating elements from each other in a solution containing several compounds.

QUESTIONS AND EXERCISES.

What is the composition and source of "*black antimony*"?—In what alloys is metallic antimony a characteristic ingredient?—What is the quantivalence of antimony as far as indicated by the formulæ of the official preparations?—By a diagram show how "*butter of antimony*" is prepared.—Write out equations or diagrams expressive of the reactions which occur in converting antimony chloride into oxide.—What is the formula of tartar emetic?—Explain the preparation of antimony oxysulphide (*Antimonium Sulphuratum*, U. S. P.) by aid of diagrams.—Give a comparative statement of the tests for arsenum and antimony.—How is antimony detected in the presence of arsenum?—How may arsenum and iron be distinguished analytically?—Describe a method by which antimony, magnesium, and iron may be separated from each other.—Draw out a chart for the analysis of an aqueous liquid containing salts of arsenum, zinc, calcium, and ammonium.

COPPER, MERCURY, LEAD, SILVER.

These metals, like arsenum and antimony, are precipitated from acidified solutions by sulphuretted hydrogen in the form of sulphides, but the sulphides, unlike those of arsenum and antimony, are soluble in alkalies. The atom of copper is usually bivalent, Cu'' ; mercury bivalent in the mercuric salts, Hg'' , and univalent in the mercurous salts, Hg' ; lead sometimes quadrivalent, Pb'''' , but generally exerting only bivalent activity, Pb'' ; and silver univalent, Ag' .

COPPER.

Symbol, Cu. Atomic weight, 63.3.

Source—The commonest ore of this metal is *copper pyrites*, a double copper and iron sulphide raised in Cornwall; Australia and Russia supply *malachite*; a mixed carbonate and hydrate; much ore is also imported from Spain and from South America. It is smelted in enormous quantities at Swansea, South Wales, a locality peculiarly fitted for the operation on account of its proximity to the coal-fields and its position as a seaport, these united advantages ensuring cheap fuel and freightage to the different metallurgical estab-

TABLE OF SHORT DIRECTIONS FOR THE ANALYSIS OF AN AQUEOUS SOLUTION OF SALTS OF **ANY** OR **ALL** OF THE METALLIC ELEMENTS HITHERTO CONSIDERED.
Acidify with HCl, and pass H_2S through the solution; filter.

Precipitate As Sb. Wash, separate by ammonium carbonate, or by strong hydrochloric acid; or dissolve in a few drops of hydrochloric and nitric acids, and examine by Hofmann's method (p. 188).	Precipitate Fe. Examine orig. solution for ferrous or ferric state.		Filtrate Al Zn. Neut. by HCl, add NH_4HO , stir, filter.		Filtrate Ba Ca Mg K Na NH_4 . Add $(\text{NH}_4)_2\text{CO}_3$, boil, filter.					
	Ppt. Al.	Sol. Zn.			Precipitate Ba.	Filtrate Ca.	Precipitate Ba Ca. Dissolve in $\text{HC}_2\text{H}_3\text{O}_2$, add K_2CrO_4 , filter.	Filtrate Mg K Na NH_4 . Add $(\text{NH}_4)_2\text{HPO}_4$, shake, filter.	Precipitate Mg.	Filtrate NH_4 Na K. (page 165).
			Precipitate† Fe Al Zn. Wash, dissolve in HCl, boil with HNO_3 (p. 163), add KHO in excess, † stir, filter.		Filtrate Fe Al Zn Ba Ca Mg K Na NH_4 . Add NH_4HO ; * NH_4HS ; stir, filter.					

* It is not usually necessary here to add NH₄Cl to keep Mg in solution, the HCl with some of the NH₄HO commonly giving sufficient NH₄Cl for this purpose.

† Much time may sometimes be saved by carefully remembering the color of the various hydrates and sulphides precipitated. Thus if the NH₄HS precipitate is white, iron cannot be present, and NH₄HO, for Al and Zn, may be at once added to the hydrochloric solution of the precipitate. The group-tests in this Table are H₂S, NH₄HS, and (NH₄)₂CO₃.

‡ The pupil should vary the process sometimes, as described in the foot-notes on pages 163 and 165. The treatment of other groups should occasionally be varied. By remembering the reactions on which analytical processes are based a pupil may often devise modified processes, and thus frequently save time, avoid working in one groove only, and acquire broader chemical views.

ishments. By Hollway's economical method of smelting copper pyrites and other sulphides, after the sulphide is once melted air is driven, not over, as usual, but through the mass; the combustion of the sulphur then becomes self-supporting, and is greatly accelerated.

Alchemy.—The alchemists termed this metal *Venus*, perhaps on account of the beauty of its lustre, and gave it her symbol ♀, a compound hieroglyphic also indicating a mixture of gold, ☉, and a certain hypothetical substance called acrimony, ✚, the corrosive nature of which was symbolized by the points of a Maltese cross. To this day the blue show-bottle in the shop-window of the pharmacist is occasionally ornamented by such a symbol, indicating, possibly, that the blue liquid in the vessel is a preparation of copper.


Coinage.—The material of the British "copper" coinage is now a bronze mixture, composed, in 100 parts by weight, of 95 copper, 4 tin, and 1 zinc, the same as in the copper coinage of France. The penny is coined at the rate of 48 pence in 1 pound avoirdupois of 7000 grains, or 453.6 grammes; the halfpenny at 80 in the pound avoirdupois, and the farthing at 160. These British bronze coins or tokens are a legal tender in payments to the amount of 1s.

Metallic Copper, U. S. P. (*Cuprum*, B. P.) is in the form of fine wire, about No. 25 of the Birmingham wire gauge, or 0.02 of an inch in diameter, and is used in preparing *Spiritus Ætheris Nitrosi*, B. P., or in the form of "pure metallic copper, thin and bright" (*Copper-foil*, B. P.).

Quantivalence.—Copper forms two classes of salts; in one the atom is bivalent (Cu''), in the other it exerts univalent activity (Cu'). The former are of primary importance to the student, the latter being for the most part unstable and wanting in technical interest. Their compounds are distinguished as *cupric* and *cuprous*. Cuprous iodide (Cu_2I_2) will be subsequently referred to as a convenient form in which to remove iodine from solution, while the formation of cuprous oxide (Cu_2O) under given circumstances will come under notice as an indicator of the presence of sugar in a liquid.

REACTIONS HAVING SYNTHETICAL INTEREST.

The processes for the following salts include the only synthetical reactions having any medical or pharmaceutical interest: (1) cupric oxide, black copper oxide, prepared by heating fragments of copper to low redness on a piece of earthenware in an open fire; (2) cupric sulphate, the common copper sulphate, prepared by boiling black oxide and about an equal weight of sulphuric acid in water, filtering and setting aside the solution, so that crystals may form on cooling; and (3) copper ammonio-sulphates, for the preparation of which see pp. 177–193.




Copper Sulphate.

Synonyms.—Sulphate of Copper ; Cupric Sulphate ; Blue Vitriol ; Blue Stone.

Copper Sulphate (*Cupri Sulphas*, U. S. P., $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$) is the only copper salt much used in pharmacy. It is a by-product in silver-refining ($2\text{Ag}_2\text{SO}_4 + \text{Cu}_2 = 2\text{CuSO}_4 + 2\text{Ag}_2$). A little is formed in roasting copper pyrites. In the latter case some ferrous sulphide and copper sulphide are oxidized to sulphates ; but the low red heat finally employed decomposes ferrous sulphate, while copper sulphate is unaffected ; it is purified by crystallization from a hot aqueous solution, though frequently much ferrous sulphate remains in the crystals. Copper sulphate results on dissolving in diluted sulphuric acid the black oxide (CuO) obtained in annealing copper plates (see the foregoing equation) ; it may also be prepared by boiling copper with three times its weight of sulphuric acid ($2\text{H}_2\text{SO}_4 + \text{Cu} = \text{CuSO}_4 + \text{SO}_2 + 2\text{H}_2\text{O}$), diluting, filtering, evaporating and crystallizing. In this process a little black copper sulphide is formed.

Anhydrous Sulphate of Copper, B. P. (CuSO_4), is a yellowish-white powder prepared by depriving the ordinary blue crystals of copper sulphate of their water of crystallization by exposing to a temperature of about 200°C . It is used in testing alcohol and similar spirituous liquids for water, becoming blue if the latter be present.

Nitrate of Copper (*Cupri Nitras*, B. P.), *copper nitrate* or *cupric nitrate*. Digest copper in diluted nitric acid. When action has ceased, evaporate and crystallize. If the crystals form at a temperature of 73° to 80°F ., they are prismatic and trihydrous, $\text{Cu}(\text{NO}_3)_2 \cdot 3\text{H}_2\text{O}$; at lower temperatures, tabular and hexahydrous, $\text{Cu}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$.



Verdigris (from *verdi-gris*, Sp., green-gray) is a subacetate or oxyacetate of copper (B. P.) ($\text{Cu}_2\text{O} \cdot 2\text{C}_2\text{H}_3\text{O}_2$), obtained by exposing alternate layers of copper and fermenting refuse grape-husks to the action of air. Digested with twice its weight of acetic acid and a little water, the mixture being evaporated to dryness and the residue dissolved in water, it forms the official (B. P.) solution of acetate of copper ($\text{Cu}_2\text{C}_2\text{H}_3\text{O}_2$).

The modes of forming *Cupric Sulphide*, *Hydrate*, *Oxide*, *Ferrocyanide* and *Arsenite*, as well as *Metallic Copper*, are incidentally alluded to in the following analytical paragraphs.

REACTIONS HAVING ANALYTICAL INTEREST (TESTS).

First Analytical Reaction.—Pass sulphuretted hydrogen through an acidified solution of a copper salt (sulphate, for example) ; a black precipitate (cupric sulphide, CuS) falls.

Second Analytical Reaction.—To an aqueous copper solution

add ammonium sulphhydrate; by this reagent also cupric sulphide is precipitated, insoluble in excess.

Note.—Cupric sulphide is not altogether insoluble in ammonium sulphhydrate if free ammonia or much ammoniacal salt be present; it is quite insoluble in the fixed alkaline sulphhydrates.

Third Analytical Reaction.—Immerse a piece of iron or steel, such as the point of a penknife or a piece of wire, in a few drops of copper solution; the copper is deposited of characteristic color, an equivalent quantity of iron passing into solution. By this reaction copper may be recovered on the large scale from waste solutions, old hoop or other scrap iron being thrown into the liquors.

Note.—This reaction furnishes another illustration of direct chemical substitution. As a matter of *fact*, 56 parts of the iron displace 63.3 of copper. As a matter of *theory*, in the whole mass of copper sulphate each of the (theoretical) ultimate particles or *atoms* of copper, weighing 63.3, is displaced by, or substituted by, one of the (theoretical) ultimate particles or *atoms* of iron, weighing 56, in the mass of iron. *Why* should the fact be as stated? We do not *know*, but the best—nay, the only—suggested explanation is that by Dalton, the one just applied—namely, “the atomic theory,” the theory that matter is not infinitely divisible, but composed of finite particles conveniently termed *atoms*. Just as the bricks in a house might be displaced, one by one, by similarly shaped pieces of stone, without the structure, quâ structure or architecture, being altered, so the copper atoms in the chemical structure termed copper sulphate may be displaced, one by one, by iron atoms, the essential *structure* (at first CuSO_4 , afterward FeSO_4) not being altered. (In this particular case some other buildings or wings—namely, $2\text{H}_2\text{O}$ —are added to the structure at the same time, but this need not complicate the leading idea just offered.)

Fourth Analytical Reaction.—Add ammonia to a cupric solution; cupric hydrate (Cu_2HO) of a light blue color is precipitated. Add excess of ammonia; the precipitate is redissolved, forming a blue solution of copper ammonio-salt, so deep in color as to render ammonia an exceedingly delicate reagent for this metal.

A copper ammonio-sulphate may be obtained in large crystals by adding strongest solution of ammonia to powdered copper sulphate until the salt is dissolved, placing the liquid in a test-glass or cylinder, cautiously pouring in twice its volume of strong alcohol or methylated spirit, taking care that the liquids do not become mixed, tying over the vessel with bladder, and setting aside for some weeks in a cool place (Wittstein). The constitution of copper ammonio-sulphate and other ammonio-salts and corresponding salts of silver will be alluded to in connection with “white precipitate,” the official “ammoniated mercury.”

Fifth Analytical Reaction.—Add solution of potash or soda to a cupric solution; cupric hydrate (Cu_2HO) is precipitated, insoluble in excess. Boil the mixture in the test-tube; the hydrate is decomposed, losing the elements of water and becoming the black anhydrous oxide (CuO).

Sixth Analytical Reaction.—Add solution of potassium ferrocyanide (K_4Fcy) to an aqueous cupric solution; a reddish-brown precipitate (cupric ferrocyanide, Cu_2Fcy) falls. This is an extremely delicate test for copper.

Note.—Of course this reaction, like most chemical reactions, offers a case of chemical *substitution* (four univalent atoms of potassium, K_4 , by two bivalent atoms of copper, Cu_2); it only is not quite so direct as that of the third reaction.

Seventh Analytical Reaction.—To a cupric solution add solution of arsenic and cautiously neutralize with alkali; green cupric arsenite (CuHAsO_3) falls.

Note.—This precipitate has been mentioned already under Arsenum. An arsenum salt is thus a test for copper, as a copper salt is for arsenum—a remark that may obviously be extended to most analytical reactions, for the *body acted upon characteristically by a reagent is as good a test for the reagent as the reagent is for it*; indeed, it becomes a reagent when the other body is the object of search.

Most copper salts color flame green, the chloride blue.

Antidotes.—In cases of poisoning by compounds of copper, iron filings should be administered, the action of which has just been explained. (See Third Analytical Reaction.) Potassium ferro-cyanide may also be given. (See Sixth Analytical Reaction.) Albumen forms with copper a compound insoluble in water; hence raw eggs should be swallowed, vomiting being induced or the stomach-pump or stomach-siphon applied as speedily as possible.

QUESTIONS AND EXERCISES.

What are the analytical relations of copper, mercury, lead, and silver to each other and to arsenum and antimony?—Name the sources of copper.—What proportion of copper is contained in English and French "copper" coins?—Give diagrams showing how copper sulphate is prepared on the small and large scales.—Work out a sum showing how much crystallized copper sulphate may be obtained from 100 parts of sulphide. *Ans.* 261 $\frac{1}{2}$.—How may copper oxide be prepared?—Mention the formula of verdigris.—Name a good chemical test for copper.—What is the analytical position of copper?—Mention the chief tests for copper.—How may copper be separated from arsenum?—Why is finely-divided iron an antidote in poisoning by copper?

MERCURY.

Symbol, Hg. Atomic weight, 200.

Molecular weight, 200 (*not* double the atomic weight).

Source.—Mercury occurs in nature as sulphide (HgS), forming the ore *cinnabar* (an Indian name expressive of something red), and is obtained from Spain, California, Eastern Hungary, China, Japan, and Peru.

Preparation.—The metal is separated by roasting off the sulphur and then distilling, or, better, distilling with lime, which combines with and retains the sulphur.

Properties.—Mercury (*Hydrargyrum*, U. S. P.) is a silver-white, lustrous metal, liquid at common temperatures. It boils at 662° F., and at -40° F. solidifies to a malleable mass of octahedral crystals. When quite free from other metals it does not tarnish, and its globules roll freely over a sheet of white paper without leaving any streak or losing their spherical form.

Formula.—The formula of the mercury molecule is Hg, and not Hg₂, because (at all events at the high temperature at which alone the weight of its vapor can be determined) two volumes, which, if hydrogen would weigh 2 parts (H₂) or oxygen 32 parts (O₂), in the case of mercury vapor weigh only 200 parts (Hg); that is, only once the atomic weight, not twice. That 200, and not 100, is the atomic weight of mercury, is shown by the fact that 200 is the minimum proportion, relative to 1 of hydrogen, in which mercury combines, and by its relation to heat. Still, it is difficult to imagine an atom existing in a free state in nature; and the suggestion has been made that (as is proved to be the case with sulphur) mercury, as we know it, is in an abnormal condition, and that if the weight of its vapor could be taken at a lower temperature or under some other conditions, its molecular weight might be found to be 400. Similar remarks may be made respecting zinc and cadmium, the molecular weights of which, so far as we know, are identical with their atomic weights. At a *very* high temperature the weight of the vapor of iodine indicates a uniatomic molecule (I), at less elevated temperature two atoms in the molecule (I₂)—a fact strongly supporting the inference that the real molecular weight of mercury is 400.

Medicinal Compounds.—The compounds of mercury used in medicine are all obtained from the metal. The metal itself, rubbed with chalk or with confection of roses and powdered liquorice-root, or with lard and suet, until globules are not visible to the *unaided eye*, is often used in medicine. The preparations are: the *Hydrargyrum cum Creta*, U. S. P., or "gray powder;" *Massa Hydrargyri*, U. S. P., or "blue pill;" and *Unguentum Hydrargyri*, U. S. P., or "blue ointment." There is also a compound ointment, a plaster of mercury, a plaster of ammoniacum and mercury, a liniment, and a suppository. Their therapeutic effects are probably due, not to the large quantity of metallic mercury in them, but to the small quantities of black and red oxide which occur in them through the action of the oxygen of the air on the finely-divided metal. The

proportion of oxide or oxides varies according to the age of the specimen. All these medicinal preparations of metallic mercury are indefinite and unsatisfactory, and that through no fault of the pharmacist. They much need investigation by pharmacologists and therapeutists. Here, as in many similar cases, if Medicine would first ascertain her own requirements, and then make them known, her handmaid Pharmacy would be found to be quite capable of supplying them.

Mercurous and Mercuric Compounds.—Mercury combines with other elements and radicals in two proportions: those compounds in which the acidulous radicals are in the lesser amount are termed *mercurous*, the higher being *mercuric*. Thus calomel (HgCl) * is mercurous chloride, while corrosive sublimate (HgCl_2) is mercuric chloride. In every pair of mercury compounds the mercuric contains *twice* as much complementary radical, in proportion to the mercury, as the mercurous.

Notes on Nomenclature.—The remarks made concerning the two classes of iron salts, ferrous and ferric (p. 143), apply in the main to the two series of mercury salts. The latter are systematically distinguished in most modern works by the terms *mercurous* and *mercuric*. But the alteration of the names of well-known substances in medicine and pharmacy, especially if they are poisonous substances, is liable to be attended by serious consequences to patients; therefore in the British and United States Pharmacopœias, which include only a few in comparison with the whole number of mercury salts, older and more strongly contrasted names are employed, thus:

Systematic Names.	Official Names.
Mercurous iodide . .	Yellow mercurous iodide.
Mercuric iodide . . .	Red mercuric iodide.
Mercurous nitrate . .	Not mentioned.
Mercuric nitrate . .	Mercuric nitrate.
Mercurous sulphate . .	Not mentioned.
Mercuric sulphate . .	Basic mercuric sulphate.
Mercurous chloride . .	Mild mercurous chloride.
Mercuric chloride . .	Corrosive mercuric chloride.
Mercurous oxide . . .	Not mentioned.
Mercuric oxide . . .	{ Red mercuric oxide, or red precipitate. { Yellow mercuric oxide, or yellow precipitate.

Specific Gravity.—Mercury is 13.6 times as heavy as water.

Amalgams.—The compound formed on fusing metals together is usually termed an *alloy* (*ad* and *ligo*, I bind), but if mercury is a constituent, an *amalgam* (μάλαγμα, *malagma*, from μαλάσσω, *malasso*, I soften, the presence of mercury lowering the melting-point of such a mixture). Most metals, even hydrogen, according to Loew, form amalgams. "Electric amalgam" consists of 1 part each of tin and zinc and 3 parts mercury.

* The specific gravity of the vapor of calomel, and the fact that the salt is *not* decomposed at the temperature at which its specific gravity is taken, indicate that the formula of calomel is HgCl , and not Hg_2Cl_2 .

REACTIONS HAVING SYNTHETICAL INTEREST.

The Two Iodides.

First Synthetical Reaction.—Rub together a small quantity of mercury and iodine, controlling the rapidity of combination by adding previously, and afterward occasionally, a few drops of spirit of wine, which by evaporation absorbs heat and thus keeps down the temperature. The product is either mercuric iodide, mercurous iodide, or a mixture of the two, as well as mercury or iodine if excess of either has been employed. If the two elements have been previously weighed in single atomic proportions, 200 of mercury to 127 of iodine (about 8 to 5, or 1 ounce of mercury to 278 grains of iodine), the mercurous or greenish (grayish-green) iodide results (HgI); if in the proportion of one atom of mercury to two atoms of iodine (200 to twice 127, or about 4 to 5), the mercuric or red iodide, HgI_2 , results—an iodide that is official, but made in another way (see p. 198). The mercurous iodide should be made and dried (without heat) with as little exposure to light as possible. Red iodide may be removed from it by well washing with alcohol, and, if pure, the mercurous iodide is yellow.

Mercurous iodide is decomposed slowly by light, and quickly by heat, into mercuric iodide and mercury. Mercuric iodide occurring as an impurity in mercurous iodide may be detected by digesting in ether (in which mercurous iodide is insoluble), filtering, and evaporating to dryness; mercuric iodide remains. Mercuric iodide is stable, and may be sublimed in scarlet crystals without decomposition. (For the mechanical details of the method by which a specimen of the crystals may be obtained, and the precautions to be observed, *vide* "Corrosive Sublimate," p. 202.)


Relation of Mercuric Iodide to Light.—In condensing, mercuric iodide is at first yellow, afterward acquiring its characteristic scarlet color. This may be shown by smearing or rubbing a sheet of white paper with the red iodide, and then holding the sheet before a fire or over a flame for a few seconds. As soon as the paper becomes hot the red instantly changes to yellow, and the salt does not quickly regain its red color, even when cold, if the paper is carefully handled. But if a mark be made across the sheet by anything at hand, or the salt be pressed or rubbed in any way, the portions touched immediately return to the scarlet condition. According to Warington, this change is consequent upon rhomboidal crystals being converted into octahedra with a square base, and will serve as an excellent illustration of the influence of physical structure in causing color. The yellow modification so acts on the rays of white light shining on its particles as to absorb the violet and reflect the complementary hue, the yellow, which, entering the eye of the observer, strikes his retina, and thus conveys to the brain the impression of yellowness; and the red modification, though actually the same

chemical substance, is sufficiently different in the structure of its crystals to absorb the green constituent of white light and reflect the complementary ray, the red.

Illustration of the Chemical Law of Multiple Proportions (p. 49).—Applying the atomic theory to the above iodides, it will at once be apparent why mercury and iodine should combine in the proportion of 200 of mercury with either 127 or 254 of iodine, and not with any intermediate quantity. For it is part of that theory that masses are composed of atoms, and that atoms are indivisible, and that the weight of the atom of mercury is to that of iodine as 200 is to 127. Mercury and iodine can only combine, therefore, in atomic proportions, atom to atom (which is the same as 200 to 127), or one atom to two atoms (which is the same as 200 to 254). To attempt to combine them in any intermediate proportion would be useless; a mere mixture of the two iodides would result. A higher proportion of mercury than 200 to 127 of iodine gives but a mixture of mercurous iodide and mercury; a higher proportion of iodine than 254 to 200 of mercury gives but a mixture of mercuric iodide and iodine. Or, for example, 200 grains of mercury mixed with, say, 200 of iodine would yield 139 grains of mercurous iodide and 261 of mercuric iodide; for the 200 grains of mercury uniting with 127 of the iodine give, for the moment, 327 grains of mercurous iodide and 73 of iodine still free. The 73 grains of iodine will immediately unite with 188 of the mercurous iodide (for if 127 of I require 327 of HgI to form HgI_2 , 73 will require 188) and form 261 grains of mercuric iodide, diminishing the 327 grains of mercurous iodide to 139.

Note on Atomic Weight.—But the student will ask, *Does* 127 represent the atomic position of iodine, relative, of course, to 1 of hydrogen, 16 of oxygen, and so on? Yes; because all synthetical and analytical operations with iodine show that 127 parts by weight is the quantity in which iodine migrates from compound to compound, or either displaces or is displaced by chlorine or bromine; secondly, because equal gaseous volumes of such elements contain equal numbers of atoms, and such equal volumes weighing 1, 16, 127, etc., each of the atoms of the respective elements must weigh 1, 16, 127, etc.; and thirdly, because similar numerical relationships are met with when the specific heats of the elements are determined or when their electrical, optical, isomorphic, and other relationships are studied. (*Vide* Index under the words “Atomic,” “Constitution,” “Molecular,” “Structure,” etc.).

Preparation of Red Mercury Iodide by Precipitation.—To a few drops of a solution of a mercuric salt (corrosive sublimate, for example) add solution of potassium iodide, drop by drop; a precipitate of mercuric iodide forms, and at first redissolves, but is permanent when sufficient iodide has been added. Continue the addition of potassium iodide; the precipitate is redissolved.



Notes.—When first precipitated mercuric iodide is yellowish-red, but soon changes to scarlet. Its solubility either in solution of the mercuric salt or in solution of potassium iodide renders the detection of a small quantity of a mercuric salt by potassium iodide, or a small quantity of an iodide by a mercuric solution, difficult, and hence lessens the value of the reaction as a test. But the reaction has synthetical interest, the method of precipitation being that adopted in the British and United States Pharmacopœias (*Hydrargyri Iodidi Rubrum*, U. S. P., or *Red Mercuric Iodide*). Mercuric iodide thus made has the same composition as that prepared by direct combination of its elements. Equivalent proportions of the two salts must be used in making the preparation ($\text{HgCl}_2 = 271$; $2\text{KI} = 332$). About 4 parts of corrosive sublimate are dissolved in 50 or 60 of water (warmth quickens solution) and 5 of potassium iodide in 15 or 20 of water, the solutions mixed and the precipitate collected on a filter, strained, washed twice with distilled water, and dried on a plate over a water-bath. Mercuric nitrate, which is more soluble, and therefore somewhat more convenient for use on the large scale, may be used instead of the mercuric chloride. The mercury in mercuric or mercurous iodide is set free, and sublimes in globules on heating either powder with dried sodium carbonate in a test-tube; the iodine may be detected by digesting with solution of soda, filtering, and to the solution of sodium iodide thus formed adding starch-paste and acidulating with nitrous acid, when blue iodide of starch results. Mercuric iodide is insoluble in water, slightly soluble in alcohol, tolerably soluble in ether. Precipitated red mercury iodide mixed with white wax, lard, and oil forms the *Unguentum Hydrargyri Iodidi Rubri*, B. P.

“Mercuric Potassium Iodide Test-solution,” U. S. P., is made by dissolving in a little water 13.5 grms. of mercuric iodide, and mixing with a solution of 49.8 grms. of potassium iodide, and, after well shaking, making up to 1000 cc.

The Two Nitrates.

Second Synthetical Reaction.—Place a globule of mercury about half the size of a pea in a test-tube; add 20 or 30 drops of nitric acid; boil slowly until red fumes no longer form; set aside. On cooling, if a globule of mercury still remains in the tube, crystals of mercurous nitrate separate. These may be dissolved in water slightly acidulated by nitric acid. The solution may be retained for subsequent analytical operations.



Third Synthetical Reaction.—Place mercury in excess of strong nitric acid and warm the mixture; mercuric nitrate is formed, and will be deposited in crystals as the solution cools; or to crystals of mercurous nitrate add nitric acid, and boil until red fumes cease to form. Retain the product for a subsequent experiment.


When mercury and nitric acid are boiled together, mercurous nitrate is formed if the mercury be in excess, while mercuric nitrate is produced if the acid preponderate.

The mercuric nitrates vary somewhat in composition, according to the proportion, strength, and temperature of the acid used in their formation. A mercuric nitrate may be obtained having the formula Hg_2NO_3 .



Mercuric Oxynitrates.—From the normal mercuric nitrate several oxynitrates may be obtained. Thus on merely evaporating a solution of mercuric nitrate and cooling, crystals having the formula $\text{Hg}_6\text{O}_36\text{NO}_3$ are deposited. The latter, by washing with cold water, yield a yellow pulverulent oxynitrate, $\text{Hg}_6\text{O}_44\text{NO}_3$: mixed with lard, this has sometimes been used as an ointment. Boiled in water, the yellow gives a brick-red oxynitrate, $\text{Hg}_6\text{O}_52\text{NO}_3$.


The pharmacopoeial preparations of mercuric nitrate are *Solution of Mercuric Nitrate*, *Liquor Hydrargyri Nitratis*, U. S. P., containing about 60 per cent. of mercuric nitrate and 11 of free acid, sp. gr. 2.100, and *Unguentum Hydrargyri Nitratis*, U. S. P. The former is made by placing 40 parts of red mercuric oxide in 45 parts of nitric acid diluted with 15 parts of water.



The *unguentum* or "citrine ointment" is made by oxidizing lard oil with nitric acid, and then adding a solution of mercury in nitric acid. It is sometimes diluted with soft paraffin (*Unguentum Hydrargyri Nitratis Dilutum*, B. P.).

The Two Sulphates.

Fourth Synthetical Reaction.—Boil 2 or 3 grains of mercury with a few drops of strong sulphuric acid in a test-tube, or, better, small dish, in a fume-chamber; sulphurous acid gas (SO_2) is evolved, and mercuric sulphate or mercury persulphate (*Hydrargyri Persulphas*, B. P., HgSO_4) results—a white, heavy, crystalline powder.



Between 2 and 3 ounces of mercuric sulphate may be prepared from a fluidrachm of mercury and a fluidounce of sulphuric acid, boiled together in a small dish. These equal the official proportions. The operation is completed and any excess of acid removed by evaporating the mixture of metal and liquid to dryness, either in the open air or in a fume-chamber, sulphuric acid vapors being excessively irritating to the mucous membrane of the nose and throat; dry crystalline mercuric

sulphate remains. If residual particles of mercury are observed, the mass should be damped with sulphuric acid and again heated.

By-products.—In chemical manufactories secondary products, such as the sulphurous gas of the above reaction, are termed *by-products*, and if of value are utilized. In the present case the gas is of no immediate use, and is therefore allowed to escape. When very pure sulphurous acid gas is required for experiments on the small scale, this would be the best method of making it, a delivery-tube being adapted by a cork to the mouth of a flask containing the acid and metal. The mercury sulphate would then become the by-product.

Mercuric Oxsulphate.—Water decomposes mercuric sulphate into a soluble acid salt and an insoluble yellow oxsulphate ($\text{Hg}_3\text{O}_2\text{SO}_4$). The latter is called *turpeth mineral*, from its resemblance in color to *vegetable turpeth*, the powdered root of *Ipomœa turpethum*, an Indian substitute for jalap. The yellow mercury sulphate was formerly official in the British Pharmacopœia, but now only in the United States Pharmacopœia—*Hydrargyri Subsulphas Flavus*, or *Yellow Mercuric Subsulphate*. It should be entirely soluble in 10 parts of hydrochloric acid.

Fifth Synthetical Reaction.—Rub a portion of the dry mercuric sulphate of the previous reaction with as much mercury as it already contains; the product, when the two have thoroughly blended, is mercurous sulphate (Hg_2SO_4): it may be retained for a subsequent experiment.

Molecular Weight.—The exact proportion of mercury to sulphate is merely a matter of calculation; for the combining proportion of a compound (if it possess any combining power), or its proportion for interchange or transposition (metathesis), is the sum of the combining proportions, or interchanging or transposing proportions, of its constituents. In other words, *the molecular weight of a compound is the sum of the atomic weights of its elements*. In accordance with this rule (deduced from the first law of chemical combination, p. 47), 296 of mercuric sulphate and 200 of mercury (about 3 to 2) are the proportions necessary to the formation of mercurous sulphate, Hg (200) + HgSO_4 (296) = Hg_2SO_4 (496).


The Two Chlorides.

Sixth Synthetical Reaction.—Mix thoroughly a few grains of dry mercuric sulphate with about four-fifths its weight of sodium chloride, and heat the mixture, slowly, in a test-tube in a fume-chamber; mercuric chloride (HgCl_2), or *corrosive sublimate*, mercury bichloride or perchloride (*Hydrargyri Perchloridum*, B. P.), *Hydrargyri Chloridum Corrosivum*, U. S. P. (*Corrosive Mercuric Chloride* or *Corrosive Chloride of Mercury*),

sublimes and condenses in the upper part of the tube in heavy colorless crystals or as a crystalline mass.

Somewhat larger quantities (in proportion of 20 of sulphate to 16 of salt, and, *vide infra*, 1 of black manganese oxide) may be sublimed in a pair of 2-ounce or 3-ounce round-bottomed gallipots, the one inverted over the other, and the joint luted

Fig. 36.



Sublimation.

by moist fireclay (the powdered clay kneaded with water to the consistence of dough). The luting having been allowed to dry (somewhat slowly to avoid cracks), the pots are placed upright on a sand-tray (plate-shape answers very well), sand piled round the lower and a portion of the upper pot, and the whole heated over a good-sized gas flame for an hour or more in a fume-chamber (see Fig. 36—pots raised to show joint). Red mercury iodide and calomel may be sublimed in the same way. The former requires less, the latter more, heat than corrosive sublimate.



Note.—If the mercuric sulphate contain any mercurous sulphate, some calomel may be formed. This result will be avoided if 2 or 3 per cent. of black manganese oxide be previously mixed with the ingredients, the action of which is to eliminate chlorine from the excess of sodium chloride used in the process, the chlorine converting any calomel into corrosive sublimate. Sodium manganate and a lower manganese oxide are simultaneously produced.

Precaution.—The operation is directed to be conducted with care in a fume-chamber, because the vapor of corrosive sublimate, which might possibly escape, is very acrid and highly poisonous. Mercuric chloride volatilizes, though extremely slowly and slightly, at warm temperatures.

10 grains of mercury perchloride and the same quantity of ammonium chloride in 1 pint of water form the *Liquor Hydrargyri Perchloridi*, B. P. A dilute aqueous solution of mercury perchloride is liable to decomposition, calomel being precipitated, water decomposed, hydrochloric acid formed, and oxygen gas evolved. The presence of excess of ammonium chloride, with a portion of which the mercuric chloride forms a stable double salt, prevents the decomposition.

Seventh Synthetical Reaction.—Mix a few grains of the mer-

curous sulphate of the fifth reaction with about a third of its weight of sodium chloride, and sublime in a test-tube; crystalline mercurous chloride (HgCl) or calomel (*Hydrargyri Subchloridum*, B. P., *Hydrargyri Chloridum Mite*, U. S. P., the *Mild Chloride of Mercury* or *Mild Mercurous Chloride*) results. Larger quantities may be prepared in the manner directed for corrosive sublimate, a somewhat higher temperature being employed: similar precautions must also be observed. The official proportions are 10 of mercuric sulphate to 7 of mercury and 5 of dry sodium chloride. "Moisten the mercury sulphate with some of the water, and rub it and the mercury together until globules are no longer visible; add the sodium chloride, and thoroughly mix the whole by continued trituration. When dry, sublime by a suitable apparatus into a chamber of such size that the calomel, instead of adhering to its sides as a crystalline crust, shall fall as a fine (dull-white) powder on the floor. Wash this powder with boiling distilled water until the washings cease to be darkened by a drop of ammonium sulphhydrate. Finally, dry at a temperature not exceeding 212° F. "




The term *calomel* ($\kappa\alpha\lambda\delta\varsigma$, *kalos*, good, and $\mu\acute{\epsilon}\lambda\alpha\varsigma$, *melas*, black) probably was simply indicative of the esteem in which black mercury sulphide was held, the compound to which the name calomel was first applied.

Test for Corrosive Sublimate in Calomel.—If the mercurous sulphate contain mercuric sulphate, some mercuric chloride will also be formed. Corrosive sublimate is soluble in water, calomel insoluble; the presence of the former may therefore be proved by boiling a few grains of the calomel in distilled water, filtering and testing by sulphuretted hydrogen or ammonium sulphhydrate, as described hereafter. If corrosive sublimate be present, the whole bulk of the calomel must be washed with hot distilled water till the filtrate ceases to give any indications of mercury. Corrosive sublimate is more soluble in alcohol, and still more in ether, calomel insoluble. Ether in which calomel has been digested should therefore, after filtration, yield no residue on evaporation. Calomel is converted by hydrocyanic acid into mercuric salt and a black powder readily yielding metallic mercury. Powell and Bayne have shown that a certain proportion of hydrochloric acid arrests this action.


Note.—The above process is that of the British Pharmacopœia, but calomel may also be made by other methods. Calomel mixed with lard forms the *Unguentum Hydrargyri Subchloridi*, B. P., and with sulphurated antimony, guaiacum resin, and mucilage of tragacanth the *Pilulæ Antimonii Compositæ*, U. S. P., or "Plummer's Pills," and with colocynth, jalap, and gamboge the *Pilulæ Catharticæ Compositæ*, U. S. P.

The Two Oxides.

Eighth Synthetical Reaction.—Evaporate the mercuric nitrate of the third reaction to dryness in a small dish, in a fume-chamber or in the open air, if more than a few grains have been prepared, and heat the residue till no more fumes are evolved; red mercuric oxide (HgO), “red precipitate,” the red mercury oxide (*Hydrargyri Oxidum Rubrum*, U. S. P., or *Red Mercuric Oxide*) remains.




The nitric constituents of the salt may be partially economized by previously thoroughly mixing with the dry mercuric nitrate as much mercury as is used in its preparation, or as much as it already contains (ascertained by calculation from the atomic weights and the weight of nitrate under operation, as in making mercurous sulphate), and well heating the mixture. In this case the free mercury is also converted into mercuric oxide. This is the official process, the pharmacopoeial quantities being 4 ounces of mercury dissolved in $4\frac{1}{2}$ fluidounces of nitric acid diluted with 2 ounces of water, the solution evaporated to dryness, the residue thoroughly mixed with 4 more ounces of mercury, and the whole heated until acid vapors cease to be evolved. (Mercuric oxide is tested for nitrate by heating a little of the sample in a test-tube, when orange nitrous vapors are produced, and are visible in the upper part of the tube if nitrate be present.)



Mercuric oxide is an orange-red powder, more or less crystalline according to the extent to which it may have been stirred during preparation from the nitrate, much rubbing giving it a pulverulent character. Mixed with hard and soft paraffin, it yields the *Unguentum Hydrargyri Oxidi Rubri*, B. P. (1 part in 8). Mercuric oxide in contact with oxidizable organic matter is liable to reduction to black or mercurous oxide.

Ninth Synthetical Reaction.—To a solution of potash or soda or lime-water, in a test-tube or larger vessel, add solution of corrosive sublimate or of mercuric nitrate; yellow mercury oxide, or yellow mercuric oxide (HgO), is precipitated (*Hydrargyri Oxidum Flavum*, U. S. P.).




18 grains of corrosive sublimate to 10 ounces of lime-water form the *Lotio Hydrargyri Flava*, B. P. The precipitate only differs physically from the red mercuric oxide; the yellow is more

minutely divided than the red. Mercuric oxide is very slightly soluble in water, but sufficiently so to communicate a metallic taste.

Tenth Synthetical Reaction.—To calomel add solution of potash or soda or lime-water; black mercury oxide, or mercurous oxide (Hg_2O), is produced, and may be filtered off, washed, and dried. (This reaction and the formation of a white curdy precipitate on the addition of solution of silver nitrate to the filtrate from the mercurous oxide, acidified by nitric acid, form sufficient evidence of a powder being or containing calomel. The curdy precipitate is silver chloride.)

30 grains of calomel to 10 ounces of lime-water form the *Lotio Hydrargyri Nigra*, B. P.



REACTIONS HAVING ANALYTICAL INTEREST (TESTS).

(The mercury occurring as mercuric or mercurous salt.)

First Analytical Reaction.—The Copper Test.—Deposition of mercury upon and sublimation from copper: Place a small piece of bright copper, about half an inch long and a quarter of an inch broad, in a solution of any salt of mercury, mercurous or mercuric, and heat in a test-tube; the copper becomes coated with mercury in a fine state of division. (The absence of any notable quantity of nitric acid must be ensured, or the copper itself will be dissolved. See below.) Pour away the supernatant liquid from the copper, wash the latter once or twice by pouring water into, and then out of, the tube, remove the metal, take off excess of water by gentle pressure in a piece of filter-paper, dry the copper by passing it quickly through a flame, holding it by the fingers; finally, place the copper in a dry, narrow test-tube, and heat to redness in a flame, the tube being held almost horizontally; the mercury sublimes and condenses as a whitish sublimate of minute globules on the cool part of the tube outside the flame. The globules aggregate on gently pressing with a glass rod, and are especially visible where flattened between the rod and the side of the test-tube.

Notes on the Test.—This is a valuable test for several reasons: It is very delicate when performed with care; it brings before the observer the element itself—one which from its metallic lustre and fluidity cannot be mistaken for any other; it is a test for mercurous and mercuric salts; it eliminates mercury in the presence of most other substances, organic or inorganic.

In performing the test the presence of any quantity of nitric acid may be avoided by adding an alkali until a slight permanent precipitate appears, and then very slightly reacidifying with a drop or two of acetic acid, or by concentrating in an evaporating-dish after adding a little sulphuric acid, and then rediluting.

Tests continued (Mercuric Salts).


Second Analytical Reaction.—To a few drops of a solution of a mercuric salt (corrosive sublimate, for example) add solution of potassium iodide, drop by drop; a yellowish-red precipitate (mercuric iodide, HgI_2) forms, and at first redissolves, but is permanent when sufficient potassium iodide has been added. Continue the addition of potassium iodide; the precipitate is redissolved. (See Notes on p. 198.)

Third Analytical Reaction.—Add a solution of mercuric salt to solution of ammonia, taking care that the mixture, after well stirring, still smells of ammonia; a white precipitate falls.

Ammoniated Mercury.

Performed in a test-tube, this reaction is a very delicate test of the presence of a mercuric salt; performed in larger vessels, the mercuric salt being corrosive sublimate (3 ounces dissolved in 3 pints of distilled water, the solution poured into 4 fluidounces of ammonia solution, and the precipitate washed and dried over a water-bath), it is the usual and the pharmacopœial process for the preparation of "white precipitate," the old "ammonio-chloride" or "amido-chloride of mercury" (so called because then considered to be a compound of mercury with chlorine and with *amidogen*, NH_2 , or $\text{HgCl}_2 \cdot \text{Hg} \cdot 2\text{NH}_2$, chloride and *amide* of mercury), now known as ammoniated mercury (*Hydrargyrum Ammoniatum*, U. S. P.).


Constitution of Ammoniated Mercury.—This precipitate is now, however, considered to be mercuric-ammonium chloride ($\text{NH}_2\text{Hg}'\text{Cl}$); that is, ammonium chloride (NH_4Cl) in one molecule of which two univalent atoms of hydrogen are substituted by one bivalent atom of mercury.




Varieties of Ammoniated Mercury.—If the order of mixing be reversed, and ammonia be added to solution of mercuric chloride, a double mercuric-ammonium and mercury chloride results ($\text{NH}_2\text{HgCl} \cdot \text{HgCl}_2$): it contains 76.55 per cent. of mercury. Previously to the year 1826 "white precipitate" was officially made by adding a fixed alkali to a solution of equal parts of corrosive sublimate and sal-ammoniac; this gave a double mercuric-ammonium and ammonium chloride ($\text{NH}_2\text{HgCl} \cdot \text{NH}_4\text{Cl}$), containing 65.57 per cent. of mercury. This compound is now known as "*fusible white precipitate*," because at a temperature somewhat below red-

ness it fuses and then volatilizes. The "white precipitate," which has been official since 1826, contains, theoretically, 79.52 per cent. of mercury. The true compound may be distinguished as "*infusible* white precipitate," from the fact that when heated it volatilizes without fusing. An ointment of this body is official (*Unguentum Hydrargyri Ammoniati*, U. S. P.). Prolonged washing with water converts "white precipitate" into a yellowish compound (NH_2HgCl , HgO); hence the official preparation is not thoroughly freed from the ammonium chloride which is formed during its manufacture, but which, if present in larger proportion than 7 or 8 per cent., gives to it the character of partial or complete fusibility. The officially recognized "ammoniated mercury" should volatilize at a temperature below redness without fusing, and should yield 77.5 per cent. of metallic mercury. With iodine, chlorine, or bromine white precipitate may yield the highly and dangerously explosive nitrogen iodide, chloride, or bromide.

Note.—Mercuric-ammonium chloride is only one member of a large class of compounds derivable from the various salts of ammonium by substitution of atoms of hydrogen in the molecules by atoms of other radicals. The composition of mercurous-ammonium chloride (see page 208) and of silver ammonio-nitrate (B. P.) is consistent with this view. In these formulæ ammonium is symbolized by NH_4 or Am indifferently. The use of the latter promotes clearness in the formulæ, but it must only be employed when the ammonium acts like an elementary radical.



The composition of the copper ammonio-sulphates (pp. 177 and 193) is consistent with the second and third of the following formulæ, the first being that of ammonium sulphate:



The iodide of dimercuric-ammonium ($\text{NH}_2\text{Hg}_2\text{I}$) is formed in testing for ammonia by the "Nessler" reagent. (*Vide* Index.) Troost has obtained NHAm_3Cl .

Fourth Analytical Reaction.—Pass sulphuretted hydrogen through a mercuric solution; a black precipitate (mercuric sulphide, HgS) falls.

Note 1.—Sulphuretted hydrogen also precipitates mercurous sulphide (Hg_2S) from mercurous solutions, and in appearance the precipitates are alike; hence this reagent does not distinguish between

mercurous and mercuric salts. But in the course of systematic analysis mercuric salts are thrown down from solution as sulphide after mercurous salts have been otherwise removed. Both sulphides are insoluble in ammonium sulphhydrate.

Note 2.—An insufficiency of gas gives a white or colored precipitate of oxysulphide. Prolonged contact with sulphuretted hydrogen or a sulphhydrate, especially if warm, converts the black into a red sulphide.

Ethiops mineral, the old *Hydrargyri Sulphuretum cum Sulphure*, is a mixture of mercury sulphide and sulphur, obtained on triturating the elements in a mortar till globules are no longer visible. Its name is probably in allusion to its similarity in color to the skin of the *Æthiop*. It was formerly official.

Vermilion is mercuric sulphide prepared by sublimation. For a description of the Chinese method of manufacturing it, see the *Pharmaceutical Journal* for December 17, 1881.

Tests continued (Mercurous Salts).

Fifth Analytical Reaction.—To a solution of a mercurous salt (the mercurous nitrate obtained in the second synthetical reaction, for example) add hydrochloric acid or any soluble chloride; a white precipitate (calomel, HgCl) occurs. This reaction was formerly official in the Dublin Pharmacopœia as a process for the preparation of calomel.

Sixth Analytical Reaction.—To solution of a mercurous salt add potassium iodide; a green precipitate results (mercurous iodide, HgI).

Seventh Analytical Reaction.—To a mercurous salt, dissolved or undissolved (*e. g.* calomel), add ammonia; a black salt (*e. g.* mercurous-ammonium chloride ($\text{NH}_2\text{Hg}_2\text{Cl}$)) is formed. (See page 206 for explanation of composition.)

Other Tests for Mercury.

The elimination of mercury in the actual state of metal by the copper test, coupled with the production or non-production of a white precipitate on the addition of hydrochloric acid to the original solution, is usually sufficient evidence of the presence of mercury and its existence as a mercurous or mercuric salt. But other tests may sometimes be applied with advantage. Thus metallic mercury is deposited on placing a drop of the solution on a plate of gold (sovereign or half-sovereign) and touching the drop and the edge of the plate simultaneously with a key: an electric current passes under these circumstances from the gold to the key, and thence through the liquid to the gold, decomposing the salt, the mercury of which forms a white metallic spot on the gold, while the other elements go

to the iron. This is called the *galvanic test*, and is useful for clinical purposes. Solution of stannous chloride (SnCl_2) (see Index) from the readiness with which it forms stannic salts (SnCl_4 , SnO_2 , etc.) gives a white precipitate of mercurous chloride in mercuric solutions, and quickly still further reduces this mercurous chloride (and other mercury salts) to a grayish mass of finely-divided mercury; this is the old *maggie test*, probably so called from the white and gray appearance of the precipitate. The reaction may even be obtained from such insoluble mercury compounds as "white precipitate."—Confirmatory tests for mercuric and mercurous salts will be found in the action of solution of potash, solution of soda, lime-water, solution of ammonia, and solution of potassium iodide. (*Vide* pages 205, 206, and 208.)—Normal alkaline carbonates produce yellowish mercurous carbonate and brownish-red mercuric carbonate, both of them unstable.—Alkaline bicarbonates give yellowish mercurous carbonate with mercurous solutions, and with mercuric salts white (becoming red) mercuric oxysalt.—Yellow potassium chromate (K_2CrO_4) gives with mercurous salts a red precipitate (mercurous chromate, Hg_2CrO_4).—Mercury and all its compounds are volatile, many of them being decomposed at the same time, and yielding globules of condensed metal; the experiment is most conveniently performed in a test-tube.—All dry compounds of mercury are decomposed when heated in a dry test-tube with dried sodium carbonate, mercury subliming and condensing in visible globules, or as a whitish deposit yielding globules when rubbed with a glass rod.

Antidote.—Albumen gives a white precipitate with solution of mercuric salts; hence the importance of administering white of egg, while waiting for a stomach-pump or stomach-siphon, in case of poisoning by corrosive sublimate.

QUESTIONS AND EXERCISES.

Name the chief ore of mercury, and describe a process for the extraction of the metal.—Give the properties of mercury.—In what state does mercury exist in "gray powder"?—What other preparations of metallic mercury itself are employed in medicine?—State the relation of the mercurous to the mercuric compounds.—Distinguish between an *alloy* and an *amalgam*.—State the formulæ of the two mercury iodides.—Under what circumstances has mercuric iodide different colors?—Illustrate the chemical law of multiple proportions as explained by the atomic theory, employing for that purpose the stated composition of the two mercury iodides.—Write down the formulæ of mercurous and mercuric nitrates and sulphates.—How is mercuric sulphate prepared?—What is the formula of "turpeth mineral"?—Describe the processes necessary for the conversion of mercury into calomel and corrosive sublimate, using diagrams.—Why is black manganese oxide sometimes mixed with the other ingre-

dients in the preparation of corrosive sublimate?—Give the chemical and physical points of difference between calomel and corrosive sublimate.—How may calomel in corrosive sublimate be detected?—Work out a sum showing how much mercury will be required in the manufacture of 1 ton of calomel. *Ans.* 17 cwt. nearly.—Mention the official preparations of the mercury chlorides.—Give the formulæ and mode of formation of the red, yellow, and black mercury oxides, employing diagrams.—Explain the action of the chief general test for mercury.—How are mercurous and mercuric salts analytically distinguished?—Give a probable view of the constitution of *Hydrargyrum Ammoniatum*, and an equation showing how it is made.—State the best temporary antidote to poisoning by mercury.

LEAD.

Symbol, Pb. Atomic weight, 206.4.

Source.—The ores of lead are numerous, but the one from which the metal is chiefly obtained is lead sulphide (PbS), or *galena* (from γαλήνη, *galēnē*, tranquillity, perhaps from its supposed effect in allaying pain).

Preparation.—The ore is first roasted in a current of air; much sulphur is thus burnt off as sulphurous acid gas, while some of the metal is converted into oxide and a portion of the sulphide oxidized to sulphate. Oxidation being stopped when the mass presents certain appearances, the temperature is raised, and the oxide and sulphate, reacting on undecomposed sulphide, yield the metal and much sulphurous acid gas: $2\text{PbO} + \text{PbS} = \text{Pb}_3 + \text{SO}_2$ and $\text{PbSO}_4 + \text{PbS} = \text{Pb}_2 + 2\text{SO}_2$.

Uses.—The uses of lead are well known. Alloyed with some arsenum, it forms common *shot*; with antimony gives *type-metal*; with tin, *solder*; and in smaller quantities enters into the composition of *Britannia metal*, *pewter*, and other alloys. Lead is so slightly attacked by acids that chemical vessels and instruments are sometimes made of it. Hot hydrochloric acid slowly converts it into lead chloride, with evolution of hydrogen. Sulphuric acid by aid of air only very slightly attacks it, with formation of lead sulphate and water. Even nitric acid very slowly converts it into nitrate, with evolution of nitric oxide and nitrous oxide gases and water.

The salts of lead used in pharmacy and all other preparations of lead are obtained, directly or indirectly, from the metal itself. Heated in a current of air, lead combines with oxygen and forms lead oxide (PbO), a yellow powder (*massicot*), or, if fused and solidified, a brighter, reddish-yellow, heavy mass of bright scales (*Plumbi Oxidum*, or *Lead Oxide*, U. S. P.), termed *litharge* (from λίθος, *lithos*, a stone, and ἀργυρος, *argyros*, silver). It is from this oxide that the chief lead compounds are obtained. Lead oxide, by further roasting in a current of air, yields *red lead* (or *minium*), Pb_3O_4 or $\text{PbO}_2 \cdot 2\text{PbO}$. Both oxides are much used by painters, paper-stainers, and glass-manufacturers. *White lead* is a mixture of lead carbonate (PbCO_3) and hydrate ($\text{Pb}2\text{HO}$) (commonly 2 molecules of the former to 1 of the latter), usually ground up with about 7 per cent. of linseed oil;

it is made by exposing lead, cast in spirals or little gratings, to the action of air, acetic fumes, and carbonic acid gas, the latter generated from decaying vegetable matter, such as spent tan: lead oxyacetate slowly but continuously forms, and is as continuously decomposed by the carbonic acid gas, with production of hydrate and carbonate, or *dry white lead*. The grating-like masses, when ground, form the heavy white pulverulent official *Plumbi Carbonas*, U. S. P. The latter is the active constituent of *Unguentum Plumbi Carbonatis*, U. S. P., the old *Unguentum Cerussæ*. It is also made by bringing carbonic acid gas and litharge together in a solution of lead acetate.


Lead compounds are poisonous, producing saturnine colic, and even paralysis. These effects are termed *saturnine* from an old name of lead, *saturn*. The alchemists called lead saturn, first, because they thought it the oldest of the seven then known metals, and it might therefore be compared to Saturn, who was supposed to be the father of the gods; and, secondly, because its power of dissolving other metals recalled a peculiarity of Saturn, who was said to be in the habit of devouring his own children.

Quantivalence.—The atom of lead is sometimes quadrivalent ($Pb^{''''}$), but in most of the compounds used in medicine it exerts bivalent activity only ($Pb^{''}$).

REACTIONS HAVING SYNTHETICAL INTEREST.

Lead Acetate, or Acetate of Lead.

First Synthetical Reaction.—Place a few grains of lead oxide in a test-tube, add about an equal weight of water and two and a half times its weight of acetic acid, and boil; the oxide dissolves (or, rather, disappears—dissolves with simultaneous decomposition) and forms a solution of lead acetate ($Pb2C_2H_3O_2 \cdot H_3O_2$). When cold or on evaporation, if much water has been used (the solution being kept faintly acid), crystals of lead acetates ($Pb2C_2H_3O_2 \cdot 3H_2O$), *Plumbi Acetas*, U. S. P., are deposited. Larger quantities are obtained by the same method.



The salt is termed *Sugar of Lead*, from its sweet taste. Besides its direct use in pharmacy, it forms three-fourths of the *Pilula Plumbi cum Opio*, B. P., is the chief constituent of *Unguentum Plumbi Acetatis*, B. P., and an ingredient in *Suppositoria Plumbi Composita*, B. P.

Lead Subacetate or Oxyacetate.




Second Synthetical Reaction.—Boil lead acetate with four times its weight of water and rather more than two-thirds its weight of lead oxide; the filtered liquor is solution of lead

oxyacetate, *Liquor Plumbi Subacetatis*, U. S. P., solution of subacetate of lead.

The official liquor is made by boiling 170 parts of acetate and 100 of oxide in 800 of distilled water for half an hour (constantly stirring), filtering, and making up for any loss during evaporation by diluting the filtrate with boiled and cooled distilled water until it weighs 1000 parts. Sp. gr. about 1.195.

A similar solution was used by M. Goulard, who called it *Extractum Saturni*, and drew attention to it in 1770. It is now frequently termed *Goulard's Extract*. A more dilute solution, 3 of liquor and 97 of boiled and cooled *distilled* water, is also official in the Pharmacopœia under the name of *Diluted Solution of Lead Subacetate* (*Liquor Plumbi Subacetatis Dilutus*). The latter is commonly known as *Goulard Water* or "lead-water." The stronger solution is the chief ingredient in *Ceratum Plumbi Subacetatis*, U. S. P., cerate of lead subacetate, a slight modification of the old *Goulard's cerate*. A strong solution of oxyacetate of lead in glycerin constitutes the *Glycerinum Plumbi Subacetatis*, B. P., and this with a mixture of soft and hard paraffin gives the *Unquentum Glycerini Plumbi Subacetatis*, B. P., a modification of the old *Goulard's cerate*.

Lead Oxyacetates.—The official "subacetate of lead" is not a definite chemical salt. It is probably a mixture of two lead subacetates, which are well-known crystalline compounds, and which the author is disposed to regard as having a constitution similar to that he has already indicated for some other salts. (See Iron and Antimony, also Bismuth.) Exposed to air, it absorbs carbonic acid gas, and lead hydrato-carbonate is deposited.



Lead Nitrate, Red Lead, Lead Peroxide.

Third Synthetical Reaction.—Digest a few grains of red lead in nitric acid and water; lead nitrate (Pb2NO_3) is formed, and remains in solution, while a puce-colored lead peroxide (PbO_2) is precipitated. *Lead Nitrate* (*Plumbi Nitras*, U. S. P.) is more directly made by dissolving litharge (PbO) in nitric acid.




The former reaction serves to bring before the student two other lead oxides—namely, red lead (Pb_3O_4) and lead peroxide (PbO_2). In the latter oxide the quadrivalent character of lead is obvious.

Lead nitrate is used officially in preparing lead iodide. For the latter purpose the above mixture may be filtered, the precipitate of lead peroxide purified from adhering nitrate by passing hot water through the filter, the filtrate and washings evaporated to dryness to remove excess of nitric acid, the residual lead nitrate redissolved by ebullition with a small quantity of hot water, and the solution set aside to crystallize or a portion at once used for the following experiment. Lead nitrate forms white crystals derived from octahedra.

Lead peroxide dissolved in cold strong hydrochloric acid apparently yields an unstable perchloride (PbCl_4).

Lead Iodide, or Iodide of Lead.

Fourth Synthetical Reaction.—To a neutral solution of lead nitrate add solution of potassium iodide; a precipitate of lead iodide (PbI_2) falls (*Plumbi Iodidum*, U. S. P.). Equal weights of the salts may be used in making large quantities.




Lead iodide is the chief ingredient in *Emplastrum Plumbi Iodidi*, B. P., and *Unguentum Plumbi Iodidi*, B. P. Lead iodide is soluble in solution of ammonium chloride.

Crystals of Lead Iodide.—Heat lead iodide with the supernatant liquid, and, if necessary, filter; the salt is dissolved, and again separates in golden crystalline scales as the solution cools.

Lead Oleate (Lead Plaster).

Fifth Synthetical Reaction.—Boil together in a small dish a few grains of very finely powdered lead oxide, with twice its weight of olive oil and two or three times as much water, well stirring the mixture, and from time to time replacing water that has evaporated; the product is a white mass of lead oleate ($\text{Pb2C}_{18}\text{H}_{33}\text{O}_2$) (*Emplastrum Plumbi*, B. P.), glycerin remaining in solution. Larger quantities are prepared in the same manner.



The action between the lead oxide and olive oil is slow, requiring several hours for its completion. Lead plaster is a constituent of seven of the thirteen plasters (*Emplastrum*) mentioned in the United States Pharmacopœia.

The *glycerin* may be obtained by treating the aqueous product of the above reaction with sulphuretted hydrogen to remove a trace of lead, then digesting with animal charcoal, filtering, and evaporating. But on the large scale glycerin is produced as a by-product

in the manufacture of candles, for its elements are found in nearly all vegetable and animal fats. (*Vide Index*.) If in making lead plaster the mixture be evaporated to dryness (*Emplastrum Plumbi*, U. S. P.), some of the glycerin will escape with the steam and some remain with the plaster.

Modes of formation of *Lead Chloride, Sulphide, Chromate, Sulphate, Hydrate*, and other salts are incidentally described in the following analytical paragraphs.

REACTIONS HAVING ANALYTICAL INTEREST (TESTS).

First Analytical Reaction.—To solution of a lead salt (acetate, for example) add hydrochloric acid; a white precipitate (lead chloride, PbCl_2) is obtained. Boil the precipitate with much water; it dissolves, but on the solution cooling is redeposited in small acicular crystals. Filter the cold solution, and pass sulphuretted hydrogen through it; a black precipitate (lead sulphide, PbS) shows that the lead chloride is soluble to a slight extent in cold water.

Note.—A white precipitate on the addition of hydrochloric acid, soluble in hot water and blackened by sulphuretted hydrogen, sufficiently distinguishes lead salts from those of other metals; but the non-production of such a precipitate does not prove the absence of a small quantity of lead, lead chloride being slightly soluble in cold water.

Second Analytical Reaction.—Through a dilute solution of a lead salt acidulated with hydrochloric acid pass sulphuretted hydrogen; a black precipitate (lead sulphide, PbS) occurs.

Lead in Water.—The foregoing is a very delicate test. Should a trace of lead be present in water used for drinking purposes, sulphuretted hydrogen will detect it. On passing the gas through a pint of such (acidulated) water a brownish color is produced. If the tint is scarcely perceptible, set the liquid aside for a day; the gas will become decomposed, and a thin layer of sulphur be found at the bottom of the vessel—white if no lead be present, but more or less brown if it contain lead sulphide. Hygienists regard one-twentieth grain of lead per gallon as dangerous, while a lesser quantity may do harm. Water commonly used for drinking purposes should not contain a trace.

Third Analytical Reaction.—To solution of a lead salt add ammonium sulphhydrate; a black precipitate (lead sulphide) falls, insoluble in excess.

Fourth Analytical Reaction.—To solution of a lead salt add solution of potassium chromate (K_2CrO_4); a yellow precipitate (lead chromate, PbCrO_4) is formed, insoluble in weak acids or in solution of ammonium chloride.

Chromes.—This reaction has technical as well as analytical interest. The precipitate is the common pigment termed *chrome yellow* or *lemon chrome*. Boiled with lime and water, a portion of the chromic radical is removed as a soluble calcium chromate, and a lead oxychromate, of a bright red or orange color (*orange chrome*), is produced.

Fifth Analytical Reaction.—To solution of a lead salt add dilute sulphuric acid or solution of a sulphate; a white precipitate (lead sulphate, PbSO_4) falls.

Lead sulphate is slightly soluble in strong acids and in solutions of alkaline salts; it is insoluble in acetic acid. It is readily dissolved, and, indeed, decomposed, by solution of ammonium acetate, the liquid yielding the ordinary reactions with soluble chromates and iodides.

In dilute solutions the above sulphuric reaction does not take place immediately; the precipitate, however, falls after a time: its appearance may be hastened by evaporating the mixture nearly to dryness and then rediluting.

The white precipitate generally noticed in the vessels in which diluted sulphuric acid is kept is lead sulphate, derived from the leaden chambers in which the acid is made: solubility in strong acid and insolubility in weak explains its appearance.

Antidotes.—From the insolubility of lead sulphate in water, the best *antidote* in a case of poisoning by the acetate or other soluble lead salt is a soluble sulphate, such as Epsom salt, sodium sulphate or alum, vomiting being also induced or the stomach-pump or stomach-siphon applied as quickly as possible.

Other tests for lead will be found in the reaction with *potassium iodide* (*vide* p. 213); with *alkaline carbonates*, a white precipitate ($2\text{PbCO}_3 + \text{Pb}_2\text{HO}$) insoluble in excess; with *alkalies*, a white precipitate (Pb_2HO) more or less soluble in excess; with alkaline *phosphates*, *arsenates*, *ferrocyanides*, and *cyanides*, precipitates mostly insoluble, but of no special analytical interest. Insoluble salts of lead may be decomposed by solution of potash (KHO) or soda (NaHO).

The metal is precipitated in a beautifully crystalline state by metallic zinc and some other metals; the *lead tree* is thus formed.—The *blowpipe flame* decomposes solid lead compounds placed in a small cavity in a piece of charcoal, a soft malleable bead of metal being produced and a yellowish ring of oxide deposited on the charcoal.

QUESTIONS AND EXERCISES.

Write down questions descriptive of the smelting of galena.—Mention some of the alloys of lead.—How is litharge produced?—Give the formulæ of white lead and red lead.—Describe the manufacture of white lead.—

What is the quantivalence of lead?—Draw a diagram showing the formation of lead acetate.—Describe the preparation and composition of *Liquor Plumbi Subacetatis*.—What is the action of nitric acid on red lead, litharge, and lead?—How is the official lead iodide prepared?—Describe the reaction between lead oxide, water, and olive oil at the temperature of boiling water, and give chemical formulæ explanatory of the constitution of the products.—Mention the chief tests for lead.—How would you search for lead in potable water?—What is the composition of chrome yellow?—State a method whereby lead, barium, and silver, in solution, may be separated from each other.—Name the best antidote in cases of poisoning by lead salts.

SILVER.

Symbol, Ag. Atomic weight, 108.

Source.—This element occurs in nature in the metallic state and as ore, the common variety of the latter being silver sulphide (Ag_2S) in combination with much lead sulphide, forming *argentiferous galena*.

Preparation.—The lead from such galena (p. 210) is melted and slowly cooled; crystals of lead separate and are raked out from the still fluid mass, and thus an alloy very rich in silver is finally obtained; this is roasted in a current of air, whereby the lead is oxidized and removed as litharge, pure silver remaining. Other ores undergo various preparatory treatments according to their nature, and are then shaken with mercury, which amalgamates with and dissolves the particles of metallic silver, the mercury being subsequently removed from the amalgam by distillation. Soils and minerals containing metallic silver are also treated in this way. An important improvement in the amalgamation process, by which the mercury more readily unites with the silver, consists in the addition of a small proportion of sodium to the mercury.—Silver chloride may be dissolved from ores by solution of sodium hyposulphite.

Silver is not readily affected by the weak acid or other fluids of food, though it is rapidly tarnished by sulphur or sulphur compounds. It does not perceptibly attack hydrochloric acid; reduces strong nitric acid to nitrous anhydride (N_2O_3), and a weaker acid to nitric oxide (NO); it reduces hot sulphuric acid to sulphurous anhydride (SO_2), silver sulphate (Ag_2SO_4) being formed. The latter salt is crystalline and slightly soluble in water.

REACTIONS HAVING SYNTHETICAL INTEREST.

Impure Silver Nitrate, or Nitrate of Silver.

First Synthetical Reaction.—Dissolve a silver coin in nitric acid; nitric oxide gas (NO) and nitrous anhydride (N_2O_3) are evolved, and a solution of silver and copper nitrates is obtained.

Silver Coinage.—Pure silver is too soft for use as coin; it is therefore hardened by alloying with copper. The silver money of England contains 7.5, of Prussia 25, and of France 10 and 16.5 per

cent. of copper, for the fineness of the French standard silver is 0.900 in the five-franc piece, while an inferior alloy of 0.835 is used for the lower denominations. The single-franc piece, composed of the latter alloy, is still made to weigh 5 grammes, the weight originally chosen for the franc as the unit of the monetary scale when the fineness of the coin was 0.900. It has now become a token, like the British shilling, of which the nominal value exceeds the metallic value. 1 pound troy of British standard silver is coined into 66 shillings, of which the metal is worth from 50s. to 60s., or less or more, according to the market price of silver. The standard fineness of this silver is 0.925—*i. e.* 3 of copper in 40 of the alloy. British silver coins are a legal tender in payments to the amount of 40s only.

Silver Chloride, or Chloride of Silver.

Second Synthetical Reaction.—To the product of the foregoing reaction add water and hydrochloric acid or a soluble chloride; white silver chloride (AgCl) is precipitated, copper still remaining in solution. Collect the precipitate on a filter and wash with water; it is pure silver chloride.

Note.—Copper may also be separated by evaporating the solution of the metals in nitric acid to dryness and gently heating the residue, when the copper nitrate is decomposed, but the silver nitrate is unaffected. The latter may be dissolved from the residual copper oxide by water.

Silver chloride may be obtained in crystals by evaporation of its solution in ammonia.

Pure Silver.


Third Synthetical Reaction.—Place the silver chloride of the previous reaction in a dish, wet it with dilute sulphuric acid, and float a piece of sheet zinc on the mixture; metallic silver is precipitated, and after about one day wholly removed from combination. Collect the precipitate on a filter and wash with water; it is pure metallic silver, and is readily fusible into a single button, especially if mixed with a little borax and nitre.

Note.—Any considerable quantity of silver chloride may be reduced to a lump of the metal by fusion in a crucible, with about half its weight of sodium carbonate. The chloride is also reduced by boiling with caustic alkali and grape-sugar until a trial sample is entirely dissolved by nitric acid.

Pure Silver Nitrate, or Nitrate of Silver.

Fourth Synthetical Reaction.—Dissolve the pure silver of the previous reaction in nitric acid (3 of silver require about 2 or $2\frac{1}{2}$ of strong acid diluted with 5 of water), and remove


excess of acid by evaporating the solution to dryness, slightly heating the residue; the product is pure silver nitrate. Dissolve by heating with a small quantity of water; on the solution cooling or on evaporation colorless tabular crystals of silver nitrate are obtained.



Notes.—The solution of pure or refined silver (*Argentum Purificatum*, B. P.) in nitric acid, evaporation, and crystallization constitutes the usual process for the preparation of the *Silver Nitrate*, or *Nitrate of Silver*, *Argenti Nitras*, U. S. P. The salt fused with 4 per cent. of hydrochloric acid (yielding about 5 per cent. of interlacing silver chloride), and poured into proper moulds, yields the white cylindrical sticks or rods, *Moulded Silver Nitrate*, or *Fused Nitrate of Silver*, *Argenti Nitras Fusus*, U. S. P., commonly termed *caustic* (from *καίω*, *kaio*, I burn) or *lunar caustic*. (The alchemists called silver *Diana* or *Luna*, from its supposed mysterious connection with the moon.) “Caustic points” commonly contain potassium nitrate, which imparts toughness, the *Diluted Silver Nitrate*, or *Diluted Nitrate of Silver*, *Argenti Nitras Dilutus*, U. S. P., or *Mitigated Caustic*, or *Toughened Caustic*, containing 1 part of silver nitrate to 2 parts of potassium nitrate.

The specimen of silver nitrate obtained in the foregoing reaction, dissolved in water, will be found useful as an analytical reagent. Silver nitrate is soluble in rectified spirit, but after a time reaction and decomposition occur.

Silver salts are decomposed when in contact with organic matter, especially in the presence of light or heat, the metal itself being liberated or a black insoluble compound formed. Hence the value of the nitrate in the manufacture of indelible ink for marking linen; hence, too, the reason of the practice of rendering silver solution clear by subsidence and decantation, rather than by filtration through paper; and hence the cause of those cases of actual combustion which have been known to occur in preparing pills containing silver oxide and essential oil or other organic matter. Linen marked with such ink should not be cleansed by aid of bleaching liquor, as the marked parts are then apt rapidly to be oxidized into perfectly rotten matter, holes resulting. Paul says the reaction is as follows:



Silver Oxide, or Oxide of Silver.

Fifth Synthetical Reaction.—To solution of silver nitrate add solution of potash or soda or lime-water; an olive-brown precipitate of silver oxide (Ag_2O) occurs. The washed and dry oxide, like most silver compounds, is decomposed by heat with production of metal. It is also reduced by contact with organic matter. (See the previous paragraph.)

Argenti Oxidum, U. S. P., may be thus made, lime-water being the precipitant employed, soda or potash not being so readily removed by washing. $3\frac{1}{2}$ pints of good lime-water will decompose $\frac{1}{2}$ an ounce of silver nitrate.



Silver oxide is also precipitated on adding ammonia to solution of silver nitrate, but it is rapidly taken up by the ammonium nitrate formed at the same time, argentammonium nitrate, NH_3AgNO_3 , being, doubtless, produced. More ammonia then yields argentammon-ammonium nitrate (see p. 207). The direct solution of silver oxide in ammonia may give the highly explosive substance known as Berthollet's fulminating silver (? NH_2Ag). Ordinary *fulminating silver*, $\text{C}_2\text{N}_2\text{O}_2\text{Ag}_2$, results from the interaction of silver nitrate, nitric acid, and alcohol. The mercury compound, $\text{C}_2\text{N}_2\text{O}_2\text{Hg}$, is used in percussion caps.

Methods of forming several other salts of silver are incidentally mentioned in the following analytical paragraphs.

REACTIONS HAVING ANALYTICAL INTEREST (TESTS).

First Analytical Reaction.—To a solution of a silver salt add hydrochloric acid or other soluble chloride; a white curdy precipitate (silver chloride) falls. Add nitric acid and boil; the precipitate does not dissolve. Pour off the acid and add solution of ammonia; the precipitate dissolves. Neutralize the ammoniacal solution by an acid; the white curdy precipitate (silver chloride, AgCl) is reproduced.

This is the most characteristic test for silver. The precipitated chloride is also soluble in solutions of sodium hyposulphite or potassium cyanide—facts of considerable importance in photographic operations.

Other analytical reagents than the above are occasionally useful.—Sulphuretted hydrogen, or ammonium sulphhydrate, gives a black precipitate (silver sulphide, Ag_2S), insoluble in alkalis.—Solutions of potash or soda give a brown precipitate (silver oxide, Ag_2O), converted into a fulminating compound by prolonged contact with ammonia.—Sodium phosphate gives a pale yellow precipitate (silver phosphate, Ag_3PO_4), soluble in nitric acid and in ammonia.—Ammonium arsenate gives a chocolate-colored precipitate (silver arsenate, $\text{Ag}_3\text{As}_4\text{O}$), already noticed in connection with arsenic acid.—Potassium

iodide or bromide gives a yellowish-white precipitate, silver iodide (*Argenti Iodidum*, U. S. P., or silver iodide) or bromide (AgI or AgBr), insoluble in acids and only slightly soluble in ammonia.—Potassium cyanide gives a white precipitate (silver cyanide, AgCy), soluble in excess, somewhat soluble in ammonia, insoluble in dilute nitric acid, soluble in boiling concentrated nitric acid. *Argenti Cyanidum*, U. S. P., or silver cyanide, may be made by distilling a mixture of potassium ferrocyanide and dilute sulphuric acid, and passing the resulting hydrocyanic acid into a solution of silver nitrate: $\text{HCy} + \text{AgNO}_3 = \text{AgCy} + \text{HNO}_3$, the precipitate being well washed and dried.—Yellow potassium chromate (K_2CrO_4) gives a red precipitate (silver chromate, Ag_2CrO_4).—Red potassium chromate also gives a red precipitate (acid silver chromate, $\text{Ag}_2\text{CrO}_4, \text{CrO}_3$).—Many organic acids afford insoluble salts of silver.—Several metals displace silver from solution, mercury forming in this way a crystalline compound known as the silver tree, or *arbor Dianæ*.—In the blowpipe flame silver salts, placed on charcoal with a little sodium carbonate, yield bright globules of metal, accompanied by no incrustation, as in the corresponding reaction with lead salts; the experiment may be performed with the nitrate, which first melts, and then, like all nitrates, deflagrates, yielding a white metallic coating of silver which slowly aggregates to a button.

Antidotes.—Solution of common salt, sal-ammoniac, or any other inert chloride should obviously be administered where large doses of silver nitrate have been swallowed. A quantity of sea-water or brine would convert the silver into insoluble chloride, and at the same time produce vomiting.

DIRECTIONS FOR APPLYING SOME OF THE FOREGOING REACTIONS TO THE ANALYSIS OF AN AQUEOUS SOLUTION OF A SALT OF ONE OF THE METALS COPPER, MERCURY (MERCUROUS OR MERCURIC), LEAD, SILVER.

Add hydrochloric acid:

Silver is indicated by a white curdy precipitate, soluble in ammonia.

Mercurous salts also by a white precipitate, turned black by ammonia.

Lead by a white precipitate, insoluble in ammonia. Confirm by boiling a small portion of the hydrochloric precipitate in water; it dissolves.

If hydrochloric acid gives no precipitate, silver and mercurous salts are absent. Lead can only be present in very small quantity. Mercuric salts may be present. Copper may be present. Divide the liquid into three portions, and apply a direct test for each metal as follows:

Lead is best detected by the sulphuric test, the tube being set aside for a time if the precipitate does not appear at once.

Mercury is best detected by the copper test. If present here, it occurs as mercuric salt.

Copper betrays itself by the blue color of the liquid under examination. Confirm by the ammonia test.

If the above reactions are not thoroughly conclusive, confirmatory evidence should be obtained by the application of other reagents for copper, mercury, lead, or silver.

TABLE OF SHORT DIRECTIONS FOR APPLYING SOME OF THE FOREGOING REACTIONS TO THE ANALYSIS OF AN AQUEOUS SOLUTION OF SALTS OF ANY OR ALL OF THE METALS COPPER, MERCURY (EITHER MERCUROUS OR MERCURIC SALT, OR BOTH), LEAD, SILVER.

Add hydrochloric acid, filter, and wash the precipitate with a small quantity of cold water.

Ppt. Pb Hg (ous) Ag. Wash with boiling water.			Filtrate. Cu Hg (ic) Pb. Divide into three portions. Test for		
Ppt. Hg (ous) Ag. Add NH_4HO .		Filtrate Pb. Add H_2SO_4 , white ppt.*	Cu by NH_4HO ; blue sol. Hg (mercuric) by Cu; globules. Pb by H_2SO_4 ; white ppt.*		
Precipitate Hg (mercurous), black.	Filtrate Ag. Add an acid, white ppt.				

* Liquids containing only a small quantity of lead do not readily yield lead sulphate on the addition of sulphuric acid. Before lead can be said to be absent, therefore, the liquid should be evaporated to dryness with one drop of sulphuric acid, and the residue digested in water; any lead sulphate then remains as a heavy, white, insoluble powder.

OUTLINE OF THE FOLLOWING TABLES FOR **ANY** OR **ALL** OF
THE COMMON METALS.

HCl.	H ₂ S.	NH ₄ HS.	(NH ₄) ₂ CO ₃ .	(NH ₄) ₂ HAso ₄ .	
Hg (as mercurous salt)	Cu	Zn	Ba	Mg	K
Pb (partially)	Hg (as mer- curic salt)	Al	Ca		Na
Ag	Pb* (entirely)	Fe			NH ₄ .
	As				
	Sb				

The practical student should examine solutions containing these common metals until he is able to analyze with facility and accuracy. In this way he will best perceive the peculiarities of each element and their general relations to each other. As the rarer metals are not included here, the tables are not complete analytical schemes; only general memoranda respecting them will, therefore, now be given. For *complete* memoranda, see notes to subsequent complete tables.

MEMORANDA RELATING TO THE GENERAL ANALYTICAL TABLES
FOR **ANY** OR **ALL** OF THE COMMON METALS (p. 224).

The *group-reagents* adopted in the table are, obviously, *hydrochloric acid*, *sulphuretted hydrogen*, *ammonium sulphhydrate*, *ammonium carbonate*, and *ammonium arsenate*. If a group-reagent produces no precipitate, it is self-evident that there can be no member of the group present. At first, therefore, add only a small quantity of a group-reagent, and if it produces no effect add no more; for it is not advisable to overload a solution with useless reagents: substances expected to come down as precipitates are not infrequently held in the liquid by excess of acid, alkali, or strong aqueous solution of some group-reagent thoughtlessly added. Indeed, experienced manipulators make preliminary trials with group-reagents on a few drops only of the liquid under examination: if a precipitate is produced, it is added to the bulk of the original liquid, and the addition of the group-reagent continued; if a precipitate is not produced, the few drops are thrown away, and the unnecessary addition of a group-reagent thus avoided altogether—an advantage fully making up for the extra trouble of making a preliminary trial. While shunning

* See note on p. 221.

SHORT DIRECTIONS FOR THE ANALYSIS OF AN AQUEOUS SOLUTION OF ORDINARY SALTS OF ONE OF THE METALS HITHERTO CONSIDERED.

Add hydrochloric acid.

Pt. Hg(ous) Pb. Ag. Add NH_4HO. Hg(ous), black ppt. Pb, ppt. still white. Ag, ppt. dissolved.	If HCl gave no precipitate, the metal is still in the liquid; pass H_2S through it.	
Note 1.—If HCl gave no precipitate, neither Hg(ous) nor Ag is present; and Pb only in minute amount, if at all.	Pt. Cu Hg(ic) Pb As Sb. As, yellow ppt. Sb, orange ppt. Cu Hg(ic) } black ppt. Pb	If H_2S gave no precipitate, the metal is still in the liquid; add NH_4Cl, NH_4HO, and NH_4HS.
Note 2.—Hg obtained here must have existed in the solution as a mercurous salt.	Test original solution for Cu by NH_4HO; blue sol. Hg by Cu; globules. Pb by H_2SO_4; white ppt.	Pt. Fe Al Zn Fe, black ppt. Test original solution for ferric salt by K_4Fey (dark blue ppt.); and for ferrous salt by K_6Fecl (dark blue ppt.).
Note 3.—Sb is also precipitated by HCl, but is dissolved on adding more HCl; the Hg, Pb, and Ag precipitates are not soluble in excess of HCl.	Note 1.—If H_2S gave no precipitate, neither Cu, Hg, Pb, As, nor Sb is present.	Al Zn } white ppt. Test original solution by NH_4HO. Al, white ppt. insoluble in excess. Zn, white ppt. soluble in excess.
	Note 2.—Hg and Pb may give colored precipitates (oxysulphides, etc.) with H_2S if too little of the latter has been passed through the solution.	If NH_4HS, etc. gave no precipitate, the liquid may still contain either Ba, Ca, Mg, K, Na, or NH_4; add successively K_2CrO_4 for Ba, $(\text{NH}_4)_2\text{C}_2\text{O}_4$ for Ca, Na_2HPO_4 for Mg. If neither Ba, Ca, nor Mg is found, examine the original solution for NH_4 by KHO, Na by the flame-test, and K by PcCl_4.

TABLE OF SHORT DIRECTIONS FOR APPLYING SOME OF THE FOREGOING ANALYTICAL REACTIONS TO THE ANALYSIS OF AN AQUEOUS SOLUTION OF ORDINARY SALTS OF ANY OR ALL OF THE METALS HITHERTO CONSIDERED.

(Read the "MEMORANDA" on page 222.)

Add hydrochloric acid, and filter.

Precipitate Hg(ous) Ag Pb. Wash, boil with water, filter.		Filtrate Cu Hg(ic) Pb As Sb Fe Al Zn Ba Ca Mg K Na NH ₄ . Pass H ₂ S through the liquid until it ceases to cause any alteration; filter.									
Precipitate Hg(ous) Ag. Add NH ₄ HO to the well-washed pre- cipitate on the filter.	Filtrate Pb. Add H ₂ SO ₄ white precipi- tate. (See footnote, p. 221.)	Precipitate Cu Hg(ic) Pb As Sb. Wash, digest in NH ₄ HS, filter.					Filtrate Fe Al Zn Ba Ca Mg K Na NH ₄ . Add NH ₄ HO, NH ₄ HS, and filter.				
		Precipitate Cu Hg(ic) Pb. Wash, dis- solve in a few drops of HNO ₃ and HCl; evap. nearly to dryness; redissolve in H ₂ O, divide into three, and test for Cu by NH ₄ HO; blue solution. Hg by Cu; globules. Pb by H ₂ SO ₄ ; white precip.					Filtrate As Sb. Add HC ₂ H ₃ O ₂ and boil; digest the precipitate in strong HCl; boil, dilute, filter.				
Precip. Hg(ous). Black precipi- tate.	Filtrate Ag. Add HNO ₃ ; white pre- cipita- tate.	Precipitate Fe. ? Ferric or ferrous. Test origi- nal solution by yellow and red prussiates.					Filtrate Al Zn. Neutralize by HCl; add NH ₄ HO, stir, filter.				
		Precip. As. Yel- low. Confirm by test- ing original solution by the H tests.					Filtrate Sb. Add H ₂ S. Orange precip. Confirm by test- ing original solution by the H tests.				
		Precipitate Ba Ca. Wash, dissolve in H ₂ O, H ₂ SO ₄ , boil; add K ₂ CrO ₄ , filter.					Filtrate Ba Ca Mg K Na NH ₄ . Add (NH ₄) ₂ CO ₃ ; boil, filter.				
		Precip. Ba. Yellow.					Filtrate Ca. Add (NH ₄) ₂ C ₂ O ₄ ; white precip.				
		Ppt. Mg. White.					Filtrate K Na NH ₄ . Evap., ignite, dis- sol. in H ₂ O; test for K by PtCl ₄ ; yel. ppt. Na by flame; yellow. NH ₄ in orig. sol. by NaHO.				

excess, however, care must be taken to avoid deficiency: a substance only partially removed from solution through the addition of an insufficient amount of a reagent will appear where not expected, be constantly mistaken for something else, and cause much trouble; this will not occur if the appearance, odor, or reaction of the liquid on test-paper be duly observed. It is also a good plan, when a group-reagent has produced a precipitate and the latter has been filtered out, to add a little more of the reagent to the clear filtrate; if more precipitate is produced, an insufficient amount of the group-reagent was introduced in the first instance, but the error is corrected by simply refiltering; if no precipitate occurs, the mind is satisfied and the way cleared for further operations.

Group-precipitates, or any precipitates still requiring examination, should, as a rule, be well washed before further testing: this is to remove the aqueous solution of other substances adhering to the precipitate (the mother-liquor, as it is termed), so that subsequent reactions may take place fairly between the reagent used and the precipitate only. A precipitate is sometimes in so fine a state of division as to retard filtration by clogging the pores of the paper, or even to pass through the filter altogether; in these cases the mixture may be warmed or boiled (or a fresh quantity of the original solution may be warmed *before* the group-reagent is added), which usually causes aggregation of the particles of a precipitate, and hence facilitates the passage of liquids.

Division of Work.—It is immaterial whether a solution be first divided into group-precipitates or each precipitate be examined as soon as produced: if the former method be adopted, confusion will be avoided by labelling or marking the funnels or papers holding the precipitate "the HCl ppt.," "the H_2S ppt.," and so on.

The colors and general appearance of the various sulphides and hydrates precipitated should be borne in mind, as the absence of other bodies, as well as the presence of those thrown down, is often at once thus indicated.

Application of confirmatory tests must be frequent.

Results of analyses should be recorded neatly in a memorandum-book—*a*, for correction and endorsement by the teacher; *b*, for future reference by the student, or, *c*, by those who may need evidence respecting his labors; and, *d*, promote mental orderliness.

The various reactions which occur in an analysis have already come before the reader in going through the tests for the individual metals or in other analytical operations; it is unnecessary, therefore, again to draw out equations or diagrams. But the reactions should be thought over, and, if not perfectly clear to the mind, be written out again and again till thoroughly understood.

QUESTIONS AND EXERCISES.

By what process is silver obtained from argentiferous galena?—What weight of English silver coin will yield one pound of pure silver nitrate?—How may the metal be recovered from impure silver salts?—Give a diagram showing the formation of silver nitrate.—Describe the reaction

of lime-water and silver nitrate.—Mention the chief test for silver, and state how silver salts may be distinguished from those of lead and mercury.—Name the antidote for silver.—Give processes for the qualitative analysis of liquids containing the following substances: *a.* Antimony and Mercurous salt. *b.* Lead and Calcium. *c.* Silver and Mercurous salt. *d.* Lead and Mercuric salt. *e.* Copper and Arsenum. *f.* Arsenum and Antimony. *g.* Aluminium and Zinc. *h.* Iron and Copper. *i.* Magnesium, Calcium, and Potassium. *j.* Silver, Antimony, Zinc, Barium, and Ammonium.—Enumerate the so-called group-tests for the metals.—Give a general sketch of the method of analyzing a solution suspected to contain two or more salts of common metals. Classify the common metals according to their analytical relations.

METALS OF MINOR PHARMACEUTICAL IMPORTANCE.

Thus far has been considered, somewhat in detail, the chemistry of the common metals, salts of which are frequently used in medicine or in testing medicinal substances. These are, exclusive of hydrogen,

Potassium,	Barium,	Zinc,	Arsenum,	Mercury,
Sodium,	Calcium,	Aluminium,	Antimony,	Lead,
Ammonium (?),	Magnesium,	Iron,	Copper,	Silver.

Of the remaining metals, eight, exclusive of molybdenum, are mentioned in the British and United States Pharmacopœias—namely:

Lithium,	Manganese,	Tin,	Platinum,
Cerium,	Chromium,	Gold,	Bismuth.

Compounds of four more are sufficiently common to occasionally come under notice:

Strontium,	Cobalt,	Nickel,	Cadmium.
------------	---------	---------	----------

These twelve metals of minor pharmaceutical interest may now be shortly studied, a few only of the reactions of each (just those mentioned in the following pages) being performed. When all have been thus treated their respective positions in the analytical groups will be indicated, and a tabular scheme be given by which an analysis of a solution containing any metal may be effected. Thus, step by step, we may learn how to analyze almost any substance that may occur, and know to what extent the presence of a rarer will interfere with the ordinary tests for a common element; additional illustrations of the working of chemical laws will be acquired and the store of chemical and pharmaceutical facts be increased. The opportunity thus afforded for improvement in habits of neatness of manipulation, in precision, and in power of classification furnishes another and no mean reason why such experiments should be prosecuted, the direct value of which may not be considerable to medical and pharmaceutical learners.

LITHIUM.

Symbol, Li. Atomic weight, 7.

Lithium is widely distributed in nature, but usually in minute proportions compared with other elements. A trace of it may be found in most soils and waters, a Cornish spring containing even considerable quantities as chloride.

One salt used in medicine is the *Citrate* ($\text{Li}_3\text{C}_6\text{H}_5\text{O}_7$) (*Lithii Citras*, U. S. P., lithium citrate), occurring in white deliquescent crystals or powder, prepared by dissolving 50 grains of the carbonate (Li_2CO_3) and 90 of citric acid (50 to 95 if both are quite pure) in 1 ounce of water, evaporating to a low bulk, and setting aside in a dry place to crystallize, or at once evaporating to dryness and powdering the residue. The crystals have the formula $\text{Li}_3\text{C}_6\text{H}_5\text{O}_7 \cdot 4\text{H}_2\text{O}$; dried at 212°F. , $\text{Li}_3\text{C}_6\text{H}_5\text{O}_7 \cdot \text{H}_2\text{O}$ (Umney). It is very soluble in water, but almost insoluble in alcohol and ether.



The effervescent lithium citrate is also official (*Lithii Citras Effervescens*, U. S. P.). It is prepared similarly to other effervescing preparations.

The benzoate (*Lithii Benzoas*, $\text{LiC}_7\text{H}_5\text{O}_2$, U. S. P.), bromide (*Lithii Bromidum*, LiBr , U. S. P.), and salicylate (*Lithii Salicylas*, $\text{LiC}_7\text{H}_5\text{O}_3$, U. S. P.) may be similarly prepared from the respective acids.

The above-named lithium salts are officially tested as follows, the bromide being acted upon directly without ignition:

"On dissolving the residue left on ignition (of either salt) in diluted hydrochloric acid, and evaporating the filtered solution to dryness, 1 part of the residue should be completely soluble in 5 parts of absolute alcohol, and the addition of an equal volume of ether to the alcoholic solution should produce no precipitate (salts of alkalis). On dissolving another portion of the residue in a small quantity of water, the solution should produce no precipitate with test-solution of ammonium oxalate (salts of alkaline earths). The aqueous solution should remain unaffected by hydrosulphuric acid or ammonium sulphide (abs. of metals)."—U. S. P.

The carbonate (*Lithii Carbonas*, U. S. P.) is a white granular powder obtained from the minerals which contain lithium—namely, lepidolite (from *λεπίς*, *lepís*, a scale, and *λίθος*, *lithos*, a stone; it has a scaly appearance), triphane (from *τρεῖς*, *treis*, three, and *φαῖνα*, *phainō*, I shine) or spodumene (from *σποδῶω*, *spodōō*, I reduce to ashes, in allusion to its exfoliation in the blowpipe-flame), and petalite (from *πέταλον*, *petalon*, a leaf; its character is leafy and laminated). Each contains aluminium silicate, with potassium fluoride and lithium in the case of Austrian lepidolite, which is the most abundant source, and sodium and lithium silicate in the others. The lepidolite is decomposed by sulphuric acid, alumina, etc. precipitated by ammonia; the filtrate evaporated and the residue

ignited; the resulting sulphates dissolved in water, and the lithium precipitated by a carbonate. The preparation of common alum is sometimes made a part of the factory processes, and other obvious modifications may be introduced. "If 0.5 gm. of lithium carbonate be dissolved in 2 cc. of hydrochloric acid, and the clear solution be evaporated to dryness, the dry residue should completely dissolve in 3 cc. of absolute alcohol, and an addition of 3 cc. of ether should not render the solution turbid (limit of other alkalies). 0.5 gm. of the dry carbonate, mixed with 20 cc. of water, should require for complete neutralization not less than 13.4 cc. of normal sulphuric acid (corresponding to at least 98.98 per cent. of the pure salt), methyl-orange being used as indicator."—*U. S. P. Liquor Lithiæ Effervescens*, B. P., is a solution of 10 grains of lithium carbonate in 1 pint of water charged with five times its volume of carbonic acid gas and kept in ordinary aerated-water bottles. "Half a pint, evaporated to dryness, yields 5 grains of a white solid residue, answering to the tests for lithium carbonate. . . . 10 grains of the latter salt, neutralized with sulphuric acid and afterward heated to redness, leaves 14.86 grains of dry lithium sulphate, which, when redissolved in distilled water, yields no precipitate with ammonium oxalate or solution of lime," indicating absence of calcium and aluminium salts. Lithium citrate should yield by incineration 52.8 per cent. of white lithium carbonate. According to C. N. Draper, lithium carbonate is soluble in 68 parts of water at 15° C. and 131 at 100° C.

*Lithium urate** is more soluble than sodium urate; hence lithium preparations are administered to gouty patients in the hope that sodium urate, with which such systems are loaded, may become lithium urate and be removed.

In chemical position lithium stands between the alkaline and the alkaline-earth metals, its hydrate, carbonate, and phosphate being slightly soluble in water. The double platinum and lithium chloride also is soluble in water. The atom of lithium is univalent, L'.

Analytical Reaction.—Moisten the end of a platinum wire with solution of a minute particle of solid lithium salt, and introduce it into the flame of a Bunsen burner or other almost colorless flame (spirit-lamp or blowpipe flame); a magnificent crimson tinge is imparted.

The light thus emitted by ignited lithium vapor is of a purer scarlet than that given by strontium, the next element. When the flames are examined by spectral analysis (physically analyzed by a prism), the red rays are, in the case of strontium, found to be associated with blue and yellow, neither of which is present in the lithium light, blue lithium rays only appearing at temperatures much higher than those of the air-gas flame, or, indeed, any other ordinary flame.

* Urates will be considered subsequently in connection with uric acid.

STRONTIUM.

Symbol, Sr. Atomic weight, 87.4.

Source.—Strontium is not widely distributed in nature; but the carbonate (SrCO_3), known as *strontianite*, and the sulphate (SrSO_4), known as *celestine* (from *cælum*, the sky, in allusion to its occasional bluish color), are by no means rare minerals.

Strontium Bromide, $\text{SrBr}_2 \cdot 6\text{H}_2\text{O}$ (*Strontii Bromidum*, U. S. P.), occurs as colorless deliquescent hexagonal crystals, soluble in water and alcohol, insoluble in ether. It may be prepared by dissolving the carbonate (*strontianite*) in hydrobromic acid or roasting the sulphate with coal (celestine), and treating the resulting sulphide with hydrobromic acid.



Strontium Iodide, $\text{SrI}_2 \cdot 6\text{H}_2\text{O}$ (*Strontii Iodidum*, U. S. P.), occurs in crystals similar to the bromide; soluble in water and alcohol, but only slightly in ether. It may be prepared similarly to bromide, using hydriodic instead of hydrobromic acid.

Strontium Lactate. (See Index.)

Strontium salts are used by firework-manufacturers in preparing red fire. The color they impart to flame is a beautiful crimson—ignited strontium vapor emitting red rays, as already explained. Strontium nitrate (Sr_2NO_3) is best for pyrotechnic compositions, its oxygen enabling it to burn freely when mixed with charcoal, sulphur, etc. It or any salts may be obtained, as described above, by dissolving the carbonate or sulphide in an acid.

The position of strontium among the chemical elements is between barium and calcium; its sulphate is extremely sparingly soluble in water. Its atom, like those of barium and calcium, is bivalent (Sr').

ANALYTICAL REACTIONS (TESTS).

First Analytical Reaction.—To a solution of a strontium salt (Sr_2NO_3 or SrCl_2) add ammonium carbonate; a white precipitate (strontium carbonate, SrCO_3) falls.

Second Analytical Reaction.—To a solution of a strontium salt add sulphuric acid previously so diluted that it will not precipitate calcium salts, or add an equally dilute solution of any sulphate—*e. g.* that of calcium itself; a white precipitate (strontium sulphate, SrSO_4) falls. The formation of this precipitate is promoted by stirring and by setting the liquid aside for some time. (Barium is precipitated immediately under similar circumstances.)

Third Analytical Reaction.—To a dilute solution of a strontium salt add yellow chromate (K_2CrO_4 or K_2HCrO_4); no precipitate falls.

Barium may be separated from strontium by potassium chromate, that reagent at once precipitating barium from aqueous or acetic

solutions. The value of the reaction is enhanced if acetic acid or ammonium acetate be present, strontium chromate being far more soluble in such fluids than in water (Ransom). It is also more soluble in cold than in hot fluids.

Fourth Analytical Reaction.—Insert a fragment of a strontium salt in the blowpipe flame or other equally colorless flame, or hold the end of a platinum wire dipped into a strontium solution in the flame; a crimson color is imparted.

Other Analytical Reactions.—Alkali-metal phosphates, arsenates, and oxalates give white insoluble precipitates with strontium, as with barium and calcium.

CERIUM, Ce; At. wt., 141.—This element occurs in the mineral cerite (a silicate of iron, calcium, and the three rare metals cerium, lanthanum, and didymium); also occasionally as impure fluoride, carbonate, and phosphate. Cerium oxalate, a white granular powder, is the only official salt; it may be obtained from cerite by boiling the powdered mineral in strong hydrochloric acid for several hours, evaporating, diluting, and filtering to separate silica; adding ammonia to precipitate hydrates of all the metals except calcium; filtering off, washing, redissolving in hydrochloric acid, and adding oxalic acid to precipitate cerium oxalate. The preparation will still contain lanthanum and didymium oxalates; it is therefore strongly calcined, the resulting lanthanum and didymium oxides dissolved out to some extent by boiling with a concentrated solution of ammonium chloride, the residual cerium oxide dissolved in boiling hydrochloric acid, and ammonium oxalate added to precipitate cerium oxalate ($\text{Ce}'''_2\text{C}_2\text{O}_4 \cdot 9\text{H}_2\text{O}$). According to Hartley, the precipitated hydrates should be treated with chlorine, by which ceric hydrate is left insoluble and the other hydrates converted into soluble hypochlorites.

Cerium oxalate (*Cerii Oxalas*, *Oxalate of Cerium*, or *Cerium Oxalate*, U. S. P.) is decomposed at a dull-red heat, 48 per cent. of a yellow, or, more generally, salmon-colored, mixture of oxides remaining. Usually the didymium present gives the ignited residue a reddish or reddish-brown color. The oxides are soluble in boiling hydrochloric acid (without effervescence, indicating, indirectly, absence of earthy and other carbonates or oxalates), and the solution gives, with excess of a saturated solution of potassium sulphate, a crystalline precipitate of cerium and potassium sulphate. Alumina mixed with cerium oxalate may be detected by boiling with solution of potash, filtering, and adding excess of solution of ammonium chloride, when a white flocculent precipitate (aluminium hydrate) will be obtained. The oxalic radical is recognized by neutralizing the potash solution by acetic acid and adding calcium chloride; a white precipitate (calcium oxalate) falls: this precipitate, though insoluble in acetic, should be wholly dissolved by hydrochloric acid. Acid or neutral cerium solutions give with sodium acetate and hydrogen peroxide a brownish-red color (Hartley).

According to H. G. Greenish, most samples of the oxalate contain as impurities traces of lead, iron, and magnesium.

MANGANESE.

Symbol, Mn. Atomic weight, 55.

Source.—Manganese is a constituent of many minerals, and, as black oxide (MnO_2) (*Mangani Dioxidum*, U. S. P., "containing at least 66 per cent. of the pure dioxide, MnO_2 ; *Mangani Oxidum Nigrum*, 1880), or *pyrolusite* (from $\pi\upsilon\rho$, *pur*, fire, and $\lambda\upsilon\sigma\iota\varsigma$, *luisis*, a loosing or resolving, in allusion to the readiness with which it is split up by heat into a lower oxide and oxygen), is met with in abundance in the south-west of England, Aberdeenshire, and most countries of Europe. It occurs as a steel-gray mass of prismatic crystals or in black shapeless lumps.

The chemical position of manganese is close to iron and three other metals still to be considered—cobalt, nickel, and chromium. Its atom apparently has sexivalent affinities, as seen in potassium manganese (K_2MnO_4), but commonly it is quadrivalent (Mn'''') or bivalent (Mn'').

Uses.—Metallic manganese, which may be isolated by aid of sodium, is used in alloy with iron in the manufacture of some varieties of steel. The black oxide is an important agent in the production of chlorine and in the preparation of green and purple disinfecting manganates, purple glass, and black glaze for earthenware.

REACTIONS HAVING BOTH SYNTHETICAL AND ANALYTICAL INTEREST.

First Reaction.—Boil a little black manganese oxide with hydrochloric acid in a fume-chamber until chlorine ceases to be evolved; filter; the filtrate is a solution of manganous chloride (MnCl_2).



This is the reaction commonly applied in the preparation of chlorine gas. It is also a ready method of preparing a manganous salt for analytical experiments. Coupled with the application of reagents to the filtrate, the reaction is that by which a black powder or mineral would be recognized as black manganese oxide. Black manganese oxide dissolves in *cold* hydrochloric acid, forming a dark-brown solution of a higher chloride or chlorides, MnCl_3 , Mn_2Cl_7 , or, possibly, MnCl_4 .

Second Reaction.—Heat a manganese compound with a grain or two of potassium carbonate and hydrate and a fragment of potassium nitrate or chlorate on platinum-foil in the blowpipe flame; a green mass containing *potassium manganate* (K_2MnO_4) results. Boil the foil in water; the green manganate dissolves,


the fluid soon changing to solution of the purple *potassium permanganate* ($K_2Mn_2O_8$).

Carefully performed, this is a delicate test for manganese.

The reaction is similar to that by which potassium permanganate (*Potassii Permanganas*, U. S. P.) is directed to be prepared for use in volumetric analysis. *Liquor Potassii Permanganatis*, B. P., is a solution of 88 grains of potassium permanganate in 1 pint of distilled water. Equations showing the action which occurs in making the salt have already been given in connection with the compounds of potassium (*vide* p. 77). The proportions of ingredients and details of the operation are as follows:

Reduce $3\frac{1}{2}$ parts (for experiment each "part" may be $\frac{1}{8}$ of an ounce or 4 grammes) of potassium chlorate to fine powder, and mix it with 4 of black manganese oxide; put the mixture into a porcelain basin, and add to it 5 parts of solid caustic potash previously dissolved in 4 parts of water. Evaporate to dryness, stirring diligently to prevent spurting. Pulverize the mass, put it into a covered Hessian or Cornish crucible, and expose it to a dull-red heat (not higher) for an hour (twenty or thirty minutes for quantities of 1 or 2 ounces), or till it has assumed the condition of a semi-fused mass. Allow to cool, pulverize, and boil with about 30 parts of water. Let the insoluble matter subside, decant the fluid, boil again with about 10 parts of water, again decant, neutralize the united liquors accurately with diluted sulphuric acid (or, better, carbonic acid gas), and evaporate till a pellicle forms. Set aside to cool and crystallize. Drain the crystalline mass, boil it in 6 parts of water, and strain through a funnel the throat of which is lightly obstructed by a little asbestos or gun-cotton. Let the fluid cool and crystallize, drain the dark-purple, slender, prismatic crystals, and dry them by placing under a bell-jar over a vessel containing sulphuric acid.

Instead of converting the manganate by ebullition and neutralizing the free alkali by acid, by which one-third of the manganese is lost, chlorine may be passed through the cold solution until the green color is entirely changed to purple.



Solutions of the potassium and sodium manganates are in common use as green and purple disinfecting fluids. They act by oxidizing organic matter, the manganic or permanganic radical being reduced to black manganic oxide or even a lower oxide. The reason for using asbestos instead of paper in filtering the solutions will now be understood.

The changes in color which the green mass of the above process undergoes when dropped into warm water produced for it the old name of *mineral chameleon*.

Third Reaction.—Make a borax bead by heating a fragment of the salt on the looped end of a platinum wire in the blow-pipe flame until a clear transparent globule is obtained. Place

on the bead a minute portion of a manganese compound or touch it with a drop of solution. Again fuse the borax; a bead of a violet or amethystine tint is produced.

This is a good analytical reaction. It has also synthetical interest, illustrating the use of black manganese oxide in producing common purple-tinted glass.

Expose the bead to the reducing part of the flame (p. 135), the part nearer to the blowpipe, where there are highly-heated hydrocarbon gases greedy of oxygen; the color disappears.

This is owing to the reduction of the manganic compound to a manganous condition, in which it no longer possesses peculiar coloring-power. This action also illustrates the use of black manganese oxide in glass-manufacture. Glass, when first made, is usually of a green tint, owing to the presence of ferrous impurities: the addition of manganic oxide to the materials converts the ferrous into ferric compounds, which have comparatively little colorific power, it itself being thereby reduced to manganous oxide, which also gives but little color. If excess of manganic oxide be added, a purple tint is produced.

REACTIONS HAVING ANALYTICAL INTEREST (TESTS).

Fourth Reaction.—Through a solution of a manganous salt acidified by hydrochloric acid pass sulphuretted hydrogen; no decomposition occurs. Add ammonia; the ammonium sulphhydrate thus formed causes the precipitation of a yellowish-pink or flesh-tinted precipitate (manganous sulphide, MnS , in a hydrous state).

This reaction is characteristic, manganese sulphide being the only flesh-colored sulphide known. The salt used may be the manganous chloride obtained in the first reaction, but such crude solutions usually give a black precipitate with ammonium sulphhydrate, owing to the presence of iron. The latter element may be precipitated, however, on adding excess of ammonia (and rapidly filtering, or oxygen will be absorbed and most of the manganese also be precipitated), or on boiling the manganous solution with a very little sodium carbonate, which attacks the ferric salt in preference to the manganous. *Pure* manganous chloride may be similarly obtained on boiling the impure solution with manganous carbonate; the latter decomposes the ferric chloride, with production of ferric hydrate and more manganous chloride and evolution of carbonic acid gas.

To the recently precipitated manganous sulphide add acetic acid; it is dissolved.

This solubility enables manganese to be separated from nickel, cobalt, and zinc, whose sulphides are insoluble in weak acetic acid. To express the fact in another way: manganese is not precipitated by sulphuretted hydrogen from a solution containing free acetic acid.

Fifth Reaction.—To solution of a manganous salt add ammonia; a white precipitate (manganous hydrate, Mn_2HO) falls. Add excess of ammonia; some of the precipitate is dissolved, and may be detected in the quickly-filtered solution by ammonium sulphhydrate. But both precipitate and solution rapidly absorb oxygen, the manganese passing into a more highly oxidized condition, in which it is insoluble in ammonia. The fixed alkalies give a similar precipitate *insoluble* in excess. The precipitate rapidly absorbs oxygen, becomes brown, and gradually passes into a higher oxide.

Sixth Reaction.—To a solution of a manganous salt add diluted nitric acid, and either red lead or the puce-colored lead oxide or peroxide, and then boil; a red tint, said to be due to permanganic acid, is imparted to the liquid. If chlorides are present, the manganese, etc. should be separated by potash or soda, the precipitate be well washed, dissolved in nitric acid, and then the oxide be added (Crum).

Seventh Reaction.—Heat a little manganese dioxide in a test-tube with sulphuric acid; oxygen is evolved and manganese sulphate formed (*Mangani Sulphas*, $\text{MnSO}_4 \cdot 4\text{H}_2\text{O}$, U. S. P.); add water, boil, filter, evaporate, and set aside to crystallize. Larger quantities are made in a similar manner.

Manganese sulphate ($\text{MnSO}_4 \cdot 5\text{H}_2\text{O}$) occurs in colorless or pale rose-colored, transparent crystals, which, when deposited from a solution at a temperature between 68° and 86° , have the form of right rhombic prisms and contain four molecules of water (U. S. P.). This salt is very soluble in water. Other sulphates containing 1, 2, 3, and 9 of water are known. The solution is not colored by tincture of nutgall (a black shows iron), but affords with caustic alkalies a white precipitate (Mn_2HO), which, by exposure to the air, soon absorbs oxygen and becomes brown. Ammonium sulphhydrate throws down a flesh-colored precipitate (MnS), and potassium ferrocyanide a white one (Mn_2Fcy).

Many other reactions occur between manganese salts and various reagents, but are of no particular synthetical or analytical interest.

COBALT AND NICKEL.

Krüss and Schmidt state that these very closely allied metals, as hitherto known, are not true elements, but contain a third element, the oxide of which resembles, yet distinctly differs from, alumina and zinc oxide.

COBALT.

Symbol, Co. Atomic weight, 58.6.

Source.—Cobalt occurs sparingly in nature as the arsenide

(CoAs_2), or *tin-white cobalt*, and occasionally as a double arsenide and sulphide ($\text{CoAs}_2, \text{CoS}_2$), or *cobalt-glance* (from *glanz*, brightness, in allusion to its lustre).

Uses.—Its chief use is for coloring glass blue. Cobalt is also the coloring constituent of *smalt* (from *smelt*, a corruption of *melt*), a finely-ground sort of glass used as a blue pigment by paper-stainers and others, and employed also by laundresses to neutralize the yellow tint of washed linen.

The salts of cobalt may be obtained from the oxide (CoO), and the oxide from *zaffre*, a mixture of sand and roasted ore.

Quantivalence.—The atom of cobalt often exhibits quadrivalent affinities, but still more often exerts only bivalent powers (Co''). Cobalt has analytical relations with zinc, nickel, and manganese.

ANALYTICAL REACTIONS (TESTS).

First Analytical Reaction.—Pass sulphuretted hydrogen through an acidified solution of a salt of cobalt—the chloride (CoCl_2) or nitrate (Co_2NO_3), for example; no decomposition occurs. Add ammonia; the ammonium sulphhydrate thus formed causes a black precipitate (cobalt sulphide, CoS). (The moist precipitate slowly absorbs oxygen from the air, yielding some cobalt sulphate, CoSO_4 .)

Second Analytical Reaction.—Add ammonia gradually to a cobalt solution; a blue precipitate (impure cobalt hydrate, Co_2HO) falls. Add excess of ammonia; the precipitate is dissolved, yielding a liquid somewhat more reddish-brown than the original solution. A similar precipitate is given by the fixed alkalis, *insoluble* in excess.

Third Analytical Reaction.—Make a borax bead by heating a fragment of the salt on the looped end of a platinum wire in the blowpipe flame until a clear transparent globule is obtained. Place on the bead a minute portion of any cobalt compound or touch it with a drop of solution. Again fuse the borax; a blue bead results.

This is a delicate test for cobalt. From what has been said previously it will be seen that this experiment has also considerable synthetical interest.

Fourth Analytical Reaction.—To a solution of a salt of cobalt add a few drops of hydrochloric acid, then excess of solution of potassium cyanide, and boil for ten minutes; oxygen is absorbed, and potassium cobalticyanide ($\text{K}_6\text{Co}_2\text{Cy}_{12}$) formed. Add hydrochloric acid, and boil the mixture (in a fume-cupboard, to avoid inhalation of any hydrocyanic acid); the excess of potassium cyanide is thus decomposed, the cobalticyanide unaffected. Now add *excess* of solution of potash;

the potassium cobalticyanide is probably decomposed, but the cobalt remains dissolved in the alkaline liquid. Nickel under similar circumstances is precipitated, the reaction thus affording means of separating these closely-allied metals from each other.

Other reactions between the cobalt solution and different reagents may be performed, and various precipitates obtained, but these have no special analytical interest.

Invisible Ink.—Many salts of cobalt containing water of crystallization are light red, the anhydrous more or less blue. Prove this by writing some words on paper with a solution of chloride of cobalt sufficiently dilute for the characters to be invisible when dry: hold the sheet before a fire or over a flame; the letters at once become visible, distinct, and of a blue color. Breathe on the words or set the sheet aside for a while; the characters are once more invisible, owing to absorption of moisture. Hence solution of cobalt chloride forms one of the so-called *sympathetic inks*.

NICKEL.

Symbol, Ni. Atomic weight, 58.6.

The ores of nickel and cobalt are commonly associated in nature. Indeed, it is from *speiss*, an arsenio-sulphide of nickel obtained in the manufacture of a pigment of cobalt, smalt, already mentioned, that much of the nickel met with in commerce has hitherto been obtained. *Garnierite*, silicate of magnesium and of nickel, containing no cobalt, is also a valuable source of nickel. Nickel is used in the preparation of the white alloy known as German or nickel silver and for plating iron.

Quantivalence.—Nickel exerts bivalent activity (Ni'') in its ordinary compounds. Its salts and their solutions are usually green. They are chiefly made, directly or indirectly, from the metal itself.

ANALYTICAL REACTIONS (TESTS).

First Analytical Reaction.—Pass sulphuretted hydrogen through an acidified solution of a salt of nickel—chloride (NiCl_2), nitrate (Ni_2NO_3), or sulphate (NiSO_4); no decomposition occurs. Add ammonia; the ammonium sulphhydrate formed causes a black precipitate (nickel sulphide, NiS).

Note.—When nickel sulphide is precipitated by the direct addition of the common yellow solution of ammonium sulphhydrate, which always contains free sulphur, there is much difficulty in filtering the mixture, owing to the slight solubility of nickel sulphide in the reagent and the formation of some nickel sulphate (NiSO_4), oxygen being

absorbed from the air by the sulphide. This may be avoided by warming the mixture and using freshly-made ammonium sulphhydrate, in which the nickel sulphide is insoluble; or, when practicable, the salt of nickel may be precipitated from an ammoniacal solution by sulphuretted hydrogen.

Second Analytical Reaction.—Add ammonia, drop by drop, to a nickel solution; a pale-green precipitate (nickel hydrate, $\text{Ni}(\text{OH})_2$) falls, especially on boiling the mixture. Add excess of ammonia; the precipitate dissolves, yielding a bluish rather than the original green-colored solution. A similar precipitate is given by the fixed alkalies, *insoluble* in excess.

Third Analytical Reaction.—Nickel salts color a borax bead, when hot, a reddish-yellow tint; the reaction is not very serviceable analytically.

Fourth Analytical Reaction.—To a solution of a salt of nickel add solution of potassium cyanide; nickel cyanide (NiCy_2) is precipitated. Add excess of solution of potassium cyanide; the precipitate is dissolved with formation of double nickel and potassium cyanide ($\text{NiCy}_2 \cdot 2\text{KCy}$). Next add hydrochloric acid, and boil the mixture (in a fume-cupboard, adding a little hydrochloric acid from time to time until all smell of hydrocyanic acid has disappeared). Lastly, add *excess* of solution of potash; pale-green nickel hydrate is precipitated.

Qualitative Separation of Cobalt and Nickel.

(This process requires much practice for its successful performance, and need not be attempted by pupils whose studies are restricted to medicine or pharmacy.)

The foregoing reaction serves for the separation of nickel from cobalt. On adding excess of potassium cyanide to a very slightly acidified solution containing the two metals, and well boiling, a solution of potassium cobalticyanide and double nickel and potassium cyanide results. On boiling with excess of hydrochloric acid the nickel salt is converted into chloride, and this, with the potassium cobalticyanide, gives a milky-looking precipitate of nickel cobalticyanide ($\text{Ni}_3\text{Co}_2\text{Cy}_{12}$), insoluble in the acid. On then adding excess of potash the nickel cobalticyanide is decomposed, nickel hydrate remaining as a green flocculent precipitate and potassium cobalticyanide going into solution. After filtering off the nickel, cobalt is detected in the filtrate by evaporating to dryness and testing the residue with borax in the blowpipe flame.

Other Reactions between a nickel solution and various reagents give, in many cases, insoluble precipitates which, from their green color, are occasionally useful in distinguishing nickel from allied elements.

CHROMIUM.

Symbol, Cr. Atomic weight, 52.5.

Source.—The chief ore of chromium is chrome ironstone (a compound of the metallic oxides FeO , Cr_2O_3), occurring chiefly in the United States and Sweden. In constitution it seems to resemble magnetic iron ore (FeO , Fe_2O_3). The metal may be isolated by aid of sodium.

Preparation of Red Potassium Chromate.—On roasting the powdered ore with potassium carbonate and nitre, yellow potassium chromate (K_2CrO_4) is obtained; the mass, treated with acid, yields red or bichromate (K_2CrO_4 , CrO_3) (*Potassii Bichromas*, U. S. P.), the old dichromate or bichromate of potash; from this other chromates are prepared, and by reduction, as presently explained, the salts of chromium itself. The yellow and orange lead chromates are used as pigments.

Notes.—Red potassium chromate is a somewhat abnormal salt, containing, possibly, neutral chromate associated with chromic anhydride, and hence termed *potassium anhydrochromate* (K_2CrO_4 , CrO_3), or *potassium pyrochromate* ($\text{K}_2\text{Cr}_2\text{O}_7$). The value of chromates as chemical reagents is alluded to in connection with chromate of barium (pp. 105–117). Heated strongly in a crucible, red potassium chromate splits up into yellow chromate, glistening chromium oxide, and oxygen. Red ammonium chromate by heat yields several times its volume of bluish-green chromium oxide, water, and nitrogen: $(\text{NH}_4)_2\text{CrO}_4$, $\text{CrO}_3 = \text{Cr}_2\text{O}_3 + 4\text{H}_2\text{O} + \text{N}_2$.

Quantivalence.—Chromium stands in close chemical relation to iron, aluminium, and manganese. Its atom is sexivalent if the formula of the fluoride (CrF_6) be correct. Like iron and aluminium, it is trivalent, as seen in chromic chloride (Cr_2Cl_6), but sometimes exerts only bivalent activity, as in chromous chloride (CrCl_2).

Passage of Chromium from the Acidulous to the Basylous Side of Salts.—Through an acidified solution of red potassium chromate pass sulphuretted hydrogen; sulphur is deposited, and a green salt of chromium remains in solution—chloride (Cr_2Cl_6) if hydrochloric acid be used, and sulphate ($\text{Cr}_2\text{3SO}_4$) if sulphuric be the acid employed. Boil the liquid to expel excess of sulphuretted hydrogen, filter, and reserve the solution for subsequent experiments. (For an equation explanatory of this reaction see p. 239). Alcohol, sugar, or almost any substance which is tolerably liable to oxidation will answer as well as sulphuretted hydrogen.

Chromium Sulphate ($\text{Cr}_2\text{3SO}_4$), like aluminium sulphate ($\text{Al}_2\text{3SO}_4$), unites with alkali-metal sulphates to form *alums*, which resemble common alum both in crystalline form and in structure; they are of a purple color.


REACTIONS.

Chromium as Chromic Acid or other Chromate.—This is the

state in which chromium will usually be met with, the most common salt being the red chromate of potassium or bichromate. Mix four volumes of a cold saturated aqueous solution of red potassium chromate with five of oil of vitriol; on cooling, *chromic anhydride* or *anhydrous chromic acid* (CrO_3) (*Acidum Chromicum*, U. S. P.) separates in crimson needles. After well draining, the crystals may be freed from adhering sulphuric acid by washing once or twice with nitric acid: the latter may be removed by passing dried and slightly warmed air through a tube containing the crystals. It may also be freed from sulphuric acid by one or two recrystallizations. In contact with moisture chromic anhydride takes up water and forms solution of true chromic acid (H_2CrO_4), 1 part of the anhydride and 3 of water forming the *Liquor Acidi Chromici*, B. P. Chromic anhydride is a powerfully corrosive oxidizing agent; it melts between 356° and 374° F., and at a higher temperature decomposes, yielding chromium oxide and oxygen; it oxidizes organic matter with great violence, spontaneous ignition sometimes resulting.

The oxygen in chromic acid and other chromates, and in manganates, permanganates, black manganese oxide, and puce-colored lead oxide, is in a physically different state to that in hydrogen peroxide, barium peroxide, and similar compounds. On bringing chromic acid or the above acidified solution of red potassium chromate into contact with solution of hydrogen peroxide a strong effervescence of oxygen ensues. According to Schönbein and Brodie, the oxygen in chromic acid is in the negative or ozonic state, while that of hydrogen peroxide is in the positive or so-called antozonic condition. Both are equally active, but neutralize each other, forming neutral or ordinary oxygen.

In the analytical examination of solutions containing chromates the chromium will always come out in the state of green chromic hydrate along with ferric hydrate and alumina, the prior treatment by sulphuretted hydrogen reducing the chromium in the molecule to the lower state, thus:



Chromium having been found in a solution, its condition as chromate may be ascertained by applying to the original solution salts of barium, mercury, lead, and silver. (See the various paragraphs relating to those metals.)

Ba_2NO_3 gives yellow BaCrO_4 with chromates.

HgNO_3 " red Hg_2CrO_4 "

AgNO_3 " red Ag_2CrO_4 "

AgNO_3 " red $\text{Ag}_2\text{CrO}_4, \text{CrO}_3$ with bichromates.

$\text{Pb}_2\text{C}_2\text{H}_3\text{O}_2$ " yellow PbCrO_4 with both.

Barium nitrate does not completely precipitate bichromates, barium bichromate being soluble in water; barium chromate is insoluble in water or acetic acid, but soluble in hydrochloric or nitric acid. Mercurous nitrate does not wholly precipitate bichromates: mercuric nitrate or chloride only partially precipitates chromates, and does not precipitate bichromates. The mercurous chromate is insoluble or nearly so in diluted nitric acid. The silver chromates are soluble in acids and alkalies. Lead acetate precipitates chromates and bichromates, acetic acid being set free in the latter case.

A delicate reaction for dry chromates will be found in the formation of *chlorochromic anhydride* (CrO_2Cl_2). A small portion of the chromate is placed in a test-tube with a fragment of dry sodium chloride and a drop or two of sulphuric acid, and the mixture heated; red irritating fumes of chlorochromic anhydride are evolved and condense in dark-red drops on the side of the tube.

Larger quantities are obtained by the same reaction, the operation being conducted in a retort with thoroughly dry materials, for the compound is decomposed by water. It may be regarded as chromic anhydride in every molecule of which an atom of oxygen is displaced by an equivalent quantity (two atoms) of chlorine. It is not used in medicine, but is of interest to the chemical student as being an illustration of a class of similar bodies—*chloro-acidulous* or *chloro-anhydro compounds*. The reaction is also occasionally serviceable for the detection of chlorides.

ANALYTICAL REACTIONS OF CHROMIUM SALTS (TESTS).

First Analytical Reaction.—To solution of a salt of chromium (chloride, sulphate, or chrome alum) add ammonium sulphhydrate; a bulky green precipitate (chromic hydrate, $\text{Cr}_2\text{6HO}$), containing a large quantity of water (seven molecules, $7\text{H}_2\text{O}$), is precipitated.



Second Analytical Reaction.—To solution of a chromium salt add ammonia; green chromic hydrate is precipitated, insoluble in excess.

Third Analytical Reaction.—To solution of a chromium salt add solution of potash or soda, drop by drop; green chromic hydrate is precipitated. Add excess of the fixed alkali; the precipitate is dissolved. Well boil the solution; the green chromic hydrate is reprecipitated.

Iron, chromium, and aluminium salts, chemically so alike, may be separated by this reaction. Ferric hydrate is insoluble in solutions of the fixed alkalies, cold or hot; chromium hydrate soluble in cold, but not in hot; aluminium hydrate in both. To a solution contain-

ing all three metals, therefore, add potash or soda, stir, and filter; the iron is thrown out: boil the filtrate and filter; the chromium is thrown out; neutralize the latter filtrate by acid, and then add ammonia; the aluminium is thrown out. Note, however, that ferric hydrate will prevent chromium hydrate being dissolved by potash or soda if the ferric hydrate is in considerable excess. Before concluding that chromium is entirely absent, the fourth reaction should be performed. Iron, chromium, and aluminium hydrates are insoluble in ammonia, and may therefore easily be separated from the hydrates of the somewhat analogous metals, zinc, cobalt, nickel, and manganese.

Fourth Analytical Reaction.—Add a salt of chromium (either of the above precipitates of chromic hydrate or the dry residue of the evaporation of a few drops of a solution of a chromium salt) to a few grains of nitre and sodium carbonate on platinum-foil, and fuse the mixture in the blowpipe flame; a yellow mass (potassium and sodium chromate, KNaCrO_4) is formed. Dissolve the mass in water, add acetic acid to decompose excess of carbonate, and apply the reagents for chromates. This is a delicate and useful reaction if carefully performed.

TIN.

Symbol, Sn. Atomic weight, 118.

Source.—The chief ore of tin is stannic oxide (SnO_2), occurring in veins under the name of *tin-stone* or in alluvial deposits as *stream-tin*. The oldest mines are those of Cornwall. Much tin is now imported from Australia.

Preparation.—The metal is obtained by reducing the roasted and washed ore by charcoal or anthracite* coal at a high temperature, and is purified by slowly heating, when the pure tin, fusing first, is run off, a somewhat less fusible alloy of tin, with small quantities of arsenum, copper, iron, or lead, remaining. The latter is known as *block-tin*; the former, heated till brittle and then hammered or let fall from a height, splits into prismatic fragments resembling starch or columnar basalt, and is named *dropped* or *grain-tin*. Good tin emits a crackling noise in bending, termed the “cry” of tin, caused by the friction of its crystalline particles on each other.

Uses.—Tin is an important constituent of such alloys as pewter, Britannia metal, solder, speculum-metal, bell-metal, gun-metal, and bronze. It is very ductile, and may be rolled into plates or leaves, known as *tin-foil*, varying from $\frac{1}{250}$ to $\frac{1}{1000}$ of an inch in thickness. Common tin-foil, however, usually contains a large proportion of lead. The reflecting surface of *looking-glasses* was formerly always

* *Anthracite* (from $\alpha\nu\theta\rho\alpha\acute{\varsigma}$, *anthrax*, a burning coal), or *stone coal*, differs from the ordinary *bituminous* or *caking coal* in containing less volatile matter, and therefore in burning without flame. It gives a higher temperature, and, from its non-caking properties is, in furnace operations, more manageable than bituminous coal.

an amalgam of tin and mercury, produced by carefully sliding a plate of glass over a sheet of tin-foil on which mercury had been rubbed and then excess of mercury been poured; but pure silver, deposited from a solution, is now largely employed. *Pins* are made of brass wire on which tin is deposited. *Tin-plate*, of which common utensils are made, is iron alloyed with tin by dipping the acid-cleansed sheet into vessels of melted tin covered with melted zinc chloride in the one case and oil in the other—fluids which, by dissolving any trace of oxide or by preventing oxidation, enable the tin more completely to alloy with the iron. *Tin tacks* are in reality tinned iron tacks: a tin nail would be too soft to drive into wood. Tin may be granulated by melting and triturating briskly in a hot mortar, by shaking melted tin in a box on the inner sides of which chalk has been rubbed, or in thin little bells or corrugated fragments (Granulated Tin, B. P.), by melting in a ladle and, immediately it is fluid, pouring from the height of a few feet into water. Powdered tin has been used medicinally as a mechanical irritant to promote expulsion of worms. The hairs of the pod of *kiwach* (Hindustani) or *cowhage* (*Mucuna pruriens*) (P. I.) is almost the only other medicine (excluding diluents and dentifrices) which acts in such a directly mechanical manner.

The chemical position of tin among the metals is close to that of arsenum and antimony. Its atom is quadrivalent, Sn'''' , and bivalent, Sn'' . The two classes of salts are termed stannic and stannous respectively.

REACTIONS HAVING SYNTHETICAL INTEREST.

Stannous Chloride, or Lower Chloride of Tin.

First Synthetical Reaction.—Warm a fragment of tin with hydrochloric acid; hydrogen escapes and solution of stannous chloride (SnCl_2 , perhaps Sn_2Cl_4) is formed. It may be retained for future experiments.

1 ounce of tin dissolved in 3 fluidounces of hydrochloric acid and 1 of water, and the resulting solution diluted to 5 fluidounces, constitutes the "Solution of Stannous Chloride," B. P.

Solid Stannous Chloride.—By evaporation of the above solution stannous chloride is obtainable in crystals ($\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$). It is a powerful reducing agent, even a dilute solution precipitating gold, silver, and mercury from their solutions, converting ferric and cupric into ferrous and cuprous salts, and partially deoxidizing arsenic, manganic, and chromic acids. It absorbs oxygen from the air, and is decomposed when added to a large quantity of water unless some acid be present. It is used as a mordant in dyeing and calico-printing.

Stannic Chloride, or Perchloride of Tin.


Second Synthetical Reaction.—Through a portion of the solution of the stannous chloride of the previous reaction pass chlorine gas; solution of stannic chloride (SnCl_4) is formed. Or

add hydrochloric acid to the stannous solution, boil, and in a fume-chamber slowly drop in nitric acid until no more fumes are evolved; again stannic chloride results. Reserve the solutions for subsequent experiments.

Stannic Oxide, or Anhydride and Stannates.

Third Synthetical Reaction.—Boil a fragment of tin with nitric acid, evaporate to dryness, and strongly calcine the residue; light buff-tinted stannic anhydride (SnO_2) is produced. Heat the stannic anhydride with excess of solid caustic potash or soda; stannate of the alkali-metal (K_2SnO_3 or Na_2SnO_3) results. Dissolve the stannate in water and add hydrochloric acid; white gelatinous *stannic acid* (H_2SnO_3) is precipitated. Stannic acid is also obtained on adding an alkali to solution of stannic chloride; it is soluble in excess of acid or alkali.

The product of the action of nitric acid on tin is also an acid, but from its insolubility in hydrochloric and other acids is different from ordinary stannic acid. It is termed *metastannic acid* (from *meta*, *meta*, beyond), and its molecule probably has a composition expressed by the formula $\text{H}_{10}\text{Sn}_5\text{O}_{15}$ (*vide* Index, "Isomerism"). It is also produced on gently heating stannic acid:



Metastannates have the general formula $\text{M}'_2\text{H}_8\text{Sn}_5\text{O}_{15}$.

Both acids yield buff-colored stannic oxide or anhydride (SnO_2) when strongly heated. The latter is employed in polishing plate under the name of *putty powder*. *Sodium stannate* ($\text{Na}_2\text{SnO}_3 \cdot 4\text{H}_2\text{O}$) is used as a mordant by dyers and calico-printers under the name of *tin prepare-liquor*.

REACTIONS HAVING ANALYTICAL INTEREST (TESTS).

Stannous or Stannic Salts.—Heat any solid compound of tin with a mixture of potassium cyanide and sodium carbonate on charcoal by the inner flame of the blowpipe. Hard globules of tin separate, having, when cut by a knife, characteristic brightness and whiteness.

Stannous Salts.

First Analytical Reaction.—Through a diluted solution of a stannous salt (stannous chloride, for example) pass sulphuretted hydrogen gas; a brown precipitate (stannous sulphide, SnS) results. Pour off the supernatant liquid, add ammonia to the moist precipitate (to neutralize acid), and, lastly, yellow ammonium sulphhydrate solution; the precipitate is dissolved.

Aqueous solution of ammonium sulphhydrate becomes yellow when a day or two old, and then contains excess of sulphur, some of that element having become displaced by oxygen absorbed from the air; hence in the above reaction the stannous sulphide (SnS), in dissolving, becomes stannic sulphide (SnS_2); for the latter is precipitated on decomposing the alkaline liquid by an acid.

Second Analytical Reaction.—To solution of a stannous salt add solution of potash or soda: a white precipitate falls (stannous hydrate, $\text{Sn}2\text{HO}$). Add excess of the alkali; the precipitate dissolves. Boil the solution; some of the tin is reprecipitated (as blackish stannous oxide, SnO). Ammonia gives a similar white precipitate, insoluble in excess. The alkaline carbonates do the same, carbonic acid gas escaping.

Stannic Salts.

Third Analytical Reaction.—Through solution of a stannic salt (stannic chloride, for example) pass sulphuretted hydrogen gas; a yellow precipitate results (stannic sulphide, SnS_2). Pour off the supernatant liquid, and to the moist precipitate add ammonia (to neutralize acid), and then ammonium sulphhydrate; the precipitate dissolves.

Note.—In this reaction the presence of much hydrochloric acid must be avoided; the formation of the precipitate is also facilitated if the solution be warmed. Stannic sulphide, like arsenum and antimony sulphides, dissolves in a solution of alkaline sulphide or sulphhydrate, with formation of definite crystallizable *sulphostannates* ($\text{M}'_2\text{SnS}_3$).

Anhydrous stannic sulphide, prepared by sublimation, has a yellow or orange lustrous appearance, and is known as *mosaic gold*. It was formerly used by decorators as *bronzing-powder*, but the latter is now commonly powdered *bronze-leaf*.

Fourth Analytical Reaction.—To solution of a stannic salt add potash or soda; a white precipitate appears (stannic acid, H_2SnO_3). Add excess of the alkali; the precipitate dissolves. Boil the mixture; no reprecipitation occurs—a fact enabling stannic to be distinguished from stannous salts.

Ammonia gives a similar precipitate slowly soluble in excess. The fixed alkali-metal carbonates do the same, carbonic acid gas escaping; after a time the stannic salt is again deposited, probably as stannate of the alkali-metal. Ammonium carbonate and all the bicarbonates give a precipitate of stannic acid insoluble in excess.

Antidotes.—In cases of poisoning by tin salts (dyers' tin-liquor, *e.g.*) solution of ammonium carbonate should be given, and white of egg is also said to form an insoluble precipitate. Vomiting should be induced, and the stomach-pump or stomach-siphon applied.

GOLD.

Symbol, Au. Atomic weight, 196.85.

Source.—Gold occurs in the free state in nature, occasionally in nodules or *nuggets*, but commonly in a finer state of division termed *gold-dust*.

Preparation.—Gold is separated from sand, crushed quartz, or other earthy matter with which it may be associated by agitation with water, when the gold, from its relatively greater specific gravity, falls to the bottom of the vessel first, the lighter mineral matter running off with the water. From this rich sand the gold is dissolved out by mercury, the amalgam filtered and afterward distilled, when the mercury volatilizes and gold remains. The amalgamation may be facilitated by the use of sodium, as already described in treating of silver. From even the poorest ores gold may be dissolved by solution of potassium cyanide. (See Faraday on gold-leaf, 1857.) $4\text{KCy} + \text{Au}_2 + \text{O} + \text{H}_2\text{O} = 2\text{AuKCy}_2 + 2\text{KOH}$. Elkington deposited the metal by the aid of a battery, of which the anode or positive pole was a plate of gold, in 1840, deposition taking place at the cathode or negative pole (*ava, ana*, upward or onward; *kata, kata*, downward; *ὁδος, odos*, way). The gold solution was made, as now, from the gold chloride and potassium cyanide, potassium chloride and auro-cyanide (AuKCy_2) being formed, but Elkington in 1857 made it direct from a gold anode with a small cathode in solution of potassium cyanide.

Pure gold is too soft for general use as a circulating medium. *Gold coin* is an alloy of copper and gold, that of Great Britain containing 1 of the former to 11 of the latter, or $8\frac{1}{2}$ per cent. of copper, that of France, Germany, and the United States about 10 per cent. *Jewellers' gold* varies in quality, every 24 parts containing 18, 15, 12, or 9 parts of gold, the alloys being technically termed 18, 15, 12, or 9 *carat fine*, the reckoning being in the old "parts per 24," instead of the more usual parts per cent. Articles made of the better qualities are usually stamped by authority. Trinkets of inferior intrinsic worth are commonly thinly coated with pure gold by electro-deposition or otherwise. The so-called *mystery gold* is an alloy of about 1 part platinum and 2 parts copper with a little silver. It resists the action of strong nitric acid. The action of aqua regia and then ammonia reveals its cupric character. *Gold-leaf* is nearly pure gold passed between rollers till it is about $\frac{1}{800}$ of an inch in thickness, and then hammered between sheets of animal membrane, termed gold-beaters' skin, and calf-skin vellum, till it is $\frac{1}{160000}$ or $\frac{1}{200000}$ of an inch in thickness. It may even be hammered till 280,000 leaves would be required to form a pile an inch thick.

Gold Coinage.—The weight of gold is expressed in England in ounces troy and decimal parts of an ounce, and the metal is always taken to be of standard fineness (11 gold and 1 alloy) unless otherwise described. The degree of "fineness" of gold, as ascertained by assay, is expressed decimally, fine pure gold ("gold free from metallic impurities," B. P.) being taken as unity, or 1.000. Thus gold of British standard is said to be 0.9166 fine, of French standard 0.900 fine. The legal weight of the sovereign is 0.2568

ounce of standard gold; or 123.274 grains, the weight coming from 1 pound of standard gold (5760 grains) being coined into 44½ guineas. Gold coins are legal tender to any amount, provided that the weight of each sovereign does not fall below 122.5 grains, or in the case of a half-sovereign 61.125 grains; these are the "least current" weights of the coins.


In the United States the weight of gold is expressed in ounces troy and decimal parts thereof. 900 parts of gold are alloyed with 100 parts of copper. The weight of the eagle or ten-dollar gold piece is 258 grains and 900 fine.

Note.—In analysis gold comes out among the sulphides of metals precipitated by sulphuretted hydrogen, and of those sulphides, it, like the sulphides of tin, antimony, and arsenum, is soluble in yellow ammonium sulphhydrate solution.

Quantivalence.—Gold is trivalent (Au''') in the auri- or auric compounds, univalent (Au') in the auro- or aurous salts.

REACTIONS.

Synthetical Reaction.—Place a fragment of gold (*e. g.* gold-leaf) in 10 or 20 drops of aqua regia (a mixture of 3 parts of nitric and 4 or 5 of hydrochloric acid), and set aside in a warm place; solution of gold perchloride or auric chloride (AuCl_3) results. Evaporate nearly to dryness to remove most of the excess of acid, dilute with water, and retain the solution for subsequent experiments. 60 grains of gold treated thus, and the resulting chloride dissolved in 5 ounces of distilled water constitutes "Solution of Perchloride of Gold," B. P. The salt itself is very deliquescent. A compound of gold and sodium chlorides, in molecular proportions, crystallizes readily and is more stable.



This reaction has analytical interest also, for in examining a substance suspected to be or to contain metallic gold, solution would have to be effected in the above way before reagents could be applied. Gold is insoluble in hydrochloric, nitric, and the weaker acids.

Gold and Sodium Chloride (*Auri et Sodii Chloridum*, U. S. P.) is "a mixture of equal parts, by weight, of dry auric chloride and sodium chloride."

ANALYTICAL REACTIONS (TESTS).

First Analytical Reaction.—Through a few drops of solution of an auric salt (the chloride, AuCl_3 , is the only convenient one) pass sulphuretted hydrogen; a brown precipitate results (auric sulphide, Au_2S_3). Filter, wash, and add yellow ammonium sulphhydrate solution; the precipitate dissolves.

Second Analytical Reaction.—To solution of a salt of gold add a ferrous salt and set the tube aside; metallic gold, having its characteristic lustrous appearance, is precipitated, a ferric

salt remaining in solution. Oxalic acid also and most free metals precipitate the gold.

This is a convenient way of preparing pure gold—or *fine gold*, as it is termed—or of working up the gold residues of laboratory operations. The precipitate, after boiling with hydrochloric acid, washing and drying, may be obtained in a button by mixing with an equal weight of borax or acid potassium sulphate and fusing in a good furnace.

Third Analytical Reaction.—Add a few drops of dilute solutions of stannous and stannic chloride to a considerable quantity of distilled water; pour the liquid, a small quantity at a time, into a very dilute solution of auric chloride (AuCl_3), well stirring; the mixture assumes a purple tint, and flocks of a precipitate known as the *Purple of Cassius* (from the name of the discoverer, M. Cassius) are produced.

The same compound is formed on immersing a piece of tin-foil in solution of auric chloride; it is said to be a mixture of auric, aurous, stannic, and stannous oxides. It is the coloring agent in the finer varieties of ruby glass.

PLATINUM.

Symbol, Pt. Atomic weight, 194.4.

Source.—Platinum, like gold, occurs in nature in the free state, the chief sources of supply being Mexico, Brazil, and Siberia. It is separated from the soil by washing.

Uses.—The chief use of platinum is in the construction of foil, wire, crucibles, spatulas, capsules, evaporating-dishes, and stills for the use of the chemical analyst or manufacturer. It is tolerably hard, fusible with very great difficulty, not dissolved by hydrochloric, nitric, or sulphuric acid, and only slightly affected by alkaline substances. It is attacked by aqua regia, with production of platinum perchloride or platonic chloride ($\text{PtCl}_4 \cdot 5\text{H}_2\text{O}$). It forms fusible alloys with lead and other metals, and with phosphorus a phosphide which easily melts. Neither of these substances, therefore, nor mixtures which may yield them, should be heated in platinum vessels. Hammered or chased, not drawn, vessels are the most durable. They are best cleaned by aid of a little fine water-worn (not “sharp”) sea-sand. They should not be very suddenly heated or very suddenly cooled. They should only be heated by the outer portions of flames, exposure to strongly heated carboniferous or siliconiferous surfaces being avoided, for at high temperatures platinum has a tendency to unite with carbon or silicon.

The chemical position of platinum among the elements is close to that of gold. Its atom is quadrivalent ($\text{Pt}^{'''}$) in some compounds, in others apparently bivalent (Pt''). The higher salts are termed *platinic*, the lower *platinous*.

The specific gravity of platinum is 21.5, and that of *iridium*, an allied metal, 22.4.

REACTIONS.

Platinic Chloride, or Platinum Perchloride.

Synthetical Reaction.—Place a fragment of platinum in a little aqua regia and set the vessel aside in a warm place, adding more acid from time to time if necessary; solution of platinum perchloride (PtCl_4) results. Evaporate the solution to remove excess of acid, and complete the desiccation over a water-bath. Dissolve the residue in water, and retain the solution for subsequent experiments and as a reagent for the precipitation of salts either of potassium or ammonium. 1 part of platinum treated in the above manner, and the resulting chloride dissolved in 20 parts of water, constitutes "Solution of Perchloride of Platinum," B. P. and U. S. P., or 1.7 grms. of pure platinic chloride ($\text{PtCl}_4 \cdot 5\text{H}_2\text{O}$) dissolved in 20 cc. of distilled water gives "Platinic Chloride Test-solution," U. S. P.

This reaction has analytical interest, for, in testing a substance suspected to contain metallic platinum, solution would have to be thus effected before reagents could be applied.

ANALYTICAL REACTIONS (TESTS).

First Analytical Reaction.—Through a few drops of a solution of a platinic salt (PtCl_4 is the only convenient one), to which an equal quantity of solution of sodium chloride has been added, pass sulphuretted hydrogen; a dark-brown precipitate results (platinic sulphide, PtS_2). Filter, wash, and add ammonium sulphhydrate; the precipitate dissolves.

If sodium chloride be not present in the above reaction, the precipitated sulphide will contain platinous chloride, and may detonate if heated.

Second Analytical Reaction.—Add excess of sodium carbonate and some sugar to solution of platinum perchloride, and boil; a black precipitate (metallic platinum) falls.


Platinum black (B. P.) is the name of this precipitate. It possesses in a high degree a quality common to many substances, but largely possessed by platinum—namely, that of absorbing or *occluding* gases. In its ordinary state, after well washing and drying, it absorbs from the air and retains many times its bulk of oxygen. A drop of ether or alcohol placed on it is rapidly oxidized, the platinum becoming hot. This action may be prettily shown by pouring a few drops of ether into a beaker (one having portions of the top and sides broken off answers best), loosely covering the vessel with a card, and suspending within the beaker a platinum wire, one end being attached to the card by passing through its centre, the other terminating in a short coil or helix near the surface of the ether: on

now warming the helix in a flame and then rapidly introducing it into the beaker, it will become red hot and continue to glow. In this experiment partial combustion goes on between the ether vapor and the concentrated oxygen of the air, the products of the oxidation revealing themselves by their odor.

Third Analytical Reaction.—To solution of platinum perchloride add solution of ammonium chloride; a yellow granular precipitate (the double chloride, $\text{PtCl}_4 \cdot 2\text{NH}_4\text{Cl}$) falls. When slowly formed in dilute solutions the precipitate is obtained in minute orange prisms.

Potassium chloride (KCl) gives a similar precipitate ($\text{PtCl}_4 \cdot 2\text{KCl}$). Platinic chloride having been stated to be a test for potassium and ammonium salts, the reader is prepared to find that potassium and ammonium salts are tests for platinic salts. The double sodium compound ($\text{PtCl}_4 \cdot 2\text{NaCl}$) is soluble in water.

Collect the precipitate, dry, and heat in a small crucible; it is decomposed, and metal in a finely-divided gray state (*spongy platinum*) remains.



Heat decomposes the salt of potassium into $\text{Pt} + 2\text{KCl} + \text{Cl}_2$, the chlorine escaping and the potassium chloride remaining with the platinum.

In working up the platinum residues of laboratory operations the mixture should be dried, burnt, boiled successively with hydrochloric acid, water, nitric acid, water, then dissolved in aqua regia, excess of acid removed by evaporation, ammonium chloride added, the precipitate washed with water, dried, ignited, and the resulting spongy platinum retained or converted into perchloride for use as a reagent for alkali-metals. It is by such processes that the native platinum is treated to free it from the rare metals palladium, rhodium, osmium, ruthenium, and iridium. The spongy platinum is converted into the massive condition by a refinement on the blacksmith's process of *welding* (German *wellen*, to join), or by fusing in a flame of pure oxygen and hydrogen gases—the oxyhydrogen blowpipe.

Occlusion by Spongy Platinum.—Spongy platinum has great power of occlusion. A small piece held in a jet of hydrogen causes ignition of the gas, owing to the close approximation of particles of oxygen (from the air) and hydrogen. Dobereiner's lamp is constructed on this principle, the apparatus being essentially a vessel in which hydrogen is generated by the action of diluted sulphuric acid on zinc, and a cage for holding the spongy platinum.

CADMIUM.

Symbol, Cd. Atomic weight, 112.

In most of its chemical relations cadmium resembles zinc. In nature it occurs chiefly as an occasional constituent of the ore of that metal. In distilling zinc containing cadmium the latter, being the

more volatile, passes over first. In analytical operations cadmium, unlike zinc, comes down among the metals precipitated by sulphuretted hydrogen; that is, its sulphide is insoluble in diluted hydrochloric acid, while zinc sulphide is soluble. It is a white malleable metal, nearly as volatile as mercury. Sp. gr. 8.7.

Beyond the occasional employment of the sulphide as a pigment (*jaune brillant*), and the iodide in photography, cadmium and its salts are but little used. The atom of cadmium is bivalent (Cd'').

REACTIONS.

Cadmium Iodide.

First Synthetical Reaction.—Digest together, in a flask, metallic cadmium, warm water, and iodine until the color of the iodine disappears; solution of cadmium iodide (CdI_2) remains. Pearly micaceous crystals may be obtained on evaporating the solution.

This salt is employed with other iodides in iodizing collodion for photographic use. It readily melts, and is soluble in water or spirit, the solution reddening litmus.

Cadmium Sulphate.

Second Synthetical Reaction.—Dissolve cadmium in nitric acid; pour the resulting solution of cadmium nitrate ($\text{Cd}2\text{NO}_3$) into a solution of sodium carbonate; dissolve the precipitate of cadmium carbonate (CdCO_3) in dilute sulphuric acid, separate and crystallize. Cadmium sulphate (CdSO_4) is a white crystalline salt soluble in water.

ANALYTICAL REACTIONS (TESTS).

First Analytical Reaction.—Through solution of a cadmium salt (CdI_2 or CdCl_2) pass sulphuretted hydrogen; a yellow precipitate (cadmium sulphide, CdS) falls, resembling in appearance arsenous, arsenic, and stannic sulphides. Add ammonium sulphhydrate; the precipitate, unlike the sulphides just mentioned, does not dissolve.

Cadmium and copper sulphides may be separated by solution of potassium cyanide, in which copper sulphide is soluble and cadmium sulphide insoluble.

Second Analytical Reaction.—To a cadmium solution add solution of potash; a white precipitate results (cadmium hydrate, $\text{Cd}2\text{HO}$), insoluble in excess of the potash.

Zinc hydrate ($\text{Zn}2\text{HO}$), precipitated under similar circumstances, is soluble in solution of potash; the filtrate from the cadmium hydrate may therefore be tested for any zinc occurring as an impurity by applying the appropriate reagent—ammonium sulphhydrate.

Before the blowpipe flame, on charcoal, cadmium salts give a brown deposit (cadmium oxide, CdO).

BISMUTH.

Symbol, Bi. Atomic weight, 208.

Source.—Bismuth occurs in the metallic state in nature. It is freed from adherent quartz, etc. by simply heating, when the metal melts, runs off, and is collected in appropriate vessels (*Bismuthum*, B. P.). It is also met with in combination with other elements. Bismuth is grayish-white, with a distinct pinkish tinge.

Purification.—Arsenum may be removed from melted bismuth by a rod of iron, iron arsenide rising to the surface of the mass: antimony, by stirring in some bismuth oxide, when antimony oxide separates. Other metals in bismuth, especially copper, are converted into sulphides, while bismuth is not affected, on fusing the crude metal with about 5 per cent. of potassium cyanide and 2 per cent. of sulphur, the whole being well stirred for a quarter of an hour with a clay rod (stem of a tobacco-pipe). On pouring off the metal from the flux, and melting and stirring it with about 5 per cent. of a mixture of potassium and sodium carbonates, sulphur and traces of other impurities are removed, and the metal is obtained pure (*Bismuthum Purificatum*, B. P.).—*Tamm*.

Uses.—Beyond the employment of some of its compounds in medicine bismuth is but little used. Melted bismuth expands considerably on solidifying, and hence is valuable in taking sharp impressions of dies. It is a constituent of some kinds of type-metal and of pewter solder.

The position of bismuth among the metals is close to that of arsenum and antimony. Its atom is trivalent (Bi''') and rarely quinquivalent (Bi^v).

REACTIONS HAVING SYNTHETICAL INTEREST.

Bismuth Nitrate, or Nitrate of Bismuth.

First Synthetical Reaction.—To a few drops of nitric acid and an equal quantity of water, in a test-tube, add a little powdered bismuth, heating the mixture if necessary; nitric oxide (NO) escapes, and solution of *bismuth nitrate* (Bi₃NO₃) results.



The solution evaporated gives crystals (Bi₃NO₃·5H₂O), any arsenum which the bismuth might contain remaining in the mother-liquor.

To make bismuth *nitrate* and other salts on a larger scale, 2 ounces of the metal, in small fragments, are gradually added to a mixture of 4 fluidounces of nitric acid and 3 of water, and when effervescence (due to escape of nitric oxide) has ceased the mixture is heated for ten minutes, poured off from any insoluble matter, evaporated to 2 fluidounces to remove excess of acid, and then either

set aside for *crystals* to form, or poured into half a gallon of water to form bismuth *oxynitrate*, or into a solution of 6 ounces of ammonium carbonate in a quart of water to form the *oxycarbonate*, as described in the following reactions.

The precipitates should be washed with cold water, and dried at a temperature not exceeding 150° F. Exposed in the moist state to 212° F. (100° C.) for any length of time, they undergo slight decomposition.

Bismuth Subnitrate or Oxynitrate.

Second Synthetical Reaction.—Pour some of the above solution of nitrate into a considerable quantity of water; decomposition occurs, and bismuth oxynitrate (BiONO_3) in a hydrous state ($\text{BiONO}_3 \cdot \text{H}_2\text{O}$) (*Bismuthi Subnitratis*, *Subnitrate of Bismuth*, or *Bismuth Subnitrate*, U. S. P.), is precipitated:



Filter, and test the filtrate for bismuth by adding excess of sodium carbonate; a precipitate shows that some bismuth remains in solution. The following equation, therefore, probably more nearly represents the decomposition:



Decomposition of bismuth nitrate by water is the process of the Pharmacopœia for the preparation of bismuth oxynitrate or "subnitrate" for use in medicine. For this purpose the original metal must contain no arsenum. In manufacturing the compound, therefore, before pouring the solution of nitrate into water the liquid should be tested for arsenum by one of the hydrogen tests; if that element be present, the solution must be evaporated, and only the deposited crystals be used in the preparation of the oxynitrate. For on pouring an arsenical solution of bismuth nitrate into water the arsenum is not wholly removed in the supernatant liquid, unless the oxynitrate be redissolved and reprecipitated several times, according to the amount of arsenum present.

Bismuth subnitrate is gradually decomposed by solution of alkaline carbonates; also by the bicarbonates, with production of carbonic acid gas, bismuth oxycarbonate and nitrate of the alkali-metal being formed. It is sometimes administered in the form of a lozenge (*Trochisci Bismuthi*, B. P.).


Oxysalts of Bismuth.—It will be noticed that the formula for bismuth subnitrate (BiNO_3) does not accord with that of other nitrates, the characteristic elements of which are NO_3 . Analogy would seem to indicate, however, that the fourth atom of oxygen has different functions to the three in the NO_3 ; for on pouring solution of bismuth chloride (BiCl_3) into water, oxychloride is produced (BiOCl) (a white powder used as a cosmetic, "pearl-white" (*Blanc de Perle*), also in enamels and in some varieties of sealing-wax). The

bromide (BiBr_3) and iodide (BiI_3), similarly treated, yield oxybromide (BiOBr) and oxyiodide (BiOI). The subnitrate (BiNO_3) is therefore probably an analogous compound, an oxynitrate (BiONO_3). The sulphate (Bi_2SO_4) also decomposes when placed in water, giving what may be termed an oxysulphate, the formula of which is $\text{Bi}_2\text{O}_2\text{SO}_4$.

It is difficult to prove whether or not the water in the "subnitrate" or hydrous bismuth oxynitrate ($\text{BiONO}_3 \cdot \text{H}_2\text{O}$) is an integral part of the salt. If it is, the compound is simply bismuth hydratonitrate ($\text{BiNO}_3 \cdot 2\text{H}_2\text{O}$).


Bismuth Oxide.

Third Synthetical Reaction.—Boil bismuth subnitrate with solution of soda for a few minutes; it is converted into yellowish bismuth oxide (Bi_2O_3) (*Bismuthi Oxidum*, B. P.).



Bismuth Subcarbonate or Oxycarbonate.

Fourth Synthetical Reaction.—To solution of bismuth nitrate add solution of ammonium carbonate; a white precipitate of hydrous oxycarbonate ($2\text{Bi}_2\text{O}_2\text{CO}_3 \cdot \text{H}_2\text{O}$) (*Bismuthi Subcarbonas*, or *Bismuth Subcarbonate*, U. S. P.) falls.



This compound may be regarded as similar in constitution to the oxysalts just described. In Bi_2CO_3 one scarcely recognizes the characteristic elements of carbonates, but, considering the preparation to be an oxycarbonate ($\text{Bi}_2\text{O}_2\text{CO}_3$), its relations to carbonates and oxides are evident. These subsalts may all be viewed as normal bismuth salts in one molecule of which an atom of oxygen displaces an equivalent proportion of other acidulous atoms or radicals:

Chloride	Bi_3Cl	Oxychloride . .	BiOCl
Bromide	Bi_3Br	Oxybromide . .	BiOBr
Iodide	Bi_3I	Oxyiodide . .	BiOI
Nitrate	Bi_3NO_3	Oxynitrate . .	BiONO_3
Sulphate	Bi_2SO_4	Oxysulphate . .	$\text{Bi}_2\text{O}_2\text{SO}_4$
Carbonate (unknown) .	Bi_2CO_3	Oxycarbonate . .	$\text{Bi}_2\text{O}_2\text{CO}_3$


They may be viewed, in short as salts in process of conversion to oxide; continue the substitution further, and each yields bismuth oxide (Bi_2O_3). They are also regarded as salts of a hypothetical radical bismuthyl (BiO).

Bismuth Citrate, or Citrate of Bismuth.


Fifth Synthetical Reaction.—Heat 10 parts of bismuth oxynitrate, 7 of citric acid crystals, and 30 to 40 of water together

for a few minutes, until a drop of the mixture forms a clear solution with ammonia-water. Dilute the crystalline mass with eight to ten times its volume of water, and set aside for a short time to let the citrate deposit; decant the clear liquid. Wash the crystalline sediment three or four times in a similar manner, drain and dry, either on a water-bath or by mere exposure. The yield is $13\frac{2}{3}$ parts, showing that the salt is anhydrous and that its formula is $\text{BiC}_6\text{H}_5\text{O}_7$ (Rother). This is the *Bismuthi Citras*, U. S. P.

Sixth Synthetical Reaction.—To bismuth citrate, rubbed to a paste with a very little water, add sufficient solution of ammonia to dissolve the citrate. The product, when of a strength of 800 grains of citrate in a pint, is the official Solution of Citrate of Bismuth and Ammonium (*Liquor Bismuthi et Ammonii Citratis*, B. P.), shortly termed *Liquor Bismuthi*. This solution, evaporated to the consistence of syrup, spread on glass and dried at 100°F ., yields scales of the solid salt (*Bismuthi et Ammonii Citras*, U. S. P.). In constitution it is possibly ordinary ammonium citrate, in which three atoms of hydrogen are displaced by one of bismuth. (For similar salts see p. 156.)



Hydrogen-ammonium citrate.



Bismuth-ammonium citrate.

REACTIONS HAVING ANALYTICAL INTEREST (TESTS).

First Analytical Reaction.—Through solution of a bismuth salt (a slightly acid solution of nitrate, for example) pass sulphuretted hydrogen; a black precipitate (bismuth sulphide, Bi_2S_3) falls. Add ammonia (to neutralize acid), and then ammonium sulphhydrate; the precipitate, unlike As_2S_3 and Sb_2S_3 , is insoluble.

Second Analytical Reaction.—Concentrate almost any acid solution of a bismuth salt and pour into much water (containing sodium chloride); a white precipitate results.

This reaction is characteristic of bismuth salts: it has already been amply explained. The oxychloride is specially insoluble, and is distinguished from that of antimony by being insoluble in solution of tartaric acid.

Third Analytical Reaction.—To a solution of bismuth salt add an alkali; a white precipitate results (bismuth hydrate, $\text{Bi}(\text{OH})_3$), insoluble in excess and becoming yellowish on boiling.

Fourth Analytical Reaction.—A small quantity of the following reagent, including both supernatant liquid and precipitated yellow scales, is transferred to a test-tube, and gradually heated till solution takes place. Any liquid containing, or supposed to contain, bismuth is then added, and the whole allowed to cool. The separated scales will show a distinct change in color from the original yellow to dark orange or crimson, according to the quantity of bismuth present.

The reagent may be prepared by adding to a boiling solution of lead acetate, containing about $\frac{1}{2}$ grain to the ounce, a few drops of acetic acid and solution of potassium iodide in considerable excess. On cooling, lead iodide is deposited in the characteristic yellow crystalline plates or scales.

Test for Calcium Phosphate in Bismuth Salts.—Dissolve the powder in nitric acid, add about twice its weight of citric acid and sufficient ammonia to give decided alkalinity; then boil, keeping the mixture faintly alkaline with ammonia: bismuth remains in solution and calcium phosphate is precipitated.

Official (B. P.) Tests for Other Impurities in Bismuth or its Salts.—Dissolve in nitric acid; concentrate and set aside for crystals of bismuth nitrate to form; pour off the mother-liquor, which will contain any impurities in a concentrated form. If this mother-liquor be evaporated with hydrochloric acid until all the nitric acid is dissipated, a little of the product yields no evidence of arsenum on being examined by the hydrogen test, commonly known as Marsh's test; no blue coloration on adding water and excess of ammonia (copper), and no precipitate on filtering and saturating the ammoniacal filtrate with nitric acid (silver); no white precipitate with diluted sulphuric acid (lead); no red or black precipitate with sodium sulphite (tellurium or selenium); and no blue precipitate with potassium ferrocyanide (iron).

THE READER IS AGAIN ADVISED TO TRACE OUT THE EXACT NATURE OF EACH OF THE FOREGOING REACTIONS, CHIEFLY BY AID OF EQUATIONS OR DIAGRAMS.

QUESTIONS AND EXERCISES.

Enumerate the fifteen metals salts of which are frequently employed in pharmacy.—Mention the twelve rare metals interesting to pharmacists.—Name the sources and official compounds of lithium.—Explain the formation of lithium citrate.—What is the strength of *Liquor Lithiæ Effervescens*?—On what chemical hypothesis are lithium compounds administered to gouty patients?—Describe the relation of lithium to

other metals.—What is the chief test for lithium?—Write a paragraph on strontium, its natural compounds, chemical relations, technical applications, and tests.—What are the formula and properties of cerium oxalate?—Name the commonest ore of manganese and give an equation descriptive of its reaction with hydrochloric acid.—Explain the formation of potassium permanganate, employing diagrams or equations.—How do the potassium manganates act as disinfectants?—What are the chief tests for manganese?—What are the chief uses of the compounds of cobalt?—How is cobalt analytically distinguished from nickel?—Mention applications of nickel in the arts.—What is the general color of nickel salts?—State the method of preparation of red potassium chromate.—Give the formulæ of red and yellow potassium chromates.—How is red potassium chromate obtained?—Describe the action of sulphuretted hydrogen on acidified solutions of chromates.—What is the formula of chrome alum?—Mention the chief tests for the chromic radical and for chromium.—How would you detect iron, chromium, and aluminium in a solution?—Define *tin-stone*, *stream-tin*, *block-tin*, *grain-tin*, *tin plate*.—Describe the position occupied by tin in relation to other metals.—What is the difference between stannic acid and metastannic acid?—State the applications of tin in the arts.—Mention the chief tests for stannous and stannic salts.—Name the best antidote in cases of poisoning by tin solutions.—How is gold-dust separated from the earthy matter with which it is naturally associated?—How much pure gold do English coin and jewellers' gold contain?—State the average thickness of gold-leaf.—What is the weight of a sovereign?—Explain the term "fineness" as applied to gold. What effect is produced on gold by hydrochloric, nitric, and nitrohydrochloric acids, respectively?—By what reagents may metallic gold be precipitated from solution?—How is "purple of Cassius" prepared?—Whence is platinum obtained?—Why are platinum utensils peculiarly adapted for use in chemical laboratories?—How is platinum perchloride prepared?—Name the chief tests for platinum.—What is "platinum black"?—Describe an experiment demonstrative of the large amount of attraction for gases possessed by metallic platinum.—How is "spongy platinum" produced?—By what process may platinum be recovered from residues?—What is *occlusion* in Chemistry?—In what condition does cadmium occur in nature?—By what process may cadmium iodide be prepared?—Mention the chief test for cadmium.—Distinguish cadmium sulphide from sulphides of similar color.—How is cadmium separated from zinc?—How does bismuth occur in nature?—What is the quantivalence of bismuth?—Write down equations descriptive of the action of nitric acid on bismuth, and water on bismuth nitrate.—How may arsenum be excluded from bismuth salts?—Give a diagram of the official process for bismuth carbonate.—Write formulæ showing the accordance in molecular constitution of the official subnitrate and carbonate with the other salts of bismuth, and with ordinary nitrates and carbonates.—How is *Liquor Bismuthi et Ammonii Citratis* prepared?—Mention the tests for bismuth.

The Analytical Classification of Metals.

Practical Analysis.

Bismuth is the last of the metals whose synthetical or analytical relations are of general interest. The position of the rarer among the common metals, and the influence which either has on the other during the manipulations of analysis, will now be considered. These objects will be best accomplished, and a more intimate

TABLE OF SHORT DIRECTIONS FOR THE ANALYSIS OF AN AQUEOUS OR ONLY SLIGHTLY ACID SOLUTION OF ORDINARY SALTS OF ONE OF THE COMMON AND RARER METALS HITHERTO CONSIDERED.
Add hydrochloric acid.

Precipitate Hg(ous) Pb Ag. Collect, wash, and add NH_4HO . Hg ppt., blackened. Pb ppt., still white. Ag ppt., dissolved. Sb and Bi may also be precipitated by HCl , but are dissolved on adding more HCl .			
IF HCl gave no precipitate the metal is still in the liquid; pass H_2S through the solution.			
Precipitate Cd Cu Hg(ic) Pb Bi As Sb Sn Au Pt. Collect wash, add NH_4HS . Insoluble. Soluble. Cd, yellow. As(ous & ic) Cu Hg(ic) } black. Sn(ic) } yellow. Pb } Sp, orange. Bi } Au Pt } black.		IF H_2S gave no precipitate add NH_4Cl , NH_4HO , and NH_4HS . Precipitate Zn Mn Co Ni Al Fe Cr. Zn } white. Al } Cr, green. Mn, skin-tint. Ni } black. Co } Fe }	
Apply special tests for each to the original solution. For these, see the previous pages.		IF NH_4HS gave no precipitate, add $(\text{NH}_4)_2\text{CO}_3$. Precipitate Ba Sr Ca. Collect, wash, dissolve in HCl , H_3O_2 , add K_2CrO_4 . Ppt. Ba. Sol. Sr Ca. Add dil. H_2SO_4 . Ppt. Sol. Sr.—Ca.	
Test specially for each in original solution. See previous pages.		IF $(\text{NH}_4)_2\text{CO}_3$ gave no precipitate, add $(\text{NH}_4)_2\text{HAsO}_4$. Ppt. Mg. If no precipitate, test original solution in flame on loop of Pt wire. L, crimson. Na, yellow. K, violet. If neither, test orig. sol. for NH_4 .	

acquaintance with all the metals be obtained, by analyzing, or studying the methods of analyzing, solutions containing one or more metallic salts.

Of the foregoing tables, the first includes directions for the analysis of an aqueous or only slightly acid solution containing but one salt of any of the metals hitherto considered. Here the color of the precipitate or precipitates afforded by a metal under given circumstances must largely be relied on in attempting the detection of the various elements.

The long table is intended as a chart for the analysis of solutions containing salts of more than one of the common and rarer metals. It is a compilation from the foregoing reactions—an extension of the scheme for the analysis of salts of the ordinary metals. It often may be altered or varied in arrangement to suit the requirements of the analyst.

OUTLINE OF THE ANNEXED ANALYTICAL TABLES.

HCl	H ₂ S	NH ₄ HS	(NH ₄) ₂ CO ₃	(NH ₄) ₂ HSO ₄	
Hg (as mercurous salt)	Cd	Zn	Ba	Mg	K
	Cu	Mn*	Sr		Na
Pb (partially)	Hg (as mercuric salt)	Co	Ca		NH ₄
	Ph (entirely)	Ni			
Ag	Bi	Al			
	As (as arsenous or arsenic salt)	Fe			L
	Sb	Cr			
	Sn (as stannous or stannic salt)				
	Au				
	Pt				

Note.—The laboratory student should practice the examination of aqueous solutions of salts of the above metals until he is able to analyze with facility and accuracy.

* See p. 234.

§ OR ONLY SLIGHTLY ACID SOLUTION OF ORDINARY

Sr Mg Li K Na NH₄.

Filtrate
Cr Ba Ca Sr Mg Li K Na NH₄
O, NH₄HS, warm gently and filter.

Filtrate

Ba Ca Sr Mg Li K Na NH₄

Add (NH₄)₂CO₃, warm, filter.

	Precipitate	Filtrate
	Ba Sr Ca.	Mg Li K Na NH ₄ .
51. pass H ₂ S,	Collect, wash, dissolve in HC ₂ H ₃ O ₂ , add excess of K ₂ CrO ₄ , filter.	Add (NH ₄) ₂ HAsO ₄ , stir, filter

ate	Ppt.	Filtrate	Ppt.	Filtrate
Ni.	Ba.	Sr Ca.	Mg.	Li K Na NH ₄
and a little	Yellow.	Add dilute H ₂ SO ₄ ,	White	Evaporate to small
HO. filter.		let stand, filter.		bulk. Add NH ₄ HO.

precipitate on Ni. ve in HCl, proceed as ed on page 237.	Ppt. Sr. White	Filt. Ca. Add NH_4HO and $(\text{NH}_4)_2\text{C}_2\text{O}_4$. White ppt.	Ppt. Li. See p. 262.	Filtrate K Na NH_4 . Evaporate. ignite, dissolve. K by PtCl_4 . Na by flame. NH_4 in original solution.
---	----------------------	--	-------------------------------	---

See also p. 261.

[To face page 258.]

TABLE OF SHORT DIRECTIONS FOR APPLYING SOME OF THE FOREGOING ANALYTICAL REACTIONS TO THE ANALYSIS OF AN AQUEOUS OR ONLY SLIGHTLY ACID SOLUTION OF ORDINARY SALTS OF ANY OF THE COMMON OR RARER METALS OF GENERAL INTEREST.

Add hydrochloric acid.

Precipitate
Hg ions Pb Az
Wash, boil with water, filter

Filtrate
Cd Cu Hg(c) Pb Bi Asous(c) Sb Sn(cous(c) Zn Mn Co Ni Al Ferous(c) Cr Ba Ca Sr Mg Li K Na NH₄
Pass H₂S through the liquid, filter.

Precipitate
Hg ions Pb
Wash, add NH₄HO, filter
Filtrate
Pb
Add H₂SO₄,
White ppt.

Precipitate
Cd Cu Hg(c) Pb Bi As Sb Sn.
Collect, wash, digest in NH₄HS, filter.

Filtrate
Zn Mn Co Ni Al Fe Cr Ba Ca Sr Mg Li K Na NH₄
Add NH₄Cl, NH₄HO, NH₄HS, warm gently and filter.

Ppt.
Hg
Add HNO₃
Black.

Precipitate
Cd Cu Hg(c) Pb Bi.
Wash, boil in HNO₃, filter.

Filtrate
As Sb Sn.
Add dilute HCl, filter, drain well, add strong
HCl, boil, dilute slightly, filter.

Precipitate
Zn Mn Co Ni Al Fe Cr.
Collect, wash, dissolve in HCl with a few drops of HNO₃,
boil, add NH₄HO, stir, filter

Filtrate
Ba Ca Sr Mg Li K Na NH₄
Add NH₄CO₃, warm filter.

See also p. 259.

Ppt.
Hg
ic.
Black.
Confirm
by Cu
test in
original
solid
then

Filtrate
Cd Cu Pb Bi.
Add NH₄HO, filter.

Insol
As.
Yellow.
Confirm
by Eleit-
mann's
test.

Filtrate
Sn Sb.
Pour into H-apparatus.

Precipitate
Fe Al Cr.
Wash, dry, fuse on foil with
Na₂CO₃ and KNO₃, boil in
water and filter.

Filtrate
Zn Mn Co Ni.
Acidify with HC₂H₃O₂, pass H₂S,
filter.

Precipitate
Ba Sr Ca.
Collect, wash, dissolve in
HC₂H₃O₂, add excess of
K₂CrO₄, filter.

Filtrate
Mg Li K Na NH₄
Add NH₄HO, stir, filter

Precipitate
Pb Bi.
Wash, add a few
drops HNO₃,
dilute, filter.

Filtrate
Cd Cu.
Add K₂CrO₄ and
H₂S.

Sn
remains on Zn.
Dissolve in HCl
and apply tests.

See also pp. 259-260

Au and Pt are specially sought when necessary.

Ppt.
Bi.
White

Filt.
Pb.
Dilute,
add
H₂SO₄,
set aside,
White
ppt.

Ppt.
Cd.
Yel-
low.

Filt.
Cu.
Acidify
with
HC₂H₃O₂,
Brown
ppt.

See also p. 260.

Residue
Fe₂O₃,
brown.
Test
original
solution
for
ferrous-
or ferric
state.

Filtrate.
If yellow, Cr pre-
sent. Divide in
two parts.

Sol.
Al.
Add
NH₄Cl
and
warm,
White
ppt.

Sol.
Cr.
Add
HC₂H₃O₂
and excess
of AgNO₃;
red ppt.
Or boil
with H₂SO₄
and spirit.
Green
solution.

See also p. 261.

* Test this also for Mn, by
Crum's process, p. 261.

Sol.
Mn.
Add
NH₄HO
and
NH₄HS.
Pink,
turning
brown.

Filt.
Zn.
Add
NH₄HS
White
ppt.

Search
also for
Mn in
the
Fe Al Cr
ppt.

Precipitate
Zn Co Ni.
Boil with H₂L and a little
HNO₃; add KHO, filter.

Filt.
Zn.
Add
NH₄HS
White
ppt.

Precipitate
Co Ni.
Dissolve in HCl,
and proceed as
directed on page
237.

Ppt.
Ba.
Yellow.

Filtrate
Sr Ca.
Add dilute H₂SO₄,
let stand, filter.

Ppt.
Mg.
White

Filtrate
Li K Na NH₄
Evaporate to small
bulk. Add NH₄HO.

See also p. 261.

Ppt.
Sr.
White

Filt.
Ca.
Add
NH₄HO and
(NH₄)₂C₂O₄
White ppt.

Ppt.
Li.
See
p. 262.

Filtrate
K Na NH₄.
Evaporate,
ignite, dissolve
K by PtCl₄,
Na by flame.
NH₄ in original
solution.

That on p. 258 is a mere outline of the other two tables. It gives the position of the metals in relation to each other, and will much aid the memory in recollecting that relation.

The analysis of solutions containing only one metal will, as already stated, serve to impress the memory with the characteristic tests for the various metals and other radicals and familiarize the mind with chemical principles. Medical students and junior pharmaceutical students seldom have time to go farther than this. More thorough analytical and general chemical knowledge is only acquired by working on such mixtures of bodies as are met with in actual practice, beginning with solutions which may contain any or all of the members of a group (see previous pages), then examining solutions containing more than one group, and finally analyzing liquids in which are dissolved several salts of any of the common or rarer metals.

The author cannot too strongly recommend students thoroughly to master the art of analysis, not only on account of its direct value, but because its practice enables the learner rapidly and soundly to acquire a good knowledge of Chemistry and greatly to improve his general mental faculties.

GENERAL AND SPECIAL MEMORANDA RELATING TO THE PRECEDING ANALYTICAL TABLES.

General Memoranda.

These charts are constructed for the analysis of salts more or less soluble in water.—The student has still to learn how substances insoluble in water are to be brought into a state of solution; but, once dissolved, their analysis is effected by the same scheme as that just given. The tables, especially the longer, folded one, may therefore be regarded as fairly representing the method by which metallic constituents of chemical substances are separated from each other and recognized.—The methods of isolation of the complementary constituent of the salt (the reactions of non-metals and acidulous radicals) will form the next object of practical study.


The general memoranda given in connection with the first table (p. 224) are equally applicable to this extended second table, and should again carefully be read through.

Special Memoranda.

The hydrochloric-acid precipitate may at first include some antimony and bismuth as oxychlorides, readily dissolved, however, by excess of acid.—If either of these elements be present, the washings of the precipitate will probably be milky; in that case add a few drops of hydrochloric acid, which will clear the liquid and make way for the application of the test for lead.

The sulphuretted-hydrogen precipitate may be white, in which case it is nothing but sulphur; for, as already indicated, ferric salts are reduced to ferrous, and chromates to the lower salts of chromium,

by sulphuretted hydrogen, finely divided, and, therefore, whitish sulphur being deposited :



But the precipitate may also be colored, or even white, when only lead or mercury is present, through an insufficiency of sulphuretted hydrogen having produced oxysulphide or hydrato-sulphide, etc. The gas should be passed through the liquid until, even after well shaking, the latter smells strongly of sulphuretted hydrogen.

The portion of the sulphuretted-hydrogen precipitate dissolved by ammonium sulphhydrate may include a trace of copper, copper sulphide being not altogether insoluble in ammonium sulphhydrate.

—On adding hydrochloric acid to the ammonium sulphhydrate solution, a white precipitate of sulphur only may be precipitated, the ammonium sulphhydrate nearly always containing free sulphur.

—Strong hydrochloric acid does not readily dissolve small quantities of antimony sulphide out of much arsenum sulphide; and, on the other hand, the strong hydrochloric acid takes into solution a small quantity of arsenum sulphide if much antimony sulphide be present. The precipitates or the original solutions should therefore be examined by the other (hydrogen) tests for these elements if doubt exist concerning the presence or absence of either. Tin remains in the hydrogen bottle in the metallic state, deposited as a black powder on the zinc used in the experiment. The contents of the bottle are turned out into a dish, ebullition continued until evolution of hydrogen ceases, and the zinc is taken up by the excess of sulphuric acid employed; any tin is then filtered out, washed, dissolved in a few drops of hydrochloric acid, and the liquid tested for tin by the usual reagents.—Tin may be detected in the mixed tin, arsenum, and antimony sulphides by the blowpipe reaction (p. 243).

The portion of the sulphuretted-hydrogen precipitate not dissolved by the ammonium sulphhydrate may leave a yellow semi-fused globule of sulphur on boiling with nitric acid. This globule may be black, not only from presence of mercuric sulphide, but also from enclosed particles of other sulphides protected by the sulphur from the action of the acid. It may also contain lead sulphate, produced by the action of nitric acid on lead sulphide. In cases of doubt the mass must be removed from the liquid, boiled with nitric acid till dissolved, the solution evaporated to remove excess of acid, and the residue examined; but usually it may be disregarded.—Before testing for bismuth any considerable excess of acid should be removed by evaporation, and the residual liquid should be freely diluted. If no precipitate (bismuth oxynitrate) appear, ammonium chloride solution may be added, bismuth oxychloride more readily forming than even oxynitrate.—Or any nitric acid or sulphuric acid having been neutralized by ammonia, hydrochloric acid is added, and then potassium iodide; a rich orange color results if bismuth be present.—Bismuth may also be detected in the mixed precipitated bismuth and lead hydrates obtained in the ordinary course of analysis by dissolving a portion of the precipitate in acetic

acid, and adding the liquid to the hot lead iodide solution mentioned in the reactions for bismuth (p. 255).—In testing for lead by sulphuric acid the liquid should be diluted and set aside for some time.

Mercury may also be isolated by digesting the sulphuretted-hydrogen precipitate in sodium sulphhydrate instead of ammonium sulphhydrate. The arsenum, antimony, tin, and mercury sulphides are thus dissolved out. The mixture is then filtered, excess of hydrochloric acid added to the filtrate, and the precipitated sulphides collected on a filter, washed, and digested in ammonium sulphhydrate; mercury sulphide remains insoluble, while the arsenum, antimony, and tin sulphides are dissolved. By this method copper also appears in its right place only, copper sulphide being insoluble in sodium sulphhydrate. The other metals are then separated in the usual way.

The ammonium-sulphhydrate precipitate may, if the original solution was acid, contain barium, calcium, and magnesium phosphates, oxalates, silicates, and borates. These will subsequently come out with the iron, and, being white, give the iron precipitate a light-colored appearance; their examination must be conducted separately by a method described subsequently in connection with the treatment of substances insoluble in water.—The precipitate containing aluminium, iron, and chromium hydrates often contains some manganese. This manganese may be detected by washing the hydrates to remove all trace of chlorides, boiling with nitric acid, adding either puce-colored lead oxide or red lead, and setting the vessel aside; if manganese be present, a red or purple liquid is produced.—Nickel sulphide is not easily removed by filtration (*vide* p. 236) until most of the excess of ammonium sulphhydrate has been dissipated by prolonged ebullition.

The ammonium-carbonate precipitate may not contain the whole of the barium, strontium, and calcium in the mixture unless free ammonia be present; for the carbonates of those metals are soluble in water charged with carbonic acid. If, therefore, the liquid is not distinctly ammoniacal, solution of ammonia should be added.—Neither ammonium carbonate nor hydrate wholly precipitates magnesian salts; and, as partial precipitation is undesirable, a solvent, in the form of an alkaline salt (ammonium chloride), if not already in the liquid, should be added.—In the chart opposite p. 258 strontium is ordered to be separated from calcium by adding to the acetic solution diluted sulphuric acid. The latter, unless extremely dilute, may precipitate calcium. Any such loss of calcium is in itself of little consequence, because enough calcium sulphate remains in the filtrate to afford a calcium reaction when ammonia and ammonium oxalate are subsequently added. But the calcium precipitated by the sulphuric acid may be wrongly set down as strontium. Therefore test a little of the acetic solution for strontium by an aqueous solution of calcium sulphate, when, if no precipitate falls after setting aside for several minutes, strontium may be regarded as absent. If a precipitate occurs, strontium is present; the rest of the acetic solution is then tested for calcium, as directed in the chart, the final testing by ammonium oxalate being, of course, preceded by the addition of ammonia.—Barium may be overlooked

if oxidation happens to have converted any sulphur into sulphuric acid.

Lithium.—The search for lithium may usually be omitted. Should a precipitate, supposed to be due to lithium, be obtained, it must be tested in a flame (= scarlet tint), as a little magnesium not infrequently shows itself under similar circumstances.

Spectral Analysis.—If present only in minute proportions, the lithium may also remain with the alkalies; it can then only be detected by physical analysis (by a prism) of the light emitted from a tinged flame—by, in short, an instrument termed a spectroscope. Such a method of examination is called spectral analysis, a subject of much interest and of no great difficulty, but scarcely within the range of medical or pharmaceutical students; it will be described briefly in connection with the methods of analyzing solid substances.

QUESTIONS AND EXERCISES.

Describe a general method of analysis by which the metal of a single salt in a solution could be quickly detected.—Give illustrations of black, white, light-pink, yellow, and orange sulphides.—Mention the group-reagents usually employed in analysis.—Under what circumstances may a hydrochloric precipitate contain antimony or bismuth?—If a sulphuretted-hydrogen precipitate is white, what substances are indicated?—Give processes for the qualitative analysis of liquids containing the following substances: *a.* Arsenum and cadmium. *b.* Bismuth and antimony. *c.* Ferrous and ferric salts. *d.* Aluminium, iron, and chromium. *e.* Arsenum, antimony, and tin. *f.* Lead and strontium. *g.* Iron, sodium, and arsenum. *h.* Mercury, manganese, and magnesium. *i.* Zinc, manganese, nickel, and cobalt. *j.* Barium, strontium, and calcium.

THE ACIDULOUS RADICALS.

Introduction.—The twenty-seven radicals which have up to this point mainly occupied attention are (admitting ammonium, NH_4) metals, and they have been almost exclusively studied not in the free state, but in the condition in which they exist in salts. Moreover, these metals have been treated as if they formed the more important constituent, the stronger part, the foundation or *base* of salts. Attention has been continuously directed to the metallic or *basyulous* side of salts. And, indeed, there is still one more basyulous radical which must be mentioned, though it is usually supposed to play only a subordinate part in medicine—hydrogen. Unlike the salts of most metals, those of hydrogen (the so-called *acids*) are never, in medicine or the arts generally, professedly used for the sake of their hydrogen, but always for the other part of the salt, the *acidulous* side. And it is not for their basyulous radical that these hydrogen salts are now commended to notice,* but in order to

* It must not be forgotten that the commonest salt of any radical whatsoever is a salt of hydrogen, hydrogen oxide (H_2O), or hydrogen hydrate (HHO), *water*. In the reactions already performed the value of this compound has been constantly recognized, both for its hydrogen

study, under the most favorable circumstances, those acidulous groupings which have continually presented themselves in operations on salts, but which were for the time of secondary importance. These *acidulous radicals* may now be treated as the primary object of attention; and there is no better way of doing so than by operating on their compounds with hydrogen, the relatively inferior medicinal importance of which element, as compared with potassium, iron, and other basylous radicals, will serve to give the desired prominence to the *acidulous radicals* in question.*

Common Acids.—These salts of hydrogen (hydrogen easily displaceable, wholly or, in certain cases, in part, by ordinary metals) are the common, sharp, sour bodies termed *acids* (from the Latin root *acies*, an edge). The following table includes the formulæ and usual names of the most important: others will be noticed subsequently. A few of those mentioned are unstable or somewhat rare; in such cases a common metallic salt containing the acidulous radical may be used for reactions:

HCl	hydrochloric acid.	H ₂ S	sulphydric acid.†
HBr	hydrobromic acid.	H ₂ SO ₃	sulphurous acid.
HI	hydroiodic acid.	H ₂ SO ₄	sulphuric acid.
HCN	hydrocyanic acid.	H ₂ CO ₃ ?	carbonic acid.
HNO ₃	nitric acid.	H ₂ C ₂ O ₄	oxalic acid.
HC ₂ H ₃ O ₂	acetic acid.†	H ₂ C ₄ H ₄ O ₆	tartaric acid.
H ₃ C ₆ H ₅ O ₇	citric acid.	H ₃ PO ₄	phosphoric acid.
		H ₃ BO ₃	boric acid.

The usual names are here retained for these acids, but in studying their chemistry and chemical relations to other salts they are usefully

and for its oxygen, but most of all as the vehicle or medium by which nearly all other atoms are enabled to come into that contact with each other without which their existence would be almost useless; for some atoms are like some animals—out of water they are as inactive as fishes. It is true that both fishes and salts have usually to be removed from water to be utilized by man, but before they can be assimilated, either as food or as medicine, they must again seek the agency of water—in becoming dissolved.

* Actually, it is as difficult to determine the relative importance of the different atoms or groups of atoms in a molecule as it is of the different parts or members of an animal or vegetable, the different units or societies in a community, the different planets or solar systems of the universe; nay, the different pieces or parts of an engine or the different pigments or portions of a picture: *l'union fait la force*.

† The hydrogen on the acidulous side must not be confounded with the basylous hydrogen in all these hydrogen salts or acids; the two perform different functions. Hydrogen in the acidulous portion is like the hydrogen in the basylous radical ammonium: it has combined with other atoms to form a group which plays more or less the part of an elementary radical. Cobalt, chromium, iron, platinum, etc. resemble hydrogen in this respect, in often uniting with other atoms to form definite acidulous radicals in which the usual basylous character of the metals has for the time disappeared. In *hydrides* (p. 124) hydrogen itself is an acidulous radical.

‡ Synonyms: hydrosulphuric acid and sulphuretted hydrogen.

spoken of by such more purely chemical names, as (for hydrochloric acid) chloride of hydrogen or hydrogen chloride, hydrogen nitrate, and so on—hydric or hydrogen sulphate, dihydric tartrate, trihydric phosphate, etc.

A prominent point of difference will at one be noticed between the basylous radicals met with up to the present time and the acidulous groupings included in the above tabular list. The former are nearly all elements, ammonium only being a compound; the latter are mostly compounds, chlorine, bromine, iodine, and sulphur being the only elements. This difference will not, however, be so apparent when the chemistry of alcohols, ethers, and such bodies has been mastered, for they may be regarded as salts of *compound* basylous radicals.

Rarer Acids.—The above acids contain the only acidulous groupings that commonly present themselves in analysis or in pharmaceutical operations. There are, however, several other acids (such as hypochlorous, nitrous, hypophosphorous, valerianic, benzoic, gallic, tannic, uric, hyposulphurous, hydroferrocyanic, hydroferricyanic, lactic, etc.) with which it is desirable to be more or less familiar; reactions concerning these will therefore be described. Arsenous, arsenic, stannic, manganic, and chromic acids have already been treated of in connection with the metals they contain; in practical analysis these acids always become sufficiently altered for their metals to come out among the basylous radicals.

Quantivalence.—A glance at the foregoing table is sufficient to show the quantivalence of the acidulous radicals there mentioned. The first six are clearly univalent; then follow six bivalent, leaving three trivalent.

These all combine with equivalent amounts of basylous radicals to form various salts; hence they may be termed monobasylous, dibasylous, and tribasylous radicals. The acids themselves were formerly spoken of as monobasic, bibasic, and tribasic respectively, or monobasic and polybasic, in reference to the amount of *base* (hydrates or oxides) they could decompose; but the terms are no longer definite, and hence but little used in mineral chemistry.

Antidotes.—The antidotes in case of poisoning by the strong acids will obviously be non-corrosive alkaline substances, as soap and water, magnesia, and common washing "soda" or other carbonates. Vinegar, lemon-juice, and weak and non-corrosive acids would be the appropriate antidotes to caustic alkalies.

Analysis.—The practical study of the acidulous side of salts will occupy far less time than the basylous. Salts will then be briefly examined as a whole.

Caution.—Once more: it is only for convenience in the division of chemistry for systematic study that salts may be considered to contain basylous and acidulous radicals, or separate sides, so to speak; for we possess no absolute knowledge of the internal arrangement of the atoms (admitting that there are such things) in the molecule of the salt. We only know that certain groups of atoms may be transferred from compound to compound in mass (that is, without apparent decomposition); hence the assumption

that these groups are radicals. A salt is probably a whole, having no such sides as those mentioned.

QUESTIONS AND EXERCISES.

Mention the basylous radical of acids.—Give illustrations of univalent, bivalent, and trivalent acidulous radicals or monobasylous, dibasylous, and tribasylous radicals.—What is the difference between an elementary and a compound acidulous radical?—Name the grounds on which salts may be assumed to contain basylous and acidulous radicals.

HYDROCHLORIC ACID AND OTHER CHLORIDES.

Formula, HCl. Molecular weight,* 36.5.


The acidulous radical of hydrochloric acid and of other chlorides is the element chlorine (Cl). It occurs in nature chiefly as sodium chloride (NaCl), either solid under the name *rock-salt*, mines of which are not uncommon, or in solution in the water of all seas. Common table-salt is more or less pure sodium chloride in minute crystals. Chlorine, like hydrogen, is univalent (Cl'); its atomic weight is 35.5. Its molecule is symbolized thus, Cl₂.

REACTIONS.

Hydrochloric Acid.

Synonyms.—Hydrogen Chloride; Chloride of Hydrogen; Muriatic Acid; Chlorhydric Acid.

First Synthetical Reaction.—To a few fragments of sodium chloride in a test-tube or a small flask add about an equal weight of sulphuric acid; colorless and invisible gaseous hydrochloric acid is evolved, a sodium sulphate remaining. Adapt to the mouth of the vessel, by a perforated cork, a piece of glass tubing bent to a right angle, heat the mixture, and convey the gas into a small bottle containing a little water; solution of hydrochloric acid results.




Hydrochloric Acid.—The product of this operation is the nearly colorless and very sour liquid commonly termed hydrochloric acid. When of certain given strengths (estimated by volumetric analysis), it forms *Acidum Hydrochloricum*, U. S. P. (*Muriatic Acid*), and *Acidum Hydrochloricum Dilutum*, U. S. P. The former has a specific gravity of about 1.163 (1.1578), and contains 31.9 per cent. of real acid; the latter, sp. gr. 1.050, with 10 per cent. of the real acid, is made by diluting 6 fluid parts of the strong acid with 13 of water.

* The weight of a molecule is the sum of the weights of its atoms.

The above process is that of the manufacturer, larger vessels being employed, and the gas being freed from any trace of sulphuric acid by washing. Other chlorides yield hydrochloric acid when heated with sulphuric acid, but sodium chloride is always used because cheap and common.

Fig. 37.



Preparation of Hydrochloric Acid.

Common yellow hydrochloric acid is a by-product in the manufacture of sodium carbonate from common salt by the process in which sodium chloride is first converted into sulphate, hydrochloric acid being liberated. This impure acid is liable to contain iron, arsenum, fixed salts, sulphuric acid, sulphurous acid, nitrous compounds, and chlorine.

The process for the preparation of hydrochloric acid is as follows: it may be carried out by the student with about one-twelfth of the quantities mentioned:


"Take of sodium chloride, dried, 48 ounces, sulphuric acid 44 fluidounces, water 36 fluidounces, distilled water 50 fluidounces. Pour the sulphuric acid slowly into 32 ounces of the water, and, when the mixture has cooled, add it to the sodium chloride previously introduced into a flask having the capacity of at least one gallon; connect the flask by corks and a bent glass tube with a three-necked wash-bottle, furnished with a safety-tube, and containing the remaining 4 ounces of the water (or let the flask-tube pass loosely through a wider tube fixed in the cork of the wash-bottle, as shown in Fig. 37): then, applying heat to the flask, conduct the disengaged gas through the wash-bottle into a second bottle containing the distilled water by means of a bent tube dipping about half an inch below the surface, and let the process be continued until the product measures 66 ounces or the liquid has acquired a specific gravity of 1.16. The bottle containing the distilled water must be kept cool during the whole operation."—*B. P.*

The modification of wash-bottle shown in the figure allows of the easy insertion or removal of a delivery tube. The wider tube there shown, or an ordinary tube-funnel, also acts as a *safety-tube* by

admitting the air the moment there is any tendency in the water in the receiver to be forced back on account of a too rapid absorption of the gas. The time of the student will be saved if hydrochloric acid already in stock be placed in the wash-bottle instead of water.

Invisible gaseous hydrochloric acid forms visible grayish-white fumes on coming into contact with air. This is due to combination with the moisture of the air. The intense greediness of hydrochloric gas and water for each other is strikingly demonstrated on opening a test-tube full of the gas under water; the latter rushes into and instantly fills the tube. If the water is tinged with blue litmus, the acid character of the gas is prettily shown at the same time. The test-tube, which should be perfectly dry, may be filled from the delivery-tube direct, for the gas is somewhat heavier than, and therefore readily displaces, air. The mouth may be closed by the thumb of the operator. At low temperatures hydrochloric acid and water form a crystalline compound, $\text{HCl} \cdot 2\text{H}_2\text{O}$.

Note.—The process includes the use of as much sulphuric acid as is necessary for the production of acid sodium sulphate (NaHSO_4), which remains in the generating vessel. A hot solution of this residue, neutralized by sodium carbonate, filtered and set aside, yields normal sodium sulphate (*Sodii Sulphas* or *Sodium Sulphate*, U. S. P., the old sulphate of soda, “Glauber’s salt”), in the form of transparent, oblique, efflorescent prisms ($\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$).



Chlorine.


Second Synthetical Reaction.—To some drops of hydrochloric acid (that is, the common aqueous solution of the gas) add a few grains of black manganese oxide, and warm the mixture; *chlorine*, the acidulous radical of all chlorides, is evolved, and may be recognized by its peculiar odor or irritating effect on the nose and air-passages.




Chlorine-water.—This is the process of the Pharmacopœia for the production of chlorine-water (*Aqua Chlori*, U. S. P.), the gas being first washed and then passed into water. 1 ounce of oxide to 6 fluidounces of acid diluted with 2 of water, and the gas passed through a wash-bottle containing about 2 ounces of water, yields enough chlorine to produce about $1\frac{1}{2}$ pints of chlorine-water. (On this small scale less than half the acid is utilized through incomplete decomposition of the materials, and especially through incomplete absorption of the chlorine gas.) Chlorine slowly decomposes water with production of hydrochloric acid and oxygen gas; hence for medicinal purposes the solution should be freshly prepared; it is best preserved in a green-glass well-stoppered bottle in a cool and dark place. At common temperatures (60°F .), if fresh and thoroughly saturated, chlorine-water contains more than twice (2.3) its

bulk of chlorine, or less than 1 per cent. (about 0.75) by weight. Chlorine passed into cold water yields crystals of hydrous chlorine ($\text{Cl}5\text{H}_2\text{O}$), and these, when heated under pressure, give an upper layer of chlorine-water and a lower layer of *liquid chlorine*.

Note.—To obtain the chlorine from other chlorides sulphuric acid, as well as black manganese oxide, must be added. Hydrochloric acid is first formed. The action described in the foregoing equation then goes on, except that half instead of the whole of the oxygen from the black oxide is employed for the removal of the hydrogen from the chlorine of the hydrochloric acid, the other half being taken up by the hydrogen of the sulphuric acid. Thus, assuming common salt to be the chloride used, the following equations may represent the supposed steps of the process:



or the whole may be included in one equation.



This reaction may occasionally have analytical interest, a very small quantity of combined chlorine being recognized by its means. But the following test is nearly always applicable for the detection of this element, and leaves nothing to be desired in point of delicacy.

ANALYTICAL REACTIONS (TESTS).

To a drop of hydrochloric acid, or to a dilute solution of any other chloride, add solution of silver nitrate; a white curdy precipitate (silver chloride, AgCl) falls. Pour off the supernatant liquid, add nitric acid, and boil; the precipitate does not dissolve. Pour off the acid and add dilute ammonia; the precipitate quickly dissolves. Neutralize the solution by an acid; the curdy precipitate again appears.

The formation of this white precipitate, its appearance, insolubility in boiling nitric acid, solubility in solution of ammonia or its carbonate, and reprecipitation by an acid, form abundant evidence of the presence of chlorine. Its occurrence as a chloride of a metal is determined by testing for the metal with the appropriate reagents; its occurrence as hydrochloric acid is considered to be indicated by the odor, if strong, and the sour taste, if weak, of the liquid, and the action of the liquid on blue litmus-paper, which, like other acids, it reddens. If hydrochloric acid be present in excessive quantity, it will, in addition to the above reactions, give rise to strong effervescence on the addition of a carbonate, a chloride being formed. The chlorine in insoluble chlorides, such as calomel, "white precipitate," etc., may be detected by boiling with alkali, filtering, acidulating the filtrate by nitric acid, and then adding the silver nitrate.

Antidotes.—In cases of poisoning by strong hydrochloric acid

solution of sodium carbonate (common washing-soda) or a mixture of magnesia and water may be administered as an antidote.

QUESTIONS AND EXERCISES.

The official hydrochloric acid contains 31.8 per cent. by weight of gas, and its specific gravity is 1.16: work out a sum showing what volume of it will be required, theoretically, to mix with black manganese oxide for the production of 1 gallon of chlorine-water, 1 fluidounce of which contains 2.66 grains of chlorine. *Ans.*, $5\frac{1}{2}$ fl. ozs. nearly (5.4)—Why does hydrochloric acid gas give visible fumes on coming into contact with air?—How much sodium chloride will be required to furnish 1 pound of chlorine?—Give the analytical reactions of chlorides. —What antidotes may be administered in cases of poisoning by hydrochloric acid?

HYDROBROMIC ACID AND OTHER BROMIDES.

Formula of Hydrobromic Acid, HBr. Molecular weight, 81.

BROMINE.—*Source, Preparation, and Properties.*—The acidulous radical of hydrobromic acid and other bromides is the element bromine, Br. (*Bromum*, U. S. P.). It occurs in nature chiefly as magnesium bromide ($MgBr_2$) in sea-water and certain saline springs, and is commonly prepared from the *bittern*, or residual liquors, of salt-works. It may be liberated from its compounds by the process for chlorine from chlorides—that is, by heating with black manganese oxide and sulphuric acid (see note on p. 268). It is a dark-red, volatile liquid, emitting an odor more irritating, if possible, than chlorine—of specific gravity 2.97 to 3.14; boiling-point, $135-145^\circ F$.

If bromine be added to an excess of potassium or sodium hydrate solution, it should combine to form a permanently clear liquid, without the separation of oily drops or the development of an odor resembling that of chloroform (absence of bromoform or other organic bromine compounds).

If an aqueous solution of bromine be shaken with a slight excess of reduced iron until it becomes nearly colorless, the filtered liquid, on the addition of a small amount of ferric chloride and of starch mucilage, should not assume a blue color (absence of iodine).


If 1 cc. of a saturated aqueous solution of bromine be diluted with 9 cc. of water, then mixed with 3 cc. of ammonium carbonate T.S. and 5 cc. of decinormal silver nitrate V.S., and the whole actively shaken, the filtered liquid, when supersaturated with nitric acid, should not become more than opalescent, nor separate a flocculent precipitate within three minutes (absence of more than 3 per cent. of chlorine).

Quantivalence.—The atom of bromine, like that of chlorine, is univalent (Br'). The atomic weight of bromine is 80. Free bromine has the molecular formula Br_2 .

Hydrobromic Acid.—Hydrogen bromide, or hydrobromic acid, may be made by decomposing phosphorus bromide by water:

$\text{PBr}_3 + 4\text{H}_2\text{O} = 5\text{HBr} + \text{H}_3\text{PO}_4$. A small quantity is prepared by placing seven or eight drops of bromine at the bottom of a test-tube, putting in fragments of glass to the height of about an inch or two, then ten or eleven grains of phosphorus, then another inch of glass, and finally a couple of inches of glass fragments slightly wetted with water, a delivery-tube being fitted by a cork. The phosphorus combines readily, almost violently, with the bromine as soon as the vapor of the latter, aided by a little warmth from a flame, rises to the region of the phosphorus. The phosphorus bromide thus formed then suffers decomposition by the water of the moist glass, phosphoric and phosphorous acids being produced. The hydrobromic acid gas passes over (heat being applied in the after part of the operation) and may be condensed in water or in solution of ammonia. The latter solution on evaporation yields ammonium bromide.

Fig. 38.



Preparation of Hydrobromic Acid.

Acidum Hydrobromicum Dilutum, U. S. P.—Pass sulphuretted-hydrogen gas through bromine covered with water, and, when all bromine has disappeared, distil the mixture. The distillate, when diluted until it has a sp. gr. of 1.300, contains 33 per cent. of HBr. Diluted until the sp. gr. is 1.077, it contains 10 per cent. of HBr, and is then of official strength.



Potassium Bromide (KBr) is very largely employed in pharmacy, and is the salt, therefore, which may be used in studying the reactions of this acidulous radical. The official method of making the salt has been alluded to under the salts of potassium (p. 77).

Other bromides are seldom used; they may be prepared in the same way as, and closely resemble, the corresponding chlorides or iodides.

Sodium Bromide crystallizes in anhydrous cubes (NaBr) from solutions at 110° or 120° F., and in hygroscopic prisms ($\text{NaBr} \cdot 2\text{H}_2\text{O}$) at ordinary temperatures.

Ammonium Bromide (NH_4Br) (*Ammonii Bromidum*, U. S. P.) is prepared by agitating iron wire with a solution of bromine until the odor of bromine can be no longer perceived, adding solution of ammonia, filtering, and evaporating the filtrate to dryness. It forms

a white granular salt, which becomes slightly yellow on exposure to air, is readily soluble in water, less so in spirit, and, when heated, sublimes. Ammonium bromide or iodide may also be made by mixing equivalent quantities of strong hot, aqueous solutions of the corresponding potassium salts and of ammonium sulphate. To the cooled liquids rectified spirit is added, which precipitates the potassium sulphate. The spirit recovered by distillation of the clear liquid leaves the required salt as a residue in the retort.

Calcium Bromide, CaBr_2 (*Calcii Bromidum*, U. S. P.), may be prepared by neutralizing hydrochloric acid by calcium hydrate or carbonate, filtering, and evaporating to dryness; or by uniting bromine with iron, boiling the aqueous solution with lime until the mixture is red, filtering, and evaporating. It is a white deliquescent granular salt, soluble in water and in alcohol.

Solution of Bromine, B. P., 10 minims in 5 ounces, or *Bromine-water*, U. S. P., 1 cc. in 100 cc., is an aqueous solution, bromine being slightly soluble in water.

Hypobromites, *Bromates*, and *Perbromates*, analogous to hypochlorites, chlorates, and perchlorates, are producible.

Bromates, occurring as an impurity in bromides, are detected by dropping diluted sulphuric acid on to the salt, when a yellow color, due to free bromine, is produced *immediately* if bromates are present.

ANALYTICAL REACTIONS (TESTS).

First Analytical Reaction.—To a few drops of a solution of a bromide (KBr or NH_4Br) add solution of silver nitrate; a yellowish-white precipitate (silver bromide, AgBr) falls. Treat the precipitate successively with nitric acid and dilute ammonia, as described for the silver chloride; it is only sparingly dissolved in ammonia.

Second Analytical Reaction.—To solution of a bromide add a drop or two of chlorine-water or a bubble or two of chlorine gas; then add a few drops of chloroform or ether or carbon bisulphide, shake the mixture, and set the test-tube aside: the chlorine, from the greater strength of its affinities, displaces the bromine, which is dissolved by the chloroform, etc., the solution falling to the bottom of the tube in the case of the heavy chloroform or carbon bisulphide, or rising to the top in the case of the light ether. Either solution has a distinct yellow or reddish-yellow or red color, according to the amount of bromine present.

Notes.—This reaction serves for the isolation of bromine when mixed with many other substances. Excess of chlorine must be avoided, as colorless bromine chloride is then formed. Iodides give a somewhat similar appearance; the absence of iodine must therefore be ensured by a process given in the next section. The above solution in chloroform or ether may be removed from the tube by

drawing up into a *pipette* (small pipe—a narrow glass tube, usually having a bulb or expanded portion in the centre), the bromine fixed by the addition of a drop of solution of potash or soda, the chloroform or ether evaporated off, and the residue tested as described in the next reaction.

Third Analytical Reaction.—Liberate bromine from a bromide by the cautious addition of chlorine or chlorine-water, then add a few drops of cold “mucilage of starch” (*vide* Index); a yellow combination of bromine and starch, commonly termed “starch bromide,” is formed.


The above reaction may be varied by liberating the bromine by a little black manganese oxide and a drop of sulphuric acid, the upper part of the inside of the test-tube being smeared over with the mucilage of starch. Even sulphuric acid alone, if strong, liberates bromine from a bromide, the hydrogen of the hydrobromic acid first produced uniting with the oxygen of the sulphuric acid—the latter being reduced to sulphurous acid or even to hydrosulphuric acid.

HYDRIODIC ACID AND OTHER IODIDES.

Formula of Hydriodic Acid, HI. Molecular weight, 128.

Source.—The acidulous radical of hydriodic acid and other iodides is the element iodine (I). It occurs in nature as sodium and magnesium iodides in sea-water. Sea-weeds, sponges, and other marine organisms, which derive much of their nourishment from sea-water, store up iodides in their tissues, and it is from the ashes of these that supplies of iodine (*Iodum*, U. S. P.) are obtained. Mineral iodides also are met with, and iodates occur in crude cubic nitre.

Process.—The sea-weed ash or *kelp* is treated with water, insoluble matter thrown away, and the decanted liquid evaporated and set aside to allow of the deposition of most of the sodium and potassium sulphates, carbonates, and chlorides. The residual liquor is treated with excess of sulphuric acid, which causes evolution of carbonic and sulphurous or sulphuretted gases, deposition of sulphur and more sodium sulphate, and formation of hydriodic acid. To the decanted liquid is added black manganese oxide, and the mixture is then slowly distilled; the iodine sublimes, and is afterward purified by resublimation.



The analogy of chlorine, bromine, and iodine is well indicated by the fact that each is obtained from its compounds by the same reaction. Iodine is liberated from any iodide as bromine from bromides or chlorine from chlorides—namely, by the action of black manganese oxide and sulphuric acid.

Properties.—Iodine is a crystalline purplish-black substance; its

vapor, readily seen on heating a fragment in a test-tube, is dark-violet. Its vapors are irritating to the lungs, but a trace may be inhaled with safety (*Vapor Iodi*, B. P.). It melts at about 230° F., boils at about 392° , and is entirely volatilized, the first portions containing any iodine cyanide or cyanogen iodide that may be, though very rarely is, present. The latter body occurs in slender, colorless prisms, emitting a pungent odor.

"A solution of iodine in chloroform should be perfectly clear and limpid (absence of moisture)."

The presence of cyanogen, chlorine, or bromine is determined as follows: When shaken with distilled water it should not communicate to the latter more than a light brownish tinge, and no deep brown color (absence of iodine chloride). If the iodine be removed from this dilute aqueous solution by agitation with carbon disulphide, and after the separation of the latter some dilute solution of ferrous sulphate with a trace of ferric chloride be added, finally soda solution, and the whole supersaturated with hydrochloric acid, no blue precipitate should make its appearance (absence of iodine cyanide). If iodine be dissolved in sulphurous acid, the solution strongly supersaturated with ammonia, and completely precipitated by silver nitrate, the filtrate, on being supersaturated with nitric acid, should not at once become more than faintly cloudy (absence of more than traces of chlorine or bromine). "Triturate 0.25 grm. of finely-powdered iodine in 10 cc. of water, and filter off the solution; to the aqueous filtrate, in a test-tube, add a slight excess of silver nitrate, shake actively, allow the precipitate to subside, and, having poured off the clear, supernatant liquid completely, shake the precipitate with a mixture of 1 cc. of ammonia-water and 9 cc. of water, and filter. Upon the addition of a slight excess of nitric acid to the filtrate, not more than a slight opalescence should make its appearance (limit of chlorine or bromine)." — *U. S. P.*

This latter reaction is applied for the detection of chloride or bromide in iodides, omitting the addition of sulphurous acid.

Quantivalence.—The atom of iodine, like those of bromine and chlorine, is univalent* (I'). The atomic weight of iodine is 127; its molecular formula, I_2 .

Hydrogen Iodide, or Hydriodic Acid, is a heavy, colorless gas. Its solution in water may be made by passing sulphuretted hydrogen into water in which iodine is suspended, the chief reaction being $2H_2S + 2I_2 = S_2 + 4HI$. (See the analogous reaction for HBr , p. 270.)

* There is a compound of iodine having the formula ICl_3 . Iodine would at first sight, therefore, seem to be a trivalent element (I'''), and bromine and fluorine, from their close chemical analogy with iodine, would necessarily be regarded as trivalent also. From this aspect the position of chlorine would be anomalous. Probably, however, the compound is only a molecular combination of true iodine chloride, ICl , with added chlorine, Cl_2 . Iodine forms, with potassium iodide, a periodide or tri-iodide, KI_3 , which may be obtained in lustrous prismatic crystals. This, too, may have the formula KI, I_2 . A mercuric hexiodide (HgI_6 , perhaps HgI_2, I_2, I_2) is also known; and an ammonium periodide, NH_4I, I_2 : both, probably, only "additive" compounds. (See p. 139.)

Hydriodic acid may also be prepared by placing 20 parts of iodine and 2 of water in a retort the neck of which points upward, and the end of the neck of which is connected by a glass tube with a bottle or other vessel containing a little water. Into the tubulure of the retort is passed, at first a drop at a time, a mixture of 1 part of red phosphorus with 2 of water. Abundance of hydriodic acid is evolved on the application of a gentle heat, and falls into and dissolves in the water in the receiver. Phosphoric acid remains. $P_2 + 5I_2 + 8H_2O = 10HI + 2H_3PO_4$. (See the analogous reaction for HBr, p. 269.)

Or iodine may be dissolved in carbon bisulphide in a tall cylinder, water added, and sulphuretted hydrogen passed through the mixture. The water dissolves the hydriodic acid produced, the bisulphide retaining the separated sulphur. The aqueous solution only needs boiling for two or three minutes to remove excess of sulphuretted hydrogen.

Syrupus Acidi Hydriodici, U. S. P., contains 1 per cent. of real acid. *Potassium Iodide*, or Iodide of Potassium (KI), is largely used in medicine, and hence is the most convenient iodide on which to experiment in studying the reactions of this acidulous radical. Solid iodine itself might be taken for the purpose, but its use and action in that state having already been alluded to in describing potassium, cadmium, and mercury iodides, its analytical reactions in the combined condition are those which may now occupy attention.

Ammonii Iodidum, U. S. P., Ammonium Iodide or Iodide of Ammonium, may be made by decomposing the two bodies, potassium iodide and ammonium sulphate, which give ammonium iodide and potassium sulphate; the latter salt is separated by adding alcohol to the cooled solution, when, by reason of its insolubility in alcohol, it crystallizes out, and the separated solution of ammonium iodide is then evaporated to dryness. It occurs usually in minute white crystalline cubes.

Solution of Iodine.—Iodine is slightly soluble in water (iodine-water), and readily soluble in an aqueous solution of potassium iodide. 5 parts of iodine and 10 of potassium iodide, dissolved in 85 of distilled water, form *Liquor Iodi Compositus*, U. S. P. ("Lugol's Solution"); 4 parts of iodine and 1 of potassium iodide, rubbed with 2 parts of water and 93 of benzoated lard, form *Unguentum Iodi*, U. S. P. It is more soluble in spirit (*Tinctura Iodi*, U. S. P.) or in a spirituous solution of potassium iodide (*Tinctura Iodi*, B. P.). It combines with sulphur, forming an unstable grayish-black solid iodide (S_2I_2), having a radiated crystalline structure (*Sulphuris Iodidum*, U. S. P., or *Sulphur Iodide*). If 100 parts be thoroughly boiled with water, the iodine will pass off in vapor, and about 20 parts of sulphur remain.—*B. P. and U. S. P.*

ANALYTICAL REACTIONS (TESTS).

First Analytical Reaction.—To a few drops of an aqueous solution of an iodide (*e. g.* KI) add solution of silver nitrate;

a light yellow precipitate (silver iodide, AgI) falls. Pour away the supernatant liquid and treat the precipitate with nitric acid; it is not dissolved. Pour away the acid, and then add dilute ammonia; it is only sparingly dissolved.

This reaction is useful in separating iodine from most other acidulous radicals, but does not distinguish iodine from bromine.

The presence of chloride in silver iodide may be detected by boiling with dilute solution of ammonium carbonate, filtering off the insoluble silver iodide, and saturating the filtrate with nitric acid; any silver chloride is then precipitated.

Ammonia, it will be remembered, dissolves silver chloride readily; hence the presence of potassium chloride in bromide or iodide may be detected by dissolving in water, adding excess of silver nitrate, collecting the precipitate, washing, digesting in ammonia, filtering, and adding excess of nitric acid to the filtrate; more than a trace of white curdy precipitate indicates chloride (of potassium). Silver bromide and iodide are, however, slightly soluble in ammonia. Better processes are given on pp. 276-279.

Second Analytical Reaction.—Liberate iodine from an iodide by the cautious addition of chlorine, then add mucilage of starch; a deep-blue combination of iodine and starch, commonly termed "starch iodide," is formed.

Starch is highly sensitive to the action of iodine; this reaction is consequently very delicate and characteristic. Heat decomposes the blue compound. Excess of chlorine must be avoided or colorless iodine chloride will be produced. Nitrous acid, or a nitrite acidulated with sulphuric acid, may be used instead of chlorine. Concentrated sulphuric acid also liberates iodine from iodides, the hydrogen of the hydriodic acid first produced uniting with the oxygen of the sulphuric acid—the latter (H_2SO_4) being reduced to sulphurous acid (H_2SO_3), or even to hydrosulphuric acid (H_2S).

In testing bromine for iodine the bromine must be nearly all converted into hydrobromic acid by dilute solution of sulphurous acid, or be nearly all removed by solution of soda, before the mucilage of starch is added.


Ozone (O_3).—Papers soaked in mucilage of starch containing potassium iodide form a test for free chlorine and nitrous acid, and are also employed by meteorologists to detect an allotropic or physically polymeric and energetic form of oxygen, termed by Schönbein *ozone* (from $\delta\zeta\omega$, *ozo*, I smell). This substance liberates iodine from the potassium iodide (with formation of starch iodide), and is supposed to occur normally in the atmosphere, the salubrity or insalubrity of which is said to be dependent to some extent on the presence or absence of ozone. The possible occurrence of nitrous or chlorinoid gases in the air, however, renders the test untrustworthy. Houzeau proposes to test for ozone by exposing litmus-paper of a neutral tint soaked in a dilute solution of potassium iodide: the potash set free by action of the ozone turns the

paper blue. The same paper without iodide would indicate the extent to which the effect might be due to ammonia-vapor. Ozone, or, rather, ozonized air, is produced artificially in large quantities on passing air through a box (Beane's ozone-generator) highly charged with electricity. In the latter operation condensation of the volume of air, or rather of the oxygen in the air, occurs. Small quantities are obtained by exposing in a loosely-closed bottle a stick of phosphorus partly covered by water, but the product is mixed with hydrogen peroxide. Ozone is a powerful bleaching, disinfecting, and general oxidizing agent; insoluble in water, soluble in oils of turpentine, cinnamon, and some other liquids. From experiments that have been made by Soret on the specific gravity of ozone, its molecular formula would seem to be O_3 , that of ordinary oxygen being O_2 . Its smell is peculiar.

Third Analytical Reaction.—To a neutral aqueous solution of an iodide add a solution containing 1 part of copper sulphate to $2\frac{1}{2}$ parts of ferrous sulphate, and well shake; a dirty-white precipitate of cuprous iodide (Cu_2I_2) falls.



Or to the liquid containing an iodide add the solution of copper sulphate and some solution of sulphurous acid, and warm the mixture; cuprous iodide falls.



Separation of Chlorides, Bromides, and Iodides.—Chlorides and bromides are not affected in the above way: the reaction is useful, therefore, in removing iodine from a solution in which chlorides and bromides have to be sought. The total removal of iodine by the former of the two modifications of the process is ensured by supplementing the addition of the cupric and ferrous sulphates by a few drops of solution of potash or soda, any acid which might be keeping cuprous iodide in solution being thereby neutralized, ferric or ferrous hydrate, precipitated at the same time, not affecting the reaction. Occasionally, too, it may be necessary to repeat the process with the filtrate before the last traces of iodine are removed. The second modification of the process is, on the whole, to be preferred.


Chlorides may be separated from bromides by taking advantage of the ready solubility of silver chloride and the slow and slight solubility of silver bromide in ammonia, especially in (a fair, not a great, excess of) ammonia containing silver chloride. The presence of much silver bromide, however, considerably reduces the power of ammonia to dissolve silver chloride.

Hart's Test.—(If nitrates, chlorates, bromates, or iodates are present, it is necessary to fuse the substance with a little sodium carbonate and charcoal to reduce them. If the haloids are united with silver, it is best to fuse with sodium carbonate


and extract with water, although with iodine and bromine this is not absolutely necessary.) The substance is placed in the flask shown in the figure given in the section on the quantitative analysis of manganese oxide (*vide* Index), with some water and a few drops of solution of ferric sulphate. Into the bulbs are poured a few drops of dilute starch mucilage. The bulbs are kept cold by immersing in water in a beaker. The contents of the flask are then boiled, and if iodine is present the starch is colored blue. This test is extremely delicate. If iodine is found, the cork with the bulb-tube is removed, and the solution boiled until, on testing again in the same way, no more iodine is found. If much iodine is present, it is necessary to add more ferric sulphate solution. The bulb-tube is now cleaned, charged with a few drops of water and a drop or two of chloroform, and a very small crystal of potassium permanganate added to the solution in the flask. The contents of the flask are boiled again, and if bromine is present the chloroform becomes red. The tube is now removed, and more potassium permanganate and ferric sulphate added, little by little, the mixture being boiled between each addition until the bromine has all been driven off. A few drops of alcohol are added to the contents of the flask to decolorize any excess of permanganate, and after filtration chlorine is tested for in the filtrate with silver nitrate.

Chlorides may also be detected in bromides and iodides by taking advantage of the formation of chlorochromic anhydride (p. 240) and the non-occurrence of corresponding compounds of bromine or iodine, as follows:

To a solution of a mixture of an iodide with a bromide and a chloride add a concentrated solution of sodium sulphite, then a reagent prepared by mixing equal volumes of sulphuric acid and saturated solution of copper sulphate, until no further precipitation of cuprous iodide occurs. Next add solution of soda to remove excess of copper sulphate, filter, and evaporate to dryness. Transfer the dried residue, together with an equal bulk of red potassium chromate, to a dry test-tube fitted with a delivery-tube, or to a small retort, and cover the mixture with sulphuric acid. Distil into water. Chromic anhydride and hydrochloric and hydrobromic acids are liberated by the sulphuric acid, and, reacting one upon another, form chlorochromic anhydride, together with free bromine and chlorine.



The chlorochromic anhydride is decomposed by the excess of water into which it distils, giving rise to chromic acid, which imparts its color to the liquid, and hydrochloric acid, thus :



Chlorine gas escapes, and the bromine is dissolved by the water. The colored liquid is then shaken with chloroform, which removes the bromine, indicating bromides in the original substance. A yellow color remaining is due to chromic acid, indicating chlorides in the original substance. Or add ammonia to the distillate: the color due to bromine is thereby entirely removed, while that of the chromic compound is only slightly modified.

Instead of eliminating the iodine as cuprous iodide, it may be expelled in vapor, obvious enough by its color and odor, by fusing the dry mixture of the salts with excess of powdered red chromate. The residue, broken into small fragments, may then be distilled with the sulphuric acid for the detection of the bromine and chlorine.



Fourth Analytical Reaction.—Iodides have been shown to be useful in testing for mercuric salts (see the Mercuric Reactions, p. 206); a mercuric salt (corrosive sublimate, for example) may therefore be used in testing for iodides, a scarlet precipitate (mercuric iodide, HgI_2) being produced.

This reaction may be employed where large quantities of an iodide are present, but its usefulness in analysis is much impaired by the fact that the precipitate is soluble in excess of the dissolved iodide or in excess of the mercurial reagent. Its color and insolubility in water distinguish it from mercuric chloride, bromide, and cyanide, which are white soluble salts.

Fifth Analytical Reaction.—Iodides have also (see the Lead Reactions, p. 213) been shown to be useful in testing for lead salts; similarly a lead salt (acetate, for example) may be used in testing for iodides, in solutions which are either neutral or faintly acid with acetic acid, a yellow precipitate (lead iodide, PbI_2), soluble in hot water and crystallizing in yellow scales on cooling, being produced.

Lead chloride, bromide, and cyanide are white; hence the above reaction may occasionally be useful in distinguishing iodine from the allied radicals. But lead iodide is slightly soluble in cold water; hence small quantities of iodine cannot be detected by this reaction. (For Iodates, see Index.)

Analogies between Chlorine, Bromine, Iodine, and their Compounds.

—These elements form a natural group or family, each distinct from the other, yet closely related. Moreover, their dissimilarities are so curiously gradational as to irresistibly suggest the idea that some day we may find the differences between these bodies to be in degree rather than in kind. Thus chlorine is a gas and iodine a solid, while bromine occupies the intermediate condition. The atomic weight of bromine is nearly midway between those of chlorine and iodine. The same may be said of the weight of equal volumes of each in a gaseous state. The specific gravity of fluid chlorine is 1.33, of iodine 4.95, while bromine is nearly 3. Liquid chlorine is transparent, iodine opaque, bromine intermediate. The crystalline forms of the chloride, bromide, and iodide of a metal are commonly identical. One volume of either element in the gaseous state combines with an equal volume of hydrogen (at the same temperature) to form two volumes of a gaseous acid, very soluble in water (hydrochloric acid, hydrobromic acid, hydriodic acid). Many other analogies are traceable. (*Vide Index*, “Periodic Law.”)

QUESTIONS AND EXERCISES.

State the method by which bromine is obtained from its natural compounds—Mention the properties of bromine.—How may potassium and ammonium bromides be made?—By what reagents may bromides be distinguished from chlorides?—Whence is iodine obtained?—By what process is iodine isolated?—State the properties of iodine.—What is the nature of sulphur iodide?—Give the analytical reactions of iodides.—What three substances may indirectly be detected by a mixture of potassium iodide and mucilage of starch?—Describe a method by which iodides may be removed from a solution containing chlorides and bromides.

HYDROCYANIC ACID AND OTHER CYANIDES.

Formula of Hydrocyanic Acid, HCN or HCy . Molecular weight, 27.

History of Cyanogen.—The acidulous radical of hydrocyanic acid and other cyanides is a compound body, cyanogen (CN , or, shortly, Cy or C_2N_2 or $\text{N}\equiv\text{C}-\text{C}\equiv\text{N}$). It is so named from *κύανος*, *kuanos*, blue, and *γεννάω*, *gennao*, I generate, in allusion to its prominent chemical character of forming, with iron, the different varieties of prussian blue. It was from prussian blue that Scheele in 1782 first obtained what we now, from our knowledge of its composition, term hydrocyanic acid (HCy , or HCN , or $\text{H}-\text{C}\equiv\text{N}$), but which he called prussic acid. Cyanogen was isolated by Gay-Lussac in 1814, and was the first compound radical distinctly proved to exist.

Sources.—Cyanogen does not occur in nature, and is only formed from its elements under certain circumstances. It is found in small quantities among the gases of iron-furnaces, and is produced to a slight extent in distilling coals for gas. In the form of potassium ferrocyanide it is obtained abundantly by heating animal refuse containing nitrogen, such as the scrapings of horns, hoofs, and hides

(5 parts), with potassium carbonate (2 parts) and waste iron (filings, etc.) in a covered iron pot. The residual mass is boiled with water, the mixture filtered, and the filtrate evaporated and set aside for crystals to form. The cyanogen, produced from the carbon and nitrogen of the animal matter, unites with the potassium, and afterward, on boiling with water, with iron, to form what is often termed the yellow prussiate of potash, or potassium ferrocyanide, *Potassii Ferrocyanidum*, U. S. P. ($K'_4Fe''Cy'_6, 3H_2O$), a compound occurring in four-sided tabular yellow crystals. It contains the elements of cyanogen, yet is not a cyanide, for it is not poisonous and is otherwise different from cyanides: it will be further noticed subsequently. From this salt all cyanides are directly or indirectly prepared.

Potassium Cyanide (KCy) (*Potassii Cyanidum*, U. S. P.), the most common cyanide, may be obtained by heating the ferrocyanide to redness until gas (chiefly nitrogen) ceases to be evolved, and iron carbide settles to the bottom of the molten mass of almost pure cyanide. The product, carefully poured off and cooled, is an opaque crystalline mass containing about 95 per cent. of the salt. It also may be procured by fusing 8 parts of potassium ferrocyanide with 3 of potassium carbonate in a crucible; carbonic acid gas (CO_2) is evolved, iron (Fe) is set free, and potassium cyanate ($KCyO$), a body that will be subsequently noticed, is formed at the same time:



Double Cyanides exist, such as sodium and silver cyanide ($NaCy$, $AgCy$), formed in the process (subsequently described) of quantitatively determining the amount of hydrocyanic acid in a liquid by a standard solution of silver nitrate: these compounds have more or less the properties of their constituents. But other cyanogen compounds, not double cyanides, occur in which the cyanogen is so intimately united with a metal as to form a distinct radical: such are ferrocyanides and ferricyanides—salts which will be noticed in due course.

Cyanogen, like chlorine, bromine, and iodine, is univalent (Cy'). It may be isolated by simply heating mercuric cyanide ($HgCy_2$) or silver cyanide ($AgCy$). It is a colorless gas, burning, when ignited, with a beautiful peach-blossom-colored flame.

Mercuric cyanide is produced in crystals on dissolving 1 part of potassium ferrocyanide in 15 parts of boiling water, adding 2 parts of mercuric sulphate, keeping the whole hot for ten or fifteen minutes, and then filtering and setting aside to cool. In addition to mercuric cyanide ($HgCy_2$), mercury (Hg), ferric sulphate (Fe_3SO_4), and potassium sulphate (K_2SO_4) are formed. Any excess of ferrocyanide also gives prussian blue by reaction with the ferric sulphate. *Hydrargyri Cyanidum*, U. S. P., may also be made by dissolving red oxide of mercury in diluted hydrocyanic acid. A small flame of cyanogen may be obtained on heating a few crystals of mercuric cyanide in a short piece of glass tubing closed at one end, and applying a light to the other end as soon as evolution of gas commences; brown paracyanogen (C_3N_3) and mercury remain.

REACTIONS.

Diluted Hydrocyanic Acid, or Prussic Acid.

Synthetical Reaction.—Dissolve 2 or 3 grains of potassium ferrocyanide in five or six times its weight of water in a test-tube, add a few drops of sulphuric acid and boil the mixture, conveying the evolved gas by a bent glass tube (adapted to the test-tube by a cork) into another test-tube containing a little water; the product is a dilute solution of hydrocyanic acid. Made by this process in larger quantities and of a certain definite strength (2 per cent.), this solution is the *Acidum Hydrocyanicum Dilutum*, U. S. P., “a colorless liquid of peculiar odor. Specific gravity, 0.997.”



The following are the details of the official (U. S. P.) process:

Place 20 parts of potassium ferrocyanide in coarse powder in a tubulated retort, and add to it (40) parts of water. Connect the neck of the retort (which is to be directed upward), by means of a bent tube, with a well-cooled condenser, the delivery-tube of which terminates in a receiver surrounded with ice-cold water, and containing 65 parts of distilled water. All the joints of the apparatus, except the neck of the receiver, having been made air-tight, pour into the retort, through the tubulure, the 8 parts of sulphuric acid, previously diluted with 25 parts of water. Agitate the retort gently,* and then heat it, in a sand-bath, until the contents are in brisk ebullition, and continue the heat regularly until there is but little liquid mixed with the saline mass remaining in the retort.* (The

* This operation is peculiarly liable to those sudden and tumultuous evolutions of vapor, or “bumpings,” or “*soubresauts*,” which often interfere with successful distillation. Such phenomena occur, according to Tomlinson, whenever unaided heat has to overcome the great amount of adhesion naturally existing between certain liquids and vapors, or, rather, between the normal liquid and those particles of it which, becoming strongly heated at the heated part of the vessel, have assumed the condition of particles of dissolved vapor, and which would at once pass from this condition into that of permanent vapor but for adhesion. Ordinarily, a glass or other surface is not absolutely clean, but is more or less covered with specks, traces of materials deposited from the air, the fingers, cloths, etc. Some liquids seem to have little or no adhesion for these materials, while certain vapors have greater adhesion for the films than for the liquids. Hence, in ordinary regular ebullition the vapors accumulate on the films, and then at once become subject to the pressure of the mass of fluid, and so pass off in bubbles. But when the films are absent or have become removed during distillation, the heat accumulates until it is sufficient to overcome the adhesion of the superheated particles, and these are then, all of them at once, converted into vapor, the liquid sometimes boiling over or even bursting the vessel. “Bumping” may be prevented by using glass vessels roughened inside (by hydrofluoric acid or otherwise), or by inserting fragments of substances for which vapor-particles have adhesion, but no known substance has this property in an absolute degree. Tobacco-pipe or pumice-stone, pieces of cork, thick paper, resin, sulphur, platinum wire, etc. are useful when there is

end of the condenser or an attached tube should pass quite into the receiver.) Detach the receiver, and add to its contents so much distilled water as may be required to bring the product to the strength of 2 per cent. of absolute hydrocyanic acid. (*Vide* paragraphs on quantitative analysis.)

The residue of this reaction is acid potassium sulphate (KHSO_4), which remains in solution, and potassium and iron ferrocyanide ($\text{Fe}'\text{K}_2\text{FeCy}_6$), an insoluble powder sometimes termed Everitt's yellow salt, from the name of the chemist who first made out the nature of the reaction. The latter compound becomes bluish-green during the reaction, owing to the absorption of oxygen.

Another Process.—A 2 per cent. solution may also be made by shaking together in a glass-stoppered bottle 6 grms. of silver cyanide, 5 cc. of hydrochloric acid, and 35 cc. of distilled water, and pouring off the clear liquid.

Pure anhydrous hydrocyanic acid is a colorless, highly volatile, intensely poisonous liquid, solidifying when cooled to a low temperature.* It may be made by passing sulphuretted hydrogen over mercuric cyanide. The official solution of the acid is fairly stable, but is said to be rendered more so by the presence of a minute trace of sulphuric or hydrochloric acid. A stronger acid is liable to assimilate the elements of water and yield ammonium formate (NH_4CHO_2). Solutions of hydrocyanic acid often become brown by formation of what is, apparently, *paracyanogen* (C_3N_3). According to Williams, aqueous hydrocyanic acid containing 20 per cent. of glycerin can be kept for an almost indefinite length of time. The official acid should be preserved in well-corked bottles tied over with impervious tissue and kept inverted, when not in use, in a cool, dark place. Unless such precautions are adopted, the fluid rapidly loses strength by escape of the vapor of the acid.

Note.—A few drops of diluted hydrocyanic acid, so placed that its vapor may be inhaled, form the *Vapor Acidi Hydrocyanici*, B. P., or inhalation of hydrocyanic acid.

Hydrocyanic acid also occurs in cherry-laurel water and bitter-almond water. (*Vide* Index.) *Aqua Laurocerasi*, B. P., is made to contain 0.1 per cent. of real acid (HCy).

The methods of determining the strength of hydrocyanic-acid solutions will be given in connection with volumetric and gravimetric quantitative analysis. They are based on the formation of silver cyanide and its solubility in solution of alkaline cyanides, as described in the next reaction.

The hydrocyanic acid used in pharmacy is extremely liable to variation in strength. It should frequently be tested volumetrically.

no chemical action between them and the liquid. Tomlinson recommends coconut-shell charcoal. A slow current of gas, such as hydrogen, air, or carbonic acid gas, also promotes escape of vapor from a liquid, a few capillary tubes, sealed at one end and placed mouth downward in the fluid, answering well. A jet of steam prevents bumping, but is not always applicable.

* Traces are formed when electricity passes between carbon poles in slightly moist air (Dewar).

ANALYTICAL REACTIONS (TESTS).


First Analytical Reaction.—To a few drops of the hydrocyanic-acid solution produced in the above reaction, or to any solution of a cyanide, add excess of solution of silver nitrate; a white precipitate (silver cyanide, AgCy) falls. When the precipitate has subsided, pour away the supernatant liquid and place half of the residue in another test-tube: to one portion add nitric acid, and notice that the precipitate does not dissolve; to the other add ammonia, and observe that the precipitate, though soluble, dissolves somewhat slowly. (Silver chloride, which is also white, is readily soluble in ammonia.) Silver cyanide dissolves in solution of cyanides of alkali-metals, soluble double cyanides being formed (*e. g.* KCyAgCy).

Solubility of Precipitates in Strong Solutions of Salts.—Silver cyanide and many other precipitates insoluble in acids (similar remarks apply to precipitates insoluble in alkalies) are often soluble in the saline liquids formed by the addition of acids and alkalies to each other. Hence the precaution of adding the latter reagents to separate portions of a precipitate, or of not adding the one until the other has been poured away.

Cyanogen in an insoluble cyanide, such as silver cyanide itself, is readily recognized on heating the substance in a short piece of glass tubing closed at one end like a test-tube and drawn out at the other end so as to have but a small opening; on applying a flame the escaping cyanogen ignites and burns with a characteristic peach-blossom tint. Metallic silver remains.

Antidote.

Second Analytical Reaction.—To a dilute solution of hydrocyanic acid, or a soluble cyanide, add a few drops of solution of a ferrous salt and a drop or two of solution of a ferric salt (ferrous sulphate and ferric chloride are usually at hand); to the mixture add potash, soda, magnesia, or sodium carbonate, and then hydrochloric acid; a precipitate of prussian blue remains. The decompositions may be traced in the following equations:



The test depends on the conversion of the cyanogen into ferrocyanogen by aid of the iron of a ferrous salt and the combination of the ferrocyanogen, so produced, with the iron of a ferric salt.

Hence a mixture of a ferrous sulphide, solution of iron perchloride, and either magnesia or sodium carbonate is the recognized *antidote* in cases of poisoning by hydrocyanic acid or potassium cyanide.

In such an alkaline mixture the poisonous cyanide, by reaction with ferrous hydrate, is at once converted into innocuous potassium or sodium ferrocyanide, etc.: should the mixture become acid, the ferric salt present reacts with the soluble ferrocyanide, forming insoluble prussian blue, which is also inert. From the rapidity of the action of these poisons, however, there is seldom time to prepare an antidote. Emetics, the stomach-pump or stomach-siphon, the application of a stream of cold water to the spine, and the above antidote form the usual treatment.

Third Analytical Reaction.—To solution of hydrocyanic acid add ammonia and common yellow ammonium sulphhydrate, and evaporate the liquid nearly or quite to dryness in a small dish, occasionally adding ammonia till the excess of ammonium sulphhydrate is decomposed; add water and acidify the liquid with hydrochloric acid, and then add a drop of solution of a ferric salt; a blood-red solution of iron sulphocyanate will be formed.

This is a very delicate reaction. Some free sulphur in the yellow ammonium sulphhydrate unites with the alkaline cyanide and forms sulphocyanate ($2\text{NH}_4\text{Cy} + \text{S}_2 = 2\text{NH}_4\text{CyS}$); the ammonia combines with excess of free sulphur and forms, among other salts, ammonium sulphhydrate, the whole of which is removed by the ebullition. If the liquid has not been evaporated far enough, ammonium sulphhydrate may still be present, and give black ferrous sulphide on the addition of the ferric salt; hence prior acidification.

Hydrocyanic Acid in the Blood.—According to Buchner, the blood of animals poisoned by hydrocyanic acid, instead of coagulating as usual, remains liquid and of a clear cherry-red color for several days. In one case he obtained the reactions of the acid on diluting and distilling the blood fifteen days after death, and applying the usual reagents to the distillate. Aqueous solution of hydrogen peroxide changes such blood to a deep-brown color.

Schönbein's test for hydrocyanic acid is said to be extremely delicate. Filtering-paper is soaked in a solution of 3 parts of guaiacum resin in 100 of alcohol. A strip of this paper is dipped in a solution of 1 part of copper sulphate in 50 of water; a little of the suspected solution is placed on this paper and exposed to the air, when it immediately turns blue. Or the paper may be placed over the neck of an open bottle of medicine supposed to contain hydrocyanic acid or otherwise exposed to the suspected vapor of the acid.

QUESTIONS AND EXERCISES.

Write a paragraph on the history of cyanogen.—Mention the source of the cyanogen of cyanides.—How is potassium ferrocyanide prepared?—What is the formula of potassium ferrocyanide?—Is potassium ferrocyanide poisonous?—Write an equation expressive of the reaction which ensues when potassium ferrocyanide and carbonate are brought together at a high temperature.—What are the properties of cyanogen?—How

may it be obtained in a pure condition?—How is mercuric cyanide prepared?—What other substances and secondary products are formed at the same time?—How much real hydrocyanic acid is contained in the official *Acidum Hydrocyanicum Dilutum*?—Give details of the preparation of hydrocyanic acid and an equation of the reaction.—State the proportion of water that must be added to an aqueous solution containing 15 per cent. of hydrocyanic acid to reduce the strength to 2 per cent. *Ans.* $6\frac{1}{2}$ to 1.—What are the characters of pure undiluted hydrocyanic acid?—How may it be obtained?—Enumerate the tests for cyanogen, giving equations.—Explain the action of the best antidote in cases of poisoning by hydrocyanic acid or potassium cyanide.—Show how it acts in alkaline and acid solutions respectively.

NITRIC ACID AND OTHER NITRATES.

Formula of the acid, HNO_3 or HONO_2 . Molecular weight, 63.

Introduction.—The group of elements represented by the formula NO_3 is that characteristic of nitric acid and all other nitrates; hence it is expedient to regard these elements as forming an acidulous radical, which may be termed *the nitric radical*. Like the hypothetical basylous radical ammonium (NH_4), this supposed acidulous radical (NO_3) has not been isolated. Possibly it is liberated when chlorine is brought into contact with silver nitrate; but if so, its decomposition into white crystalline nitric anhydride (N_2O_5) and oxygen (O) is too rapid to admit of its identification.

Sources.—The nitrogen and oxygen of the air combine and ultimately form nitric acid whenever a current of electricity (as in the occurrence of lightning) passes. The nitrates found in rain may partly or wholly thus originate. The oxidation of ammoniacal matter and of the nitrogenous constituents of animal and vegetable matters in the soil, favored by the darkness, by the occurrence of calcium carbonate, and by the presence of some low form of vegetable life acting as a ferment, results in the production of nitrates. Hence nitrates are commonly met with in waters, soils, and the juices of plants. In the concentrated plant-juices, termed medicinal "Extracts," small prismatic crystals of potassium nitrate may occasionally be observed. (The cubical crystals often met with in extracts are potassium chloride.) Nitric acid and other nitrates are obtained from potassium and sodium nitrates, and these from the surface layers of the soil of tropical countries. *Potassium nitrate*, or *prismatic nitre* (from the form of its crystals), is chiefly produced in and about the villages of India. The natives simply scrape the surface of waste grounds, mud-heaps, banks, and other spots where a slight incrustation indicates the presence of appreciable quantities of nitre, mix the scrapings with wood-ashes (potassium carbonate, to decompose the calcium nitrate always present), digest the mixture in water, and evaporate the liquor. The immediate product is purified by careful recrystallizations, and is sent into commerce in the form of white crystalline masses or fragments of striated six-sided prisms. Besides its use in medicine (*Potassii Nitras*, U. S. P., Nitrate of Potassium, or *Potassium Nitrate*), it is employed in very

large quantities in the manufacture of gunpowder. *Charta Potassii Nitratis*, U. S. P., *Nitrate-of-Potassium Paper*, is made by dipping strips of white unsized paper in a solution of 1 part of the salt in 4 parts of water and drying them. *Sodium Nitrate* (*Sodii Nitrates*, U. S. P.), the old nitrate of soda, occurs in deposits from three inches to three yards in thickness on and near the surface, and at any depth down to about thirty feet, in many parts of Peru, Bolivia, and Chili, but more especially in the district of Atacama. The mineral is termed *caliche*, and commonly contains 50 per cent. of sodium nitrate. The latter is distinguished as *Chili saltpetre* or *Chili nitre*, or (from the form of its crystals—obtuse rhomboids, not true cubes) *cubic nitre*, and is chiefly used as a manure and as a source of nitric acid, its tendency to absorb moisture unfitting it for use in gunpowder. In many parts of Europe potassium nitrate is made artificially by exposing heaps of animal manure, refuse, ashes, and soil to the action of the air and the heat of the sun; in the course of a year or two the nitrogen of the animal matter becomes oxidized to nitrates; the latter are removed by washing. According to Warington, the nitrifying ferment appears capable of existing in three conditions: (1) The nitric ferment of soil, which converts both ammonium salts and nitrites into nitrates; (2) the altered ferment, which converts ammonium salts into nitrites, but fails to change nitrites into nitrates; and (3) the surface organism (a bacterium) which changes nitrites into nitrates. Similar nitrification goes on in impure well and river waters, which thereby tend to become pure.

Note.—The word *nitric* is from *nitre*, the English equivalent of the Greek *νίτρον* (nitron), a name applied to certain natural deposits of *natron* (sodium carbonate), for which potassium nitrate seems at first to have been mistaken. *Saltpetre* is simply *sal petræ*, salt of the rock, in allusion to the natural origin of potassium nitrate. *Sal prunella* (from *sal*, a salt, and *pruna*, a live coal) is potassium nitrate melted over a fire and cast into cakes or bullets.

The nitric radical is univalent (NO_3').

Constitution of Salts.

It is here necessary again to caution the reader against regarding salts as invariably possessing a known constitution, or supposing that they always possess two or more sides or contain definite radicals. The erroneous conception which, of all others, is most likely to be imperceptibly formed is that of considering salts to be binary bodies. For, first, the names of salts are necessarily binary. A student hears the names "iron sulphate," "copper sulphate," and simultaneously receives the impression that each salt has two sides, copper or iron occupying one, and something indicated by the word "sulphate" the other. Such words as "vitriol," green or blue, or "nitre," would perhaps implant unitary ideas in the mind, but it is simply impossible to give such names to all salts as will convey the impression that each salt is a whole, and therefore unitary. The name "sulphate of potash" produces binary impressions, and the less incorrect name, "sulphate of potassium" or "potassium sulphate," is in this respect no better. Secondly, it is impracticable

to study salts as a whole. Teachers are almost unanimous in the opinion that students should first master the reactions characteristic of the metals in salts, and then the residues which, with those metals, make up the salts, and *vice versâ*. It is not only impracticable, but impossible, to study salts as a whole; binary ideas concerning them are therefore almost inevitably imbibed. We come to regard a salt as a body which splits up in one direction only; look upon nitre, for instance, and all other nitrates, as containing NO_3 and a metal, M ; whereas KNO_3 may be split up into KNO_2 and O , or into K_2O , N_2 , and O_5 , or may contain K_2O and N_2O_5 . These are the chief disadvantages attending the employment of the binary hypothesis in studying chemical compounds: if they be borne in mind, the hypothesis may be freely used without much danger of permanent mental bias. Thus, in nitre let the group of elements (NO_3) which, with potassium, makes up the whole salt, be called the nitric radical, the name of the latter being directly derived from its hydrogen salt. Similarly, allow the acidulous residues of other salts of metals to be termed respectively the chloric, acetic, sulphurous, sulphuric, carbonic, oxalic, tartaric, phosphoric, citric, boracic radicals. In short, these compound radicals should be regarded as groupings common to many salts, and which may usually be transferred without any apparent breaking or splitting; at the same time we must be prepared to find that occasionally a salt divides in other directions. In this way perhaps erroneous impressions will gain least hold on the mind, and a way be left open for the easy entrance of new truths should the real constitution of salts be discovered.

Formerly, salts (such as magnesium sulphate) were regarded as containing (a) an oxide of a metal (MgO) and an anhydride (SO_3), the latter being incorrectly called an acid (sulphuric acid); or (b) as containing two simple radicals (e.g. KI , NaCl , $\text{KC}_2\text{H}_3\text{O}_2$, HgS)—the former being called *oxyacid salts* or *oxysalts*, and the latter *haloid salts* (from *ἅλς*, *als*, sea-salt, and *εἶδος*, *eidōs*, likeness). Such distinction is no longer maintained, the two classes being merged. This is an important educational gain on the side of simplicity; for whereas under the old system much time was necessarily expended before salts of a metal and salts of the oxide of that metal could be distinguished (e.g. KI and $\text{K}_2\text{O}, \text{SO}_3$), now, all salts being regarded as salts of the metals themselves (e.g. KI and K_2SO_4), no such distinction is necessary.

REACTIONS.


Nitric Acid.

Synonyms.—Hydrogen Nitrate; Nitrate of Hydrogen.

Synthetical Reaction.—To a fragment of potassium nitrate or sodium nitrate in a test-tube add a drop or two of sulphuric acid, and warm; nitric acid (HNO_3) is evolved in vapor. The fumes may be condensed by a bent tube fitted to the test-tube not by a cork, as for hydrochloric acid—because the nitric

vapors would strongly act on it—but by plaster of Paris, a paste of which sets hard on being put aside for a short time and is unaffected by the acid.

On a somewhat larger scale nitric acid may be prepared by heating, in a stoppered or plain retort, a mixture of equal weights of potassium nitrate and sulphuric acid; the acid distills over, and acid potassium sulphate remains behind:




Half the quantity of sulphuric acid may be taken; but in that case neutral potassium sulphate (K_2SO_4) is produced, which, from its hard, slightly soluble character, is removed with difficulty from the retort. On the manufacturing scale the lesser proportion is used, but instead of retorts iron cylinders are employed, from which the residual salt is removed by chisels. Moreover, the cheaper sodium salt is the nitrate from which manufacturers usually prepare nitric acid, 7 parts of sodium nitrate and 4 of sulphuric acid being employed.

Note.—The acid potassium sulphate is readily converted into neutral sulphate (*Potassii Sulphas*, Sulphate of Potassium, or *Potassium Sulphate*, U. S. P.) by dissolving in water, adding potassium carbonate until effervescence ceases to occur, filtering, and setting aside to crystallize.


Pure nitric acid (HNO_3) is a colorless liquid, somewhat difficult of preparation; its specific gravity is 1.52. The strongest acid met with in commerce has a sp. gr. of 1.5, and contains 93 per cent. of real nitric acid (HNO_3); it fumes disagreeably, is unstable, and, except as an escharotic, is seldom used. The United States Pharmacopœia contains two acids: *Acidum Nitricum*, of sp. gr. 1.414 and containing 68 per cent. of real acid (HNO_3); and another, *Acidum Nitricum Dilutum*, sp. gr. 1.057, containing 10 per cent. The stronger liquid, although containing water, is usually simply termed "nitric acid." The official nitric acid, of sp. gr. 1.414, is a definite hydrous acid ($2\text{HNO}_3, 3\text{H}_2\text{O}$); it distills at 240°F . without change. If a weaker acid be heated, it loses water; if a stronger acid be heated, it loses nitric acid until the density of 1.42 is reached. *Aqua fortis* is an old name for nitric acid (*Aqua Fortis Simplex*, sp. gr. 1.22 to 1.25; *Aqua Fortis Duplex*, 1.36). The strength of a specimen of nitric acid is determined by volumetric analysis. *Nitric Anhydride* (N_2O_5), sometimes but erroneously called *Anhydrous Nitric Acid*, is a solid crystalline substance formed on passing dry chlorine over dry silver nitrate.

Metals reduce nitric acid to nitrous acid and to the various oxides of nitrogen, or even to nitrogen itself, according to the strength of acid, temperature, and amount of nitrate present. Not infrequently ammonium nitrate is simultaneously formed. Thus with zinc:



Aqua Regia, or Nitrohydrochloric Acid.—2 parts of nitric acid

and 9 of hydrochloric acid by volume form the *Acidum Nitrohydrochloricum*, U. S. P., and the same weights, with 78 of water, give the *Acidum Nitrohydrochloricum Dilutum*. The mixture should be set aside for a week in summer or a fortnight in winter to ensure mutual decomposition and full development of the chief active product, chlorine:



In the latter stages of the reaction the decomposition expressed in the following equation also probably occurs:




The undiluted mixture of acids is known as *aqua regia*, from its property of dissolving gold, the "king of metals."

"Diluted nitrohydrochloric acid" may attack organic matter with evolution of nitrous gases, hence should not be dispensed with tinctures, etc. without further dilution.

ANALYTICAL REACTIONS (TESTS).

First Analytical Reaction.—To a solution of any nitrate (*e. g.* KNO_3) add sulphuric acid, and then copper turnings, and warm; colorless nitric oxide gas (NO) is evolved, which at once unites with the oxygen in the tube, giving *red fumes* of nitric peroxide or nitrogen peroxide (NO_2).




Performed on a larger scale in a vessel to which a delivery-tube is attached, the reaction of nitric acid on copper becomes of syn-
thetical interest, being the process for the preparation of nitric oxide gas for the purposes of chemical experiment.

Small amounts of a nitrate may be overlooked by this test, the color of the red fumes not being very intense.


Undiluted nitric acid poured on to copper turnings gives dense red vapors of *nitrous acid* (HNO_2), *nitrous anhydride* (N_2O_3), *nitric peroxide* (NO_2), *nitric oxide* (NO), and even *nitrogen* (N_2), the reaction varying somewhat according to the temperature of the mixture and (Ackworth) the amount of copper nitrate in solution. With ordinary copper diluted nitric acid gives nitric oxide, $\text{Cu}_3 + 8\text{HNO}_3 = 3(\text{Cu}_2\text{NO}_3) + 4\text{H}_2\text{O} + 2\text{NO}$.

Second Analytical Reaction.—To a cold solution of a nitrate, even if very dilute, add three or four crystals of ferrous sulphate, shake gently for a minute, in order that some of the sulphate may become dissolved, and then pour 8 or 10 drops of strong sulphuric acid down the side of the test-tube, so that it may form a layer at the bottom of the vessel; a reddish-purple or black coloration will appear between the acid and the supernatant liquid.

This is a very delicate test for the presence of nitrates. Nitrites give the reaction without the sulphuric acid. The black color is due to a solution, or perhaps combination of nitric oxide with a portion of the ferrous salt. The nitric oxide is liberated from the nitrate by the reducing action of the hydrogen of the sulphuric acid, the sulphuric radical of which is absorbed by another portion of the ferrous sulphate, the latter then becoming ferric sulphate.



The process of oxidation is one frequently employed in experimental chemistry, and nitrates, from their richness in oxygen, but more especially because always at hand, are the oxidizers usually selected for the purpose. In the operation they generally split up in one way—namely, into oxide of their basylous radical, nitric oxide gas, and available oxygen. Thus hydrogen nitrate (nitric acid) commonly yields hydrogen oxide (water) and the other bodies mentioned, as shown in the following equation :



When nitrates, other than nitric acid, are used for the purpose of oxidation, a stronger acid, generally sulphuric, is commonly added in order that nitric acid may be formed, the hydrogen nitrate splitting up more readily than any other nitrates.

The five oxides of nitrogen have now been mentioned—namely :

Nitrous oxide	N_2O	} or {	N_2O
Nitric oxide*	NO		N_2O_2
Nitrous anhydride	N_2O_3		N_2O_3
Nitric peroxide*	NO_2		N_2O_4
Nitric anhydride	N_2O_5		N_2O_5

Nitrous oxide is a colorless gas, not altered on exposure to air ; nitric oxide is also colorless, but gives red fumes in the air ; nitrous anhydride is a red vapor condensible to a blue liquid ; nitric peroxide is a red vapor condensible to an orange liquid ; nitric anhydride is a colorless crystalline solid. The two anhydrides, by absorbing water, yield respectively nitrous acid (HNO_2) and nitric acid (HNO_3). Nitrous oxide is also probably an anhydride, the acid of which would doubtless have the formula HNO , while the silver and sodium salts certainly have the formula AgNO and $\text{NaNO}_3\cdot\text{H}_2\text{O}$ (Divers ; Menke). The above series of compounds forms a good illustration of the doctrine of multiple proportions (p 49).

Third Analytical Reaction.—Direct the blowpipe flame on to charcoal until a spot is red hot ; now place on the spot a fragment of a nitrate ; deflagration ensues.

This reaction does not distinguish nitrates from chlorates. It is sufficient for the recognition of very small quantities of either class of salts (especially when they are mixed with other substances).

* The specific gravities of these gases indicate that NO and NO_2 are the correct formulæ, and not N_2O_2 and N_2O_4 .

Gunpowder is an intimate mechanical mixture of 75 parts of nitre, 15 to $12\frac{1}{2}$ parts of charcoal, and 10 to $12\frac{1}{2}$ parts of sulphur. In burning it may be said to give potassium sulphide (the white smoke, K_2S), nitrogen (N), carbonic oxide (CO), and carbonic acid (CO_2) gases, though the decomposition is seldom complete. The sudden production of a large quantity of highly heated gas from a small quantity of a cold solid is sufficient to account for all the effects of gunpowder.

Fourth Analytical Reaction.—To nitric acid or other nitrate add solution of “indigo sulphate;” the color (indigotin-disulphonic acid, $C_{16}H_8(HSO_3)_2N_2O_2$) is discharged.

Indigo Test-solution, U. S. P. (Sulphindyllic or Sulphindigotic Acid), is made by digesting 1 part of dry finely-powdered indigo in 6 parts of strong sulphuric acid in a test-tube; set the mixture aside for two days; the blue liquid is then poured into 25 parts of water, the whole well shaken, set aside, and the clear liquid decanted. Free chlorine also destroys the color of this reagent.

Indigo, U. S. P. ($C_{16}H_{10}N_2O_2$), is a blue coloring-matter deposited when infusion of various species of *Indigofera* is exposed to air and slight warmth. Under these circumstances, *indican*, a yellow, transparent, amorphous substance, soluble in water, breaks up into indigo, which is insoluble and falls as a sediment, and a sort of sugar termed *indiglucin*. The indigo is collected, drained, pressed, and dried. By action of deoxidizing agents indigo is converted into soluble colorless *indigogen*, *reduced indigo*, or *indigo white*; 1 part of powdered indigo, 2 of ferrous sulphate, 3 of slaked lime, and 200 of water, shaken together and set aside in a well-closed bottle, give this *colorless indigo* ($C_{16}H_{12}N_2O_2$). A piece of yarn, calico, or similar fabric dipped into such a solution and exposed to air becomes dyed blue, deposition of insoluble indigo-blue occurring within the cells and vessels of the fibre. This operation is readily performed on the small scale, and forms an illustration of a characteristic feature of the art of dyeing—namely, the introduction of soluble coloring-matter into a fabric by permeation of the walls of its cellular and vascular tissue, and the imprisonment of that coloring-matter within the cells and vessels by conversion into a solid and insoluble form. (*Vide* also p. 140.) Indigo is probably a derivative of benzene.

Pure indigo, or *indigotin*, may be obtained in beautiful needles by spreading a paste of indigo and plaster of Paris on a tin plate, and when quite dry placing a lamp underneath, moving the latter from place to place as the indigo sublimes and condenses on the surface of the plaster. It may also be obtained in crystals by gently boiling finely-powdered indigo with aniline, filtering while hot, and setting aside; these crystals may be washed with alcohol. Hot paraffin may be employed instead of aniline. Indigo may be produced artificially. Toluene, from coal-tar, is by Perkin's process converted into cinnamic acid, this into a nitro-derivative, and this again into orthonitrophenylpropionic acid. From the latter alkali and grape-sugar deposit crystalline indigo (Baeyer).

Distinction between Nitric Acid and Other Nitrates.—Presence of

the nitric radical in a solution having been proved by the above reactions, its occurrence as the nitrate of a metal is demonstrated by the neutral or nearly neutral deportment of the liquid with test-paper and the detection of the metal—its occurrence as nitric acid by the sourness of the liquid to the taste and the strong effervescence produced on the addition of a carbonate.

Antidote.—In cases of poisoning by strong nitric acid solution of sodium carbonate (common washing soda) or magnesia and water may be administered as an antidote.

QUESTIONS AND EXERCISES.


Trace the origin of nitrates.—In what does cubic nitre differ from prismatic nitre?—Describe a process by which potassium nitrate may be obtained artificially.—State the difference between potassium nitrate, nitre, saltpetre, and sal prunella.—What group of elements is characteristic of all nitrates? and what claim has this group to the title of radical?—Mention the usual theory regarding the manner in which atoms are arranged in reference to each other in such salts as potassium nitrate.—How is the official nitric acid prepared?—Give the properties of nitric acid.—What reactions occur when strong nitric and hydrochloric acids are mixed?—How is nitric oxide prepared?—Enumerate and explain the tests for nitrates?—Into what substances does nitric acid usually split when employed as an oxidizing agent?—How is nitrous oxide prepared? Enumerate the five oxides of nitrogen.—What is the nature of gun-powder?—Write a few sentences on the chemistry of indigo, one of the tests for nitric acid.—How is nitric acid distinguished from other nitrates?—What quantity of cubic nitre will be required to produce ten carboys of official nitric acid, each containing 114 lbs.? *Ans.* 1076½ lbs.

CHLORIC ACID AND OTHER CHLORATES.

Formula of the acid, HClO_3 or ClO_2OH . Molecular weight, 84.5. Chlorates (p. 293) are made from hypochlorites.

Hypochlorous Acid (HClO or ClOH) and other Hypochlorites.


Place a few grains of red mercury oxide in a test-tube, half fill the tube with chlorine-water, and well shake the mixture; the resulting liquid is a solution of hypochlorous acid, mercuric oxychloride remaining undissolved:



By the metathesis (double decomposition) of hypochlorous acid and oxides of hydrates other pure hypochlorites are formed: $\text{HClO} + \text{NaHO} = \text{NaClO} + \text{H}_2\text{O}$.

The direct action of chlorine on metallic hydrates and some carbonates is supposed to give a compound of chloride and hypochlorite, as described in connection with the synthetical

reactions of calcium (p. 114, *Calc Chlorata*, U. S. P.). (See also p. 87, *Liquor Sodæ Chloratæ*, U. S. P.)



The condition of the chlorine in these bodies is not satisfactorily made out, so that their constitution is not definitely determined. The action of acids on them results in the evolution of chlorine; hence the great value of the calcium compound (chlorinated lime, or chloride of lime) in bleaching operations:




The solubility of hypochlorites in water, their peculiar odor, greatly intensified on the addition of acid, and their bleaching powers (see the above calcium reaction) are the characters on which to rely in searching for hypochlorites.

Chlorates.

The group of elementary atoms represented by the formula ClO_3 is that characteristic of chloric acid and all other chlorates; hence it is expedient to regard it as being an acidulous radical, which may be termed the chloric radical. Like the nitric radical, it has not been isolated. Chloric anhydride (Cl_2O_5), unlike nitric anhydride, has not yet been obtained in the free condition.

Chlorates are artificial salts. They are formed by simply boiling aqueous solutions of the common bleaching salts (chlorinated lime, chlorinated soda, chlorinated potash). Heat thus converts




One chlorate may also be made from another by double decomposition. In making chlorates economically the chlorinated salt is, of course, at once converted into chlorate.

Potassium Chlorate.

Thus, Potassium Chlorate (*Potassii Chloras*, or *Chlorate of Potassium*, U. S. P.), the old chlorate of potash, is commercially made by saturating with chlorine gas a moistened mixture of 3 parts of potassium chloride and 10 of slaked lime, and well boiling the product. Chlorinated lime is first formed;

this, on continued boiling with water, splits up into calcium chloride and calcium chlorate; and the latter, reacting on the potassium chloride, yields calcium chloride and potassium chlorate.



The operation may be conducted on a small scale by rubbing together in a mortar the above proportions of ingredients in ounces or half ounces, adding enough water to make the whole assume the character of damp lumps, placing the porous mass in a funnel (loosely plugged with stones or pieces of glass), and passing chlorine gas (p. 29) up through the mass by attaching the tube delivering the gas to the neck of the funnel. When the whole mass has become of a slight pink tint (due to a trace of permanganate), it should be turned into a dish, well boiled with water, filtered, the filtrate evaporated if necessary, and set aside; the potassium chlorate crystallizes out in colorless rhomboidal plates, calcium chloride remaining in the mother-liquor.


In the official process potassium carbonate is alluded to as being used in place of chloride, but otherwise the method is similar to that just described. Chlorinated potash and chlorinated lime are first formed:




These on boiling with water split up into chlorates and chlorides:



the whole of the potassium chloride and calcium chlorate finally yielding potassium chlorate and calcium chloride:



Neglecting intermediate decompositions, the reactions may be represented by the following equation:



Sodium Chlorate (*Sodii Chloras*, *Chlorate of Sodium*, U. S. P.), NaClO_3 , is similarly prepared.

Potassium chlorate is soluble in water to the extent of 6 or 7 parts in 100 at common temperatures. It is usually administered medicinally in aqueous solution, sometimes also in lozenges (*Tro-*

chisci Potassii Chloratis, U. S. P.). Potassium chlorate must on no account be rubbed with sulphur or sulphides in a mortar or otherwise, friction of such a mixture resulting in violent explosion.

Potassium chlorate, when heated, yields potassium chloride and oxygen, and is the salt commonly employed in the preparation of the gas for experimental purposes. But if the action be carried on at as low a temperature as possible, and be arrested when 100 parts of the chlorate have (Teed) yielded 7.84 parts of oxygen, the residual salt will be found to contain only potassium perchlorate (KClO_4) and chloride: $10\text{KClO}_3 = 6\text{KClO}_4 + 4\text{KCl} + 3\text{O}_2$. A higher temperature causes the decomposition of the perchlorate: $\text{KClO}_4 = \text{KCl} + 2\text{O}_2$. When the chlorate is heated with manganese peroxide no perchlorate is formed.

Perchloric Acid (HClO_4).—Crude potassium perchlorate, obtained as just indicated, is boiled (in a fume-cupboard) with hydrochloric acid to decompose any chlorate that may be remaining, and then separated from chloride by washing and crystallization, chloride being far more soluble in water than perchlorate. Perchloric acid is then obtained by distilling the potassium perchlorate with sulphuric acid; it is stable, and is occasionally administered in medicine.

Chloric acid (HClO_3) may be isolated, but is unstable, quickly decomposing into chlorine, oxygen, and perchloric acid; some other chlorate (*e. g.* KClO_3) must therefore be used in studying the reactions of the chloric radical.

Table of the Chlorine Acids.

Hydrochloric acid	HCl .
Hypochlorous acid	HClO .
Chlorous acid	HClO_2 .
Chloric acid	HClO_3 .
Perchloric acid	HClO_4 .

The chloric radical is univalent (ClO_3'). The acidulous radicals of the other chlorine acids are also univalent, as indicated in the foregoing formulæ.

ANALYTICAL REACTIONS (TESTS).

First Analytical Reaction.—To solution of a chlorate (*e. g.* potassium chlorate) add solution of silver nitrate; no precipitate falls, showing that the chlorine must be performing different functions from those it possesses in chlorides. Evaporate the solution to dryness and place the residue in a small dry test-tube, or at once drop a fragment of a chlorate into a test-tube and heat strongly; oxygen is evolved, and may be recognized by its power of rekindling an incandescent match inserted in the tube. Boil the residue with water, and again add solution of silver nitrate; a white precipitate falls having all the

characters of silver chloride, as described under Hydrochloric Acid.

Second Analytical Reaction.—To a fragment of a chlorate add two or three drops of strong sulphuric acid; an explosive gas (Cl_2O_4) is evolved, somewhat resembling chlorine in odor, but possessing a deeper color than that element.




Warm the upper part of the test-tube to 150° or 200° F., or introduce a hot wire; a sharp explosion ensues, due to decomposition of the gas, chlorine peroxide, into its elements.

Third Analytical Reaction.—Heat a small fragment of a chlorate with hydrochloric acid; a yellowish-green explosive gas, termed *euchlorine*, is evolved. Its color is deeper than that of chlorine; hence the name (from $\epsilon\upsilon$, *eu*, well, and $\chi\lambda\omega\rho\acute{o}\varsigma$, *chloros*, green). In odor it resembles chlorine, and is probably a mixture of that element with one of the oxides of chlorine.

Fourth Analytical Reaction.—Direct the blowpipe flame on to charcoal until a spot is red hot, and then place on the spot a fragment of a chlorate; deflagration ensues, as with nitrates.

Bromates.


Bromates are salts closely resembling chlorates and iodates. The formula of bromic acid is HBrO_3 . The presence of bromates in bromides is shown by the production of a yellow color on the addition of diluted sulphuric acid.



Iodates.

Iodic Acid (HIO_3), or *Hydrogen Iodate*.—Iodine is warmed in a flask with five times its weight of the strongest nitric acid (sp. gr. 1.5), in a fume-cupboard, until all action ceases. On cooling, iodic acid separates in small pyramidal crystals. These are removed, the residual liquid evaporated to dryness to remove excess of nitric acid, the residue and the first crop dissolved in a small quantity of boiling water, and the solution set aside to crystallize. Neutralized with carbonates or hydrates, it yields *iodates*.

Potassium iodate and sulphurous acid decompose each other with elimination of iodine (or with formation of a blue color if starch be present). Sulphurous acid occurring as an impurity in acetic and other acids may thus be detected.



Ferric Iodate, or rather *Oxyiodate* ($\text{Fe}_2\text{O}_4\text{IO}_3, 8\text{H}_2\text{O}$), is precipitated on adding solution of ferric chloride to solution of potassium iodate.

QUESTIONS AND EXERCISES.

How may hypochlorous acid be formed?—What are the relations of hypochlorous acid to bleaching-powder?—By what reaction is chlorine eliminated from hypochlorites?—State the general reaction by which chlorates are formed.—Give details of the preparation of potassium chlorate.—Mention the properties of potassium chlorate.—What decompositions occur when potassium chlorate is heated?—Find the molecular weight of potassium chlorate.—What weight of oxygen is yielded when 1 oz. of potassium chlorate is completely decomposed? and how much potassium chloride remains?—100 cubic inches of oxygen, at 60° F. and barometer at 30 inches, weighing 34.203 grains, and 1 gallon containing 277½ cubic inches, what weight of potassium chlorate will be required to yield 10 gallons of the gas?—*Ans.* 5½ oz.—How many cubic inches of oxygen are producible from 1 oz. of potassium chlorate?—Calculate the weight of potassium chlorate theoretically obtainable from 100 parts of chloride.—How is perchloric acid prepared?—Enumerate the chlorine acids.—How may the presence of chlorides in chlorates be demonstrated?—Mention the tests for chlorates.—Give the formula of chlorine peroxide.—What is euchlorine?—How may iodic acid be made?—Describe the preparation of potassium iodate?

ACETIC ACID AND OTHER ACETATES.

Formula of Acetic Acid, $\text{HC}_2\text{H}_3\text{O}_2$. Molecular weight, 60.

Source.—Acetic Acid, the Hydrogen Acetate or Acetate of Hydrogen, is said to occur naturally in certain plant-juices and animal fluids in minute proportions, but otherwise is an artificial product. Much is furnished by the destructive distillation of wood. When first discovered this was regarded as a new acid, and was named *pyroligneous acid*, a hybrid word from $\pi\upsilon\rho$, *pŭr*, fire, and *lignum*, wood, a term still retained for the crude acid. The latter, neutralized by calcium carbonate, the whole evaporated, and the residue gently heated to drive off volatile tarry matters, gives calcium acetate, which, after conversion into sodium acetate and recrystallization, furnishes by distillation with sulphuric acid and water acetic acid in a state of purity. Diluted acetic acid is *white vinegar*, one of the now many varieties of *vinegar*. It has been known as *wood vinegar* for the past sixty years. It is generally colored brown with caramel to meet the taste of the public. In Germany and France large quantities of acetic acid are made by the spontaneous oxidation of the alcohol in inferior wines in the presence, according to Pasteur, of a plant-ferment termed *Mycoderma Aceti* (the *Bacterium Mycodermi* of Cohn); hence the *white-* and *red-wine vinegar* (*vinegar*, from the French *vin*, wine, and *aigre*, sour). Indeed, this bacterium may be propagated, and the artificial manufacture of vinegar from alcohol and water be carried out, by its aid, on a larger scale. In England also the domestic form of acetic acid (brown vinegar) commonly has an alcoholic origin: infusion of malt and unmalted grain, or sometimes the latter alone after treatment with sulphuric acid, is fermented; and the resulting alteration of its sugar, instead of being arrested when the product is an alcoholic liquid, a sort of beer, is allowed to go on to the next stage, acetic acid; it usually


contains from 3 to 6 per cent. of real acetic acid ($\text{HC}_2\text{H}_3\text{O}_2$). Different strengths of vinegar are sold under the numbers 16, 18, 20, 22, and 24, corresponding to the number of grains of anhydrous sodium carbonate neutralized by 1 Imperial fluidounce of the vinegar. All of these "brewed vinegars" are further colored with caramel to suit the popular taste. *Vinegar* is a generic term applicable to any one or to all varieties. Its essential component is acetic acid.

Official Vinegars.—The British official "vinegar" (*Acetum*, B. P.) contains $5\frac{1}{2}$ (5.41) per cent. of acid. The so-called vinegar of cantharides (*Acetum Cantharidis*, B. P.) is a solution of the active principle of cantharides in very strong acetic acid, not in official vinegar. The vinegars of squill (*Acetum Scillæ*, U. S. P.) and of ipecacuanha (*Acetum Ipecacuanhæ*, B. P.) contain diluted acetic acid; that is, *wood vinegar*, not the official "vinegar," which is that made from malted and unmalted grain, commonly 1 part of the former to 3 of the latter. In the British Pharmacopœia "vinegar," except rare occasional use for its own sake, is only employed in the preparation of *Emplastrum Saponis Fuscum*.

The *Acetum Opii*, U. S. P., or *Black Drop* of America, is made from nutmeg, saffron, and sugar, as well as opium and diluted acetic acid.

The Acetic Radical.—The group $\text{C}_2\text{H}_3\text{O}_2$ is that characteristic of acetic acid and other acetates. It is univalent.

Acetyl.—The characteristic group in acetates ($\text{C}_2\text{H}_3\text{O}_2$) is considered to contain, rather than to be, a radical—the radical $\text{C}_2\text{H}_3\text{O}$, termed *acetyl*. Acetates yield a body having the composition $\text{C}_2\text{H}_3\text{OCl}$, which is regarded as acetyl chloride; from this may be obtained acetic anhydride ($\text{C}_4\text{H}_6\text{O}_3$), which, by absorbing water, becomes acetic acid.



The quantitative relation of acetic acid to alcohol will be evident from the following equation, representing, empirically, the formation of the acid:



Acetates in aqueous solution are liable to decomposition. In solution of acetate of morphine a myceloid growth occasionally forms, acetic acid disappears, and morphine is deposited. Solution of ammonium acetate is liable to a similar change, gradually becoming alkaline.


SYNTHETICAL REACTION.

Acetic Acid.


To a few grains of sodium acetate in a test-tube add a little water and some sulphuric acid, and heat the mixture; acetic acid is evolved, and may be condensed by a bent tube adapted to the test-tube by a cork in the usual way.

Acetic Acid, or *Hydrogen Acetate*, or *Acetate of Hydrogen*.—This is the process by which sodium or calcium acetate (the neutralized products of the distillation of wood) is made to yield acetic acid on the large scale. As with nitric and hydrochloric acids, the loose term “acetic acid” is that usually applied to aqueous solutions of acetic acid. The *Acidum Aceticum*, U. S. P., contains 36 per cent. of real acid—that is, of $\text{HC}_2\text{H}_3\text{O}_2$; for it contains only 30.6 per cent. of *acetic anhydride* ($\text{C}_4\text{H}_6\text{O}_3$), still occasionally though somewhat obscurely termed *anhydrous acetic acid*. Its specific gravity is 1.048. *Acidum Aceticum Dilutum*, U. S. P., contains 6 per cent. of $\text{HC}_2\text{H}_3\text{O}_2$. Sp. gr. 1.0083. Glacial acetic acid ($\text{HC}_2\text{H}_3\text{O}_2$) contains no water. It solidifies to a crystalline mass at and below 15°C . (59°F .), hence the appellation *glacial* (from *glacies*, ice). Sp. gr. from 1.056 to 1.058. Good commercial glacial acetic acid (*Acidum Aceticum Glaciale*, U. S. P.) does not contain more than 1 per cent. of water. Although water is lighter than this acetic acid, yet addition of water at first renders the acid heavier; evidently, therefore, condensation, or contraction in bulk, occurs on mixing the liquids: after 10 per cent. has been added the addition of more water produces the usual effect of dilution of a heavy liquid by a lighter—namely, reduction of relative weight. This matter will be better understood after the subject of specific gravity has been studied. Glacial acetic acid mixes readily with most oils.


The following equation shows the reaction :



or, assuming the existence of acetyl ($\text{C}_2\text{H}_3\text{O}$) in acetic acid (united with hydroxyl HO), and the radical sulphuryl (SO_2) in sulphuric acid (united with 2HO),



or, on the assumption that salts contain the oxide of a basylous radical united with the anhydride of an acid (the old view under which such names as acetate of soda were formed),



Note on the Constitution of Salts.

Which of these three equations—or, more broadly, which of the three views of the constitution of salts illustrated by the equations—is correct is questionable. Whether it is $\text{C}_2\text{H}_3\text{O}_2$, $\text{C}_2\text{H}_3\text{O}$ or $\text{C}_4\text{H}_6\text{O}_3$ which migrates from one acetic compound to another, whether it is SO_4 , SO_2 , or SO_3 which migrates from one sulphuric compound to another, and so on with other acidulous groupings, cannot at present be determined. There are some objections to each view, and possibly neither is right. Either the given radicals cannot be isolated, or application of the forces of heat, light, and electricity do not confirm views arrived at by the results of operations with the chemical force; or a salt comes to be regarded as having so large a number

of constituent parts that the view breaks down from the sheer inability of students, at all events at the present stage of study, to grasp the complicated analogies involved. Yet for the purposes of description, study, and conversation some system must be adopted. Let the first, then, be taken for the present, over-reliance on it being checked by the use of general instead of special names for the hypothetical radicals, and other systems be employed in other cases. (See also p. 286.)

ANALYTICAL REACTIONS (TESTS).

First Analytical Reaction.—To an acetate add sulphuric acid and heat the mixture; a characteristic odor (acetic acid) is evolved.

Note 1.—Iodine, sulphurous acid, and other substances of powerful odor mask that of acetic acid; they must be removed, therefore, usually by precipitation or oxidation, before applying this test.

Note 2.—It will be noticed that this reaction is identical with the previous one; it has synthetical or analytical interest according to the object and method of its performance.

Second Analytical Reaction.—Repeat the above reaction, a few drops of spirit of wine being first added to the acetate; acetic ether (ethyl acetate, $C_2H_5C_2H_3O_2$), also of characteristic odor, is evolved. The basylous radical ethyl (C_2H_5) will be referred to subsequently.

Third Analytical Reaction.—Heat a fragment of a dry acetate in a test-tube, and again notice the odor of the gaseous products of the decomposition; among them is *acetone* (C_3H_6O), the smell of which is characteristic. Carbonate of the metal remains in the test-tube.

Fourth Analytical Reaction.—To a solution of an acetate, made neutral by the addition of acid or alkali as the case may be, add a few drops of neutral solution of ferric chloride; a deep-red liquid results, owing to the formation of ferric acetate ($Fe_26C_2H_3O_2$). Boil; a red precipitate (iron oxyacetate) occurs, leaving the liquid colorless.

Analytical Note.—It will be noticed that the formation of characteristic precipitates, the usual method of removing radicals from solution for recognition, is not carried out in the qualitative analysis of acetates. This is because all acetates are soluble. Silver acetate ($AgC_2H_3O_2$) and mercurous acetate ($HgC_2H_3O_2$) are only sparingly soluble in cold water, but the fact can seldom be utilized in analysis. Hence peculiarities of color and odor, the next best characters on which to rely, are adopted as means by which acetates may be detected. Acetates, like other organic compounds, char when heated to a high temperature.

Note on Anhydrides.—Up to this point the student has regarded an anhydride as a body derived from an acid by removal of the

whole of the hydrogen of the acid, together with as much of its oxygen as with the hydrogen forms water. This definition will scarcely apply to acetic anhydride, and must therefore be somewhat qualified. An *anhydride* is derived from an acid, the acid having lost the whole of its *basylous* hydrogen and so much oxygen as is necessary to form water with that hydrogen. Anhydrides are obtained by heating acids and by other methods.

QUESTIONS AND EXERCISES.

What is the formula of acetic acid?—State the relation of acetic acid to other acetates.—What is the molecular weight of acetic acid?—Name the sources of acetic acid.—What is pyroligneous acid?—From what compound is the acetic acid of most foreign and English varieties of vinegar derived?—How much real acid is contained in official vinegar?—What is the nature of the “vinegars” of pharmacy?—How may acetic acid be obtained from sodium acetate?—How much real acid is contained in the official acetic acid?—Mention the strength of commercial glacial acetic acid.—Give three or more views of the constitution of acetates, illustrating each by formulæ.—Enumerate the tests for acetates.

HYDROSULPHURIC ACID AND OTHER SULPHIDES.

Formula of Hydrosulphuric Acid, H_2S . Molecular weight, 34.

Source and Varieties of Sulphur.—The acidulous radical of *hydrosulphuric acid*, *sulphydric acid*, *hydrogen sulphide*, or *sulphuretted hydrogen* and other sulphides is the element sulphur (S). It occurs in nature in the free condition, also in combination with metals, as already stated in describing the ores of some of the metals. The coal-gas industry furnishes much sulphur as a by-product, it occurring in coal, chiefly as iron pyrites. Most of the sulphur used in medicine is imported from Sicily, where it occurs chiefly associated with blue clay. It is purified by fusion, sublimation, or distillation. Melted and poured into moulds, it constitutes a crystalline mass termed *roll sulphur*. If distilled and the vapor carried into large chambers, so that it may be rapidly condensed, the crystals are so minute as to give the sulphur a pulverulent character; this is *sublimed sulphur* (*Sulphur Sublimatum*, U. S. P.) or *flowers of sulphur*: the same washed with dilute ammonia, to remove the traces of sulphuric acid (often 0.1 per cent., resulting from very slow oxidation of sulphur in ordinary moist air) or, possibly, arsenous sulphide, constitutes *Sulphur Lotum*, U. S. P. The third common form, *milk of sulphur*, will be noticed subsequently. Sulphur also occurs in nature in combination as a constituent of animal and vegetable tissues, as sulphurous acid gas (SO_2) in volcanic vapors, and as sulphuretted hydrogen in some mineral waters. *Plastic sulphur* is one of the allotropic varieties of the element, obtained on heating sulphur considerably beyond its melting-point and pouring into cold water.

Quantivalence.—Sulphur would seem to be sexivalent in sulphuric anhydride (SO_3), a substance which will be noticed under Sulphuric

Acid, and quadrivalent in sulphurous anhydride (SO_2), while it still oftener exhibits bivalent affinities (H_2S).

Molecular Weight.—At very high temperatures sulphur follows the rule that, under similar conditions of heat and pressure, molecular weights (in grammes, grains, etc.) of volatile elements occupy equal volumes of vapor; its formula therefore is S_2 , and molecular weight 64. At lower temperatures the volume weighs three times as much as it should do if following usual laws, and then the molecule would appear to contain six atoms (S_6).

Acid Salts.—Sulphur (S') being the first acidulous radical of bivalent activity met with in these sections on acids, it is desirable here to draw attention to a new class of salts to which such a radical will give rise. These are acid salts (as KHSO_4), which are intermediate between neutral salts (as K_2SO_4) and acids (as H_2SO_4). Univalent acidulous radicals, with an atom of hydrogen, give an acid, and with an atom of other basylous radicals an ordinary or neutral salt. But bivalent acidulous radicals, inasmuch as they give with two atoms of hydrogen an acid, and with two atoms of univalent metals a neutral salt, may obviously give intermediate bodies containing one atom of hydrogen and one atom of metal; these are appropriately termed acid salts: they are neither normal acids nor neutral salts, but *acid salts*. (Examples: KHCO_3 , NaHSO_4 , KHCO_3 , Na_2HPO_4 , CuHAsO_3 , CaH_2PO_4 .) Whether or not these and other salts give an acid reaction with blue litmus-paper depends on the strength of the respective radicals. Usually they do redden the test-paper, but sometimes not; thus potassium, sodium, or ammonium acid sulphides or sulphhydrates (KHS , NaHS , NH_4HS) have alkaline properties.*

The chemical analogy between sulphur and oxygen, already once alluded to (p. 177), is further illustrated by the compounds just mentioned. Thiacetic acid ($\text{CH}_3\text{CO.SH}$) exists, and other *thio* (*θειον*, *theion*, sulphur) acids. Sulphur is also closely related to the rarer element selenium. Thus we have SeO_2 (selenious anhydride) as well as SO_2 (sulphurous anhydride), H_2SeO_3 (selenious acid) as well as H_2SO_3 (sulphurous acid), H_2SeO_4 (selenic acid) as well as H_2SO_4 (sulphuric acid). The rare element tellurium also seems to have similar analogies. The four hydrogen compounds of the group have the formulæ H_2O , H_2S , H_2Se , H_2Te . (*Vide* Index, "Periodic Law.")

SYNTHETICAL REACTIONS.

Hydrogen Sulphide, or Sulphuretted Hydrogen.

Synonyms.—Hydrosulphuric Acid; Sulphydric Acid.

First Synthetical Reaction. The preparation of sulphuretted


* Chemists regard these sulphhydrates as compounds of basylous radicals with HS , a univalent grouping termed hydrosulphyl (hydrogen persulphide, H_2S_2), just as hydrates are similarly viewed as compounds of the univalent radical hydroxyl (HO) (hydrogen peroxide, H_2O_2), H_2S becoming HHS (hydrogen hydrosulphylide), and H_2O becoming HHO (hydrogen hydroxylide).

hydrogen.—This operation was described on p. 97, and probably has already been studied by the reader.


Precipitated Sulphur.

Second Synthetical Reaction.—Prepare the variety of the radical of sulphides known as *Precipitated Sulphur* (*Sulphur Præcipitatum*, U. S. P.) or *Milk of Sulphur*, by boiling a few grains of flowers of sulphur (100 parts) with slaked lime (50 parts) and some water (500 parts) in a test-tube (larger quantities in an evaporating-basin), filtering, and (reserving a small portion of the filtrate) adding diluted hydrochloric acid until the well-stirred milk-like liquid still has a faint alkaline or scarcely acid reaction on test-paper; sulphur is precipitated, and may be collected on a filter, washed, and dried (at about 130° F.). Excess of acid must be avoided, or some hydrosulphyl, the liquid hydrogen persulphide (H_2S_2), will be formed, probably causing the particles of sulphur to aggregate to a gummy mass.


This is the process of the *pharmacopœias*. Calcium polysulphide and calcium hyposulphite are formed:



On adding the acid both salts are decomposed and, after an intermediate reaction, sulphur separates:




The calcium polysulphide yields sulphuretted hydrogen and milk-white sulphur on the addition of acid. The calcium hyposulphite then yields sulphurous acid gas as well as yellowish sulphur. The gases react, and give sulphur and water, very little sulphuretted hydrogen escaping: this is the intermediate reaction just alluded to. A little pentathionic acid (see Index) is also said to be formed.



Sulphur Lozenges (*Trochisci Sulphuris*, B. P.) contain, each, 5 grains of precipitated sulphur, 8 grains of sugar, 1 grain of gum and of cream of tartar, and 1 minim of mucilage and of tincture of orange-peel.

Calcareous Precipitated Sulphur, the old "Milk of Sulphur."—To a sulphur solution prepared as before (or to the reserved portion) add a little diluted sulphuric acid; the precipitate is in this case largely mixed with calcium sulphate:



Place a little of each of these specimens of "precipitated sulphur," with a drop of the supernatant liquid, on a strip of glass, cover each spot with a piece of thin glass, and examine the precipitates under a microscope; the pure sulphur will be found to consist of minute grains or globules, the calcareous to contain comparatively large crystals (hydrous calcium sulphate).

Note.—Some of the precipitated sulphur met with in trade is, in England, still thus mixed with calcium sulphate, most of such specimens containing two-thirds of their weight of that substance. Formerly, purchasers were so accustomed to the satiny appearance of the mixed article as to regard real sulphur with suspicion, sometimes refusing to purchase it. The mixed article is certainly somewhat more easily miscible with aqueous liquids.

Calcareous precipitated sulphur yields a white ash (anhydrous calcium sulphate) when a little is burnt off on the end of a table-knife or spatula or in a crucible. (No more damage is done to the steel than a rub on a knife-board will remove.) To ascertain exactly the amount of the sulphate, place a weighed quantity in a tared crucible and heat till no more vapors are evolved. The weight of the residual anhydrous calcium sulphate ($\text{CaSO}_4 = 136$), with one-fourth thereof added, is the amount of crystalline calcium sulphate ($\text{CaSO}_4 \cdot 2\text{H}_2\text{O} = 172$) present in the original quantity of calcareous sulphur.

ANALYTICAL REACTIONS (TESTS).

To a sulphide add a few drops of hydrochloric acid; sulphuretted hydrogen will probably be evolved, well known by its smell. If the sulphide is not acted upon by the acid or if free sulphur be under examination, mix a minute portion with a fragment of solid caustic potash or soda and fuse on a silver coin or old spoon. When cold, place a drop of dilute hydrochloric acid on the spot; sulphuretted hydrogen is evolved, and a black stain, due to silver sulphate (Ag_2S), is left on the coin.

Other sulphur reactions may be adapted as tests, but the above are sufficient for all ordinary purposes. The most convenient reagent for detecting a sulphide in solution of ammonia is copper ammonio-sulphate, which gives a black precipitate (copper sulphide) if a sulphide be present.

The *Sulphur Iodide* (S_2I_2) has been mentioned under "Iodine." A *chloride* (S_2Cl_2) and *bromide* (S_2Br_2) may also be formed from their elements. A mixture of sulphur and sulphur chloride is sometimes met with under the name of *sulphur hypochloride*.

QUESTIONS AND EXERCISES.

In what forms does sulphur occur in nature?—State the modes of preparation of the three chief commercial varieties of sulphur.—To what

extent does the atom of sulphur vary in quantivalence?—State the relations of acid salts to acids and to normal salts.—Define sulphides and sulphydrates.—Describe the preparation of sulphuretted hydrogen.—What are the characters of pure precipitated sulphur?—Give equations explanatory of the reactions which occur in precipitating sulphur according to the official process.—Describe the microscopic test for calcareous precipitated sulphur.—Mention a ready physical method of detecting calcium sulphate in precipitated sulphur.—Mention the tests for sulphides and the character by which sulphuretted hydrogen is distinguished from other sulphides.—How are sulphides insoluble in acids tested for sulphur?—How would you detect a trace of sulphur in ammonia solutions?

SULPHUROUS ACID AND OTHER SULPHITES.


Formula of Sulphurous Acid, H_2SO_3 . Formula of Sulphurous Acid Gas or Sulphurous Anhydride, sometimes termed Sulphurous Acid, SO_2 . Molecular weight of Sulphurous Acid, 82; of the Anhydride, 64.

When sulphur is burnt in the air it combines with oxygen and forms sulphurous acid gas (SO_2), more correctly termed sulphurous anhydride, or occasionally, but erroneously, sulphurous acid. It is a pungent, colorless gas, readily liquefied on being passed through a tube externally cooled by a *freezing-mixture* composed of 2 parts of well-powdered ice (or, better, snow) with 1 part of common salt. If sulphurous acid gas becomes moist or is passed into water, heat is evolved and true sulphurous acid (H_2SO_3) is formed. The latter body may be obtained in crystals by freezing a strong aqueous solution; but it is very unstable, and hence the properties of the sulphurous radical must be studied under the form of some other sulphite; calcium sulphite (CaSO_3) or sodium sulphite (Na_2SO_3) may be used for the purpose.

Quantivalence.—The radical of the sulphites is bivalent (SO_3''), and hence forms acid sulphides, such as acid potassium sulphite (KHSO_3), and neutral sulphides, such as sodium sulphite (Na_2SO_3).

Note on Nomenclature.—The sulphites are so named from the usual rule, that salts corresponding with acids whose names end in *ous* have a name ending in *ite*. They are generally made by passing sulphurous acid gas over moist oxides or carbonates; in the latter case carbonic acid gas escapes.

Synthetical Reaction.—To a few drops of sulphuric acid in a test-tube add a piece of charcoal and apply heat; sulphurous acid gas is evolved, and may be conveyed by a bent tube into a small quantity of cold water in another test-tube. Larger quantities of the gas may be made in a Florence flask. The fluid product is the *Acidum Sulphurosum*, U. S. P. It should contain not less than 6.4 of the gas (SO_2), and have a sp. gr. of not less than 1.035. The process is also that described in the Pharmacopœia, except that the gas is purified by passing through a small wash-bottle before the final collection.



Sulphurous acid gas may also be made by boiling copper, mercury, or iron with sulphuric acid, sulphate of the metal being formed; also by boiling sulphur with sulphuric acid. The gas passed into water forms sulphurous acid.



If in this process the water were replaced by solutions of, or solid, metallic oxides or carbonates, sulphites of the various metals would be formed. The formula of sodium sulphite (*Sodii Sulphis*, U. S. P., the old sulphite of soda) is $\text{Na}_2\text{SO}_3 \cdot 7\text{H}_2\text{O}$; it occurs in colorless efflorescent prisms, soluble in water or spirit: under the name of *antichlor* it is used for removing traces of chlorine from paper pulp. The formula of sodium bisulphite (*Sodii Bisulphis*, U. S. P.) is NaHSO_3 . The so-called *Bisulphite of Lime*, used by brewers for retarding or arresting fermentation and oxidation, and employed for various antiseptic purposes, is a solution of calcium sulphite (CaSO_3) in free sulphurous acid (H_2SO_3), and is made by passing sulphurous acid gas (SO_2) into thin milk of lime. Its specific gravity varies from 1.050 to 1.070, and its potential strength of anhydride (SO_2) from 4 to 6 per cent. Sulphurous acid is very soluble in alcohol.

ANALYTICAL REACTIONS (TESTS).

First Analytical Reaction.—To a sulphite (sodium sulphite, for instance, made by passing sulphurous acid gas into solution of sodium carbonate) add a drop or two of diluted hydrochloric acid; a well-known peculiarly pungent smell results (sulphurous acid).

This smell is the same as that evolved on burning lucifer-matches that have been tipped with sulphur. It is due probably not to the gas (SO_2), but to the sulphurous acid (H_2SO_3) formed by the union of sulphurous acid gas with either the moisture of the air or that on the surface of the mucous membrane of the nose. The gas is highly suffocating.

Second Analytical Reaction.—To a sulphite add a little water, a fragment or two of zinc, and then hydrochloric acid; sulphuretted hydrogen will be evolved, known by its odor and by its action on a piece of paper placed like a cap on the mouth of the test-tube and moistened with a drop of solution of lead acetate, black lead sulphide being formed. Sulphurous acid may be detected in acetic acid or in hydrochloric acid by this test: $\text{H}_2\text{SO}_3 + 6\text{H} = \text{H}_2\text{S} + 3\text{H}_2\text{O}$.

Other Analytical Reactions.

To solutions of neutral sulphites add barium nitrate or chloride, calcium chloride, or silver nitrate; in each case white precipitates result (sulphites of the various metals). The barium sulphite is soluble in weak hydrochloric acid; but if a drop or two of chlorine-water is first added, barium sulphate is formed, which is insoluble in acids. The other precipitates are also soluble in acids. The silver sulphite is decomposed on boiling, sulphuric acid being formed and metallic silver set free, the mixture darkening in color.

To recognize the three radicals in an aqueous solution of sulphides, sulphites, and sulphates add barium chloride, filter, and wash the precipitate. In the filtrate sulphide are detected by the sulphuretted hydrogen evolved on adding an acid. In the precipitate sulphites are detected by the odor of sulphurous acid, produced on adding hydrochloric acid, and sulphates by their insolubility in the acid.

QUESTIONS AND EXERCISES.

What are the differences between sulphurous acid and sulphuric acid gas, sulphites and acid sulphites?—State the characters of sulphurous acid gas.—How is the official sulphurous acid prepared?—By what tests may sulphurous acid be recognized in acetic acid?—Give a method by which sulphites may be detected in presence of sulphides and sulphates.

SULPHURIC ACID AND OTHER SULPHATES.

Formula of the acid, H_2SO_4 or $\text{SO}_2(\text{OH})_2$. Molecular weight, 98.


Many sulphates occur in nature, but the common and highly important *hydrogen sulphate*, sulphuric acid, is made artificially.

Preparation of the Acid.—General Nature of the Process.—Sulphur itself, or generally the sulphur in iron pyrites, is first converted into sulphurous acid gas by burning in air, and this gas, by moisture and oxygen, into sulphuric acid ($\text{SO}_2 + \text{H}_2\text{O} + \text{O} = \text{H}_2\text{SO}_4$).


Details of the Process.—The oxygen necessary to oxidize the sulphurous acid gas cannot directly be obtained from air, but indirectly, the agency of nitric oxide (NO) and nitrous anhydride (N_2O_3) being employed, these gases becoming nitric peroxide (NO_2) by action of the air, and the nitric peroxide again becoming reduced by the action of the sulphurous acid gas, and so on. A comparatively small quantity of nitric gases will thus act as carrier of oxygen from the air to very large quantities of sulphurous acid.

The nitric oxide and nitrous anhydride are, in the first instance, obtained from nitric acid, and this from sodium nitrate, by the action of a small quantity of the sulphuric acid of a previous operation.

The following equations represent the chief successive steps :




Possibly the nitrous anhydride may, instead of absorbing oxygen, yield oxygen and become reduced to nitric oxide, or may act in both ways. Probably also *nitrosulphonic acid* $\text{SO}_2 < \begin{smallmatrix} \text{OH} \\ \text{NO}_2 \end{smallmatrix}$ is formed both from nitrous anhydride and nitric peroxide, and is then resolved by the steam into sulphuric acid and the lower oxides. And so on, over and over again.



On the large scale the sulphurous acid gas is carried, together with the nitric vapors, by flues into leaden chambers, where jets of steam supply the necessary moisture. The resulting dilute sul-

Fig. 38.





Experimental Manufacture of Sulphuric Acid.

phuric acid is concentrated by evaporation in leaden, and finally in glass or platinum, vessels.

Other Processes.—Sulphuric acid may be obtained by other processes, as by distilling ferrous sulphate resulting from the natural oxidation of iron pyrites by air; but it is not so made at the present day. The ferrous sulphate was formerly called *green vitriol* (p. 144), and the distilled product *oil of vitriol*, the latter in allusion to its consistence and origin.

Experiment.—For purposes of practical study a small quantity may be made, as shown in Fig. 39, by passing (a) sulphurous acid gas (p. 305), (b) nitric oxide in small quantity (p. 289), (c) air (forced through by aid of bellows or a gas-holder, or drawn through a *water-aspirator*, e), and occasionally (d) steam (generated in a Florence flask), through glass tubes nearly to the bottom of a two- or three-quart flask.

A slow current of sulphurous acid gas, air, and steam and a small quantity of nitric oxide will furnish, in the course of a few minutes, enough sulphuric acid for recognition by the first of the following analytical reactions. The manufacturing process may be more exactly imitated by burning sulphur in a tube placed where the flask *a* is represented in the foregoing figure, or by burning it under a funnel there attached; but in either case strong aspiration must be maintained. An instructive result also will follow the cessation of the steam-current for a short time—namely, the growth on the inner surface of the large flask of “chamber crystals,” which are either nitro-sulphonic acid or nitrosyl-sulphuric acid:



Purification.—Sulphuric acid may contain arsenic, nitrous compounds, and salts (lead sulphate, etc.). Arsenic may be detected by the hydrogen test (p. 173) or the stannous-chloride test (p. 175), nitrous compounds by powdered ferrous sulphate (which acquires a violet tint if they are present), and salts by the residue left on boiling a little to dryness in a crucible in a fume-chamber. If only nitrous compounds are present, the acid may be purified by heating with about $\frac{1}{2}$ per cent. of ammonium sulphate, water and nitrogen being produced (Pelouze). If arsenic occurs, heat with a little nitric acid (or sodium nitrate), which converts arsenous anhydride (As_2O_3) into arsenic anhydride (As_2O_5), then add ammonium sulphate, and distil in a retort containing pieces of quartz and heated by an annular-shaped burner (to prevent “bumping”—see p. 282). This is the usual process of purification. The arsenic anhydride remains in the retort. (Arsenous anhydride would be carried over with the sulphuric acid vapors.) The distillation frees the acid from other salts (such as NaHSO_4) which are not volatile.

Quantivalence.—The sulphuric radical being bivalent (SO_4''), acid as well as neutral sulphates may exist. Acid potassium sulphate (KHSO_4) is an illustration of the former, sodium sulphate (Na_2SO_4) of the latter; *double sulphates* may also occur, such as potassium and magnesium sulphate ($\text{K}_2\text{SO}_4, \text{MgSO}_4, 6\text{H}_2\text{O}$). Sulphates generally contain water of crystallization.

Pure sulphuric acid (H_2SO_4) is of specific gravity 1.848. The best "oil of vitriol" of commerce, a colorless liquid of oily consistence, is of specific gravity 1.843, and contains about 98 per cent. of real acid (H_2SO_4). This is the *Acidum Sulphuricum*, B. P. A variety less pure than this "white" acid is known as "brown acid." The *Acidum Sulphuricum*, U. S. P., should not contain less than 92.5 per cent. of the pure acid (H_2SO_4) and a sp. gr. not below 1.835. The *Acidum Sulphuricum Dilutum*, U. S. P., contains 10 per cent. of the pure acid and has a sp. gr. of about 1.070, the dilution commonly resulting in the separation, as a slight white sediment, of lead sulphate usually present in the strong acid. *Acidum Sulphuricum Aromaticum*, U. S. P., an acid greatly diluted with alcohol, in which are dissolved oil of cinnamon and tincture of ginger, contains about 20 per cent. by weight of official acid; sp. gr. about 0.939. It has been stated that in this preparation the acid and alcohol form some sulphovinic acid or ethyl-sulphuric acid, but the author has been unable to detect the latter. There are some definite compounds of sulphuric acid with water; the first ($\text{H}_2\text{SO}_4, \text{H}_2\text{O}$) may be obtained in crystals.

Sulphuric anhydride (SO_3) occurs in white crystals. It was formerly called *anhydrous sulphuric acid*, but it has no acid properties. It may be made by distilling sulphuric acid with phosphoric anhydride ($\text{H}_2\text{SO}_4 + \text{P}_2\text{O}_5 = 2\text{HPO}_3 + \text{SO}_3$). On the large scale sulphuric acid is dissociated by heat, and the dried sulphurous anhydride and oxygen made to recombine. It appears to unite with sulphuric acid and some other sulphates to form compounds ($\text{R}'_2\text{SO}_4, \text{SO}_3$) resembling in constitution red potassium chromate or *pyrochromate*. The fuming sulphuric acid ($\text{H}_2\text{SO}_4, \text{SO}_3$), sometimes termed *pyrosulphuric acid* ($\text{H}_2\text{S}_2\text{O}_7$), formerly made at Nordhausen in Saxony, seems to be such a body.

Note.—Sulphuric acid is a most valuable compound to all chemists and manufacturers of chemical substances. By its agency, direct or indirect, many, if not most, chemical transformations are effected. To describe all its uses would be to write a work on Chemistry.

ANALYTICAL REACTIONS (TESTS).

First Analytical Reaction.—To solution of a sulphate add solution of a barium salt; a white precipitate (barium sulphate, BaSO_4) falls. Add nitric acid and boil the mixture; the precipitate does not dissolve.

This reaction is as highly characteristic of sulphates as it has been stated to be of barium salts. (*Vide* p. 104.) The only error likely to be made in its application is that of overlooking the fact that barium nitrate and chloride are less soluble in strong acid than in water. On adding the barium salt to the acid liquid, therefore, a white precipitate may be obtained, which is simply barium nitrate or chloride. The appearance of such a precipitate differs considerably from that of the barium sulphate; hence a careful operator will not be misled. Should any doubt remain, water should be

added, which will dissolve the nitrate or chloride, but not affect the sulphate.

Second Analytical Reaction.—Mix a fragment of an insoluble sulphate (BaSO_4 , *e. g.*) with potassium or sodium carbonate, or, better, with both carbonates, and fuse the mixture in a small crucible. Digest the residue, when cold, in water, and filter; the filtrate may be tested for the sulphuric radical.

This is a convenient method of qualitatively analyzing insoluble sulphates, such as those of barium and lead.

Third Analytical Reaction.—Mix a fragment of an insoluble sulphate with a little alkaline carbonate on a piece of charcoal, taking care that some of the charcoal-dust is included in the mixture. Heat the little heap in the blowpipe flame until it fuses, and when cold add a drop of acid; sulphuretted hydrogen is evolved, recognized by its odor.

This is another process for the recognition of insoluble sulphates. Other preparations of sulphur, and sulphur itself, give a similar result. It is, therefore, rather a test for sulphur and its compounds than sulphates only, but the absence of other salts can be determined previously.

Note.—The presence of the sulphuric radical in a solution having been proved by the above reactions, its occurrence as a neutral sulphate of a metal is demonstrated by the neutral or nearly neutral deportment of the liquid with test-paper and the detection of the metal; its occurrence as sulphuric acid or an acid sulphate by the sourness of the liquid to the taste and the abundant effervescence produced on the addition of a carbonate.

Antidote.—In cases of poisoning by strong sulphuric acid, solution of sodium carbonate (common washing soda), magnesia and water, etc. may be administered as antidotes.

QUESTIONS AND EXERCISES.

State the formula and molecular weight of sulphuric acid.—How is it related to other sulphates?—Write a short article on the manufacture of sulphuric acid, giving either diagrams or equations.—How may nitrous compounds be detected in and eliminated from sulphuric acid?—State the method for detecting arsenic in sulphuric acid, and explain the process by which it may be removed.—Define sulphates, acid sulphates, and double sulphates.—What percentage of real acid is contained in oil of vitriol?—State the strength of “diluted” and “aromatic” sulphuric acid.—By what process is sulphuric anhydride obtained from the ordinary sulphuric acid?—Explain the reactions which occur in testing for sulphates.—Ascertain by calculation the weight of oil of vitriol (of 96.8 per cent.) necessary for the production of 1 ton of dry ammonium sulphate. *Ans.* 1718 lbs.—Name the antidotes in cases of poisoning by strong sulphuric acid.

CARBONIC ACID AND OTHER CARBONATES.

Formula of the acid, H_2CO_3 or $\text{CO}(\text{OH})_2$. Molecular weight, 62. Formula of Carbonic Acid Gas or Carbonic Anhydride, commonly termed Carbonic Acid, CO_2 . Molecular weight, 44.

Sources.—Carbonates (compounds containing the grouping CO_3) are very common in nature, the calcium carbonate (CaCO_3) being widely distributed as chalk, limestone, or marble. The hydrogen carbonate, true carbonic acid, is not known, unless, indeed, carbonic acid gas assumes that condition on dissolving in water. Such a solution (see p. 86) changes the color of blue litmus-paper, and the gas does not; this may be because only the true acid (H_2CO_3) affects the litmus, or because the gas (CO_2) cannot come into real contact with the litmus without a medium. From the commonest natural carbonate, calcium carbonate, is derived the carbonic constituent of the one most frequently used in medicine and in the arts generally, sodium carbonate.


Sodium Carbonate, the old carbonate of soda, is prepared from the chief natural salt, the chloride, by two different processes. The first is "the Leblanc process." After the chloride has been converted into sulphate (salt cake) by sulphuric acid (or by sulphurous acid, air, and steam—Hargreave's modification), $2\text{NaCl} + \text{H}_2\text{SO}_4 = \text{Na}_2\text{SO}_4 + 2\text{HCl}$, the sulphate is roasted with limestone and small coal, sodium carbonate and calcium sulphide being formed: $\text{Na}_2\text{SO}_4 + \text{C} + \text{CaCO}_3 = \text{CaS} + \text{Na}_2\text{CO}_3 + 2\text{CO}_2$. The residual mass (black ash) is digested in water, in which the sodium carbonate dissolves, the calcium sulphide remaining insoluble. The solution is evaporated to dryness, and yields crude sodium carbonate. This is roasted with a small quantity of sawdust to convert any caustic soda, resulting from the action of the lime on the carbonate, into neutral carbonate. The product is *soda-ash*. Dissolved in water and crystallized, it constitutes the ordinary "soda" used for washing purposes: recrystallized, and sometimes ground, it forms the official sodium carbonate (*Sodii Carbonas*, U. S. P.) ($\text{Na}_2\text{CO}_3, 10\text{H}_2\text{O}$.) The reaction will be more intelligible if regarded as occurring in two stages: 1st, reduction of the sodium sulphate to sulphide by the carbon of the coal, $\text{Na}_2\text{SO}_4 + \text{C} = \text{Na}_2\text{S} + 2\text{CO}_2$; 2d, reaction of the sodium sulphide and calcium carbonate, giving soluble sodium carbonate; thus, $\text{Na}_2\text{S} + \text{CaCO}_3 = \text{Na}_2\text{CO}_3 + \text{CaS}$.



The sulphur in the residual calcium sulphide may be recovered by exposure to the action of the waste carbonic acid gas of lime-kilns, calcium carbonate being formed, and the diluting nitrogen passing off, more carbonic acid afterward causing sulphuretted hydrogen to be set free. The latter is either burnt and converted into sulphuric acid, or is caused to react with air on ferric oxide, sulphur being set free.

The other process is "the ammonia process," so called because ammonia (NH_3) is used over and over again as the chief agent—or

chemical tool, so to say—in the factory. Strong brine from salt-beds is saturated first with ammonia gas (from coal which also yields gaseous fuel and coke), and then with carbonic acid gas (from coke-heated limestone). Sodium bicarbonate falls as a precipitate, and is afterward heated to furnish the sodium carbonate, with simultaneous temporary recovery of part of the carbonic acid gas. The ammonium chloride concurrently produced is, by the latest improvements, separated and heated, and its vapors passed over magnesia, magnesium chloride and regenerated ammonia gas resulting. Air passed over the heated magnesium chloride furnishes chlorine and recovered magnesia. The chlorine passed over lime (from the limestone) furnishes chlorinated lime. The former part of this process was discovered by Dyar and Hemming in 1838, and was applied, chiefly by Solvay, from 1861 to 1872; the working out of the chlorine-recovery by the magnesia method is largely due to Mond. The following notes will aid the student: *a.* Coal yields ammonia gas, gaseous fuel, and coke. *b.* $\text{CaCO}_3 = \text{CaO} + \text{CO}_2$. *c.* $2\text{NaCl} + 2\text{NH}_3 + 2\text{CO}_2 + 2\text{H}_2\text{O} = 2\text{NaHCO}_3 + 2\text{NH}_4\text{Cl}$. *d.* $2\text{NaHCO}_3 = \text{Na}_2\text{CO}_3 + \text{H}_2\text{O} + \text{CO}_2$. *e.* $2\text{NH}_4\text{Cl} + \text{MgO} = \text{MgCl}_2 + 2\text{NH}_3 + \text{H}_2\text{O}$. *f.* $\text{MgCl}_2 + \text{O} = \text{MgO} + \text{Cl}_2$. *g.* $\text{Cl}_2 + \text{CaO} = \text{CaOCl}_2$. (If the chlorine-recovery is not practised, then $2\text{NH}_4\text{Cl} + \text{CaO} = 2\text{NH}_3 + \text{CaCl}_2$, the latter being lost.) Rarely is a process carried on with such economy of elements as here obtains, while the salt, limestone, and fuel occur in England in close contiguity. After these raw materials, labor, wear and tear, loss, and capital are all paid for, single 1-cwt. casks of powdered bicarbonate (NaHCO_3) or of *granular monohydrous carbonate* ($\text{Na}_2\text{CO}_3 \cdot \text{H}_2\text{O}$) can be supplied in Great Britain at one penny per pound.



Carbonic Acid Gas (CO_2), termed also *Carbon Dioxide* and *Carbonic Anhydride*, is a product of the combustion of all carbonaceous matters. It is constantly exhaled by animals and inhaled by plants, its intermediate storehouse being the atmosphere, throughout which it is equally distributed by *diffusion* (*vide* p. 24) to the extent of about 4 parts in 10,000. A larger proportion than that just mentioned gives to confined air depressing effects, 4 or 5 per cent. rendering the atmosphere poisonous when taken into the blood from the lungs. Carbonic acid, however, may be taken into the stomach with beneficial sedative effects; hence, probably, much of the value of such effervescing liquids as soda-water, lemonade, and solutions of the various granulated preparations and effervescing powders. The gas liquefies on being compressed, and the liquid solidifies on being cooled. The weights of equal bulks of the gas, of air, and of hydrogen are as 22, 14.44, and 1. At common temperatures it dissolves in about its own volume of water, both being under the same pressure, the water retaining gas strongly if all air has been expelled.

Sulphocarbonates resemble carbonates in constitution, but contain sulphur in place of oxygen.

Sulphocarbonic Anhydride, CS_2 , commonly termed *Carbon Disulphide* or *Bisulphide* (*Carbonei Disulphidum*, U. S. P.), is a highly volatile and inflammable liquid, easily made from its elements. Sp. gr. 1.268 to 1.269 at 15°C .; boiling-point, 46°C . It may be rendered almost scentless by digestion with lime and then with copper turnings. Its possible impurities are dissolved sulphur, sulphur oils, and sulphuretted hydrogen. It is slightly soluble in water (about 1 in 400), forming a useful antiseptic fluid.

REACTIONS.

Synthetical and Analytical Reactions.—1. To a fragment of marble in a test-tube add water and then hydrochloric acid; carbonic acid gas (CO_2) is evolved, and may be conveyed into water or solutions of salts by the usual delivery-tube.

This is the process of the British Pharmacopœia, and the one usually adopted for experimental purposes. On the large scale the gas is prepared from chalk or marble and sulphuric acid, frequent stirring promoting its escape.

2. Pass the gas into lime-water; a white precipitate of calcium carbonate (CaCO_3) falls. Solution of lead subacetate may be used instead of, and is perhaps even a more delicate test than, lime-water.

The evolution of a gas on adding an acid to a salt, warming the mixture if necessary, the gas being inodorous and giving a white precipitate with lime-water, is sufficient evidence of the presence of a carbonate. Carbonates in solutions of ammonia, potash, or soda may be detected by the direct addition of solution of lime. Carbonates in presence of sulphites or hyposulphites may be detected by adding acid potassium tartrate, which decomposes carbonates with effervescence, but does not attack sulphites or hyposulphites.

3. Blow air from the lungs through a glass-tube into lime-water; the presence of carbonic acid gas is at once indicated.

The passage of a considerable quantity of normal air through lime-water produces a similar effect. A bottle containing lime-water soon becomes coated with calcium carbonate from absorption of atmospheric carbonic acid gas.

4. Fill a dry test-tube with the gas by passing the delivery-tube of the above apparatus to the bottom of the test-tube. Being rather more than once and a half as heavy as the air (1.529), it will displace the latter. Prove the presence of the gas by pouring it slowly, as if a visible liquid, into another test-tube containing lime-water; the characteristic cloudiness and precipitate are obtained on gently shaking the lime-water.

In testing for carbonates by bringing evolved gas into contact with lime-water, the preparation and adaptation of a delivery-tube

may often be avoided by pouring the gas from the generating-tube into that containing the lime-water in the manner just indicated.

5. Pass carbonic acid gas through lime-water until the precipitate at first formed is dissolved. The resulting liquid is a solution of calcium carbonate in carbonic acid water. Boil the solution; carbonic acid gas escapes, and the carbonate is again precipitated.

This experiment will serve to show how chalk is kept in solution in ordinary well-waters, giving the property of "hardness," and how the *fur* or stone-like deposit in tea-kettles and boilers is formed. It should be here stated that calcium sulphate produces similar hardness, and that these, with small quantities of magnesium sulphate and carbonate, constitute the hardening constituents of well-waters, a curd (calcium or magnesium oleate) being formed whenever soap is used with such waters. An enormous amount of soap is wasted through the employment of hard water for washing purposes. The hardness produced by the earthy carbonates is termed "temporary hardness," because removable by ebullition; that by the earthy sulphates, "permanent hardness," because unaffected by ebullition. The addition of lime-water or a mixture of lime and water removes temporary hardness (Reaction 2, p. 314), and sodium carbonate, "washing-soda," both temporary and permanent hardness, in the latter case sodium sulphate remaining in solution. Barium carbonate (ground witherite) also decomposes calcium and magnesium sulphates, barium sulphate being precipitated and calcium or magnesium carbonates formed; the latter and the carbonates originally in the water may then be precipitated by ebullition or by the action of lime-water. But the injurious effects of barium salts on man and the lower animals prevents the carbonate being used for purifying water for drinking purposes, as by accident or an unforeseen reaction a portion might become dissolved.

6. Add a solution of potassium or sodium carbonate to a mercuric salt; a brownish-red precipitate results. Add a solution of potassium or sodium bicarbonate to a mercuric solution; a white precipitate results, becoming red after some time.

QUESTIONS AND EXERCISES.

Name the chief natural carbonates.—What are the formulæ of carbonic acid and carbonic acid gas?—Adduce evidence of the existence of true carbonic acid.—Trace the steps by which the carbonic constituent of chalk is transferred to sodium by the two processes usually adopted in alkali-works—the manufacture of "soda."—Carbonic acid gas is constantly exhaled from the lungs of animals; why does it not accumulate in the atmosphere?—What is the effect of pressure on carbonic acid gas?—State the specific gravity of carbonic acid gas?—By what process may carbonic acid gas be obtained for experimental and manufacturing purposes?—Describe the action of carbonic acid gas on the potassium or sodium carbonate.—How may carbonic acid gas be detected in expired

air?—To what extent is carbonic acid gas heavier than air?—Work sums showing what quantity of chalk (90 per cent. pure) will be required to furnish the carbonic acid gas necessary to convert 1 ton of potassium carbonate (containing 83 per cent. of K_2CO_3) into acid carbonate, supposing no gas to be wasted. *Ans.* 1500 lbs.—Define “hardness” in water.—How may the presence of carbonates be demonstrated?

OXALIC ACID AND OTHER OXALATES.

Formula of Oxalic Acid, $H_2C_2O_4 \cdot 2H_2O$ or $(COOH)_2 \cdot 2H_2O$. Molecular weight, 126.

Synonym.—Hydrogen Oxalate.

Sources.—Oxalates occur in nature in the juices of some plants, as wood-sorrel, rhubarb, the common dock, and certain lichens, but the hydrogen oxalate (oxalic acid) and other oxalates are all made artificially. The carbon of many organic substances yields oxalic acid when those substances are boiled with nitric acid, and an alkaline oxalate when they are roasted with a mixture of potassium and sodium hydrates.

Experimental Process.—On the small scale a mixture of nitric acid 10 parts, loaf-sugar 2 parts, and water 3 parts quickly yields the acid. Abundance of red fumes are at first evolved. On cooling crystals are deposited. A more dilute acid, kept warm, acts more slowly, but yields a larger product. The following process is more economical:

Manufacturing Process.—On the large scale sawdust is roasted with alkalis, resulting sodium oxalate decomposed by lime, with regeneration of the soda and formation of calcium oxalate, the latter digested with sulphuric acid, and the liberated oxalic acid (oxalic acid of commerce, B.P.) made commercially pure by recrystallization.

Purified Oxalic Acid.—The acid made from sugar, recrystallized two or three times, is quite pure. Commercial acid should be mixed with insufficient water for complete solution, and the mixture occasionally shaken. Most of the impurities remain undissolved, and the saturated aqueous solution evaporated yields crystals which are fairly pure.

Quantivalence.—The elements represented by the formula C_2O_4 are those characteristic of oxalates. They form a bivalent grouping; hence neutral oxalates ($R'_2C_2O_4$) and acid oxalates ($R'HC_2O_4$) exist.

Salt of Sorrel is a crystalline compound of oxalic acid with acid potassium oxalate, the crystals containing two molecules of water of crystallization ($KHC_2O_4 \cdot H_2C_2O_4 \cdot 2H_2O$).

ANALYTICAL REACTIONS (TESTS).

First Analytical Reaction.—To solution of an oxalate (ammonium oxalate, *e. g.*) add solution of calcium chloride; a white precipitate falls (calcium oxalate, CaC_2O_4). Add to the precipitate excess of acetic acid; it is insoluble. Add hydrochloric acid; the precipitate is dissolved.

The formation of a white precipitate on adding a calcium or barium salt, insoluble in acetic but soluble in hydrochloric or nitric acid, is usually sufficient proof of the presence of an oxalate. The action of the liquid on litmus-paper, effervescence with sodium carbonate, and absence of metals would indicate that the oxalate is that of hydrogen, oxalic acid.

Note.—The *barium* oxalate is slightly soluble in acetic acid (Souhay and Lenssen), and enough may be dissolved by this acid from a mixed barium precipitate (produced on adding barium chloride or nitrate to a solution of mixed salts) to give the foregoing reaction on adding calcium chloride to the filtered acetic liquid—an effect sometimes useful in the analysis of mixed substances (Davies).

Antidote.—In cases of poisoning by oxalic acid, or salt of sorrel, chalk and water may be administered as a chemical antidote (with the view of producing the insoluble calcium oxalate), emetics, and the stomach-pump or stomach-siphon being used as soon as possible.

Second Analytical Reaction.—Heat a fragment of any dry common fixed metallic oxalate (a potassium oxalate, for example) in a test-tube; decomposition occurs, carbonic oxide, CO (a gas that will be noticed subsequently) is liberated, and a carbonate of the metal remains. Add water and then an acid to the residue; effervescence occurs.

This is a ready test for ordinary insoluble oxalates, and is trustworthy if, on heating the substance, no charring occurs, or not more than gives a gray color to the residue. Organic salts of metals decompose when heated, and leave a residue of carbonate, but, except in the case of oxalates, the residue is always accompanied by much charcoal. Insoluble oxalates and organic salts of such metals as lead and silver are, of course, liable to be reduced to oxide or even metal by heat. Such oxalates may be decomposed by boiling with solution of sodium carbonate, filtering and testing the filtrate for oxalates by the calcium chloride test.

Other Analytical Reactions.—Silver nitrate gives, with oxalates, a white precipitate (silver oxalate, $\text{Ag}_2\text{C}_2\text{O}_4$).—Dry oxalates are decomposed when heated with strong sulphuric acid, carbonic oxide and carbonic acid gases escaping. If much of the substance be operated on, the gas may be washed with an alkali, the carbonic acid be thus removed, and the carbonic oxide be ignited; it will be found to burn with a characteristic bluish flame.—Oxalates, when mixed with water, black manganese oxide (free from carbonates), and sulphuric acid, yield carbonic acid gas, which may be tested by lime-water in the usual manner.—Not only such insoluble oxalates as those of lead and silver above referred to, but any common insoluble oxalate, such as that of calcium or magnesium, may be decomposed by ebullition with solution of sodium carbonate; after

filtration the oxalic radical will be found in the clear liquid as soluble sodium oxalate.

QUESTIONS AND EXERCISES.

Explain the constitution of oxalates.—State how oxalates are obtained.—What is the quantivalence of the oxalic radical?—Give the formula of “salt of sorrel.”—Mention the chief test for oxalic acid and other soluble oxalates.—Name the antidote for oxalic acid, and describe its action.—By what reactions are insoluble oxalates recognized?

TARTARIC ACID AND OTHER TARTRATES.

Formula of Tartaric Acid, $\text{H}_2\text{C}_4\text{H}_4\text{O}_6$ or $\text{C}_4\text{H}_2(\text{OH})_2(\text{COOH})_2$. Molecular weight, 150.

Synonym.—Hydrogen Tartrate.

Sources.—Tartrates exist in the juice of many fruits, but it is from that of the grape that our supplies are usually obtained. Grape-juice contains much acid potassium tartrate ($\text{KHC}_4\text{H}_4\text{O}_6$), which is gradually deposited when the juice is fermented, as in making wine; for acid potassium tartrate, not very soluble in aqueous liquids, is still less so in spirituous, and hence crystallizes out as the sugar of the grape-juice is gradually converted into alcohol. It is found, with calcium tartrate, lining the vessels in which wine is kept; and it is from this crude substance, termed *argal* or *argol*, also from the albumenoid yeasty matter or “lees” deposited at the same time, as well as from what tartrate may be remaining in the marc left after the juice has been pressed from the grapes, that by rough recrystallization “tartar,” still containing 6 or 7 per cent. or more of anhydrous calcium tartrate ($\text{CaC}_4\text{H}_4\text{O}_6$), is obtained. From the tartar tartaric acid and other tartrates are prepared. In old dried grapes (raisins, *Uvæ*, B. P.) crystalline masses of tartar and of grape-sugar are constantly met with.

Cream of tartar, purified by crystallization (*Potassii Bitartras*, U. S. P.), occurs as “a gritty white powder or fragments of cakes crystallized on one surface;” of a pleasant acid taste, soluble in 180 parts of cold and 6 of boiling water, insoluble in spirit.*

“If 1.2 grm. of potassium bitartrate be repeatedly agitated, during half an hour, with a mixture of 5 cc. of acetic acid and 1 cc. of



* A boiling solution of tartar yields a floating crust of minute crystals on cooling, just as milk yields a floating layer of cream; hence the term *cream of tartar*. “It is called *tartar*,” says Paracelsus, “because it produces oil, water, tincture, and salt, which burn the patient as Tartarus does.” *Tartarus* is Latin (*Tátrapos*, Tartaros, Greek) for *hell*. The products of its destructive distillation are certainly somewhat irritating in taste and smell, and the “salt” (potassium carbonate) that is left is diuretic, and in larger quantities powerfully corrosive.

water, and the mixture be then diluted with 30 cc. of water and filtered, the clear filtrate should not be rendered turbid, within one minute, by the addition of 0.5 cc. of ammonium oxalate T. S. (limit of calcium salt).”—*U. S. P.*

Quantivalence.—The elements represented by the formula $C_4H_4O_6$ are those characteristic of tartrates. They form a bivalent grouping; hence neutral tartrates ($R'_2C_4H_4O_6$) and acid tartrates ($R'HC_4H_4O_6$) exist. Potassium tartrate, the *Potassii Tartras* of the United States Pharmacopœia ($K_2C_4H_4O_6$), and Rochelle salt, or potassium and sodium tartrate, the official *Potassii et Sodii Tartras*, *U. S. P.* (*Soda Tartrate*, *B. P.*), are illustrations of neutral tartrates, while cream of tartar is an example of acid tartrates. The only official tartrate not apparently included in these general formulæ is tartar emetic (*Antimonium Tartaratum*, *B. P.*, *Antimonii et Potassii Tartras*, *U. S. P.*), which is sometimes regarded as the double tartrate of potassium and a hypothetical radical, antimonyl (SbO); thus, $KSbOC_4H_4O_6$. Probably, however, it is but an antimony oxytartrate ($Sb_2O_2C_4H_4O_6$) with normal potassium tartrate ($K_2C_4H_4O_6$); for there are several oxycompounds of antimony analogous to the oxycompounds of bismuth that have been described (p. 254), normal salts partially decomposed by water into oxides, and many of these oxycompounds readily unite with normal salts of other basylous radicals. Tartar emetic would thus be antimony oxytartrate with potassium tartrate [$Sb_2O_2C_4H_4O_6, K_2C_4H_4O_6$ or $K''Sb'''(C_4H_4O_6)''O''$].

Tartaric Acid.

Tartaric Acid, the *Hydrogen Tartrate* (*Acidum Tartaricum*, *U. S. P.*), is obtained, according to the British Pharmacopœia, by boiling cream of tartar with water, adding chalk till effervescence ceases, and then calcium chloride so long as a precipitate falls; the two portions of calcium tartrate thus consecutively formed are thoroughly washed, treated with sulphuric acid, the mixture boiled for a short time, resulting calcium sulphate mostly separated by filtration, the filtrate concentrated by evaporation, any calcium sulphate that may have deposited removed as before, and concentration continued until the solution is strong enough to crystallize. Calcium tartrate from 9 ounces of cream of tartar requires 5 ounces by weight of sulphuric acid for complete decomposition.



Tartaric acid occurs in trade as colorless crystals or the same powdered. It is strongly acid and readily soluble in water or spirit.

1 part in 8 of water and 2 of spirit of wine forms "Solution of Tartaric Acid," B. P. Its aqueous solution is not stable.

Parcels of tartaric acid often contain crystals of an allotropic or physically isomeric modification. (*Vide* "Allotropy" and "Isomerism" in Index.) It is termed *paratartaric acid* (*παρά, para*, beside) or *racemic acid* (*racemus*, a bunch of grapes), and is a combination of ordinary tartaric acid, whose solution twists a ray of polarized light to the right hand (dextrotartaric or dextroracemic acid), and of lævotartaric or lævoracemic acid, whose solution twists a polarized ray to the left.* Racemic acid is inactive in this respect, the opposite properties of its constituents neutralizing each other. Racemic acid is less soluble in alcohol than tartaric acid.

REACTIONS.

Potassium Tartrate, or Tartrate of Potassium.


Synthetical Reactions.—To a small quantity of a strong solution of potassium carbonate add acid potassium tartrate so long as effervescence occurs; the resulting liquid is solution of neutral potassium tartrate, *Potassii Tartras*, U. S. P. ($K_2C_4H_4O_6$), crystals of which may be obtained on evaporation.

Note.—This is a common method of converting an acid salt of a bivalent acidulous radical into a neutral salt. The carbonate added need not be a carbonate of the same, but may be of a different metal; compounds like Rochelle salt ($KNaC_4H_4O_6$) are then obtained. Thus:

Potassium and Sodium Tartrate.

Synonyms.—Tartrate of Potassium and Sodium; Tartrate of Potash and Soda; Rochelle Salt.

To a strong hot solution of sodium carbonate add acid potassium tartrate until effervescence ceases; the resulting liquid is solution of potassium and sodium tartrate: on cooling it yields crystals. This is the official process (*Soda Tartarata*, B. P.; *Potassii et Sodii Tartras*, U. S. P.) ($KNaC_4H_4O_6 \cdot 4H_2O$).



Crystals of Rochelle salt are usually halves of colorless transparent right rhombic prisms, slightly efflorescent in dry air, soluble in 5 parts of boiling water. Potassium tartrate is slightly deliquescent, soluble in about 4 parts of boiling water.

* According to Van't Hoff and Le Bel, all compounds that cause such rotation contain at least one atom of carbon with which is united four different atoms or radicals. Such carbon atoms are conveniently termed *asymmetrical*.

EQUIVALENT WEIGHTS of *Tartaric Acid*, *Potassium Carbonate*, *Potassium Bicarbonate*, *Sodium Carbonate (crystallized)*, *Sodium Bicarbonate*, and *Ammonium and Magnesium Carbonates repeated for 20 parts of each* (and, incidentally, for other proportions).

Tart. acid . . .	$\text{H}_2\text{C}_4\text{H}_4\text{O}_6$	=150	20	$18\frac{1}{4}$	15	$10\frac{1}{2}$	$17\frac{3}{4}$	$28\frac{1}{2}$	$31\frac{1}{2}$
Potas. carb. . .	K_2CO_3 (of 84 per cent.) . .	=164	22	20	$16\frac{1}{2}$	$11\frac{1}{2}$	$19\frac{1}{2}$	$31\frac{1}{2}$	$34\frac{1}{2}$
Pot. bicarb. . .	$2(\text{KHCO}_3)$	=200	$26\frac{3}{4}$	$24\frac{1}{4}$	20	14	$23\frac{3}{4}$	$38\frac{1}{4}$	42
Sod. carb. . . .	$\text{Na}_2\text{CO}_3, 10\text{H}_2\text{O}$	=286	38	$34\frac{3}{4}$	$28\frac{1}{2}$	20	34	$54\frac{1}{2}$	60
Sod. bicarb. . .	$2(\text{NaHCO}_3)$	=168	$22\frac{1}{2}$	$20\frac{1}{4}$	$16\frac{3}{4}$	$11\frac{3}{4}$	20	32	$35\frac{1}{4}$
Ammon. carb. .	$(\text{N}_3\text{H}_{11}\text{C}_2\text{O}_6) \div 3 \times 2$	=105	14	$12\frac{3}{4}$	$10\frac{1}{2}$	$7\frac{1}{4}$	$12\frac{1}{2}$	20	$21\frac{3}{4}$
Magnes. carb. .	$(\text{MgCO}_3)_3\text{Mg}2\text{HO}, 4\text{H}_2\text{O} \div 4$	=95.5	$12\frac{3}{4}$	$11\frac{1}{2}$	$9\frac{3}{4}$	$6\frac{3}{4}$	$11\frac{1}{4}$	$18\frac{1}{4}$	20

Thus 20 parts (grains or other weights) of tartaric acid neutralize 22 of potassium carbonate, $26\frac{3}{4}$ of potassium bicarbonate, 38 of sodium carbonate, $22\frac{1}{2}$ of sodium bicarbonate, 14 of ammonium carbonate, or $12\frac{3}{4}$ of magnesium carbonate. Other quantities of tartaric acid ($18\frac{1}{4}$, 15, $10\frac{1}{2}$, $17\frac{3}{4}$, $28\frac{1}{2}$, $31\frac{1}{2}$) saturate the amount of salts mentioned in the other columns, and *vice versâ*. A similar table for citric acid will be found at p. 325, and for both acids in the Appendix. These tables afford good illustrations of both of the laws of chemical combination (pp. 47, 198). The reader should verify a few of the numbers by calculation from the atomic weights of the elements concerned in the reactions, remembering that the salts formed are considered to be neutral in constitution. In medical practice effervescing saline draughts are often designedly prescribed to contain an amount of acid or alkali considerably in excess of the proportions required for perfect neutrality.

Effervescent Tartarated Soda Powder (*Pulvis Sodæ Tartaratæ Effervescens*, B. P.), or *Seidlitz Powder*, consists of 3 parts of Rochelle salt (120 grains), with 1 part (40 grains) of acid-sodium carbonate (the mixture usually wrapped in blue paper), and 1 part (38 grains) of tartaric acid (wrapped in white paper). When administered, one powder is dissolved in a tumbler rather more than half full of water, the other added, and the mixture drunk during effervescence. It will be seen that the salts swallowed are potassium and sodium tartrate ($\text{KNaC}_4\text{H}_4\text{O}_6, 4\text{H}_2\text{O}$), sodium tartrate ($\text{Na}_2\text{C}_4\text{H}_4\text{O}_6, 2\text{H}_2\text{O}$), and acid sodium or potassium tartrate. The last-mentioned salt results because, for one reason, $2\frac{1}{2}$ grains of the tartaric acid is in excess of the quantity necessary for the formation of neutral sodium tartrate; and, for another reason, because while carbonic acid remains in great excess, a neutral tartrate containing potassium may be converted more or less into acid potassium tartrate and a bicarbonate. This amount of acid salt gives, according to the taste of some persons, agreeable acidity to the draught. The United States formula (*Pulvis Effervescens Compositus*, U. S. P.) includes rather less tartaric acid, so that only neutral salts are formed, and the occurrence of the gritty acid potassium tartrate

avoided. "Double" Seidlitz powders contain a double dose of Rochelle salt.

ANALYTICAL REACTIONS (TESTS).

First Analytical Reaction.—To solution of any normal tartrate, or tartaric acid made neutral by solution of soda, add solution of calcium chloride; a white precipitate (calcium tartrate, $\text{CaC}_4\text{H}_4\text{O}_6 \cdot 4\text{H}_2\text{O}$) falls. Collect the precipitate on a filter, wash, place a small quantity in a test-tube, and add solution of potash: on stirring the mixture the precipitate dissolves. Heat the solution: calcium tartrate is again precipitated.

In this reaction a fair amount of the calcium chloride solution should be added at once, and the test be performed without delay, or the calcium tartrate will assume a crystalline character and be with difficulty dissolved by the potash. The potash should be quite free from carbonate.

The solubility of calcium tartrate in cold potash solution enables the analyst to distinguish between tartrates and citrates, otherwise a difficult matter. Calcium citrate is not soluble, or only to a very slight extent, in the alkali. The absence of much ammoniacal salt must be ensured, calcium citrate as well as tartrate being soluble in solutions of salts of ammonium.

Second Analytical Reaction.—Acidulate a solution of a tartrate with acetic acid, add potassium acetate, and well stir the mixture; a crystalline precipitate (acid potassium tartrate) slowly separates.

This reaction is not applicable in testing for very small quantities of tartrates, the acid potassium tartrate being not altogether insoluble. The precipitate being insoluble in alcohol, however, the addition of spirit of wine renders the test far more delicate.

Third Analytical Reaction.—To a neutral solution of a tartrate add solution of silver nitrate; a white precipitate (silver tartrate, $\text{Ag}_2\text{C}_4\text{H}_4\text{O}_6$) falls. Boil the mixture; it blackens, owing to the reduction of the salt to metallic silver. Or, before boiling, add a drop or less of ammonia; a mirror will form on the tube, adhering well to the glass if the tube was thoroughly cleansed. Even an insoluble tartrate, placed in a dry tube with a few fragments of silver nitrate and a drop or less of ammonia added, gives a mirror-like character to each fragment of silver salt when the tube is gently rotated some inches above a flame.

Other Reactions.—Tartrates heated with strong sulphuric acid char immediately.—Tartaric acid and the soluble tartrates prevent the precipitation of ferric and other hydrates by alkalis, solutions of double tartrates being formed (which on

evaporation yield liquids that do not crystallize, but, spread on sheets of glass, dry up to thin transparent plates or scales). The iron potassio-tartrate (*Ferri et Potassii Tartras*, U. S. P.; *Ferrum Tartaratum* B. P.) is a preparation of this kind.—Tartrates decompose when heated, carbonates being formed and carbon set free, the gaseous products having a peculiar more or less characteristic smell resembling that of burnt sugar.

QUESTIONS AND EXERCISES.

State the origin of tartaric acid and other tartrates, and explain the deposition of argol, crude acid potassium tartrate, during the manufacture of wine.—What is the chemical formula, and what are the characters, of "cream of tartar"?—Mention the formula and quantivalence of the tartaric radical.—Write formulæ of various tartrates, including tartar emetic.—Give equations or diagrams illustrative of the production of tartaric acid from cream of tartar.—By what general process may normal or double tartrates be obtained from acid potassium tartrate?—Work out sums proving the correctness of some of the figures given on p. 321 as showing the saturating power of tartaric acid for various quantities of different carbonates, and give diagrams or equations of the reactions.—State the names and work sums showing quantities of the salts resulting from the admixture of 120 grains of potassium and sodium tartrate, 40 grains of acid sodium carbonate, and 38 grains of tartaric acid (Seidlitz powder).—Enumerate the tests for tartrates, and explain the effect of heat on tartrates of the metals.

CITRIC ACID AND OTHER CITRATES.


Formula of Citric Acid, $\text{H}_3\text{C}_6\text{H}_5\text{O}_7, \text{H}_2\text{O}$. Molecular weight, 210.

Synonym.—Hydrógen Citrate.

Source.—Citric acid (*Acidum Citricum*, U. S. P.) exists in the juice of the gooseberry, currant, cherry, strawberry, raspberry (*Rubus Idæus*, U. S. P.), and many other fruits, as well as in other parts of plants. The pulp of the fruit of *Tamarindus indica* (*Tamarindus*, U. S. P.) contains from 1 to 12 per cent. (in addition to 1.5 of tartaric acid, .5 of malic acid, and 3 per cent. of acid potassium tartrate). But it is from the lemon or lime that the acid of commerce is usually obtained. For this purpose concentrated lemon-juice is exported from Sicily, concentrated bergamot-juice from the Calabrian coast of South Italy, and concentrated lime-juice from the West Indies. The lime-fruit from *Citrus bergamia* is official in the Pharmacopœia of India.

Process.—The British Pharmacopœia directs that the hot lemon-juice (4 pints) be saturated by powdered chalk—that is, whiting—($4\frac{1}{2}$ ounces), the resulting calcium citrate collected on a filter, washed with hot water till the liquor passes from it colorless (by which not only the coloring matter, but the mucilage, sugar, and other constituents of the juice, is got rid of), then mixed with cold water (1 pint), decomposed by sulphuric acid ($2\frac{1}{2}$ fluidounces in $1\frac{1}{2}$ pints

of water), the mixture boiled for half an hour, filtered, the solution evaporated to a density of 1.21, set aside for twenty-four hours, then poured off from any deposit of crystalline calcium sulphate, further concentrated, and set aside to crystallize. If the quantity of calcium citrate to be decomposed is indefinite, the sulphuric acid may be added until a little of the supernatant fluid gives, after a minute or two, a precipitate with solution of calcium chloride. The concentrated citric solution generally crystallizes very slowly. Shaken violently, however, in a bottle with a granule or two of solid acid, it quickly yields its citric acid in a pulverulent form, and this, drained and redissolved in a very small quantity of hot water, yields crystals fairly quickly (Warington).



Quantivalence.—The elements represented by the formula $\text{C}_6\text{H}_5\text{O}_7$ are those characteristic of citrates. They form a trivalent grouping; hence three classes of salts may exist—one, two, or three atoms of the basylous hydrogen in one molecule of the acid, $\text{H}_3\text{C}_6\text{H}_5\text{O}_7$, being displaced by equivalent proportions of other basylous radicals. Constitutional formula, $\text{C}_3\text{H}_4(\text{OH})(\text{COOH})_3$.

Citric acid itself is the only citric compound of much direct importance to the pharmacist. It usually occurs in colorless crystals soluble in half their weight of boiling and three-fourths of cold water, less soluble in spirit, and insoluble in ether. A solution of 36 to 46 grains in 1 ounce of water forms a sort of artificial lemon-juice. Citrates heated with strong sulphuric acid to about 215°F . evolve carbonic oxide gas, and at higher temperatures acetone and carbonic acid gas.

The artificial production of citric acid has been accomplished by Grimaux and Adam, who, starting with glycerin, produce certain chloro- and cyano-derivatives, and ultimately citric acid itself.

Action of Heat on Citric Acid.—Citric acid, slowly heated, first loses its water of crystallization; afterward (347°F .) the elements of another molecule of water are evolved, and a residue obtained from which ether extracts *aconitic acid*, $\text{H}_3\text{C}_6\text{H}_3\text{O}_6$, identical with the aconitic acid (and the acid first termed *equiseti*c) in various species of *Aconitum* and *Equisetum*.

The official lemon-juice (*Limonis Succus*, U. S. P.) is to be freshly expressed from the ripe fruit, have a specific gravity not less than 1.030, and contain about 7 per cent. of citric acid ($\text{H}_3\text{C}_6\text{H}_5\text{O}_7, \text{H}_2\text{O}$). The acidity may be ascertained by adding solution of potash or soda (the strength of which has been determined previously with pure crystals of citric acid) till red litmus-paper is fairly turned blue. Before applying this test to commercial specimens of lemon-juice the absence of notable quantities of sulphuric, hydrochloric, acetic, tartaric, or other acid must be ensured by application of appropriate reagents. (See also "Lemon-juice," in Index.)

Lime-juice as imported into England contains an average of 7.84 per cent. of citric acid, rarely rising to 10 per cent. and very seldom falling to 7 per cent. Containing but little sugar and mucilage, it requires no addition of spirit to preserve it. *Lemon-juice* requires about 40 per cent. of proof spirit to prevent fermentation (Conroy).

EQUIVALENT WEIGHTS of *Citric Acid*, *Potassium Carbonate*, *Potassium Bicarbonate*, *Sodium Carbonate*, (crystallized), *Sodium Bicarbonate*, and *Ammonium and Magnesium Carbonates* repeated for 20 parts of each (and, incidentally, for other proportions).

Citric acid,	$\text{H}_3\text{C}_6\text{H}_5\text{O}_7 \cdot \text{H}_2\text{O}$	=210	20	17	14	9 $\frac{3}{4}$	16 $\frac{3}{4}$	26 $\frac{3}{4}$	29 $\frac{1}{4}$
Pot. carb., .	$(\text{K}_2\text{CO}_3; \text{ of } 84 \text{ per cent.}) \div 2 \times 3$	=246 $\frac{1}{2}$	23 $\frac{1}{2}$	20	16 $\frac{1}{2}$	11 $\frac{1}{2}$	19 $\frac{3}{4}$	31 $\frac{1}{2}$	34 $\frac{1}{4}$
Pot. bicarb.,	$3(\text{KHCO}_3)$	=300	28 $\frac{1}{4}$	24 $\frac{1}{4}$	20	14	24	38 $\frac{1}{4}$	41 $\frac{1}{4}$
Sod. carb., .	$(\text{Na}_2\text{CO}_3; 10\text{H}_2\text{O}) \div 2 \times 3$	=429	41	34 $\frac{3}{4}$	28 $\frac{1}{2}$	20	34	54 $\frac{1}{2}$	60
Sod. bicarb.,	$3(\text{NaHCO}_3)$	=252	24	20 $\frac{1}{2}$	16 $\frac{1}{2}$	11 $\frac{3}{4}$	20	32	35
Amm. carb.,	$(\text{N}_3\text{H}_{11}\text{C}_2\text{O}_5)$	=157	15	12 $\frac{3}{4}$	10 $\frac{3}{4}$	7 $\frac{1}{4}$	12 $\frac{1}{2}$	20	21 $\frac{3}{4}$
Mag. carb., .	$(\text{MgCO}_3)_3\text{Mg}2\text{HO}, 4\text{H}_2\text{O} \div 8 \times 3$	=143	13 $\frac{1}{2}$	11 $\frac{1}{4}$	9 $\frac{1}{2}$	6 $\frac{3}{4}$	11 $\frac{1}{2}$	18 $\frac{1}{4}$	20

Thus 20 parts (grains or other weights) of citric acid neutralize 23 $\frac{1}{2}$ of potassium carbonate, 28 $\frac{1}{2}$ of potassium bicarbonate, 41 of sodium carbonate, 24 of sodium bicarbonate, 15 of ammonium carbonate, or 13 $\frac{1}{2}$ of magnesium carbonate. Other quantities of citric acid (17, 14, 9 $\frac{3}{4}$, 16 $\frac{3}{4}$, 26 $\frac{3}{4}$, 29 $\frac{1}{4}$) saturate the amounts of salts mentioned in the other columns, and *vice versa*.

This table, the similar one for tartaric acid (p. 321), and that for both acids (*vide* Appendix) afford good illustrations of both of the laws of chemical combination (pp. 47, 198). The reader should verify a few of the numbers by calculation from the atomic weights of the elements concerned in the reactions, remembering that the salts formed are considered to be neutral in constitution. In medical practice effervescing saline draughts are often designedly prescribed to contain an amount of acid or alkali considerably in excess of the proportions required for perfect neutrality.

ANALYTICAL REACTIONS (TESTS).

First Analytical Reaction.—To a dilute solution of any neutral citrate, or citric acid carefully neutralized by alkali, add solution of calcium chloride and boil; a white precipitate (calcium citrate, $\text{Ca}_3\text{C}_6\text{H}_5\text{O}_7$) falls. Treat the precipitate as for calcium tartrate (p. 322); it is not perceptibly dissolved by the potash.

A mixture of citrates and tartrates can be separated by this reaction. They are precipitated as calcium salts, and the rapidly washed precipitate mixed with solution of potash, diluted and filtered; the filtrate contains the tartrate, which is shown to be

present by precipitation on boiling. The precipitate still on the filter is washed, dissolved in solution of ammonium chloride, and the solution boiled; the calcium citrate is reprecipitated. The presence of much sugar interferes with this reaction. A dilute solution of a citrate is not precipitated by calcium chloride until the liquid is heated: precipitation from a strong solution, also, is not thoroughly complete without ebullition of the mixture. This reaction is not thoroughly satisfactory, calcium citrate being *slightly* soluble in alkalies, in the solutions of salts produced in the reaction, and to a very slight extent even in cold water. It is readily soluble in acetic acid.

Second Analytical Reaction.—To a *neutral* solution of a citrate add solution of silver nitrate; a white precipitate (silver citrate, $\text{Ag}_3\text{C}_6\text{H}_5\text{O}_7$) falls. Boil the mixture; the precipitate does not blacken as silver tartrate does, or only after long boiling.

Other Analytical Reactions.—Citric acid forms no precipitate corresponding with the acid potassium tartate.——Lime-water, in excess, gives no precipitate with citric acid or citrates unless the solution is boiled, calcium citrate being slightly soluble in cold, but not in hot, water; it usually precipitates tartrates in the cold.——Citrates when heated with strong sulphuric acid do not char immediately.——Citric acid and citrates prevent the usual precipitation of iron by alkalies, soluble double compounds being formed. The *Ferri et Ammonii Citras*, U. S. P., is a preparation of this kind.——Metallic citrates decompose when heated, carbonates being formed and carbon set free; the odor of the gaseous products is not so characteristic as that of tartrates.——According to Cailletet, a cold saturated solution of red potassium chromate turns a solution of tartaric acid dark brown, carbonic acid gas being evolved, while a solution of citric acid only slowly becomes of a light brown.

Pusch's test for the detection of tartaric acid in citric acid depends on the well-known difference in the action of sulphuric acid on tartaric acid and on citric acid. It consists in adding to 1 gramme of powdered citric acid in a dry test-tube 10 grammes of strong pure (colorless) sulphuric acid, and keeping the part of the tube containing the mixture immersed in boiling water for an hour. The citric acid dissolves with evolution of gas and frothing to form a lemon-colored liquid, and if the sample be pure this color undergoes no change within half an hour; but if as much as $\frac{1}{2}$ per cent. of tartaric acid be present, the lemon color becomes brownish within that time, and in an hour the mixture is red-brown.

QUESTIONS AND EXERCISES.

What is the source of citric acid?—Describe the preparation of citric acid, giving diagrams.—Illustrate by formulæ the various classes of tar-

trates and citrates.—State the average proportion of citric acid in lemon-juice.—Work out sums proving the correctness of some of the figures given on p. 325, as showing the saturating power of citric acid for various carbonates.—What are the tests for citrates?—How are tartrates separated from citrates?

PHOSPHORIC ACID AND OTHER PHOSPHATES.

Formula of the Acid, H_3PO_4 or $\text{PO}(\text{OH})_3$. Molecular weight, 98.

Synonym.—Hydrogen Phosphate.

Source.—The source of the ordinary normal phosphates and of phosphorus itself (*Phosphorus*, U. S. P.) is the normal calcium phosphate ($\text{Ca}_3\text{2PO}_4$). It is the chief constituent of the bones and teeth of animals, being derived from the plants on which they feed, plants again obtaining it from the soil. Compounds of phosphorus are also met with in the brain, nerves, muscles, blood, saliva, and, according to Kirkes, even in tissues so simple that one must assume that the compounds are necessary constituents of the substance of the primary cell. They escape from the system both in the urine and in the fæces.


Process.—Phosphorus ($\text{P} = 31$) is obtained from bones by the following processes: The bones are burnt to remove all traces of animal matter. The resulting *bone-earth* is treated with hot and fairly strong sulphuric acid, by which phosphoric acid and calcium sulphate are produced:



The acid fluid, strained from the sulphate and concentrated, is mixed with charcoal, coke, or sawdust and dried in an iron pot. At this stage water escapes and metaphosphoric acid remains:



The mixture is then to be transferred to a fireclay retort and strongly heated; phosphorus vapor is evolved and is condensed under water:



The phosphorus is purified by melting under water containing sulphuric acid and red potassium chromate, and is filtered through canvas and cast into sticks.

Properties.—Phosphorus is “a semi-transparent, colorless, wax-like solid (in sticks or cakes), which emits white vapors when exposed to the air. Specific gravity, 1.77. It is soft and flexible at common temperatures, melts at 110°F ., ignites in the air at a temperature a little above its melting-point, burning with a luminous flame and producing dense white fumes. It is very poisonous. Insoluble in water, but soluble in ether and in boiling oil of turpen-

tine, also in carbon bisulphide. It is soluble in oil which has been previously heated for a short time to about 300°F . to expel moisture—1 part in 90 of dried almond oil constituting phosphorated oil (*Oleum Phosphoratum*, U. S. P.). A mixture, or rather a solution, of phosphorus in chloroform, mixed with althea, acacia, glycerin, coated with balsam of Tolu (previously dissolved in ether), forms the official phosphorus pills (*Pilule Phosphori*, U. S. P.)."

Granulated or pulverulent phosphorus is obtained by placing a portion under equal parts of spirit and water in a bottle, standing the bottle in warm water till the phosphorus melts, then inserting the stopper (glass, not cork), and shaking the whole till cold.

Red or Amorphous Phosphorus.—Ordinary phosphorus kept at a temperature of about 450°F . in an atmosphere from which air is excluded becomes red, opaque, insoluble in liquids in which ordinary phosphorus is soluble, oxidizes extremely slowly, and only ignites when heated to near 500°F . It is used in the manufacture of several varieties of lucifer-matches, not emitting the poisonous jaw-destroying fumes given off by ordinary phosphorus.

Quantivalence.—The atom of phosphorus is quivalent, as seen in the pentachloride (PCl_5) and oxychloride (POCl_3); but it often exhibits trivalent activity, as seen in the trichloride (PCl_3) and trihydride (PH_3).

Zinc Phosphide, Zn_3P_2 (*Zinci Phosphidum*, U. S. P.), occurs as a grayish-black powder or in crystalline fragments having a metallic lustre. It may be obtained by throwing phosphorus upon melted zinc.


Molecular Weight.—Phosphorus is an exception to the rule that the atomic weights (in grains, grammes, etc.) of elements occupy similar volumes of vapor at similar temperatures, the equivalent weight of phosphorus (31) only giving half such a volume. Hence, while the molecular weights—that is, double the atomic weights—of oxygen ($\text{O}_2 = 32$), hydrogen ($\text{H}_2 = 2$), nitrogen ($\text{N}_2 = 28$), etc., give a similar bulk of vapor at any given temperature, the double atomic weight of phosphorus ($\text{P}_2 = 62$) only gives half this bulk; that is, four times the atomic weight of phosphorus must be taken to obtain the whole bulk. It would appear, therefore, that the molecule of phosphorus contains four atoms, $\text{P}_4 = 124$. As with sulphur, however, phosphorus in the state ordinarily known to us may be abnormal, and conditions yet be found in which the molecular weight is double the atomic weight.

Phosphoric Acid.

Synonyms.—Orthophosphoric Acid; Hydrogen Phosphate.

The chief use of phosphorus in pharmacy is for the production of diluted phosphoric acid. Phosphorus is boiled with nitric acid and water until dissolved. The solution, evaporated to a low bulk to remove nitrous compounds, and diluted so as to contain 85 per cent. of acid (H_3PO_4), constitutes the *Acidum Phosphoricum*, U. S. P., a colorless liquid of specific gravity

1.710. The latter, diluted so as to contain 10 per cent. of the acid, constitutes the *Acidum Phosphoricum Dilutum*, U. S. P., a colorless, sour liquid of specific gravity 1.057. If the necessary appliances are at hand, specimens may be prepared by the official process as follows: $\frac{3}{4}$ of an ounce of the *concentrated*, or a $\frac{1}{4}$ pint of the *diluted*, is made by boiling together, in a flask attached to a vertical condenser, 103 grains of phosphorus, $1\frac{1}{2}$ fluidounces of the official nitric acid, and 2 ounces of water. By some such arrangement the condensed products are returned to the flask. The operation is continued until the phosphorus has disappeared.




The liquid remaining in the flask is then transferred to a dish (preferably of platinum), evaporated down to about $\frac{1}{2}$ an ounce, and, lastly, diluted with distilled water.

The use of the water in the former part of the process is to moderate the reaction. Strong hot nitric acid oxidizes phosphorus with almost explosive rapidity; hence the acid must be diluted in the first instance, and the dilution be maintained to prevent its becoming too strong by loss of water. Time is saved by using a strong acid, but in that case constant supervision is necessary, in order that water may be added or the temperature otherwise reduced should the action become too violent. Deficiency of nitric acid must also be avoided, or some phosphorous acid (H_2PHO_3) will be formed.

Markoe, also to economize time, modifies the process by adding for every ounce of phosphorus 4 or 5 grains of iodine, and, drop by drop, 25 or 30 drops of bromine. The iodine and bromine unite with the phosphorus with a readiness, or even violence, that would be explosive if not controlled by the presence of the cold fluids—further cooled, if necessary, by immersing the vessel in cold water. Phosphorus iodide (PI_3) and phosphorus bromide (PBr_3) are at once formed. These in the presence of water immediately yield hydriodic and hydrobromic acids (HI , HBr) and phosphoric acid. The nitric acid attacks the hydriodic and hydrobromic acids, forming the lower oxides of nitrogen, which escape as gas, water, and free iodine and bromine. The latter unite with more phosphorus, and the reactions are repeated. This carrying power of a little iodine or bromine or both would perhaps be indefinitely prolonged if no vapor of these elements or their acids escaped with the gases. The phosphorus having disappeared, excess of nitric acid is got rid of roughly by dropping in clean rags or paper (nitric oxide, carbonic acid gas, and water being formed), and the last portions by adding oxalic acid (which even still more readily yields similar products). Evaporation to a syrupy consistence finally removes all traces of iodine, bromine, oxalic acid, and moisture. The product is then diluted to any required extent.

Experimental Process.—A flask, in the neck of which a funnel is inserted and a second funnel inverted, so that its mouth rests within the mouth of the first, is an efficient and convenient arrangement of apparatus for the official process, especially if the operation be conducted slowly. (See Fig. 40.)

Fig. 40.



Solution of phosphoric acid evaporated to a sp. gr. of 1.850 yields a mass of prismatic crystals of H_3PO_4 , especially if a crystal or two be dropped into the fluid (Cooper). Further evaporation, it leaves a residue which melts at a low red heat, yielding *pyrophosphoric acid*, and, finally, *metaphosphoric acid* (*Glacial Phosphoric Acid*).

A commercial variety of phosphoric acid, containing no large amount of impurity, is prepared by well digesting a mixture of bone-ash, sulphuric acid, and water, filtering, concentrating, precipitating calcium by strong sulphuric acid, and heating until sulphuric vapors cease to escape; also by burning phosphorus to phosphoric anhydride, dissolving the latter in water, and boiling with a little nitric acid to oxidize any lower acids of phosphorus and to cause any meta- or pyro-phosphoric acid to take up the elements of water.

Prepared from bones, phosphoric acid is apt to develop fungoid deposit (Jensen). Prepared from phosphorus, it occasionally contains arsenum in the form of arsenic acid. The latter is detected and removed, together with any traces of platinum or lead, on passing sulphuretted hydrogen through the warmed acid.

Quantivalence.—The elements represented by the formula PO_4 are those characteristic of phosphates. The grouping is trivalent; hence there may exist trimetallic phosphates ($\text{M}'_3\text{PO}_4$), dimetallic acid phosphates ($\text{M}'_2\text{HPO}_4$), or monometallic acid phosphates ($\text{M}'\text{H}_2\text{PO}_4$), and, lastly, trihydric phosphate (H_3PO_4), or common phosphoric acid. These are the ordinary phosphates, or *orthophosphates*, met with in nature or used in pharmacy; the rarer pyrophosphates and metaphosphates, as well as the phosphites and hypophosphites, will be mentioned subsequently. Crude dry calcium phosphate ground with sulphuric acid yields the very largely used artificial manure termed "superphosphate." It contains acid calcium phosphate ($\text{CaH}_4\text{P}_2\text{O}_7, 2\text{H}_2\text{O}$) and calcium phosphate ($\text{CaSO}_4, 2\text{H}_2\text{O}$).

ANALYTICAL REACTIONS (TESTS).

First Analytical Reaction.—To an aqueous solution of a phosphate (*e. g.* Na_2HPO_4) add solution of magnesium sulphate with which ammonium chloride and ammonia have been mixed; a white crystalline precipitate falls (ammonio-magnesian phosphate, MgNH_4PO_4).

Ammonium chloride is added to prevent the precipitation of magnesium hydrate. Arsenates, which have close analogy to phosphates, give with the magnesium reagent a precipitate of similar character.

Second Analytical Reaction.—To an aqueous solution of a phosphate add solution of silver nitrate; light yellow silver phosphate (Ag_3PO_4) is precipitated—completely, if the mixture be neither acid nor alkaline. To a portion of the precipitate add ammonia; it dissolves. To another portion add nitric acid; it dissolves. By the former part of this reaction phosphates may be distinguished from their close allies the arsenates, silver arsenate being of a chocolate color.

Third Analytical Reaction.—To a solution (in a few drops of acid) of a phosphate insoluble in water (*e. g.* Ca_3PO_4) add the acetate of an alkali-metal (easily made by adding to soda or ammonia in a test-tube excess of acetic acid), and then a drop or two of solution of ferric chloride; a yellowish-white precipitate falls (ferric phosphate, $\text{Fe}_2\text{2PO}_4$), insoluble in acetic acid. Too much of the ferric chloride must not be added or ferric acetate will be produced, in which the ferric phosphate is to some extent soluble.

To remove the whole of the phosphoric radical from the solution add ferric chloride so long as a precipitate is produced, and boil; ferric phosphate and oxyacetate are precipitated.

To obtain confirmatory evidence of the presence of phosphate in this precipitate, and to separate the phosphoric radical as a more characteristic phosphate, collect the precipitate on a filter, wash, drop some solution of ammonia on it, then ammonium sulphhydrate, and finally wash with water; black ferrous sulphide remains on the filter, while ammonium phosphate occurs in the filtrate. To the filtrate add a mixture of solutions of magnesium sulphate and ammonium chloride, and well stir; a granular precipitate (ammonio-magnesian phosphate) appears.

Fourth Analytical Reaction.—In diluted nitric acid dissolve a little calcium phosphate (or any other phosphate), and then add solution of ammonium molybdate, and gently heat; a yellow precipitate falls.

This precipitate contains what is somewhat indefinitely termed phospho-molybdic acid—a compound of molybdic acid with phosphoric acid (about 4 per cent. of H_3PO_4) with ammonia (nearly 7 per cent.).

Ammonium molybdate is obtained by roasting native molybdenum sulphide (MoS_2 , U. S. P.; $(\text{NH}_4)_2\text{MoO}_4$)—which much resembles lead; hence the name of the metal, from $\mu\delta\lambda\upsilon\beta\delta\omicron\varsigma$, *molybdos*, lead—to molybdic oxide or anhydride (MoO_3), dissolving the latter in water, adding ammonia, evaporating, and crystallizing.

Molybdates having the following formulæ ($M = 1$ univalent atom of any metal) have been obtained: M_2MoO_4 ; $MHMoO_4$; $MHMoO_4 \cdot H_2MoO_4$. According to Carington, commercial ammonium molybdate is commonly the intermediate of the three salts.


Note.—The foregoing two reactions are useful in the analysis of bone-earth, of other earthy iron phosphate, and all phosphates insoluble in water. Only arsenates give similar appearances, but the acid solution of these may be decomposed by agitation with sulphurous acid, ebullition, and subsequent treatment with sulphuretted hydrogen—yellow arsenous sulphide, As_2S_3 , being then precipitated.

Other Analytical Reactions.—Solutions of barium and calcium salts give, with aqueous solutions of phosphates, white precipitates (of the respective phosphates $BaHPO_4$ or Ba_3-2PO_4 , and $CaHPO_4$ or Ca_3-2PO_4), all of which are soluble in acetic and the stronger acids; lead acetate a white precipitate.

QUESTIONS AND EXERCISES.

State the source of phosphorus.—Give equations or diagrams explanatory of the isolation of phosphorus from its natural compounds.—What is the composition of farmers' "superphosphate," and how is it prepared? Enumerate the properties of phosphorus.—Mention some solvents of phosphorus.—How are the official varieties of phosphoric acid made? Describe the precautions to be observed in making this acid.—What are the strengths of the official acids?—Write formulæ illustrative of all classes of orthophosphates.—Mention the chief tests for soluble and insoluble phosphates.—By what reactions may phosphates be distinguished from arsenates.

VANADIUM, V, 51.3, is a very rare element, and is here mentioned only because of its exceedingly interesting relationship to nitrogen, phosphorus, arsenum, and antimony, it with them forming five closely-connected members of one family. Discovered, but not isolated, by Sefström, and its compounds investigated by Berzelius, it has only of late years been obtained in a free state and fully studied by Roscoe.



Orthophosphates . . . R'_3PO_4	Orthovanadates . . . R'_3VO_4
Pyrophosphates . . . $R'_4P_2O_7$	Pyrovanadates . . . $R'_4V_2O_7$
Metaphosphates . . . $R'PO_3$	Metavanadates . . . $R'VO_3$

Isomorphous Minerals.

Apatite	$3(Ca_3-2PO_4), CaF_2$
Pyromorphite	$3(Pb_3-2PO_4), PbCl_2$
Mimetesite	$3(Pb_3-2AsO_4), PbCl_2$
Vanadinite	$3(Pb_3-2VO_4), PbCl_2$

BORIC ACID AND OTHER BORATES.

Formula of Boric Acid, H_3BO_3 . Molecular weight, 62.

The composition of artificial boric acid, sometimes termed *orthoboric acid*, *hydrogen borate*, and *boracic acid*, is expressed by the formula H_3BO_3 (*Acidum Boricum*, U. S. P.); but at a temperature of 212°F . this body loses the elements of water and yields metaboric acid, HBO_2 , which at higher temperature becomes boric anhydride (B_2O_3). Metaboric acid exists in the jets of steam (*fumerolles* or *suffioni*) that issue from the earth in some districts of Tuscany, and collects in the water of the *lagoni* (lagoons or little lakes) formed at the orifices of the steam-channels. This acid liquid, evaporated by aid of the waste natural steam and neutralized by sodium carbonate, gives common borax, possibly a sodium and hydrogen metaborate, with water of crystallization ($2\text{NaBO}_2, 2\text{HBO}_2, 9\text{H}_2\text{O}$), or possibly a sodium metaborate with boric anhydride ($2\text{NaBO}_2, \text{B}_2\text{O}_3, 10\text{H}_2\text{O}$); or possibly a sodium pyroborate ($\text{Na}_2\text{B}_4\text{O}_7, 10\text{H}_2\text{O}$), analogous to potassium pyrochromate ($\text{K}_2\text{Cr}_2\text{O}_7$) or pyrosulphuric acid ($\text{H}_2\text{S}_2\text{O}_7$). It is sometimes termed sodium baborate. Its official name is *Sodii Boras*, *Borate of Sodium* or *Sodium Borate*. It occurs "in transparent colorless crystals, occasionally slightly effloresced, with a weak alkaline reaction; insoluble in rectified spirit, soluble in water." Native borax, or *tincal*, and other borates are also found in Thibet, Nevada, Peru, Chili, and recently in California in the Clear Lake district. The introduction of the natural borax from California has reduced the price to about one-half its former amount. The Californian borax is represented as forming large portions of the crystalline bed of a dried-up lake. Borax is also largely made by boiling native calcium borax with sodium carbonate.

Fused borax readily dissolves metallic oxides, as will have been noticed already in testing for cobalt and manganese. Hence, besides its use in medicine, it is employed as a flux in refining and other metallurgic and ceramic operations.

Quantivalence.—The boric radical is trivalent (BO_3'''); the metaboric, univalent (BO_2').

The element boron, like carbon, occurs in the amorphous, graphitoid, and crystalline conditions. It is a trivalent element (B'''), yielding definite salts, such as the chloride (BCl_3) and fluoride (BF_3). Its atomic weight is 11.

REACTIONS.

First Synthetical Reaction.—To a hot solution of a crystal of borax add a few drops of sulphuric acid and set aside; on cooling crystalline scales of boric acid (H_3BO_3) (*Acidum Boricum*, U. S. P.) are obtained. They may be purified by collecting on a filter, slightly washing, drying, digesting in hot alcohol, filtering, and setting aside; pure boric acid is deposited. The acid may also be recrystallized from water. 50 grains dissolved in 1 ounce of rectified spirit constitute "Solution of Boric Acid," B. P. If 310 parts of boric acid be added to

glycerin at 150° C., and after being kept at that temperature for some time diluted to 1000 parts, *Glyceritum Boroglycerini*, U.S.P., is obtained. *Honey of Borax*, formed of 2 parts of borax to 16 of honey, is a very old antiseptic for the mouths of infants troubled by the growth called "thrush," the official British variety (*Mel Boracis*, B.P.) containing, in addition to the foregoing, 1 part of glycerin.

Boric acid occurs in "colorless, pearly, lamellar crystals; unctuous to the touch; taste feebly sour and bitter and leaving a sweetish after-flavor in the mouth. Soluble in 25 parts of water, 5 of glycerin, 16 of rectified spirit at 60° F. (15.5° C.), and in 3 of boiling water. It changes the color of litmus to wine-red; turmeric-paper, moistened with an aqueous solution slightly acidified with hydrochloric acid, becomes brownish-red on gently drying, and this color changes to a greenish if solution of potash be added. The alcoholic solution burns with a flame tinged with green. The crystals liquefy when warmed, and on careful ignition lose 43½ per cent. of their weight, the product solidifying on cooling to a brittle glass-like mass (B.P.).

Boric acid is a very weak compound. Indeed, the alkalinity of borax is as great as if it contained no acidulous material. The acid only slowly decomposes carbonates.

Second Synthetical Reaction.—Mix together 1 part of boric acid, 4 parts of acid potassium tartrate, and 10 or 20 of water; evaporate to a syrupy consistence, spread on plates, and set aside for dry scales to form. The resulting substance is, in water, far more readily soluble than either of its constituents, and is known as *potassium boro-tartrate* or *soluble cream of tartar*. The Prussian *tartarus boraxatus* differs from the foregoing French variety in containing 1 part of *borax* to 3 of acid potassium tartrate.

ANALYTICAL REACTIONS (TESTS).

First Analytical Reaction.—Dip a piece of turmeric-paper (paper soaked in tincture of turmeric-tubers and dried) into a solution of boric acid; it is colored brown red, as by alkalies.

The usual way of applying this test is as follows: Add to a solution of any borate a few drops of hydrochloric acid, immerse half of a slip of turmeric-paper in the liquid, then remove the hydrochloric acid by drying the paper over a flame. Concentrated hydrochloric acid and ferric chloride produce a somewhat similar effect.

Second Analytical Reaction.—To a fragment of a borate or a metaborate (borax, for example) in a small dish or watch-glass add a drop of sulphuric acid, and then a little alcohol; warm the mixture and set light to the spirit; the resulting flame will be tinged of a greenish color at its edges by the volatilized metaboric acid or boric anhydride.

The liquid should be well stirred while burning. Salts of copper and some metallic chlorides produce a somewhat similar color. The flame-test may also be applied to a little of a mixture of the borate with strong sulphuric acid on a platinum wire. Glycerin may be used in place of sulphuric acid (Hes), the reaction with borax being, according to Dunstan, the formation of glyceryl borate, $C_3H_5BO_3$, water, and sodium metaborate, the glyceryl borate and water reacting immediately to form boric acid and glycerin.

Other Analytical Reactions.—In solutions of borax barium salts give a white precipitate (barium metaborate, Ba_2BO_2), soluble in acids and alkaline salts. Silver nitrate also affords a white precipitate (silver metaborate, $AgBO_2$), soluble in nitric acid and in ammonia. Calcium chloride, if the solution is not too dilute, gives a white precipitate (calcium borate, $CaBO_2$).

QUESTIONS AND EXERCISES.

Illustrate the relation of vanadium to nitrogen by formulæ of compounds of each element.—Describe the preparation of borax.—Give the formulæ of boric acid, metaboric acid, and borax.—Mention the tests for borates or metaborates.

The foregoing acids and other salts contain the only acidulous radicals that are commonly met with in analysis or in ordinary medical or pharmaceutical operations. There are, however, many others which occasionally present themselves. The chief of these will now be shortly noticed; they are arranged in alphabetical order to facilitate reference.


SALTS OF RARER ACIDULOUS RADICALS.

Anemonic Acid.—*Pulsatilla*, U. S. P., is the official name for the herbs of *Anemone Pulsatilla* and *A. pratensis*. These, together with several species of *Ranunculus*, on distillation with water yield a heavy, yellow, acrid oil, which in contact with water yields crystalline poisonous *anemonin* ($C_{15}H_{12}O_6$) and amorphous *anemonic acid* ($C_{15}H_{14}O_7$).

BENZOIC ACID ($HC_7H_5O_2$) AND OTHER BENZOATES.—Slowly heat a fragment of benzoin (gum benjamin) (*Benzoinum*, U. S. P.) in a test-tube; benzoic acid (*Acidum Benzoicum*, U. S. P.), the hydrogen benzoate, rises in vapor and condenses in small, white, feathery plates and needles on the cool sides of the tube. If the benzoin is first mixed with twice its weight of sand or roughly powdered pumice-stone, and the heat very cautiously applied, the product will be less likely to be burnt


and a larger quantity will be yielded. By repeated sublimation 10 to 15 per cent. may be obtained.

A more economical process is to boil the benzoin with one-fourth its weight of lime, filter, concentrate, decompose the solution of calcium benzoate by hydrochloric acid, collect the precipitated benzoic acid, press between paper, dry, and sublime in a tube or other vessel.



There is always associated with the product a minute quantity of a mixture of volatile oils of agreeable odor, suggesting that of hay, and yielding, according to Jacobsen, methyl benzoate, guaiacol (methoxycatechol), catechol, acetylguaiacol, benzyl benzoate, benzophenone, and benzoylguaiacol.


Benzoic acid is also prepared on a large scale artificially from naphthalin, one of the crystalline by-products in the distillation of coal for gas. The naphthalin is oxidized by nitric acid to naphthalic or phthalic acid:



The phthalic acid is neutralized by lime and the calcium phthalate heated with calcium hydrate in a covered vessel at a temperature of about 640° F. for several hours. Calcium benzoate and carbonate are formed, and from the powder the benzoic acid is set free by action of hydrochloric acid.



The crystalline deposit formed when essential oil of almonds (benzoic aldehyde) is exposed to the air is benzoic acid.



Pure sublimed benzoic acid is also obtained from hippuric acid (p. 340). Such acid, if not thoroughly purified, may have an urinoid odor.

Jacobsen has prepared benzoic acid from benzotrichloride (trichloromethylbenzene, $\text{C}_6\text{H}_5\text{CCl}_3$), one of the trichlorotoluenes, by heating with glacial acetic acid and zinc chloride. This acid, if not very highly purified, may give a green color to flame when placed on platinum wire with a little copper oxide. In artificial benzoic acid the fragrant volatile oil characteristic of the acid from benzoin is of course absent.

Official Benzoates.—To a little benzoic acid add a few drops

of solution of ammonia or of sodium carbonate; it readily dissolves, forming corresponding benzoates (*Ammonii Benzoas*, U. S. P., *Sodii Benzoas*, U. S. P., $\text{NaC}_7\text{H}_5\text{O}_2$).



On evaporation acid crystals or, ammonia being added, neutral crystals, of ammonium benzoate are deposited.

Properties.—Benzoic acid is also soluble in other alkaline liquids, forming benzoates. It is slightly soluble in cold water, more so in hot, and readily soluble in rectified spirit. It melts at 248°F . and boils at 462° , volatilizing with only a slight residue. Heated with lime, it yields benzene. It dissolves in cold sulphuric acid without decomposition, is again deposited on dilution, and the traces of odoriferous and other substances present in the acid from benzoin only slightly color the fluid, even on gently warming.

Tests for Benzoates.—To a portion of a solution of a benzoate add a drop or two of sulphuric or hydrochloric acid; a white crystalline precipitate (benzoic acid) separates. To another portion, carefully made neutral, add a drop or two of neutral solution of ferric chloride; a reddish precipitate (ferric benzoate) results.

Cinnamic Acid ($\text{C}_8\text{H}_7\text{COOH}$).—Benzoic acid is distinguished from an allied body, cinnamic acid, the hydrogen cinnamate (occurring in balsams of Peru, Tolu, and storax, and sometimes in benzoin), by not yielding benzaldehyde ($\text{C}_6\text{H}_5\text{COH}$) (oil of bitter almonds) when distilled with chromic acid; that is, with a mixture of red potassium chromate and sulphuric acid, or when rubbed with half its weight of potassium permanganate. —Old hard balsam of Tolu yields cinnamic acid on boiling with lime and water and precipitating by hydrochloric acid. —Jacobsen makes it artificially by the prolonged reaction of glacial acetic acid and benzodichloride in the presence of zinc chloride.

CARMINIC ACID ($\text{C}_{14}\text{H}_{14}\text{O}_8$).—This is the coloring principle (about 10 per cent.) of the dried female *Coccus Cacti*, or cochineal (*Coccus*, U. S. P.). The carmine of trade, when unadulterated (*vide Pharmaceutical Journal*, 1859–60, p. 546), is carminic acid united with 2 or 3 per cent. of alumina and lime, or, occasionally, of tin oxide or albumen. It should be wholly soluble in solution of ammonia, giving an apparently clear, rich purple liquid. Carmine with French chalk or starch constitutes *face rouge* or *animal rouge*.

Merrick tests the relative value of several samples of cochineal or carmine by observing how much solution of potassium permanganate

is required to change the color of a decoction to faint pink. The silvery coating of cochineal is a wax, *coccerin*.

CETRARIC ACID is the bitter principle of Iceland "moss" (*Cetraria*, U. S. P.). In the lichen it is associated with much starch. Lichestearic also is present.

CHRYSOPHANIC ACID ($C_{15}H_{10}O_4$).—This yellow acid is found in various species of rhubarb-root (*Rheum*, U. S. P.), and, under the name of *parietinic acid*, in various common yellow lichens. Kublé considers—Dragendorff also—that the chrysophanic acid of rhubarb is only produced when a glucoside, *chrysophan*, is acted on by a ferment in the presence of water. The formation of chrysophanic acid is probably, in most if not in all cases, preceded by the occurrence of *chrysophan* or an allied body. The author found it in "chrysarobin," a name given by Kemp to the pith, etc. of a leguminous tree (*Andira araroba*), (*Chrysarobinum*, U. S. P.). Chrysarobin is also known as Araroba powder, Bahia powder, Brazil powder, Goa powder, and ringworm powder. The chrysarobin, as it occurs in the tree or when fresh, has been shown by Liebermann and Seidler to have the formula $C_{30}H_{26}O_7$; this, by oxidation and elimination of water, yields the chrysophanic acid, more or less of which occurs with the chrysarobin according to the age of the chrysarobin, and to, perhaps, the presence or absence of a ferment. A solution of chrysarobin in alkali rapidly absorbs oxygen, the fluid yielding chrysophanic acid. Chrysophanic acid may be obtained in crystals of a golden-yellow color, hence the name (from χρυσός, *chrusos*, gold, and φαίω, *phainō*, I shine). Its synonyms are *Rhaponticin*, *Rheic Acid*, *Rhein*, *Rheumin*, *Rhubarbaric Acid*, *Rhubarbarin*, *Rumicin*. Chrysophanic acid, actual or potential, black, red-brown, and red resins (*Aporetine*, *Phæoretine*, and *Erythroretine*), a bitter principle, and tannic acid are considered to be the conjoint source of the therapeutic properties of rhubarb. Chrysophanic acid may also be obtained from several species of *Rumex* or *dock*. "Rumicin" is a preparation of yellow dock (*Rumex*, U. S. P.). *Cascara Sagrada*, or sacred bark (*Rhamni Purshiani Cortex*, B. P.), according to Limousin, contains chrysophanic acid, a glucoside (?), a ferment, and various resins being also said to be present.

Emodin, $C_{15}H_{10}O_5$, is apparently closely associated, chemically, with chrysophanic acid. It is obtained with chrysophanic acid in the preparation of the latter from rhubarb. It also occurs in black alder bark (*Rhamni Frangulæ Cortex*, B. P.), according to Liebermann and Waldstein. It is said to be derived, together with glucose, from *frangulin*, $C_{21}H_{20}O_9$, the glucoside of the dried bark.

CORNIC ACID, or CORNIN.—This is, according to Geiger, the crystalline bitter principle of the bark of *Cornus florida* or *Dogwood*. A crystalline resin is also present.

CYANIC ACID (HCNO) AND OTHER CYANATES.—The reducing power of potassium cyanide (KCN) (or ferrocyanide,


$K_4FeC_6N_6$) on metallic compounds is due to the avidity with which it absorbs oxygen and forms cyanate ($KCNO$).

Process.—Fuse a few grains of potassium cyanide in a small porcelain crucible, and add powdered lead oxide; a globule of metallic lead is at once set free, excess of the oxide converting the whole of the potassium cyanide into potassium cyanate.


Urea.—Potassium cyanate ($KCNO$), or, better, lead cyanate (Pb_2CNO), treated with ammonium sulphate, yields ammonium cyanate (NH_4CNO); and solution of ammonium cyanate, when simply heated, changes to artificial *urea* (CH_4N_2O), the most important constituent of urine, and the chief form in which the nitrogen of food is eliminated from the animal system. The process will be more fully described subsequently in connection with Urea.

FORMIC ACID ($HCHO_2$).—The red ant (*Formica rufa*) and several other insects when irritated eject a strongly acid, acrid liquid having a composition expressed by the above formula, and which has appropriately received the name of formic acid; it is also contained in the leaves of the stinging-nettle. (According to Church, the sting of the wasp is *alkaline*.)

Process.—It may be artificially prepared by heating equal weights of oxalic acid and glycerin to a temperature of from 212° to 220° F. for fifteen hours, and then distilling the mixture with a considerable volume of water. The formic acid slowly passes over, glycerin being regenerated. The dilute acid may be concentrated by neutralizing with lead carbonate, filtering, evaporating to a small bulk, collecting the deposited crystalline lead formate, drying, decomposing in a current of dry sulphuretted hydrogen at 212° F., and rectifying the resulting syrupy acid from dry lead formate. It should be fluid at 48° F. and boil at 212° F. The following are the chief reactions:



Formic Acid may be instructively though not economically prepared by the oxidation of methylic alcohol (wood spirit), just as acetic acid and valerianic acid are obtained from ethylic alcohol and amylic alcohol respectively.



Tests.—Formic acid does not char when heated alone or with sulphuric acid, but splits up into carbonic oxide gas and water. It

is recognized by this property and by its reducing action on salts of gold, platinum, mercury, and silver. It is solid below 32° F.

GALLIC ACID.—See TANNIC ACID.

HEMIDESMIC ACID.—The supposed active principle of hemidesmus-root (*Hemidesmi Radix*, B. P.).

HIPPURIC ACID ($\text{HC}_9\text{H}_8\text{NO}_3$) is a constituent of human urine (much increased on taking benzoic acid), but is prepared from the urine of the horse (hence the name, from *ἵππος*, *hippos*, a horse), or, better, from that of the cow. To such urine add a little milk of lime, boil for a few minutes, remove precipitated phosphates by filtration, drop in hydrochloric acid until the liquid, after well stirring, is exactly neutral to test-paper, concentrate to about one-eighth the original bulk, and add excess of strong hydrochloric acid; impure hippuric acid is deposited. From a solution of the impure acid in hot water chlorine gas removes the color, and the liquid deposits crystals of pure hippuric acid on cooling.

Tests.—To a solution of a hippurate add neutral solution of ferric chloride; a brown precipitate (ferric hippurate) results. Salts of silver and mercury give white precipitates. Heat hippuric acid in a test-tube; it chars, benzoic acid sublimes, and vapors of characteristic odor are evolved; they contain, amongst other bodies, hydrocyanic acid and a substance smelling somewhat like Tonka bean. —The crystalline form of hippuric acid is characteristic; it will be described in connection with the subject of Urine.

QUESTIONS AND EXERCISES.


Give the preparation, composition, properties, and tests of benzoic acid, employing equations or diagrams.—What is the nature of carmine? —Name the bitter principle of Iceland "moss."—Mention the coloring principle of rhubarb.—To what is rhubarb considered to owe its medicinal activity?—How is potassium cyanate prepared, how converted into an ammonium salt, and what are the relations of the latter to urea?—Give the formulæ of cyanic acid, ammonium cyanate, and urea.—What is the chemical formula of formic acid?—Describe the artificial production of formic acid.—Describe the relation of formic acid to wood spirit.—State the sources, characters, and tests of hippuric acid.

HYDROFERROCYANIC ACID ($\text{H}_4\text{Fe}''\text{Cy}_6$ or $\text{H}_4\text{Fe}y'''$) AND OTHER FERROCYANIDES.—The ferrocyanide of most interest is that of potassium, the old "yellow prussiate of potash" (*Potassii Ferrocyanidum*, U. S. P.) ($\text{K}_4\text{FeC}_6\text{N}_6 \cdot 3\text{H}_2\text{O}$), the formation of which was alluded to in connection with hydrocyanic acid (see p. 280). It cannot be regarded as simply a double salt of potassium cyanide with ferrous cyanide ($\text{FeCy}_2 \cdot 4\text{KCy}$), its chemical properties being entirely different from either of those substances: moreover, unlike potassium cyanide, it is not poisonous. Most of the reactions point to the

conclusion that its iron and cyanogen are intimately united to form a definite quadrivalent radical appropriately termed *ferrocyanogen* (FeC_6N_6 or Fcy). 1 part of potassium ferrocyanide in 10 of water forms the official "Potassium Ferrocyanide Test-solution," U. S. P.

Tests.—Many of the ferrocyanides are insoluble, and are therefore precipitated when solution of potassium ferrocyanide is added to the various salts. Those of iron and copper, being of characteristic color, are adopted as tests of the presence of the metals or of the ferrocyanogen, as the case may be. To solution of potassium ferrocyanide add a ferric salt; a dark-blue precipitate (iron ferrocyanide, Fe_4Fcy_3 , prussian blue) falls. To another portion add solution of a copper salt; a reddish-brown precipitate (copper ferrocyanide, Cu_2Fcy) results.

Note.—The ferrocyanogen in potassium ferrocyanide is broken up when the salt is heated with sulphuric acid, *carbonic oxide* being evolved if the acid is strong (that is, ordinary oil of vitriol— H_2SO_4 with 2 or 3 per cent. of water), and *hydrocyanic acid* if weak:



Hydrocyanic Acid has already been described. (*Vide* p. 279.)


Carbonic Oxide (CO).—Heat two or three fragments of potassium ferrocyanide with eight or ten times their weight of sulphuric acid, and as soon as the gas begins to be evolved remove the test-tube from the flame, for the action, when once set up, proceeds somewhat tumultuously. Ignite the carbonic oxide at the mouth of the tube; it burns with a pale-blue flame, the product of combustion being carbonic acid gas (CO_2).

Carbonic oxide is a direct poison. It is generated whenever coke, charcoal, or coal burns with an insufficient supply of air. Hence the danger of open fires in the more or less closed apartments of ordinary dwellings.

Carbonic oxide may also be obtained from oxalic acid. (*Vide* p. 371.)

HYDROFERRICYANIC ACID ($\text{H}_6\text{Fe}^{\text{'''}}_2\text{Cy}_{12}$ or $\text{H}_6\text{Fdcy}^{\text{v}}$) AND OTHER FERRICYANIDES.—Pass chlorine gas slowly through solution of potassium ferrocyanide until the liquid, after frequent shaking, ceases to give a blue precipitate, when a minute portion is taken out on the end of a glass rod and brought into contact with a drop of dilute solution of a ferric salt; it now contains potassium ferricyanide (B. P.) ($\text{K}_6\text{Fe}^{\text{'''}}_2\text{Cy}_{12}$ or $\text{K}_6\text{Fdcy}^{\text{v}}$), *red prussiate of potash*, as it is termed from the color of its crystals. Excess of chlorine must be carefully avoided, as cyanogen chloride and other compounds are then formed. (Such a result does not ensue if bromide be used instead of

chlorine, but the process is less economical. Other processes are known.)



Note.—The removal of two atoms of potassium from the ferrocyanide molecules is the only change of composition that occurs; but the ferrocyanogen is altered in quality, its iron passing from the ferrous to the ferric condition, from bivalent to trivalent activity—altered to a condition in which it no longer precipitates ferric salts, but gives a dark-blue precipitate with ferrous salts.

The radical is distinguished as *ferricyanogen*.

Test.—To some of the solution add solution of ferrous sulphate; a dark-blue precipitate falls. This precipitate is ferric ferricyanide (Turnbull's blue), $Fe''_3Fe'''_2Cy'_{12}$ or Fe''_3Fdcy^v .



It will be noticed that the change in the condition of the iron keeps up the balance of the atomic values of the various parts of the radicals or of the salts; the quantivalential equilibrium is maintained.

A solution of 1 part of potassium ferricyanide in about 10 parts of water constitutes the "Potassium Ferricyanide Test-solution," U. S. P.

HYDROFLUORIC ACID (HF) AND OTHER FLUORIDES.—Molecular weight of HF, 20. The chief use of hydrofluoric acid is in etching on glass. The operation, performed on the small scale, also constitutes the best test for fluorine, the elementary radical of all fluorides.

Process and Test.—Warm any odd piece of window-glass, having an inch or two of surface, until a piece of beeswax rubbed on one side yields a thin oily film. When cool, make a cross, letter, or other mark on the glass by pressing a pointed piece of wood, a penknife, or file through the wax. Place a few grains of powdered fluor-spar, the commonest natural fluoride, in a porcelain crucible (or a lead cup), add a drop or two of sulphuric acid, cover the crucible with the prepared glass, waxed side downward, and gently warm the bottom of the crucible in a fume-chamber or in the open air in such a way as not to melt the wax. After a few minutes remove the glass, wash the waxed side by pouring water over it, scrape off most of the wax, then warm the glass and wipe off the remainder; the marks made through the wax will be found to be permanently etched on the glass; the acid has eaten into or *etched* (from the German *ätzen*, to corrode) the glass.

Calcium fluoride and sulphuric acid yield hydrofluoric acid, thus: $CaF_2 + H_2SO_4 = CaSO_4 + 2HF$. The hydrofluoric acid gas and the silica of the glass then yield gaseous silicon fluoride (SiF_4), which

escapes, and water, thus : $4\text{HF} + \text{SiO}_2 = 2\text{H}_2\text{O} + \text{SiF}_4$. The silica, being removed from the glass, leaves furrows or etched portions.


Note.—In the experiment just described the liberated hydrofluoric acid also attacks the siliceous glazing of the porcelain crucible; so that in important cases, where search is made for very small quantities of fluorine, vessels of platinum or lead must be employed.

Uses.—The aqueous solution of hydrofluoric acid, used by etchers, and commonly termed simply hydrofluoric acid or “fluoric” acid, is prepared in leaden stills and receivers and kept in leaden or gutta-percha bottles. Except these materials, as well as platinum and fluor-spar, hydrofluoric acid rapidly attacks any substance of which bottles and basins are usually made. It quickly cauterizes the skin, producing a painful, slow-healing sore. A mixture of hydrofluoric acid and ammonium fluoride, known as “white acid,” is also used for etching glass.

Quantivalence.—The atom of fluorine, like that of chlorine, bromine, or iodine, is univalent (F'). The great analogy existing between these radicals extends to their compounds.

Fluorine has been isolated by electrolyzing hydrofluoric acid (Moissan). It is a gas having a light greenish-yellow color and a penetrating and irritating odor, somewhat recalling that of strong hypochlorous acid. It combines with great avidity with all elements except oxygen.

HYPOPHOSPHOROUS ACID (H_3PO_2 or HPH_2O_2) AND OTHER HYPOPHOSPHITES.—Boil together in a fume-chamber, two or three grains of phosphorus, three or four grains of slaked lime, and about a quarter of an ounce of water, until phosphoretted hydrogen, a spontaneously inflammable, badly-smelling gas, ceases to be evolved. The lime must not be in great excess, or the hypophosphite will be converted into phosphate as fast as formed. The mixture, filtered and excess of lime removed by carbonic acid gas, yields solution of calcium hypophosphite ($\text{Ca}_2\text{PH}_2\text{O}_2$) (*Calcii Hypophosphis*, U. S. P.). The salt may be obtained in crystals by evaporation and slow cooling.



Phosphoretted Hydrogen (PH_3).—The above reaction is also that by which phosphoretted hydrogen, the third hydride of phosphorus, may be prepared. If the gas is to be collected, the phosphorus and water may first be boiled in a flask until a jet of spontaneously inflammable phosphorus vapor escapes, with steam, from the end of the attached delivery-tube. Strong hot solution of caustic potash or soda is next very gradually poured into the flask through a funnel tube previously fitted into the cork, the liquid being kept boiling. Phosphoretted hydrogen is then evolved, and, if the delivery-tube dip under water, may be collected, or allowed to slowly pass up through the water, bubble by bubble, so as to burst spontaneously into flame and form the peculiar vortex rings of

smoke (phosphoric anhydride) characteristic of the experiment. The spontaneous inflammability is due to the presence of a lower hydride, P_2H_4 .

Other hypophosphites ($Mg2PH_2O_2, 6H_2O$; $Fe2PH_2O_2, 6H_2O$; etc.) may thus be obtained from other hydrates or by double decompositions of the calcium salt and carbonates.

The potassium hypophosphite (*Potassii Hypophosphis*, U. S. P.) (KPH_2O_2) may be obtained in the same way from its hydrate, and many other hypophosphites ($Mg2PH_2O_2, 6H_2O$, $Fe2PH_2O_2, 6H_2O$, etc.) similarly from other hydrates, or by double decomposition of the calcium salt and carbonates.

Sodium Hypophosphite ($NaPH_2O_2, H_2O$) (*Sodii Hypophosphis*, U. S. P.), the old soda hypophosphite, is made by decomposing solution of calcium hypophosphite by sodium carbonate, filtering, and evaporating to dryness. It is a white, granular, or lamellar deliquescent substance: $Ca2PH_2O_2 + Na_2CO_3 = 2NaPH_2O_2 + CaCO_3$. When heated, the water is first evolved, then hydrogen and spontaneously inflammable phosphoretted hydrogen, and a mixture of sodium pyrophosphate and metaphosphate remains (Rammelsburg): $5NaPH_2O_2 = Na_4P_2O_7 + NaPO_3 + 2PH_3 + 2H_2$.

Hypophosphorous Acid, the hydrogen hypophosphite, may be prepared by decomposing the calcium salt by oxalic acid, or, better, the pure barium salt by sulphuric acid, filtering and evaporating; quinine hypophosphite by dissolving the alkaloid in hypophosphorous acid, or by decomposing quinine sulphate by barium hypophosphite. The latter is obtained on boiling excess of pure barium hydrate with ammonium hypophosphite until all ammonia is evolved. The ammonium salt is formed on bringing calcium hypophosphite and ammonium oxalate together in presence of a little ammonia.

Diluted Hypophosphorous Acid is official (*Acidum Hypophosphorosum Dilutum*, U. S. P.). It is a colorless, inodorous, sour liquid, containing about 10 per cent. of the real acid.

Hypophosphite of Iron ($Fe_26PH_2O_2$) (*Ferri Hypophosphis*, U. S. P., N. P.) or *Ferric Hypophosphite*, may be obtained by dissolving ferric hydrate in cold aqueous hypophosphorous acid and evaporating the solution.

The hypophosphites are often used in medicine in the form of syrups (*Syrupus Hypophosphitum*, U. S. P., and *Syr. Hypophosphitum cum Ferro*, U. S. P.). The term hypophosphite is in allusion to the smaller amount (*ἵπο*, *hupo*, under or deficiency) of oxygen in these compounds (R'_3PO_2) than in the phosphites (R_3PO_3), a class of salts having again less oxygen in their molecules than exists in those of the phosphates (R_3PO_4). The prefix *hypo* has similar significance in such words as hyposulphite and hypochlorite.

Tests.—To a portion of the above solution of calcium hypophosphite add solution of barium chloride, calcium chloride, or lead acetate; in neither case is a precipitate obtained, whereas soluble phosphates and phosphites yield white precipitates (of barium calcium or lead phosphate or phosphite). To other

portions add solutions of silver nitrate and mercuric chloride; the respective metals (HgO first, then Hg) are precipitated as by phosphites. To another small portion add zinc and dilute sulphuric acid; hydrogen and phosphoretted hydrogen are evolved as from phosphites. To another portion add sufficient oxalic acid to remove the calcium; filter; to the solution of hypophosphorous acid add solution of copper sulphate and slowly warm the mixture; a solid brown precipitate results (cuprous hydride, Cu_2H_2): increase the heat to the boiling-point; a gas (hydrogen) is evolved and metallic copper is set free. Add the ordinary nitric solution of a molybdate or tungstate to a hypophosphite solution, and then a very little sulphurous acid; a blue precipitate results, or in very dilute solutions a blue color, deepened on shaking or gently warming. Heat a small quantity of a solid hypophosphite on the end of a spatula in a flame, and note the resulting phosphorescent light and smell; for it splits up into pyrophosphate, a little metaphosphate, hydrogen, phosphoretted hydrogen, and water, the official calcium hypophosphite yielding about 80 per cent. of residue. The free acid undergoes similar but more complete decomposition.



5 grains of calcium hypophosphite, if of good quality, will almost decolorize a solution of 12 grains of potassium permanganate on boiling the mixture for about ten minutes. 5 grains of sodium hypophosphite should almost decolorize $11\frac{1}{2}$ grains of permanganate under similar conditions.

The same effect follows the addition of the permanganate to an acid solution of a phosphite, but not to that of an ortho-, meta-, or pyro-phosphate.

HYPOSULPHUROUS ACID ($\text{H}_2\text{S}_2\text{O}_3$) AND OTHER HYPOSULPHITES.—The only hyposulphite of much interest in pharmacy is the sodium salt (*Hyposulphite of Sodium*, U.S.P., the old hyposulphite of soda) ($\text{Na}_2\text{S}_2\text{O}_3 \cdot 5\text{H}_2\text{O}$). Hyposulphites are now more usually termed *Thiosulphates* (e.g. $\text{H}_2\text{SO}_3\text{S}$; $\text{Na}_2\text{SO}_3\text{S}$), being regarded as *sulphates* (e.g. Na_2SO_4) in whose molecules one atom of oxygen is displaced by one of sulphur ($\theta\epsilon\iota\omicron\nu$, *theion*, sulphur.)

Process.—Heat together gently or set aside in a warm place a mixture of solution of sodium sulphite (Na_2SO_3) and a little powdered sulphur; combination slowly takes place and sodium hyposulphite is formed. The solution, filtered from excess of sulphur, readily yields crystals. (The solution of sodium

sulphite may be made by saturating solution of soda with sulphurous acid gas.)

Use of Sodium Hyposulphite in Quantitative Analysis.—In the British Pharmacopœia sodium hyposulphite is given as a reagent for the quantitative estimation of free iodine in volumetric analysis. To a few drops of iodine-water add cold mucilage of starch; a deep-blue color (starch iodide) is produced. To the product add solution of sodium hyposulphite until the blue color just disappears. This absorption of iodine is sufficiently definite and delicate to admit of application for quantitative purposes. It depends on the combination of the iodine with half of the sodium in two molecules of the hyposulphite, the hyposulphurous radicals of the two molecules apparently coalescing to form a new radical, the tetrathionic (from τέτρας, *tetras*, four, and θεῖον, *theion*, sulphur), sodium tetrathionate ($\text{Na}_2\text{S}_4\text{O}_6$) and sodium iodide being formed.

Sulphur Oxyacids.—It will be as well here to give the formulæ of other oxyacids of sulphur, forming with the four already mentioned a series that is as useful as the series of compounds of nitrogen and oxygen in illustrating the soundness of Dalton's atomic theory. The first-named (see table below) (H_2SO_2) is more usually termed hyposulphurous acid, and the fourth ($\text{H}_2\text{S}_2\text{O}_3$) thiosulphuric acid. Moreover, there appears to be an acid ($\text{H}_2\text{S}_2\text{O}_4$) between those having the formulæ $\text{H}_2\text{S}_2\text{O}_3$ and $\text{H}_2\text{S}_2\text{O}_6$, which Bernthsen says is really Schützenberger's hydrosulphurous acid, but which the latter chemist says is probably a distinct acid:

Hydrosulphurous acid	H_2SO_2		?		$\text{H}_2\text{S}_2\text{O}_4$
Sulphurous acid	. . . H_2SO_3		Dithionic acid		$\text{H}_2\text{S}_2\text{O}_6$
Sulphuric acid	 H_2SO_4		Trithionic acid	. . .	$\text{H}_2\text{S}_3\text{O}_6$
Hyposulphurous, or			Tetrathionic acid	. .	$\text{H}_2\text{S}_4\text{O}_6$
Thiosulphuric acid	$\text{H}_2\text{S}_2\text{O}_3$		Pentathionic acid	. .	$\text{H}_2\text{S}_5\text{O}_6$

Use of "Hypo" in Photography.—The sodium hyposulphite is largely used in photography to dissolve silver chloride, bromide, or iodide off plates which have been exposed in the camera. Prepare a little silver chloride by adding a chloride (sodium chloride) to a few drops of solution of silver nitrate. Collect the precipitated chloride on a filter, wash, and add a few drops of solution of sodium hyposulphite; the silver salt is dissolved, solution of sodium and silver hyposulphite being formed. The solution of this double hyposulphite has a remarkably sweet taste, sweeter than syrup if the solution is strong. The double sodium and gold hyposulphite

has been employed for giving a pleasant tint to photographic prints.

Test.—To solution of a hyposulphite add a few drops of dilute sulphuric or other acid and smell the mixture cautiously; hyposulphurous acid is set free, but at once begins to decompose into sulphurous acid, recognized by its odor, and free sulphur ($2\text{H}_2\text{S}_2\text{O}_3 = 2\text{H}_2\text{SO}_3 + \text{S}_2$). This reaction constitutes the best test for hyposulphites. Another test of a soluble simple hyposulphite is its power of dissolving silver chloride with production of a more or less sweet solution.

QUESTIONS AND EXERCISES.

Give the formula of potassium ferrocyanide.—What is the supposed constitution of potassium ferrocyanide?—Enumerate the tests for ferrocyanogen.—What are the respective reactions of potassium ferrocyanide with strong and weak sulphuric acid?—Mention and explain a common source of carbonic oxide in households. What is the product of its combustion?—Write equations or diagrams illustrative of the changes effected in potassium ferrocyanide during its conversion into ferricyanide. By what reactions may the presence of a ferricyanide in a solution be demonstrated?—State the difference between prussian blue and Turnbull's blue.—Describe the source, mode of preparation, chief use of, and test for, hydrofluoric acid.—Illustrate by a diagram the preparation of sodium hyposulphite.—Mention the characteristic reactions of sodium hyposulphite.—Give the names and formulæ of eight acids each containing hydrogen, sulphur, and oxygen.

LACTIC ACID ($\text{HC}_3\text{H}_5\text{O}_3$) AND OTHER LACTATES.—Lactic acid occurs naturally in willow-bark (Dott). When milk turns sour, its sugar has become converted into an acid appropriately termed lactic (*lac*, *lactis*). Other saccharine and amylaceous substances also by fermentation yield lactic acid. The hydrogen lactate (lactic acid) is official (*Acidum Lacticum*, U. S. P.).

Process.—Calcium lactate and lactic acid may be prepared as follows: Mix together 8 parts of sugar, 1 of common cheese, 3 of chalk, and 50 of water, and set aside in a warm place (about 80°F.) for two or three weeks; a mass of small crystals of calcium lactate results. Remove these, recrystallize from hot water, decompose by sulphuric acid, avoiding excess, digest in alcohol, filter off the calcium sulphate, evaporate the clear solution to a syrup: this residue is ordinary lactic acid, *Acidum Lacticum* U. S. P., sp. gr. 1.213; it contains 75 per cent. of real acid. 3 fluid parts of this acid diluted with water to 20 fluid parts forms the *Acidum Lacticum Dilutum*, B. P.; sp. gr. 1.040.

Strontium Lactate, $\text{Sr}(\text{C}_3\text{H}_5\text{O}_3)_2 \cdot 3\text{H}_2\text{O}$ (*Strontii Lactas*, U. S. P.) is a permanent white crystalline powder, soluble in water

and in alcohol. It may be prepared by dissolving the carbonate in lactic acid.



Filter, concentrate, and allow to crystallize.

Ferrous Lactate (*Ferri Lactas*, U. S. P., $\text{Fe}2\text{C}_3\text{H}_5\text{O}_3, 3\text{H}_2\text{O}$) may be made by digesting iron filings in warm diluted lactic acid (1 acid to 16 water) till effervescence of hydrogen ceases, filtering, and setting aside to cool and crystallize. The crystals are collected, washed with alcohol, and dried. This ferrous lactate occurs in greenish-white crystalline crusts or grains, of a mild, sweetish, ferruginous taste, soluble in 48 parts of cold and 12 of boiling water, but insoluble in alcohol. Exposed to heat, it froths up, gives out thick white acid fumes, and becomes black, sesquioxide of iron being left. If it be boiled for fifteen minutes with nitric acid of the specific gravity 1.20, a white granular deposit of mucic acid will occur on the cooling of the liquid.

Test.—No single reaction of lactic acid is sufficiently distinctive to form a test. The crystalline form of the calcium lactate, as seen by the microscope, is characteristic. The production of this salt and the isolation of the syrupy acid itself are the only means, short of quantitative analysis, on which reliance can be placed. Lactic acid is soluble in water, alcohol, and ether, but almost insoluble in chloroform. It is only slightly colored by cold sulphuric acid. Warmed with potassium permanganate, it gives the odor of aldehyde.

A *variety of lactic acid* has been obtained from the juice of flesh; it is termed *sarcolactic acid* (from $\sigma\alpha\rho\acute{\alpha}\xi$, $\sigma\alpha\rho\chi\acute{o}\varsigma$, *sarx*, *sarcos*, flesh). Unlike lactic acid, it is precipitated by solution of copper sulphate.

MALIC ACID ($\text{C}_4\text{H}_6\text{O}_5$) AND OTHER MALATES (from *malum*, an apple).—The juice of unripe apples, gooseberries, currants, rhubarb-stalks, strawberries, grapes, etc. contains malic acid, hydrogen malate and potassium malate. When isolated it occurs in deliquescent prismatic crystals.

Tests.—Calcium malate ($\text{CaC}_4\text{H}_4\text{O}_5$) is soluble in water; hence the aqueous solution of malic acid or other malate is not precipitated by lime-water or calcium chloride; but on adding spirit of wine a white precipitate falls, owing to the insolubility of the calcium malate in alcohol. Malates are precipitated by lead salts; on warming the precipitate (lead malate) with acetic acid, it dissolves, separating out in acicular crystals on cooling. If the mixture be heated without acid, the precipitate aggluti-

nates and fuses. Hot strong sulphuric acid chars malic acid far less readily than it does nearly all other organic acids.

Asparagin ($C_4H_8N_2O_3 \cdot H_2O$).—This proximate principle of plants occurs in many vegetable juices, and doubtless plays a very important part in their nutrition. It is deposited in crystals when the fresh juices of asparagus, marsh-mallow, etc. are rapidly evaporated. It is noticed here because malic acid is readily obtained from it by oxidation, nitrogen being eliminated, and because its exact natural position among chemical substances is not yet well made out. The atoms of its molecule are those of ammonium aspartate ($NH_4C_4H_6NO_4$), into which it is converted when its solution is long boiled. Decomposed by aid of ferments, asparagin, absorbing hydrogen, yields ammonium succinate ($(NH_4)_2C_4H_4O_4$).


MECONIC ACID ($H_2C_7H_2O_7 \cdot 3H_2O$).—Opium contains meconic acid (from *μῆζων*, *mēkōn*, a poppy) partially combined with morphine. To concentrated infusion of opium, nearly neutralized with ammonia, add solution of calcium chloride; calcium meconate is precipitated. Wash the precipitate, place it in a small quantity of hot water; add a little hydrochloric acid; the clear liquid (filtered if necessary) deposits scales of meconic acid on cooling (*Acidum Meconicum*, B. P.).

Tests.—To solution of meconic acid or other meconate or to infusion of opium add a *neutral* solution of ferric chloride; a red solution (of ferric meconate) is produced. To a portion of the mixture add solution of corrosive sublimate; the color is not destroyed: to another portion add hydrochloric acid; the color is discharged. (These reagents act on iron sulphocyanate, which is of similar tint, in exactly the opposite manner.) To another portion add a drop of a diluted acid and boil; the color is not discharged. (A solution of ferric acetate, which is of similar color, *is* decomposed in boiling, giving a colorless fluid and a red precipitate—ferric oxyacetate.)



The normal potassium, sodium, and ammonium meconates are soluble in water, the acid meconates very slightly soluble; barium, calcium, lead, copper, and silver meconates are insoluble in water, but soluble in acetic acid.

METAPHOSPHORIC ACID (HPO_3) AND OTHER METAPHOSPHATES.—Prepare phosphoric anhydride (P_2O_5) by burning a small piece of phosphorus in a porcelain crucible placed on a plate and covered by an inverted test-glass, tumbler, half-pint measure-glass, or some such vessel. After waiting a few minutes for the phosphoric anhydride to fall, pour a little water on the plate and filter the liquid; the product is solution of metaphosphoric acid (from *μετά*, *meta*, a preposition denoting change) or hydrogen metaphosphate, $P_2O_5 + H_2O = 2HPO_3$.

Tests.—To solution of metaphosphoric acid add silver ammonio-nitrate, or to a neutral metaphosphate add solution of silver nitrate; a white precipitate (AgPO_3) is obtained. This reaction sufficiently distinguishes metaphosphates from the ordinary phosphates or orthophosphates (from *ὀρθός*, *orthos*, straight), as the common phosphates may, for distinction, be termed (which give, it will be remembered, a *yellow* precipitate with silver nitrate). Another variety of phosphates shortly to be considered, the pyrophosphates, also give a white precipitate with silver nitrate. To the solution of metaphosphoric acid obtained as above, or by the action of acetic acid on a metaphosphate, add an aqueous solution of white of egg; coagulation of the albumen ensues. Neither orthophosphoric nor pyrophosphoric acid coagulates albumen. Boil the aqueous solution of metaphosphoric acid for some time; on testing the solution the acid will be found to have been converted into orthophosphoric acid:



The ordinary medicinal phosphoric acid is made from phosphorus and nitric acid, the liquid being evaporated to a syrupy consistence to remove the last traces of nitric acid. It may contain pyrophosphoric and metaphosphoric acids if the heat employed be high enough to remove the elements of water:



On redilution the metaphosphoric acid only slowly reabsorbs water. If, therefore, on testing, metaphosphoric acid be found to be present, the solution should be boiled until conversion to orthophosphoric acid has occurred.

NITROUS ACID (HNO_2) AND OTHER NITRITES.—Strongly heat a fragment of potassium or sodium nitrate on a piece of platinum-foil; oxygen is evolved and potassium or sodium nitrite remains.


Test.—Dissolve the residue in water, add a few drops of dilute sulphuric acid, then a little weak solution of potassium iodide, and, lastly, some starch mucilage; the deep-blue “starch iodide” is produced: $2\text{HI} + 2\text{HNO}_2 = 2\text{H}_2\text{O} + 2\text{NO} + \text{I}_2$. Repeat this experiment, using potassium nitrate instead of nitrite; no blue color is produced.

Test for Nitrites in Water.—This liberation of iodine by nitrites, and not by nitrates, is a reaction of considerable value in searching for nitrites in ordinary drinking waters, the occurrence of such

salts, except in very deep-seated springs, being held to indicate the presence of nitrogenous organic matter in a state of oxidation or decay. The sulphuric acid used in the operation must be pure and the potassium iodide free from iodate. If much organic matter is present, however, the nitric acid liberated by the sulphuric may be reduced to nitrous acid. It is perhaps best, therefore, to add acetic acid, and (Fresenius) distil over 10 or 20 per cent. of the water and apply the test to this distillate. Very dilute solutions of nitrous acid may thus be distilled without the slightest decomposition.

Commercial Nitrous Acid.—The liquid commonly termed in pharmacy “nitrous acid” is simply nitric acid impure from the presence of nitrous acid.

Other nitrites used in medicine are nitrites of organic basylous radicals: ethyl nitrite ($C_2H_5NO_2$), or nitrous ether, is the most important constituent of *Spiritus Ætheris Nitrosi*, U. S. P. (*Vide* Index.) Amyl nitrite ($C_5H_{11}NO_2$) is also official (*Amyl Nitris*, U. S. P.). Ammonium nitrite, on being heated, yields pure nitrogen gas:



Sodium Nitrite, $NaNO_2$ (*Sodii Nitris*, U. S. P.). It yields ruddy nitrous fumes on the addition of sulphuric acid, and gives the usual deep-brown color with iron sulphate (p. 289). It is in the form of white fused masses or colorless hexagonal crystals, slightly deliquescent, and gradually oxidizes in the air to nitrate; very soluble in water, but only slightly so in alcohol. The official salt should conform to the following tests: It should readily dissolve in 20 parts of water, forming a colorless solution, and leaving no insoluble residue (absence of insoluble mucilage impurities). If 1 one drop of hydrochloric acid and a few drops of starch be added to 5 cc. of the aqueous solution, no blue coloration should appear (absence of iodide). 5 cc. of the aqueous solution, mixed with an equal volume of hydrogen sulphide, should produce no coloration or precipitate (absence of lead, arsenic, copper, etc.). 0.15 grm. of sodium nitrite dissolved in 5 cc. of water and introduced into a nitrometer, followed by a solution of 1 grm. of potassium iodide in 6 cc. of water and 15 cc. of normal sulphuric acid, should liberate nitrogen dioxide gas, which should measure not less than 50 cc. at $15^\circ C.$ ($59^\circ F.$) or 51.7 cc. at $25^\circ C.$ ($77^\circ F.$), corresponding to not less than 97.6 per cent. of the pure salt.

OPHELIC ACID ($C_{13}H_{20}O_{10}$).—This is one of the principles to which the herb *Ophelia chirata*, or Chiretta (*Chirata*, B. P.), owes its bitterness. It is an amorphous yellow body. Another is *Chiratin* ($C_{26}H_{48}O_{15}$), decomposable by hydrochloric acid into *Chiratogenin* ($C_{13}H_{24}O_8$) and ophelic acid (Höhn).

PHOSPHOROUS ACID (H_3PO_3 , or $P(OH)_3$, or H_2PHO_3).—The three acids of phosphorus—namely, phosphoric acid (H_3PO_4), phosphorous acid (H_2PHO_3), and hypophosphorous acid ($H_2P_2O_2$)—differ from each other in the proportion of oxygen, the molecules containing four, three, and two atoms respectively. In constitution they

differ by the hypothetical phosphoric radical or grouping being trivalent, the phosphorous radical bivalent, and the hypophosphorous radical univalent. These three acids and corresponding salts must not be confounded with pyrophosphoric and metaphosphoric acids and salts: the former are acids of phosphorus; the latter varieties of phosphoric acid: the former differ in proportion of oxygen, the latter by the elements of water:

Acids of Phosphorus.

H_3PO_4 , phosphoric acid.

H_2PHO_3 , phosphorous acid.

HPH_2O_2 , hypophosphorous acid.

Varieties of Phosphoric Acid.

H_3PO_4 , (ortho)phosphoric acid.

$\text{H}_4\text{P}_2\text{O}_7$, pyrophosphoric acid.

$\text{HP}\ddot{\text{O}}_3$, metaphosphoric acid.

When hypophosphorous acid is exposed to the air, oxygen is absorbed and phosphorous acid results; by prolonged exposure more oxygen is absorbed and phosphoric acid is obtained. When phosphoric acid—or rather, for distinction, orthophosphoric acid—is heated, every two molecules yield the elements of a molecule of water, and pyrophosphoric acid results; by prolonged exposure to heat more water is evolved and metaphosphoric acid is obtained. These differences will be further evident if the formulæ be written empirically, nearly all being doubled, thus:

$\text{H}_6\text{P}_2\text{O}_4$, hypophosphorous acid.


$\text{H}_6\text{P}_2\text{O}_6$, phosphorous acid.

$\text{H}_6\text{P}_2\text{O}_8$, { phosphoric acid, or
orthophosphoric acid.

$\text{H}_4\text{P}_2\text{O}_7$, pyrophosphoric acid.

$\text{H}_2\text{P}_2\text{O}_6$, metaphosphoric acid.

Or thus:



From the central compound, ordinary phosphoric acid, the acids of phosphorus differ by regularly diminishing proportions of the element oxygen, the varieties of phosphoric acid by regularly diminishing proportions of the elements of water.

Prepare phosphorous acid by exposing a moist stick of phosphorus to the air; a thin stream of heavy white vapor falls, which contains the acid in question. A simple method of collection is to place the stick in an old test-tube having a hole in the bottom, to support this tube by a funnel or otherwise, the neck of the funnel being supported in a bottle, test-glass, or tube, at the bottom of which is a little water. Or phosphorous oxide, P_4O_6 , may first be obtained by heating phosphorus in a

tube, through which a slow current of air is drawn, condensing the fumes in a U-tube surrounded by a freezing mixture, and then decomposing the oxide by water. Or chlorine is passed through phosphorus melted under water: $\text{PCl}_3 + 3\text{H}_2\text{O} = \text{P}(\text{HO})_3 + 3\text{HCl}$. Having collected some phosphorous acid, apply the various tests already alluded to under *Hypophosphorous Acid*, first carefully neutralizing the phosphorous acid by an alkali. The means by which the varieties of phosphoric acid are distinguished have been given under *Metaphosphoric Acid*.

Other soluble phosphites are prepared by neutralizing phosphorous acid with alkalis, and the insoluble phosphites by double decomposition.

Associated with phosphorous acid prepared as above stated there is said to be an acid of the formula H_2PO_3 , termed *hypophosphoric acid*. Its anhydride would be P_2O_4 .

It is interesting to note that during the oxidation of phosphorus in moist air, not only are phosphoric, hypophosphoric, and phosphorous acids formed, but also ozone (O_3), hydrogen peroxide (H_2O_2), and a small quantity of ammonium nitrate (NH_4NO_3).

PYROGALLIC ACID.—See TANNIC ACID.

PYROPHOSPHORIC ACID ($\text{H}_4\text{P}_2\text{O}_7$) AND OTHER PYROPHOSPHATES.—Heat ordinary sodium phosphate ($\text{Na}_2\text{HPO}_4 \cdot 12\text{H}_2\text{O}$) in a crucible; water of crystallization is first evolved, and dry phosphate (Na_2HPO_4) remains. Further heat to redness; two molecules yield one of water, while a salt having new properties is obtained: $2\text{Na}_2\text{HPO}_4 - \text{H}_2\text{O} = \text{Na}_4\text{P}_2\text{O}_7$. It is termed sodium pyrophosphate, in allusion to its origin ($\pi\tilde{\nu}\rho$, $p\tilde{u}r$, fire). From its solution in water it may be obtained in prismatic crystals ($\text{Na}_4\text{P}_2\text{O}_7 \cdot 10\text{H}_2\text{O}$), *Sodii Pyrophosphas*, U. S. P., *Pyrophosphate of Sodium*, or *Sodium Pyrophosphate*. Phosphoric acid itself is similarly affected by heat: $2\text{H}_3\text{PO}_4 - \text{H}_2\text{O} = \text{H}_4\text{P}_2\text{O}_7$ (pyrophosphoric acid), though metaphosphoric acid is also formed. Other pyrophosphates are similarly produced, or by double decomposition and precipitation, or by neutralizing pyrophosphoric acid by an oxide, hydrate, or carbonate. *Ferri Pyrophosphas Solubilis*, U. S. P., is official.

Tests.—To solution of a pyrophosphate add solution of silver nitrate; a dense white precipitate falls (silver pyrophosphate, $\text{Ag}_4\text{P}_2\text{O}_7$), differing much in appearance from the white gelatinous silver metaphosphate or the yellow orthophosphate. To pyrophosphoric acid or to a pyrophosphate mixed with acetic acid add an aqueous solution of albumen (white of egg); no

precipitate occurs. Metaphosphoric acid, it will be remembered, gives a white precipitate with albumen.

QUESTIONS AND EXERCISES.

What are the sources of lactic acid?—How is lactic acid usually prepared?—Name some of the plants in which malic acid is found.—Whence is meconic acid derived?—By what process may meconic acid be isolated?—Which is the best test for the meconic radical?—Distinguish meconates from sulphocyanates.—Give the mode of manufacture of hypophosphites.—How is phosphoretted hydrogen prepared?—By what ready method may metaphosphoric acid be obtained for experimental purposes?—Name the tests for metaphosphates.—How may meta- or pyro-phosphoric acid be converted into orthophosphoric acid?—Describe the preparation of phosphorous acid.—State the relations of the acids of phosphorus to each other.—How are the pyrophosphates prepared?—Define, by formulæ, metaphosphates, pyrophosphates, orthophosphates, phosphites, and hypophosphites.—Mention the tests by which meta-, pyro-, and ortho-phosphates are analytically distinguished.—How are hypophosphites and phosphites detected?

SILICIC ACID (H_4SiO_4) AND OTHER SILICATES.—Silicates of various kinds are among the commonest of minerals. The various *clays* are aluminium silicates; the volcanic substance termed *pumice-stone* is a porous aluminium and alkali-metal or alkaline-earth metal silicate; *meerschaum* is an acid magnesium silicate; the ordinary *sandstones* are chiefly silica; *sand*, *flint*, *quartz*, *agate*, *chalcedony*, and *opal* are silicic anhydride or *silica* (SiO_2). *Tripoli*, a polishing powder now found in many other countries than Tripoli, and consisting of infusorial skeletons, is nearly pure silica. *Bath brick*, used in knife-polishing, is a silico-calcareous deposit found in the estuary at Bridgewater and other places. *Tourmalines*, plates of which, cut parallel to the axis of a crystal, are used as polarizers or analyzers in microscopy, are all aluminium silicates with varying iron, copper, manganese, etc. silicates. *Asbestos*, or *amianth*, is a fibrous calcium and magnesium silicate, the length of the fibres being from less than one inch to five feet. A single silk-like fibre can easily be fused, but even in very small masses and for all practical purposes asbestos is infusible, and of course incombustible. It is also a bad conductor of heat. It is already largely used in packing piston-rods and joints and for steam apparatus generally; as a covering for boilers to prevent loss of heat by radiation; and for so lining ceilings, floors, and other partitions as to render rooms, etc. fireproof. Artificial acid, and therefore insoluble, silicates are familiar under the forms of *glass* and *earthenware*. Common English window-glass is usually calcium, sodium, and aluminium silicate; French glass, calcium and sodium silicate; Bohemian, chiefly potassium and calcium silicate; English flint or crystal glass for ornamental, table, and optical purposes is mainly potassium and lead silicate. Earthenware is mostly aluminium silicate (clay), with more or less of the more fusible silicates—namely, those of

calcium, sodium, and potassium, and in the commoner forms, iron silicate. The various kinds of *porcelain* (China, Sèvres, Meissen, Berlin, English), *Wedgwood-ware*, and *stoneware* are varieties of earthenware. *Kaolin*, or *China clay*, which is disintegrated *felspar*, not more common in China than in Devonshire and Cornwall, is the clay which yields the finest translucent porcelain. *Crucibles*, *bricks*, and *tiles* are clay-silicates. *Fire-clay* contains excess of silica and very small proportions of the fusible silicates; hence its refractory character. *Mortar* is essentially calcium silicate. *Portland*, *Roman*, and other "hydraulic" cements* are calcium silicates with more or less aluminium silicate.

Mix together a few grains of powdered flint or sand with about five or six times its weight of sodium carbonate and an equal quantity of potassium carbonate, and fuse a little of the mixture on platinum-foil in the blowpipe flame; the product is an alkaline, and therefore soluble, silicate, a kind of *soluble glass*. Boil the foil in water for a few minutes; filter; to a portion add excess of hydrochloric acid, evaporate the solution to dryness, and again boil the residue in water and acid; silicon oxide, silicic anhydride, or *silica* (SiO_2) remains as a light, flaky, insoluble powder.

The *soluble glass* or *glass liquor* of trade commonly contains 10 or 12 per cent. of soda (NaHO) to 20 or 25 per cent. of silica (SiO_2). When of sp. gr. 1.300 to 1.400 it satisfies official requirements (*Liquor Sodii Silicatis*, *Solution of Sodium Silicate*, U. S. P.).

The foregoing operation constitutes the *test* for silicates. By fusion with alkali the silicate is decomposed and a soluble alkaline silicate formed. On addition of acid, silicic acid (H_4SiO_4) is set free, but remains in solution if the fluid is not too strong. The heat subsequently applied eliminates water and reduces the silicic acid to silica (SiO_2), which is insoluble in water or acid. By the addition of hydrochloric acid to soluble glass, and removal of the resulting alkaline chloride and excess of hydrochloric acid by dialysis (a process to be subsequently described), a pure aqueous solution of silicic acid may be obtained; it readily changes into a gelatinous mass of silicic acid. Possibly some of the natural crystallized varieties of silica may have been obtained from the silica contained in such an aqueous solution, nearly all waters yielding a small quantity of silica when treated as above described.

A variety of silicic acid (H_2SiO_3), sometimes termed *diabasic* to distinguish it from the normal or *tetrabasic* acid (H_4SiO_4), results when the aqueous solution of the latter is evaporated *in vacuo*.

Siliciuretted hydrogen, or silicon hydride (SiH_4), is a spontaneously inflammable gas formed on treating magnesium silicide with hydrochloric acid. It is the analogue of light carburetted hydrogen or methane (CH_4). A liquid silicon chloride (SiCl_4)

* For an article on "Cements" generally see *Pharmaceutical Journal* for April 17, 1880.

analogous to carbon tetrachloride (CCl_4) and a gaseous fluoride (SiF_4) also exist.

Many other analogies are traceable between the elements silicon and carbon, including, perhaps, boron, especially amongst their organic compounds.

SUCCINIC ACID ($\text{H}_2\text{C}_4\text{H}_4\text{O}_4$).—Amber (*Succinum*) is a resin usually occurring in association with coal and lignite. From the fact that fragments of coniferous fruit are frequently found in amber, and impressions of bark on its surface, it is considered to have been an exudation from a species of *Pinus* now probably extinct. Heated in a retort, amber yields, first, a sour aqueous liquid containing acetic acid, and another characteristic body appropriately termed *succinic acid*; secondly, a volatile liquid known as *oil of amber* (*Oleum Succini*, U. S. P. 1880), resembling the oil yielded by most resinous substances under similar circumstances; and, thirdly, a pitchy residue allied to asphalt. The succinic acid is a normal constituent of the amber; the acetic acid is produced during distillation. Succinic acid has also been found in wormwood, in several pine resins, and in certain animal fluids, such as those of hydatid cysts and hydrocele. It may be obtained artificially from butyric, stearic, or margaric acid by oxidation. Tartaric, malic, and succinic acids are also convertible the one into the other.

Succinates are neutral ($\text{R}'_2\text{C}_4\text{H}_4\text{O}_4$) and acid ($\text{R}'\text{HC}_4\text{H}_4\text{O}_4$); a potassium and hydrogen succinate ($\text{KHC}_4\text{H}_4\text{O}_4, \text{H}_2\text{C}_4\text{H}_4\text{O}_4, \text{H}_2\text{O}$), analogous to the superacid oxalate, salt of sorrel, also exists. Soluble succinates give a bulky brown precipitate with neutral ferric chloride, only less voluminous than ferric benzoate; a white precipitate with lead acetate, soluble in excess of either reagent; with silver nitrate a white precipitate after a time; with barium chloride no precipitate at first, but a white one (barium succinate) on the addition of ammonia and alcohol. Succinates are distinguished from benzoates by the last-named reaction and by not yielding a precipitate on the addition of acids. (*Vide* p. 337.)

SULPHOCYANIC ACID (HCNS) AND OTHER SULPHOCYANATES.—Boil together sulphur and solution of potassium cyanide; solution of potassium sulphocyanate (KCNS) is formed. Warm the liquid, add hydrochloric acid till it faintly reddens litmus-paper, and filter; any potassium sulphide is thus decomposed, and the solution may then be used for the following reactions. The salt readily crystallizes.

Tests.—To a small portion of the solution add a ferric salt (Fe_2Cl_6); a deep blood-red solution (ferric sulphocyanate) is formed. To a portion of the red liquid add a little hydrochloric acid; the color is not discharged (ferric meconate, a salt of similar tint, is decomposed by hydrochloric acid). In the acid liquid place a fragment or two of zinc; sulphuretted hydrogen is evolved and the red color disappears.

To another portion of the ferric sulphocyanate add solution of corrosive sublimate; the color is at once discharged (ferric meconate is unaffected by corrosive sublimate). The ferric is the best test of the presence of a sulphocyanate, and indirectly is also a good test of the presence of hydrocyanic acid or cyanogen (see p. 284). Solutions of pure ferrous salts are not colored by the solution of sulphocyanate. Red ferric acetate is decomposed by ebullition. Neither the ferric acetate nor the meconate yields its color to ether, but on shaking ferric sulphocyanate solutions with ether the latter takes up the salt and becomes of a purple color.

To solution of a sulphocyanate add solution of mercuric nitrate; mercuric sulphocyanate is precipitated as a white powder.

Pharaoh's Serpents.—Mercuric sulphocyanate, thoroughly washed and made up into a little cone, forms the toy termed Pharaoh's serpent. It readily burns when ignited, the chief product being a light solid matter (mellon, C_9N_{13} , and melam, $C_3H_6N_6$) which issues from the cone in a snake-like coil of extraordinary length. The other products are mercuric sulphide (of which part remains in the snake and part is volatilized), nitrogen, sulphurous and carbonic acid gases, and vapor of metallic mercury. (For details concerning the economical manufacture of sulphocyanates see *Pharmaceutical Journal*, 2d series, vol. vii. p. 581 and p. 152.)

The sulphocyanic radical (CNS) is often termed sulphocyanogen (Scy), and its compounds regarded as *sulphocyanides*. Saliva contains sulphocyanates.

TANNIC ACID, GALLOTANNIC ACID, OR TANNIN.—*Digallic Acid* (*Acidum Tannicum*, U. S. P., $C_{14}H_{10}O_9$).—This is a very common astringent constituent of plants, but is contained in largest quantity in galls (excrescences on the oak formed by the puncture and deposited ova of an insect). English galls contain from 14 to 28 per cent. of tannic acid; Aleppo galls (*Galla*, U. S. P.), 25 to 65. It is present also in the white oak (*Quercus Alba*, U. S. P.).

Gallotannic acid, $C_6H_2(OH)_3CO \cdot O \cdot C_6H_2(OH)_2COOH$.

Gallic acid (p. 360), $C_6H_2(OH)_3COOH$.

Process.—"Expose powdered galls [about an ounce is sufficient for the purpose of study] to a damp atmosphere for two or three days, and afterward add sufficient ether to form a soft paste. Let this stand in a well-closed vessel for twenty-four hours; then, having quickly enveloped it in a linen cloth, submit it to strong pressure, so as to separate the liquid portion, which contains the bulk of the tannin in solution. Reduce the pressed cake to powder, mix it with sufficient ether, to which one-sixteenth of its bulk of water has been added,

to form again a soft paste, and press this as before. Mix the expressed liquids, and expose the mixture to spontaneous evaporation until, by the aid subsequently of a little heat, it has acquired the consistence of a soft extract; then place it on earthen plates or dishes and dry it in a hot-air chamber at a temperature not exceeding 212° F."

The resulting tannic acid occurs in "pale yellow vesicular masses or thin glistening scales, with a strongly astringent taste and an acid reaction, readily soluble in water and rectified spirit, very sparingly soluble in pure ether, though soluble in the ethereal fluid used in the foregoing process—a fluid which is really a mixture of true ether, water, and alcohol (both the latter contained in the common "ether") and a little added water also.

Medicinal Uses.—Tannic acid is very soluble in water, and in this form is usually administered in medicine. Its official (British) preparations are *Glycerinum Acidi Tannici*, *Suppositoria Acidi Tannici*, *Suppositoria Acidi Tannici cum Sapone*, and *Trochisci Acidi Tannici*. The preparations of the United States Pharmacopœia are *Collodium Stypticum*, 20 grm. in 100 cc.; *Glyceritum Acidi Tannici*, 20 per cent.; *Trochisci Acidi Tannici*, 0.06 grm. in each; and *Unguentum Acidi Tannici*, 20 per cent.

Tests.—To an aqueous solution of tannic acid add aqueous solution of gelatin; a yellowish-white flocculent compound of the two substances is precipitated.

Tanning.—The above reaction also serves to explain the chemical principle involved in *tanning*—the operation of converting skin into leather. In that process the skin is soaked in infusion of oak-bark (*Quercus Cortex*, B. P.), the tannic acid of which, uniting with the gelatinous tissues of the skin, yields a compound very well represented by the above precipitate. The outer bark of the oak contains little or no tannic acid, and is commonly shaved off from the pieces of bark which are large enough to handle; useless coloring-matter is thus also rejected. Other infusions and extracts besides that of oak-bark (chiefly catechu, sumach, and valonia) are largely used by tanners; if used alone, these act too quickly and give a harsh, hard, less durable leather. The tannic acid of these preparations is probably slightly different from that of oak-bark.

To an aqueous solution of tannic acid add a neutral solution of a ferric salt; dark bluish-black ferric tannate is slowly precipitated. This is an excellent test for the presence of tannic acid in vegetable infusions. The precipitate is the basis of nearly all black *writing-ink*. Ferrous salts give at first only a slight reaction with tannic acid, but the liquid gradually darkens: characters written with a liquid of this kind, of

proper strength, become quite black in a few hours, and are very permanent.

To an aqueous solution of tannic acid add solution of tartar emetic; antimony tannate is precipitated. This reaction and that with gelatin are useful in the quantitative estimation of the amount of tannic acid in various substances, the separation of the gelatin tannate being much promoted by previously adding some heavy neutral powder, such as barium sulphate, and well stirring while adding the gelatin.

Tannic acid, as it occurs in oak-bark, is said to be a glucoside; that is, like several other substances, it yields glucose (grape-sugar) when boiled with dilute sulphuric or hydrochloric acid, the other product being gallic acid. Possibly, however, the sugar is not a necessary constituent of the tannin, certainly not of all of them.

Catechu (*Catechu*, B. P.), *Gambier* or *Terra Japonica*, an extract of the *Uncaria Gambier*; as well as the true *Catechu Cutch*, or *Terra Japonica*, an extract from the *Acacia Catechu* (*Catechu*, U. S. P., *Catechu nigrum*, P. I.) and *A. Suma*; *East Indian Kino* (*Kino*, U. S. P.), from the *Pterocarpus marsupium*; also *Bengal or Butea Kino*, from the *palas* or *dhak* tree, *Butea frondosa* (*Butea Gummi vel Kino Bengalensis*, P. I.); Botany Bay or Australian kinos from various species of *Eucalyptus* or blue-gum trees; and some other vegetable products—contain a variety of tannic acid (*mimotannic acid*), which gives a greenish precipitate with neutral solutions of ferric salts. According to Paul and Kingzett, it yields, when decomposed, unfermentable sugar and an acid different to ordinary gallic acid. *Catechu* and *gambier* also contain *catechuic acid* or *catechin*, $C_{13}H_{12}O_5$, a body occurring in minute colorless acicular crystals, and, like *mimotannic acid*, affording a green precipitate with ferric salts.

Bael-fruit (*Belæ Fructus*, U. S. P.), from the *Ægle Marmelos*, is said to owe its power as a remedy for dysentery and diarrhoea to a variety of tannic acid, but this is questionable. About 10 per cent. of tannic acid is contained in the leaves of *Castanea dentata* (*Castanea*, U. S. P.), the tree yielding edible chestnuts. The rind of the fruit of the *pomegranate* (*Punica granatum*) (*Granati Cortex*, P. I.) contains tannic acid. The astringency of pomegranate-root bark (*Granati Radicis Cortex*, B. P. and P. I., *Granatum*, U. S. P.) is due to a tannic acid (its anthelmintic properties probably to a resinoid matter, or possibly to what Tanret states to be a liquid alkaloid, *pelletierine*, $C_{16}H_{30}N_2O_2$). A tannic acid also probably gives the astringency to logwood (*Hæmatoxylin*, U. S. P.). *Rhatany-root bark* (*Krameria*, U. S. P.) contains about 20 per cent. of tannic acid, its active astringent principle; *rhubarb-root* about 9 per cent. *Bearberry-leaves* (*Uva Ursi*, U. S. P.) owe most of their therapeutic power to about 35 per cent. of tannic acid. (The cause of their influence on the kidneys is not yet traced.) They also contain *arbutin*, a crystalline glucoside. *Larch-bark* (*Laricis Cortex*, B. P.), the inner bark of *Pinus Larix* or *Larix Europæa*, contains, accord-

ing to Stenhouse, a considerable amount of a tannic acid giving olive-green precipitates with iron salts, and *larixin* and *larixinic acid* ($C_{10}H_{10}O_5$), a somewhat bitter substance. *Areca-nuts* or *betelnuts*, from the Areca palm (*Areca Catechu*), besides the alkaloid *arekane* (Bombelon), contain a very active alkaloid, *arecoline*, $C_8H_{13}NO_2$ (Jahns), said to be the vermifugal principle; *arecaine*, an inert alkaloid (Jahns), and, according to Flückiger and Hanbury, about 16 per cent. of "tannic matter." The extract of the fruit of *gab*, or *Diospyros embryopteris* (*Diospyri Fructus*, P. I.) is a powerful astringent containing tannic acid. The rhizome (*Geranium*, U. S. P.) of *Geranium maculatum*, spotted cranesbill or arum-root, contains both tannic and gallic acids. *Sumac*, or *shumac*, or *sumach*, the leaves and stalks of various species of *Rhus*, chiefly *Rhus coriaria*, contains ordinary tannic acid and gallic acid. The fruit of sumach (*Rhus glabra*, U. S. P.) contains tannic and much malic acid. The bark of *Prinos verticillatus*, the black alder or *winterberry*, contains tannin and a bitter principle. The principal constituent of the bark of the root of *Rubus villosus*, or *high blackberry*, and of *R. canadensis* and *R. trivialis* (*Rubus*, U. S. P.), is tannic acid.

Gallic Acid ($H_3C_7H_3O_5, H_2O$) (*Acidum Gallicum*, U. S. P.) occurs in small quantity in oak-galls and other vegetable substances, but is always prepared from tannic acid. Gallic acid forms slender acicular, fawn-colored crystals, soluble in one hundred and eighteen times their weight of cold or three of boiling water, freely in spirit, sparingly in ether.

"Boil 1 part of coarsely powdered galls with 4 fluid parts of diluted sulphuric acid for half an hour, then strain through calico while hot; collect the crystals that are deposited on cooling, and purify these with animal charcoal and repeated crystallization."

Test.—To an aqueous solution of gallic acid add a neutral solution of a ferric salt; a bluish-black precipitate of ferric gallate falls, similar in appearance to ferric tannate. Ferrous salts also are blackened by gallic acid. To more of the solution add an aqueous solution of gelatin; no precipitate occurs. By the latter test gallic is distinguished from tannic acid.

Pyrogallic Acid, or Pyrogallol, $C_6H_3(OH)_3$.—This substance sublimes in light feathery crystals when gallic acid is heated. It is very soluble in water, alcohol, and ether. Or it may be formed by heating gallic acid with three or four times its weight of glycerin to a temperature of 190° or 200° C. for a short time, until carbonic acid ceases to be evolved. Longer heating at lower temperatures is not equally effective, and below 100° C. probably no pyrogallol is produced (Thorpe). To an aqueous solution add a neutral solution of a ferric salt; a red color is produced which turns dark bluish on the addition of ammonia. To another portion add a ferrous salt; a

deep-blue color results. The aqueous solution of pyrogallol reduces solutions of silver, mercury, and gold salts in the cold.

Test for the Three Acids.—To three separate small quantities of milk of lime in test-tubes add, respectively, tannic, gallic, and pyrogallic acids; the first slowly turns brown, the second more rapidly, while the pyrogallic mixture at once assumes a beautiful purplish-red color changing to brown. These reactions are characteristic; they are accompanied by absorption of oxygen from the air.

Use of Pyrogallic Acid in Gas Analysis.—A mixture of pyrogallic acid and solution of potash absorbs oxygen with such rapidity and completeness that a strong solution of each, passed up successively by a pipette into a graduated tube containing air or other gas, forms an excellent means of estimating free oxygen. The value of this method may be roughly proved by pouring a small quantity of each solution into a phial, immediately and firmly closing its mouth with a cork, thoroughly shaking the mixture, and then removing the cork under water: the water rushes in and occupies about one-fifth of the previous volume of air, indicating that the atmosphere contains one-fifth of its bulk of oxygen. The small amount of carbonic acid gas present in the air is also absorbed by the alkaline liquid; in delicate experiments this should be removed by the alkali before the addition of pyrogallic acid.

Toxicodendric Acid is the volatile, excessively acrid, and poisonous principle of the poison oak or poison ivy, the fresh leaves of which are official (*Rhus toxicodendron*, U. S. P.).

URIC ACID ($\text{H}_2\text{C}_5\text{H}_2\text{N}_4\text{O}_3$) AND OTHER URATES.—Acidulate a few ounces of human urine with hydrochloric acid, and set aside for twenty-four hours: a few minute crystals of uric acid will be found adhering to the sides and bottom of the vessel and floating on the surface of the liquid.

Microscopical Test.—Remove some of the floating particles by a slip of glass, and examine by a powerful lens or microscope; the chief portion will be found to be in yellowish semi-transparent crystals, more or less square, two of the sides of which are even and two very jagged; but other forms are common. (See the lithographs in the section on Urinary Sediments.)

Chemical Test.—Collect more of the deposit, place in a watch-glass or small white evaporating-dish, remove adherent moisture by a piece of blotting- or filter-paper, add a drop or two of strong nitric acid, and evaporate to dryness; the residue will be red. When the dish is cold add a drop of solution of ammonia; a purplish-crimson color results. The color is deepened on the addition of a drop of solution of potash.

Notes.—Uric acid (or lithic acid) and sodium, potassium, calcium, and ammonium urates (or lithates) are common constituents of animal excretions. Human urine contains about 1 part of urate (usually sodium urate) in 1000. When more than this is present the urate is often deposited as a sediment in the excreted urine, either at once or after standing a short time. Uric acid or other urate is also occasionally deposited before leaving the bladder, and, slowly accumulating there, forms a common variety of urinary calculus.—Some urates are not definitely crystalline, but, when treated with dilute nitric acid or a drop of solution of potash and then a drop or two of acetic acid, jagged microscopic crystals of uric acid are usually formed. All urates yield the crimson color when treated as above described. This color is due to a definite substance, *murexid*, $C_8H_8N_6O_6$ (from the *murex*, a shell-fish of similar tint, and from which the ancient and highly valued purple dye seems to have been prepared), and the test is known as the *murexid* test. The formation of murexid is due to the action of ammonia on *alloxan* ($C_4H_2N_2O_4 \cdot 4H_2O$) and other white crystalline products of the oxidation of uric acid by nitric acid. Murexid is a good dye; it may be prepared from *guano* (the excrement of sea-fowl), which contains a large quantity of ammonium urate. The excrement of the serpent is almost pure ammonium urate.

Uric acid and the urates will again be alluded to in connection with the subject of Morbid Urine.

Constitution of Uric Acid.—The physiological and pathological importance of uric acid has obtained for it great attention from chemists, a knowledge of its constitution being rightly regarded as amongst the most prominent of chemical desiderata. (For accounts of what has been done in this direction in recent years students of organic chemistry may consult the *Pharmaceutical Journal*, 3d Series, vol. xiv. p. 771; vol. xv. pp. 119 and 411; and vol. xviii. p. 69. See also page 390.)




VALERIANIC ACID, OR VALERIC ACID ($HC_5H_9O_2$), AND OTHER VALERIANATES.—In a test-tube place a few drops of amylic alcohol (fousel oil) with a little dilute sulphuric acid and a grain or two of red potassium chromate; cork the tube, set aside for a few hours, and then heat the mixture; valerianic acid, the hydrogen valerianate, of characteristic valerian-like odor, is evolved.

Valerianic acid occurs in valerian-root in association with the essential oil from which it is apparently derived (*vide Index*), but is usually prepared artificially, by the foregoing process, from amylic alcohol, to which it bears the same relation as acetic acid does to common alcohol:





Sodium Valerianate ($NaC_5H_9O_2$) (*Sodii Valerianas*, B. P.), the old valerianate of soda, is prepared from the valerianic acid

and amyl valerianate obtained on distilling the mixture of amylic alcohol (4 fl. oz.), sulphuric acid ($6\frac{1}{2}$ fl. oz. with 10 of water), and red potassium chromate (9 oz. in 70 of water): The mixture should stand for several hours before heat is applied.



The distillate (70 or 80 oz.) is saturated with soda, which not only yields sodium valerianate with the free valerianic acid, but decomposes the amyl valerianate produced at the same time, more sodium valerianate being formed and some amylic alcohol set free, according to the following equations:



From the solution of sodium valerianate (which should be made neutral to test-paper by careful addition of soda solution) the solid white salt is obtained by evaporation to dryness and cautious fusion of the residue. The mass obtained on cooling should be broken up and kept in a well-closed bottle. It should be entirely soluble in spirit.

Other Valerianates, as zinc valerianate (*Zinci Valerianas*, U. S. P.) and *Ferric Valerianate* (*Ferri Valerianas*, U. S. P., $\text{Fe}_2\text{6C}_5\text{H}_9\text{O}_2$), may be made by double decomposition of sodium valerianate with the sulphate or other salt of the metal the valerianate of which is desired, the new valerianate either precipitating or crystallizing out. A hot solution of zinc sulphate ($5\frac{3}{4}$ parts) and sodium valerianate (5 parts) in water (40 parts) gives a crop of crystals of zinc valerianate on cooling.

Tests.—Heated with diluted sulphuric acid, valerianates of the metals give valerianic acid, which has a highly characteristic smell. Sodium valerianate thus treated, and the resulting oily acid liquid purified by agitation with sulphuric acid and distillation, furnishes valerianic acid. Dry ammonia gas passed into valerianic acid gives white lamellar crystals of *Ammonium Valerianate* (*Ammonii Valerianas*, $\text{NH}_4\text{C}_5\text{H}_9\text{O}_2$, U. S. P.).

The amylic alcohol ($\text{C}_5\text{H}_{11}\text{HO}$) from which valerianates are pre-

pared may contain the next lower homologue, *butylic alcohol* (C_4H_9HO). (*Vide* "Homology" in the Index.) This, during oxidation, will be converted into *butyric acid* ($HC_4H_7O_2$), the next lower homologue of valerianic acid ($HC_5H_9O_2$), and hence the various valerianates be contaminated by some *butyrates*. These are detected by distillation with diluted sulphuric acid and addition of solution of copper acetate to the distillate, which at once becomes turbid if butyric acid be present. In this reaction valerianic and butyric acids are produced by double decomposition of the valerianate and butyrate by the sulphuric acid, and distil over on the application of heat. On the addition of copper acetate ($Cu_2C_2H_3O_2$) copper butyrate ($Cu_2C_4H_7O_2 \cdot H_2O$) is formed, and, being almost insoluble in water, is at once precipitated or remains suspended, giving a bluish-white opalescent liquid. Copper valerianate ($Cu_2C_5H_9O_2$) is also formed after some time, but is far more soluble than the butyrate, and only slowly collects in the form of greenish oily drops, which gradually pass into greenish-blue hydrous crystalline copper valerianate (Larocque and Huralt).

VANILLIC ACID ($HC_8H_7O_3$), or VANILLIN ($C_8H_8O_3$), or METHYLPROTOCATECHUIC ALDEHYDE ($C_7H_5CH_3O_3$), is the body to which is due the odor and flavor of vanilla. It also occurs in Siam benzoin, *Rosa canina*, etc. The white crystals commonly found on vanilla (the prepared unripe pods of *Vanilla planifolia*), previously termed *vanillin*, were found by Carles to be a weak acid. It occurs in vanilla to the extent of from $1\frac{1}{2}$ to 3 per cent. Vanillin has in recent years been prepared artificially by Tiemann and Haarman from coniferin, a glucoside existing in the sapwood of pines. The body remaining after the removal of glucose from coniferin, or, indeed, coniferin itself, by action of a mixture of red potassium chromate and sulphuric acid, yields the vanillin. It also may be obtained by a series of reactions starting from that of carbonic acid on potassium carbolate; also from the eugenol of oil of cloves. By action of hydrochloric acid vanillin yields methyl chloride and protocatechuic aldehyde. Such reactions will be better understood when the pupil has studied succeeding sections on what is commonly termed *Organic Chemistry*. Artificial vanilla is less stable than natural vanillin, perhaps because with the latter is associated a preservative resin.

QUESTIONS AND EXERCISES.

What is the constitution of nitrites?—Mention a test for nitrites in potable waters.—Which nitrites are official?—Give the names of some natural and artificial silicates.—What is "soluble glass"?—Distinguish between silica and silicic acid.—How are silicates detected?—What is the quantivalence of silicon?—Mention the sources, formulæ, and analytical reactions of succinates.—State the mode of manufacture and tests of sulphocyanates.—What proportion of tannic acid is contained in galls?—Describe the official process for the preparation of tannic acid.—Explain the chemistry of "tanning."—Enumerate the tests for tannic acid.—What is the assumed constitution of tannic acid?—Mention official substances other than galls whose astringency is due to tannic acid.—

How is gallic acid prepared?—By what reaction is gallic distinguished from tannic acid?—Mention the characteristic properties of pyrogallie acid.—Explain the murexid test for uric acid.—Describe the artificial preparation of valerianic acid and other valerianates, giving diagrams or equations.—What is the formula of valerianic acid?—How are buty-rates detected in presence of valerianates?

DETECTION OF THE ACIDULOUS RADICALS OF SALTS SOLUBLE IN WATER.

Analytical operations may now be resumed, the detection of acidulous radicals being practised for two or three days, and then full analyses made, both for basylous and acidulous radicals. To this end a few compounds of stated metals (potassium, sodium, or ammonium) should be placed in the hands of the practical student for examination according to the following paragraphs and tables. Mixtures in which both basylous and acidulous radicals may be sought should then be analyzed.

In examining salts soluble in water, and concerning which no general information is obtainable, search must first be made for any basylous radicals by the appropriate methods. (*Vide* pp. 224 or 257–262.) Certain metals having been thus detected, a little reflection on the character of their salts will at once indicate what acidulous radicals may be and what cannot be present. Thus, for instance, if the substance under examination is freely soluble in water and lead is found, only the nitric and acetic radicals need be sought, none other of the lead salts than nitrate or acetate being freely soluble in water. Moreover, the salt is more likely to be lead acetate than nitrate, for two reasons: the former is more soluble than the latter, and is by far the commoner salt of the two. Medical and pharmaceutical students have probably, in dispensing, already learned much concerning the solubility of salts, and whether a salt is rarely employed or is in common use. And, although but little dependence can be placed on the chances of a salt being present or absent according to its rarity, still the point may have its proper weight. If in a mixture of salts, ammonium, potassium, and magnesium have been found associated with the sulphuric, nitric, and hydrochloric radicals, and we are asked how we suppose these bodies may exist in the mixture, it is far more in accordance with common sense to suggest that sal-ammoniac, nitre, and Epsom salt were originally mixed together than to suppose any other possible combination. Such appeals to experience regarding the solubility or rarity of salts cannot be made by any one not previously acquainted, or insufficiently acquainted, with the characters of salts; in such cases the relation of a salt to water and acids can be ascertained by referring to the following table (p. 368) of the solubility or insolubility of about five hundred of the common and rarer salts met with in chemical operations.

The opposite course to the above (namely, to ascertain what acidulous radicals are present in a mixture, and then to appeal to experi-

ence to tell what basylous radicals may be and what cannot be present) is impracticable; for acidulous radicals cannot be separated out, one after the other, from one and the same quantity of substance by a similar treatment to that already given for basylous radicals. Indeed, such a sifting of acidulous radicals could scarcely be accomplished at all or only by a vast deal of labor. The basylous radicals must therefore first be detected.

Even when the basylous radicals have been found, the acidulous radicals which may be present must be sought for singly, the only additional aid which can be brought in being the action of sulphuric acid, a barium salt, a calcium salt, silver nitrate, and ferric chloride on *separate* small portions of the solution under examination, as detailed in the second of the following tables.

Practical Analysis.

Commence the analysis of an aqueous solution of a salt or salts, the basylous radicals in which are known, by writing out a list of the acidulous radicals which may be, or, if more convenient, of those which cannot be, present. To this end consult the following table (p. 368) of the solubility of salts in water. Look for the name of the metal of the salt in the vertical column; the letters S and I indicate which salts are soluble and which insoluble in water, an asterisk attached to the S meaning that the salt is slightly soluble. The acidulous part of the name is given in the top line of the table. All the names are in alphabetical order, for facility of reference.

Some of the salts marked as insoluble in water are soluble in aqueous solutions of soluble salts, a few forming soluble double salts. To characterize salts as soluble, slightly soluble, or insoluble only roughly indicates their relation to water: on the one hand, very few salts are absolutely insoluble in water; on the other, there is a limit to the solubility of every salt.

If only one, two, or perhaps three given acidulous radicals can be in the liquid, test directly for it or them according to the reactions given in the previous pages. If several may be present, pour small portions of the solution, rendered neutral if necessary by ammonia, into five test-tubes, and add respectively sulphuric acid, barium nitrate or chloride, calcium chloride, silver nitrate, and ferric chloride; then consult the table on p. 369 in order to correctly interpret the effects these reagents may have produced.

REMARKS ON THE TABLE, p. 369.

The first point of value to be noticed in connection with this table is one of a negative character—namely, if either of the reagents gives no reaction, it is self-evident that the salts which it decomposes with production of a precipitate must be absent. Then, again, if

the action of one of the reagents indicates the absence of certain acidulous radicals, those radicals cannot be amongst those precipitated by the other reagents; thus, if the action of sulphuric acid points to the absence of sulphides, sulphites, carbonates, cyanides, and acetates, these salts may be struck out of the other lists, and the examination of subsequent precipitates be so far simplified. Or if the barium precipitate is soluble in hydrochloric acid and the calcium precipitate in acetic acid, neither sulphates nor oxalates can be present. Observing these and other points of difference, which will be seen on careful and thoughtful reflection, and remembering the facts suggested by a knowledge of what basylous radicals are present, one acidulous radical after the other may be struck off as absent or present, leaving only one or two as the objects of special experiment. Among the chief difficulties to be encountered will be the separation from each other of chlorides, bromides, iodides, and cyanides, or of tartrates from citrates, and confirmatory tests of the presence of certain compounds. These may all be surmounted on referring back to the reactions of the various radicals, as described under their hydrogen salts, the acids.

In rendering a solution neutral for the application of the various group-tests, the necessary employment of any large amount of acid or of alkali must be noted, the presence of actual alkalies (that is, *hydrates*) or of acids themselves—*free acids*, so called—respectively being thereby indicated. Free acids also betray themselves by the *abundant* effervescence which results on the addition of a carbonate.

Sulphuric acid, the first group-reagent, may itself yield, especially when heated with some solid substances, sulphurous acid or hydrosulphuric acid (see p. 275); hence the production of the latter acids from a *diluted* solution only is evidence of the presence of a sulphide or sulphite.

In the precipitate produced by barium chloride, the second group-reagent, the oxalic radical may be specially sought by the test described in the "note" on p. 275.

Calcium chloride does not precipitate citrates readily or completely in the cold; therefore the mixture should be filtered and the filtrate boiled; calcium citrate then falls. Calcium tartrate is soluble in solution of ammonium chloride when quite freshly precipitated, but not after it has become crystalline. From their solution in ammonium chloride, calcium tartrate is mostly precipitated by ammonia, and citrate on boiling.

The rarer acidulous radicals will very seldom be met with. *Benzoates*, *hippurates* (which give benzoic acid), *hypochlorites*, *hyposulphites*, *nitrites*, and *valerianates* show themselves under the sulphuric treatment. *Ferrocyanides*, *ferricyanides*, *meconates*, *succinates*, *sulphocyanates*, *tannates*, and *gallates* appear among the salts whose presence is indicated by ferric chloride; *formates*, *hypophosphites*, *malates*, and others by silver nitrate. *Urates* char when heated, giving an odor resembling that of burnt feathers.

In actual practice the analyst nearly always has some clue to the nature of rarer substances placed in his hands.

TABLE OF THE SOLUBILITY OR INSOLUBILITY OF SALTS IN WATER.

	Acetate.	Arseniate.	Arsenite.	Carbonate.	Chloride.	Citrate.	Chromate.	Cyanide.	Hydrate.	Iodide.	Nitrate.	Oxalate.	Oxide.	Phosphate.	Sulphate.	Sulphide.	Sulphite.	Tartrate.
Aluminium.....	S	I	I	I	S	S	S	I	I	S	S	I	I	I	S	I	I	S
Ammonium.....	S	I	I	I	S	S	S	I	I	S	S	I	I	I	S	I	I	S
Antimony.....	S	I	I	I	S	S	S	I	I	S	S	I	I	I	S	I	I	S
Barium.....	S	I	I	I	S	S	S	I	I	S	S	I	I	I	S	I	I	S
Bismuth.....	S	I	I	I	S	S	S	I	I	S	S	I	I	I	S	I	I	S
Cadmium.....	S	I	I	I	S	S	S	I	I	S	S	I	I	I	S	I	I	S
Calcium.....	S	I	I	I	S	S	S	I	I	S	S	I	I	I	S	I	I	S
Chromium.....	S	I	I	I	S	S	S	I	I	S	S	I	I	I	S	I	I	S
Cobalt.....	S	I	I	I	S	S	S	I	I	S	S	I	I	I	S	I	I	S
Copper.....	S	I	I	I	S	S	S	I	I	S	S	I	I	I	S	I	I	S
Ferric.....	S	I	I	I	S	S	S	I	I	S	S	I	I	I	S	I	I	S
Ferrous.....	S	I	I	I	S	S	S	I	I	S	S	I	I	I	S	I	I	S
Gold.....	S	I	I	I	S	S	S	I	I	S	S	I	I	I	S	I	I	S
Lead.....	S	I	I	I	S	S	S	I	I	S	S	I	I	I	S	I	I	S
Magnesium.....	S	I	I	I	S	S	S	I	I	S	S	I	I	I	S	I	I	S
Manganese.....	S	I	I	I	S	S	S	I	I	S	S	I	I	I	S	I	I	S
Mercuric.....	S	I	I	I	S	S	S	I	I	S	S	I	I	I	S	I	I	S
Mercurous.....	S	I	I	I	S	S	S	I	I	S	S	I	I	I	S	I	I	S
Nickel.....	S	I	I	I	S	S	S	I	I	S	S	I	I	I	S	I	I	S
Platinum.....	S	I	I	I	S	S	S	I	I	S	S	I	I	I	S	I	I	S
Potassium.....	S	I	I	I	S	S	S	I	I	S	S	I	I	I	S	I	I	S
Silver.....	S	I	I	I	S	S	S	I	I	S	S	I	I	I	S	I	I	S
Sodium.....	S	I	I	I	S	S	S	I	I	S	S	I	I	I	S	I	I	S
Stannic.....	S	I	I	I	S	S	S	I	I	S	S	I	I	I	S	I	I	S
Stannous.....	S	I	I	I	S	S	S	I	I	S	S	I	I	I	S	I	I	S
Strontium.....	S	I	I	I	S	S	S	I	I	S	S	I	I	I	S	I	I	S
Zinc.....	S	I	I	I	S	S	S	I	I	S	S	I	I	I	S	I	I	S

* The asterisk indicates *slight* solubility.

TABLE TO AID IN THE DETECTION OF CHLORIDES, BROMIDES, IODIDES, CYANIDES, NITRATES, CHLORATES, BORATES, ACETATES, SULPHIDES, SULPHITES, SULPHATES, CARBONATES, OXALATES, TARTRATES, PHOSPHATES, AND CITRATES, IN A NEUTRAL AQUEOUS SOLUTION.

(For remarks concerning this Table see page 367.)

Sulphuric Acid decomposes	Barium Chloride precipitates	Calcium Chloride precipitates	Silver Nitrate precipitates	Ferric Chloride precipitates	Not precipitated.
Sulphides. Sulphites. Carbonates, with effervescence—hydrosulphuric and sulphurous acid gases, known by their smell and carbonic acid gas, which has no special odor, being evolved. Cyanides, with production of the odor of hydrocyanic acid. Acetates, with production of the odor of acetic acid when the solution is warmed. (For other salts see p. 367.)	Borates. Sulphites. Sulphates. Carbonates. Oxalates. Tartrates. Phosphates. Citrates. Of these white barium precipitates, the sulphate is the only one insoluble in hydrochloric acid; the tartrate and citrate char when heated on platinum foil; the sulphite and carbonate are decomposed with effervescence by acids.	Borates. Oxalates. Sulphites. Sulphates. Carbonates. Citrates. Of these white calcium precipitates, the sulphate only is sol. in much water; the borate, carbonate, and citrate are sol. in solution of ammonium chloride: all are sol. in acetic acid except oxalate and some tartrate and sulphate; all are sol. in hydrochloric acid, much sulphate excepted; the dry tartrate and citrate char when heated; the sulphite and carbonate effervesce with acids.	Chlorides, white. Bromides, white. Iodides, yellow. Cyanides, white. Borates, white. Sulphides, black. Sulphites, white. Carbonates, white. Oxalates, white. Tartrates, white. Phosphates, yellow. Citrates, white. Of these silver precipitates, the chloride, bromide, iodide, cyanide, and sulphide are insoluble in dilute nitric acid; the rest soluble.	Borates, yellowish. Sulphides, black. Carbonates, reddish. Oxalates, yellow. Phosphates, yellow. Gives red color with acetates, if neutral. (For other salts see p. 367.)	Nitrates. Chlorates. Apply special tests.

Note.—The student should practise the examination of aqueous solutions of salts until able to detect acidulous radicals with facility and precision. For this purpose he may finish the analyses of salts or solutions already examined for common and rarer metals, or have aqueous solutions of salts, or the salts themselves, specially prepared for present use, the metals of the salts being stated. He will then be in a position to effectively study the analysis of salts which may or may not be soluble in water, examining them for both basyious and acidulous radicals, (see next page).

If chromium and arsenum have been detected among the basylous radicals, those elements may be present in the form of *chromates*, *arsenates*, and *arsenites*, yielding with barium chloride yellow barium chromate and white barium arsenate and arsenite, and with silver nitrate red silver chromate, brown silver arsenate, and yellow silver arsenite.

QUESTIONS AND EXERCISES.

In analyzing an aqueous solution of salts, for which radicals would you first search, the basylous or the acidulous? and why?—In an aqueous solution there have been found magnesium (Mg) and potassium (K), with the sulphuric radical (SO_4) and iodine (I); state the nature of the salts which were originally dissolved in the water, and mention the principles which guide you to the conclusions.—Give a sketch of the method by which to analyze a neutral or only faintly acid aqueous liquid for the acidulous radicals of salts. In what stage of the process would the following salts be detected? *a.* Carbonates and sulphates. *b.* Oxalates. *c.* Tartrates and nitrates. *d.* Acetates and sulphites. *e.* Bromides and cyanides. *f.* Borates. *g.* Iodides and phosphates. *h.* Chlorates, oxalates, and acetates. *i.* Chlorides and iodides. *j.* Sulphites. *k.* Sulphides, carbonates, and nitrates. *l.* Citrates and sulphates.—Silver nitrate gives no precipitate in an aqueous solution; what acidulous radicals may be present?—Barium chloride gives no precipitate in neutral solution, but silver nitrate a white; what acidulous radicals are indicated?—Ferric chloride produces a deep-red color in a solution, calcium chloride yielding no precipitate; what salts may be present? and how may they be distinguished from each other?—Ferric chloride gives a black precipitate in a solution in which sulphuric acid develops no odor; to what is the effect due?

ANALYSIS OF SALTS.

SINGLE OR MIXED, SOLUBLE OR INSOLUBLE.

Thus far, all material substances, especially those of pharmaceutical interest, have been regarded as being definite compounds, and as having certain well-defined parts, termed, for convenience, basylous and acidulous respectively; moreover, attention has been designedly restricted to those definite compounds which are soluble in water. But there are many substances having no definite or known composition, and of those having definite composition there are many having no definite or ascertained parts. Again, of those having definite composition, and whose constitution admits of the entertainment of theory, there are many insoluble in water.

Chemical substances of whose composition or constitution little or nothing is at present known are chiefly of animal and vegetable origin, and figure in tables of analyses under the convenient collective title of "extractive matter;" they are not of immediate importance, and may be omitted from consideration.

Of substances which are definite in composition, but whose parts or radicals, if they have any, are unknown or imperfectly known,

there are only a few (such as the alkaloids, amylaceous and saccharine matters, the glucosides, and the albumenoid, resinoid, and colorific substances) which have any considerable amount of medical or pharmaceutical interest; these will be noticed subsequently.

Definite compounds most frequently present themselves; and of these by far the larger proportion (namely, the salts soluble in water) have already been fully studied. There remain, however, many salts which are insoluble in water, but which must be brought into a state of solution before they can be effectively examined. The next subject of laboratory work is, therefore, the analysis of substances which may or may not be soluble in water. This will involve no other analytical schemes than those which have been given, and will in only one or two cases increase the difficulty of the analysis of a precipitate produced by a group-reagent, but will give roundness, completeness, and a practical bearing to the reader's analytical knowledge. Such a procedure will at the same time bring into notice the methods by which substances insoluble in water are manipulated for pharmaceutical purposes or made available for use as food by plants or as food and medicine by man and animals generally.

Preliminary Examination of Solid (chiefly Mineral) Salts.

Before attempting to dissolve a salt for analysis, its appearance and other physical properties should be noted, and the influence of heat and strong sulphuric acid be ascertained. If the operator knows how to interpret what is thus observed, and to what extent to place confidence in the observations, he may more certainly obtain a high degree of precision in analysis, and will always gain some valuable negative information. But if he has only slight experience of the appearance and general properties of bodies, or has the habit of turning what should be inferences from tentative processes into foregone conclusions, he should omit the preliminary examination altogether, or only follow it out under the guidance of a judicious tutor; for it is impracticable here to do more than hint at the results which may be obtained by such an examination, or to so adapt descriptions as to prevent a student allowing unnecessary weight to preconceived ideas.

Whatever be the course pursued, short memoranda describing results should invariably be entered in the note-book.

1. Examine the physical characters of the salt in various ways, but never, or only rarely and cautiously, by the palate, on account of the danger to be apprehended.

If the salt is white, colored substances cannot be present; if colored the tint may indicate the nature of the substance or of one of its constituents, supposing that the learner is already acquainted with the colors of salts. Close observation, aided perhaps by a lens, may reveal the occurrence in a pulverulent mixture of small crystals or pieces of a single substance; these should be picked out by a needle and examined separately. In a powder or roughly-

divided mixture of substances the process of *sifting* (through such sieves as muslin of different degrees of fineness) often mechanically separates substances, and thus greatly facilitates analysis. The body may present an undoubted metallic appearance, in which case only the metals existing under ordinary atmospheric conditions need be sought. Peculiarity in smell reveals the presence of ammonia, hydrocyanic acid, hydrosulphuric acid, etc. Between the fingers a substance is perhaps hard, soft, or gritty; consequent inferences follow. Or the matter may be heavy, like the salts of barium or lead; or light, like magnesium carbonate and hydrate; or may be one of the pharmaceutically well-known class of "scale" preparations.

2. Place a grain or two of the salt in a small dry test-tube or in a piece of ordinary tubing closed at one end, and heat it, at first gently, then more strongly, and finally, if necessary, by the blowpipe.

Gases or vapors of characteristic appearance or odor may be evolved, such as iodine, nitrous fumes, sulphurous, hydrocyanic, or ammoniacal gases. Much steam given off by a dry substance indicates either hydrates or salts containing water of crystallization. (A small quantity of interstitial moisture often causes heated crystalline substances to *decrepitate* (from *decrepo*, I crackle); that is, break up with slight explosive violence, owing to the expansive force of the steam suddenly generated.) A sublimate may result, due to salts of ammonium, mercury, or arsenum, to oxalic or benzoic acid, or to sulphur free or as a sulphide—a salt wholly volatile containing such substances only. The compound may blacken, pointing to the presence of organic matter, which, in common definite salts, will probably be in the form of acetates, tartrates, and citrates, or as common salts of the alkaloids morphine, quinine, strychnine, or as a starch, sugar, salicin, or in other definite or indefinite forms common in pharmacy, and for which tests will be given in subsequent pages. If no charring occurs, the important fact is established that no organic matter is present, except cyanides, formates, or oxalates, which do not char. The residue may change color from presence or development of zinc oxide, iron oxide, etc., or melt from the presence of a fusible salt and absence of any large proportion of infusible salt, or be unaltered, showing the absence of any large amount of such substances.

3. Place a grain or two of the salt in a test-tube, add a drop or two of strong sulphuric acid, *cautiously* smelling any gas that may be evolved; afterward slowly heat the mixture, noticing the effect, and stopping the experiment when any sulphuric fumes begin to escape.

Iodine, bromine, and nitrous or chlorinoid fumes will reveal themselves by their color, indicating the presence of iodides, bromides, iodates, bromates, nitrates, and chlorates. The evolution of a colorless gas fuming on coming into contact with air and having an

irritating odor points to chlorides, fluorides, or nitrates. Gaseous products having a greenish color and odor of chlorine indicate chlorates, hypochlorites, or chlorides mixed with other substances. Slight sharp explosions betoken chlorates. Evolution of colorless gas may proceed from cyanides, acetates, sulphides, sulphites, carbonates, or oxalates. Charring will be due to citrates, tartrates, or other organic matter. If neither of these effects is produced, most of the bodies are absent or only present in minute quantity. The substances apparently unaffected by the treatment are metallic oxides, borates, sulphates, and phosphates.

4. Exposure of the substance to the blowpipe flame—on platinum wire, with or without a bead of borax or of microcosmic salt* (sodium, ammonium, and hydrogen phosphate, $\text{NaNH}_4\text{HPO}_4$); on platinum-foil or in a porcelain crucible or on a crucible lid, with or without sodium carbonate; or on charcoal, alone or in conjunction with sodium carbonate, potassium cyanide, or cobalt nitrate—will sometimes yield important information, especially to one who has devoted much attention to reactions producible by the blowpipe flame. The medical or pharmaceutical student, however, will seldom have time to work out this subject to an extent sufficient to make it a trustworthy guide in analysis. (See Plattner and Muspratt, "On the Use of the Blowpipe," and a chapter in Galloway's "Manual of Qualitative Analysis.")

Methods of Dissolving and Analyzing Single or Mixed Solid Substances.

Having submitted the substance to preliminary examination, proceed to dissolve and analyze by the following methods. These operations consist in treating a well-powdered substance consecutively with cold or hot water, hydrochloric acid, nitric acid, nitro-hydrochloric acid, or fusion with alkaline carbonates and solution of the product in water and acid. Resulting liquids are analyzed in the manner already described or by slightly modified processes, as detailed in the following paragraphs.

Solution in Water.—Boil about a grain of the salt presented for analysis in about a third of a test-tubeful of water. If it dissolves, prepare a solution of about 20 or 30 grains in $\frac{1}{2}$ an ounce or more of water, and *proceed with the analysis in the usual way*, testing first for the basylous radical or radicals by the proper group-reagents (HCl , H_2S , NH_4HS , $(\text{NH}_4)_2\text{CO}_3$, $(\text{NH}_4)_2\text{HPO}_4$), pp. 224 or 258, and then for the acidulous rad-

*So called because formerly obtained from the urine of man, who was called the *microcosmos*, or little world.

ical or radicals, directly or by aid of the prescribed reagents (H_2SO_4 , BaCl_2 , CaCl_2 , AgNO_3 , Fe_2Cl_6), p. 369.

If the salt is not *wholly* dissolved by the water, ascertain whether or not any has entered into solution by filtering, if necessary, and evaporating a drop or two of the clear liquid to dryness on platinum-foil; the presence or absence of a residue gives the information sought. If anything is dissolved, prepare a sufficient quantity of solution for analysis and proceed as usual, reserving the insoluble portion of the mixture, after thoroughly exhausting with water, for subsequent treatment by acids.

Solution in Hydrochloric Acid.—If the salt is insoluble in water, digest about a grain of it (or of the insoluble portion of a mixed salt) in a few drops of hydrochloric acid, adding water, and boiling if necessary. If the salt wholly dissolves, prepare a sufficient quantity of the liquid, noticing whether or not any effervescence (due to the presence of sulphides, sulphites, carbonates, or cyanides) occurs, and proceed with the analysis as before, except that the first step, the addition of hydrochloric acid, may be omitted.

The analysis of this solution will in most respects be simpler than that of an aqueous solution, inasmuch as the majority of salts (all those soluble in water) will be absent. This acid solution will, in short, only contain chlorides produced by the action of the hydrochloric acid on sulphides, sulphites, carbonates, cyanides, oxides, and hydrates, and certain borates, oxalates, phosphates, tartrates, and citrates (possibly silicates and fluorides), which are insoluble in water, but soluble in acids without apparent decomposition. The first four—sulphides, sulphites, carbonates, and cyanides—will have revealed themselves by the occurrence of effervescence during solution, and the presence of oxides and hydrates may often be inferred by the absence of compatible acidulous radicals. The borates, oxalates, phosphates, tartrates, and citrates alluded to will be reprecipitated in the general analysis as soon as the acid of the solution is neutralized; that is, will come down in their original state when ammonia and ammonium sulphydrate are added in the usual course. Of these precipitates, only the calcium oxalate and the calcium and magnesium phosphates need occupy attention now; for barium oxalate and phosphate seldom or never occur, and the borates, tartrates, and citrates met with in medicine or general analysis are all soluble in water. These phosphates and oxalates, then, will be precipitated in the course of analysis along with iron, their presence not interfering with the detection of any other metal. If from the unusually light color of the ferric precipitate phosphates and oxalates are suspected, it is treated according to the following table (reference to which should be inserted in the table for metals, under Fe, p. 224, and in the long table opposite p. 258).

PRECIPITATE OF PHOSPHATES, OXALATES, FERRIC HYDRATE, ETC.

Dissolve in HCl, add citric acid, then NH_4HO , and filter; then follow the table below:

Filtrate Fe. Add HCl and K_4Fcy . Blue ppt.	Precipitate $\text{Ca}_3\text{2PO}_4$, CaC_2O_4 , $\text{Mg}_3\text{2PO}_4$. Boil in acetic acid and filter.		
	Insoluble CaC_2O_4 . White. (CaF_2 may occur here.)	Filtrate $\text{Ca}_3\text{2PO}_4$, $\text{Mg}_3\text{2PO}_4$. Add $\text{Am}_2\text{C}_2\text{O}_4$, stir, filter.	
		Precipitate white, indicating $\text{Ca}_3\text{2PO}_4$.	Filtrate, add AmHO . White ppt. MgNH_4PO_4 .

In analyzing phosphates and oxalates advantage is also frequently taken of the facts that the phosphoric radical is wholly removed from solution of phosphates in acid by the addition of an alkaline acetate, ferric chloride, and subsequent ebullition, as described under "Phosphoric Acid" (p. 331), and that dry oxalates are converted into carbonates by heat, as mentioned under "Oxalic Acid" (p. 317). (See also p. 331, Fourth Analytical Reaction.)

Certain arsenates and arsenites, insoluble in water but soluble in hydrochloric acid, may accompany the above phosphates and oxalates if from any cause hydrosulphuric acid gas has not been previously passed through the solution or passed for an insufficient length of time.

If the substance insoluble in water does not *wholly* dissolve in hydrochloric acid, ascertain if any has entered into solution by filtering, if necessary, and evaporating a drop of the clear liquid to dryness on platinum-foil; the presence or absence of a residue gives the information sought. If anything is dissolved, prepare a sufficient quantity of solution for analysis, and proceed as usual, reserving the insoluble portion of the mixture, after thoroughly exhausting with hydrochloric acid and well washing with water, for the following treatment by nitric acid:

Solution in Nitric Acid.—If the salt is insoluble in water and hydrochloric acid, boil it (or that part of it which is insoluble

* Most oxalates, after being heated, effervesce on the addition of acid; fluorides may be detected by the "etching" test.

in those menstrua) in a few drops of nitric acid. If it wholly dissolves, remove excess of acid by evaporation, dilute with water, and proceed with the analysis.

This nitric solution can contain only a few substances, for nearly all salts soluble in nitric acid are also soluble in hydrochloric acid, and therefore will have been removed previously. Some of the metals, however (Ag, Cu, Hg, Pb, Bi), as well as amalgams and alloys, unaffected or scarcely affected by hydrochloric acid, are readily attacked and dissolved by nitric acid. Many of the sulphides, also, insoluble in hydrochloric acid, are dissolved by nitric acid, usually with separation of sulphur. Calomel is converted, by long boiling with nitric acid, into mercuric chloride and nitrate. The nitrates here produced are soluble in water.

This nitric solution, as well as the hydrochloric and aqueous solutions, should be examined separately. Apparently, time would be saved by mixing the three solutions together and making one analysis. But the object of the analyst is to separate every radical from every other; and when this has been partially accomplished by solvents, it would be unwise to again mix and separate a second time. Moreover, solvents often do what the chemical reagents cannot—namely, separate *salts* from each other. This is important, inasmuch as the end to be attained in analysis is not only an enumeration of the radicals present, but a statement of the actual condition in which they are present: the analyst must, if possible, state of what salts a given mixture was originally formed—how the basylous and acidulous radicals were originally distributed. In attempting this much must be left to theoretical considerations, but a process by which the salts themselves are separated is of trustworthy practical assistance; hence the chief advantage of analyzing separately the solutions resulting from the action of water and acids on a solid substance.

Solution in Nitro-hydrochloric Acid.—If the salt or any part of a mixture of salts is insoluble in water, hydrochloric acid, and nitric acid, digest it in nitro-hydrochloric acid, warming, or even boiling gently if necessary; evaporate to remove excess of acid, dilute, and proceed as before.

Mercury sulphide and substances only slowly attacked by hydrochloric or nitric acid—as, for example, calomel and ignited ferric oxide—are sufficiently altered by the free chlorine of aqua regia to become soluble.

Analysis of Insoluble Substances.

If the substance is insoluble in water and acids, it is one or more of the following substances: Sand and certain silicates, such as pipeclay and other clays; fluor-spar; cryolite ($3\text{NaF} \cdot \text{AlF}_3$); barium, strontium, and possibly calcium sulphates; tin-stone; antimonie oxide; glass; felspar (double silicate of

aluminium and other metals); silver chloride, bromide, or iodide; lead sulphate. It may also be or contain carbon or carbonaceous matter, in which case it is black and combustible, burning entirely or partially away when heated in the air, or be or contain sulphur, in which case sulphurous gas is evolved, detected by its odor when the substance is heated in air. A drop of solution of ammonium sulphhydrate, added to a little of the powder, will at once indicate the presence or absence of salts of such metals as lead and silver. For the other substances proceed according to the following (Bloxam's) method:

4 or 5 grains of the dry substance are intimately mixed with twice the quantity of dried sodium carbonate, and this mixture well rubbed in a mortar with five times its weight of *deflagrating flux* (1 of finely-powdered charcoal to 6 of nitre). The resulting powder is placed in a thin porcelain dish or crucible or clean iron tray, and a lighted match applied to the centre of the heap. Deflagration ensues and decomposition of the various substances occurs, the acidulous radicals going to the alkali metals to form salts soluble in water, the basylous radicals being simultaneously converted into carbonates or oxides. The mass is boiled in water for a few minutes, the mixture filtered, and the residue well washed. The filtrate may then be examined for acidulous radicals and aluminium, and the residue be dissolved in diluted hydrochloric acid and analyzed by the ordinary method.

The only substance which resists this treatment is chrome iron ore.

To detect alkali in felspar, glass, or cryolite Bloxam recommends deflagration of the powdered mineral with 1 part of sulphur and 6 of barium nitrate. The mass is boiled in water, the mixture filtered, ammonium hydrate and carbonate added to remove barium, the mixture again filtered, and the filtrate evaporated and examined for alkalies by the usual process.

Hydrates and Oxides,

If no acidulous radical can be detected in a substance under analytical examination, or if the amount found is obviously insufficient to saturate the quantity of basylous radical present, the occurrence of oxides or hydrates, or both, may be suspected. Confirmation of their presence will be found in the general rather than in any special behavior of the substances. Some hydrates yield water when heated in a dry test-tube held nearly horizontally in a flame, so that moisture may condense on the cool part of the tube. Some oxides yield oxygen, detected by heating in a test-tube and inserting the incandescent end of a strip of wood. Soluble hydrates cause abundant evolution of ammonia gas when heated with solution of ammonium chloride. Soluble hydrates also give character-

istic precipitates with the various metallic solutions. Hydrates and oxides insoluble in water not only neutralize much nitric acid or acetic acid, but are thereby converted into salts soluble in water. Most oxides and hydrates have a characteristic appearance. In short, some one or more properties of an oxide or a hydrate will generally betray its presence to the student who not only has knowledge respecting chemical substances, but has cultivated the faculties of observation and perception.

Fractional Operations.

Not only is the common process of *sifting* (p. 372) through sieves of varying degrees of fineness a useful fractional operation or separatory adjunct in analytical as in other work, but also fractional *elutriation* (p. 134), fractional *solution* of a mixed mass by *lixivation* (p. 89) of the substance with successive small quantities of solvents, and fractional *precipitation* with filtration after each addition of successive small quantities of a precipitant. Fractional *distillation* (*vide* Index) is often very useful, fractional *sublimation* (p. 93) and fractional *crystallization* (p. 78) occasionally, fractional *fusion* less often.

QUESTIONS AND EXERCISES.

Describe the preliminary treatment to which a salt may be subjected prior to systematic analysis.—Mention substances which might be recognized by smell.—Which classes of salts are heavy, and which light?—Name some bodies detectable by their color.—What inference may be drawn from the appearance of steam when dry substances are heated?—Why do certain crystals decrepitate?—If a powder sublimes on being heated, to what classes of compounds may it belong?—When heat causes charring, what conclusion is drawn?—No change occurring by heat, which substances cannot be present?—Give examples of salts which are identified by their reaction with strong sulphuric acid, and by their comportment in the blowpipe flame, with or without borax or microcosmic salt.—What are the solvents usually employed in endeavoring to obtain a substance in a state of solution? and what is the order of their application?—Name a few salts which may be present in an aqueous solution.—Mention some common compounds insoluble in water, but soluble in hydrochloric acid.—What substances are only attacked by nitric acid or nitro-hydrochloric acid?—At what stage of analysis do arsenites and arsenates show themselves?—Sketch out a method for the complete analysis of a liquid suspected to be an aqueous solution of neutral salts.—How can earthy phosphates and oxalates with ferric oxide be separated from each other?—How would you proceed to analyze an alloy?—By what process may substances insoluble in water or acids be analyzed?—How would you qualitatively analyze glass?

RECAPITULATORY AND OTHER NOTES ON THE CONSTITUTION OF THE DEFINITE CHEMICAL COMPOUNDS COMMONLY TERMED SALTS.

The molecules of a salt (p. 37) contain *radicals* (p. 67) which may be either elementary or compound (p. 67.)

Each radical has a definite exchangeable value (p. 123).

The definite exchangeable values of radicals differ in different series of radicals (pp. 123 to 125.)

In one and the same molecule of a salt two or more different atoms of the same element may possess the two distinct functions of being—(a) a single definite distinct radical, and (b) one member of a group of atoms which together form a single definite distinct radical (pp. 124 and 264.)

The relation to each other either of the elementary or the compound radicals in *organic* substances or salts is apparently far more complex than the relation to each other of the elementary or compound radicals in inorganic or mineral salts, as will be apparent when the section on Organic Chemistry has been studied.

The properties of salts are regarded as depending on (a) the nature, (b) the number, and (c) the position in relation to each other, of the elementary and compound radicals in a molecule, as will be apparent when the subject of isomerism has been studied.

Dumas, afterward Laurent, and then Gerhardt, attempted the classification of salts under such *types* as the following:



The hydrogen type.



The water type.



The ammonia type.

Other chemists have extended the number of such types of salts. Further, by writing the typical formulæ in the above and other manners a mode of indicating the facts assumed to be dependent on the position of the atoms in a molecule has been sought to be obtained. Finally, the natural development of this train of thought and of practice has produced the *graphic formulæ* of Kekulé, Frankland, and others, and has given rise to the doctrine of *valency*.

Caution.—The conjectural or theoretic character of our ideas respecting masses of matter being formed of molecules and molecules of atoms, and that molecules contain radicals consisting of one or more atoms, must never be lost sight of, highly valuable and practically useful though the hypotheses be (see pages 36, 51–57, 139, 264, 286, 299, 387.)

Berthollet's Laws.—"When we cause two salts to react by means of a solvent, if, in the course of double decomposition, a new salt can be produced less soluble than those which we have mixed, this salt will be produced." "When we apply dry heat to two salts, if, by double decomposition, a new salt can be produced more volatile than the salts previously mixed, this salt will be produced."

Malaguti's Law.—When solutions of two different salts are mixed and metathesis occurs, and four salts result, the proportions of the salts to each other are dependent on the strength or intensity of force with which the respective basylous and acidulous radicals are united.

The state of equilibrium just mentioned may be permanent or temporary. The latter condition obtains when one of the salts which may possibly be produced is insoluble, for as soon as precipi-

tation occurs the equilibrium is upset, and is re-established only to be upset again, and so on, until from the four salts there result one in solution and one out of solution. This would seem to be the way in which the laws termed "Berthollet's" work.

The Periodic Law.—Observations by Newlands, elaborated by Mendelejeff and Lothar Meyer, point to a law thus expressed by the latter chemist: "If the elements are arranged in order of increasing atomic weights [by which the student will understand those actual combining weights conveniently termed atomic weights], the properties of these elements vary from member to member of the series, but return more or less nearly to the same value at certain fixed points in the series." For example:

Li Be B C N O F Na Mg Al Si P S Cl K Ca, etc.

The periodicity of properties here alluded to occurs at about every eighth member of the series, irresistibly suggesting the periodicity of the musical scale. Thus sodium becomes the octave to lithium, silicon to carbon, phosphorus to nitrogen, sulphur to oxygen, chlorine to fluorine, potassium to sodium, calcium to magnesium, etc. Each chemical note (element) is distinct from the other, yet there is this curious harmony between any one and the eighth on either side. There are gaps in some of the series suggesting elements yet to be discovered, and irregularities suggesting the desirability of reconsidering some of the present weights; and other difficulties occur. Clearly the properties of the elements are in some way dependent on their "atomic" weights: the properties would indeed seem to be mere functions of these weights; the properties would seem to be determined by the weights; at all events, there is some such relationship between the elements,—all of which facts and considerations, by the way, irresistibly suggest the actual existence of *atoms* having fixed weight, and, *pro tanto*, very strongly support Dalton's atomic theory. Evidently there is less of fundamental difference between the so-called elements than we assume when we regard them as *distinct elements*. They would seem to be not so distinct as we commonly imagine. Whence the observed relationship of lithium, sodium, and potassium; of nitrogen, phosphorus, arsenum, and antimony (pp. 166, 332); of oxygen, sulphur, selenium, and tellurium (p. 302); of carbon, silicon, and perhaps boron (p. 356); of chlorine, bromine, iodine (p. 278), and fluorine (p. 343)? Are the so-called elements one and the same matter, differing only in the weight of their ultimate particles? Is the difference only, say, rate of vibration? They have not *yet* been transmuted. The subject is not developed sufficiently to warrant further consideration in this manual for ordinary medical and pharmaceutical students. Other students will find it fully considered in more advanced works. Papers and general statements will also be found in most journals of pharmacy. (See *Pharmaceutical Journal*, 3d Ser., vol. xviii. p. 882.)

ADVICE TO STUDENTS

RESPECTING THE METHOD OF STUDYING THE FOLLOWING PAGES ON
ORGANIC CHEMISTRY.

Both medical and pharmaceutical students of organic chemistry may be divided into two classes—namely, *junior students*, or those who, in the first instance at all events, desire to obtain only a general acquaintance with the subject; and *senior students*, or those who, having some general information, desire more complete and thorough knowledge of this branch of the science. To the members of each of these classes who use this Manual some advice concerning the kind and extent of work they may hope to accomplish in this department of the science will perhaps be acceptable.

Junior Students.—The whole of the following section on organic chemistry should be read through carefully once or twice, with the object not so much of remembering all that is stated as of acquiring (*a*) a general view of the scope of the subject, (*b*) a clear notion of the modes of classifying organic substances, and (*c*) an intelligent perception of their broad relationships to one another. (*d*) Special attention should be given to the methods of preparing and testing the particular substances officially recognized in the British Pharmacopœia, the student of practical chemistry preparing actual specimens of most of these substances, as well as going through tests for them and testing for impurities in them. He should make small quantities of chloroform, iodoform, spirit of nitrous ether, acetic ether, and a volatile oil; should extract gum from a gum-resin; purify some benzene, test aloin, and examine methylated spirit for its methylic constituent; prepare some spirit of wine by fermentation, concentrating the product until it will burn; make ether; convert amylic alcohol into valerianic acid; test carbolic acid and glycerin; manufacture a little soap; extract mannite from manna; go through the analytical reactions of cane- and grape-sugar; obtain starch from wheat flour, maize flour, and a potato, and examine each product with the microscope; make dextrin, pyroxylin, and collo-dion; prepare and test aldehyde, and try the action of lime on chloral hydrate; prepare and test acetic, oxalic, and citric acids; emulsify sweet and bitter almonds: prepare elaterin and test jalap resin and salicin; extract morphine or quinine or both, and perform the tests for the chief alkaloids of opium, cinchona, and nux vomica; test albumen and pepsin. Having gone through these operations, he should again read through the whole section.

Senior Students, having done all that junior students are, in the previous paragraph, advised to do, should thoughtfully study every page, reading what one other author, at least, has to say on each subject. More especially they should actually prepare, or test, or otherwise *experiment* with, one or more typical members of most of the series or sub-series of organic bodies. For example, they should prepare the *hydrocarbon* methane (from sodium acetate), convert it into a *haloid derivative* (by one of the given methods), transform this into the *alcohol* (by the agency of silver oxide and water), and this again

into the *acid* (by oxidation). The preparation of acetylene and ethylene and some of their derivatives should be tried; the differences between turpentine and petroleum spirit be experimentally proved; nitro-benzene be made, this be converted into aniline, and this again into "mauve;" aloin should be prepared; methylic alcohol be extracted from crude wood spirit and absolute alcohol be obtained from spirit of wine; alcohol and acetic acid be regenerated from the acetic ether previously prepared (by ebullition with a strong aqueous solution of potash); ethyl iodide or bromide and perhaps zinc-ethyl be made; glycol be prepared, and then be oxidized; glycerin be examined; starch be converted into dextrin and into sugar; malt extract be examined for diastase; trinitrocellulin be made; acetic aldehyde be fully examined and aldehyde-ammonia be prepared; lactic acid be made; benzoic and salicylic acids and aldehydes be obtained; natural urea be extracted and artificial urea be made; the glucosides be examined; and one or two artificial alkaloids be prepared; etc. Melting-points and boiling points of pure substances should be taken; and fractional distillation should be applied, either to acetic acid with a view to separate glacial acid on the one hand from water or weak aqueous acid on the other, to mixed alcohol and water with the object of attempting their re-separation as far as possible, or to some such mixture. Especially must the operations of quantitative analysis of organic compounds in due time be fully and thoroughly performed.

Other Students.—Students who have no occasion to apportion their periods of study in the manner contemplated in the previous paragraphs are recommended to go through the succeeding sections as they have gone through the foregoing—namely, page by page.

Note.—Students will find that in working at organic chemistry, so called, they are not departing from the method of study hitherto pursued. Hitherto they have concentrated attention on the chief elements, one at a time; they are now about to investigate the compounds of another of those elements; that is all. But it is an element having a greater range of combining powers than any yet examined. Organic chemistry is the chemistry of the element *carbon*.

ORGANIC CHEMISTRY ;*

OR

THE CHEMISTRY OF CARBON COMPOUNDS.

INTRODUCTION.

EXCEPT alcohol and a few acids, the large number of compounds which have hitherto engaged notice in this Manual have been of mineral origin. But the two other kingdoms of nature, the animal and vegetable, furnish still larger numbers of definite substances. These latter compounds, indeed, when discovered, were producible *only* by organized living structures ; hence were termed, more than two hundred years ago, *organic* (from ὄργανον, *organon*, an organ), and their study was termed *organic chemistry*. A very large number of organic compounds can now, however, be obtained artificially—without the aid of a living organism ; hence the particular distinction formerly drawn between organic and inorganic compounds, “organic” and “inorganic” chemistry, no longer fully obtains. Another definition, or additional definition, of organic chemistry or the chemistry of animate nature, the laws of which do not differ from those of inanimate nature, is now generally adopted—namely, *the chemistry of carbon compounds*. No doubt two or three kinds of compounds of carbon—carbonic acid gas and carbonates, for example—are met with in the mineral kingdom, and are therefore inorganic compounds, but they are met with in the organic kingdoms too, and therefore are organic compounds also.

Practically, all carbon compounds are organic compounds, and all of the so-called organic compounds are carbon compounds ; hence the old term, *organic chemistry*, no longer being etymologically and fully applicable, that of *the chemistry of the carbon compounds* seems natural as well as useful. Moreover, the carbon atoms possess in an altogether exceptional degree a property either not possessed or only to a much slighter extent possessed by those of other elements—namely, the property of combining with one another, and forming a sort of chain, to every link of which atoms of other elements can be attached, the result being obviously molecules of almost infinite variety and complexity—a fact which alone suffices to secure for carbon special and separate consideration by chemists. In short, the chemistry of the carbon compounds includes what was formerly as well as what is now known as organic chemistry ; or, in other words, the chemistry of organized or animate nature is included in

* Read the two previous pages of *Advice to Students*.

the chemistry of the carbon compounds. Of course, so old and historically interesting a term as *organic chemistry* will continue to be used; and there is no objection to such use, provided students remember that when the term is used as the equivalent of *the chemistry of the carbon compounds*, it is only conventionally and not etymologically accurate. Moreover, the chemistry of a carbon compound includes the chemistry of every element in that compound and the chemistry of the compound as a whole—facts which obtain even more prominence if the chemistry be spoken of by a general word, such as *organic*, rather than by the specific word *carbon*. Indeed, those organic compounds which contain nitrogen seem to be conditioned as much by their *nitrogen* as their *carbon*. So that, the word *organic* having now in chemistry lost its original specific signification, and having acquired the general signification described, it becomes, by its associations, perhaps the best word that can be chosen as the title of the great division of chemistry now under consideration.

COMPOSITION OF ORGANIC COMPOUNDS.

(a) *Qualitative Composition*.—The presence of carbon in a compound is at once shown if the compound blackens when a little is heated on a knife or platinum-foil in a flame. If the substance is heated in a dry, narrow test-tube and much moisture is condensed on the upper cool part of the tube, the presence of hydrogen and oxygen is reasonably inferred. Nitrogen may be detected by the odor of ammonia emitted on strongly heating the substance with a dry mixture of soda and lime, or it may be sought by carefully but strongly heating in a test-tube a small portion of the substance with a very small piece of sodium, and, after all action ceases, digesting the residue in water, filtering, and adding to the filtrate a ferrous salt, a ferric salt, and hydrochloric acid; a precipitate of prussian blue indicates nitrogen. Chlorine, bromine, iodine, sulphur, and phosphorus may be detected by heating the substance with nitric acid and silver nitrate in a very carefully and strongly sealed tube (in a fume-chamber, with such precautions that if the tube burst no harm to the operator shall ensue), and testing the product for chlorides, bromides, iodides, sulphates, and phosphates by methods already described.

(b) *Quantitative Composition*.—The qualitative composition of an organic substance being ascertained, the quantities of each element are then determined by methods which the student will practise when he is sufficiently advanced to work at the sections on quantitative analysis. The principles of the methods, are, however, simple, and may at once be described. For the quantitative estimation of *carbon* and *hydrogen* a carefully weighed portion of the substance is completely burned; the products, which are, of course, carbonic acid gas and water, are collected and accurately weighed. Of every 44 parts of the carbonic acid gas ($\text{CO}_2 = 44$), 12 will be carbon, and of every 18 parts of the water ($\text{H}_2\text{O} = 18$), 2 will be hydrogen; in other words, three-elevenths of the weight of carbonic acid

gas obtained will be the carbon of the original substance, while one-ninth of the water obtained will be the hydrogen of the original substance. If *nitrogen* be present, another weighed portion of the substance is so burned as to yield all its nitrogen as gas, which is carefully collected and measured, or it is heated with a mixture of soda and lime, when the nitrogen takes up hydrogen and becomes ammonia, which is collected and accurately estimated; of every 17 parts of ammonia ($\text{NH}_3 = 17$), 14 will be nitrogen. The amounts of *chlorine*, *sulphur*, etc., elements not often present, are obtained by subjecting carefully weighed portions of the original substance to the nitric treatment already alluded to, collecting and weighing the products, and calculating what proportions of the products are chlorine, sulphur, etc. The amount of *oxygen* is ascertained by difference; that is to say, the difference between the sum of the weights of carbon, hydrogen, nitrogen, etc. and the original weight of substance will be the weight of the oxygen in that original weight of substance.

For example, a fluid having well-marked definite properties, and known to contain only carbon, hydrogen, and oxygen, is so burned that 0.3 of a gramme* of it yields 0.5738 of a gramme of carbonic acid gas, and 0.3521 of water. As three-elevenths of the carbonic acid gas is carbon, and one-ninth of the water is hydrogen, it follows that the 0.3 of substance contains 0.1565 of carbon and 0.0391 of hydrogen; and the difference between these two figures and 0.3 being 0.1044, it follows that 0.1044 is the amount of oxygen in the 0.3 of original substance. For, $0.5738 \times 3 \div 11 = 0.1565$; and $0.3521 \div 9 = 0.0391$; $0.3 - (0.1565 + 0.0391) = 0.1044$.

(c) *Centesimal Composition*.—It is usual to make at least two such analyses of any organic compound; and, as different weights of the original substance will almost necessarily be subjected to combustion (for it is easier to counterpoise by weights a selected quantity than it is to counterpoise with the substance any selected weights), the results of the combustion can best be compared by converting the numbers first obtained into percentages; that is to say, by assuming in, for instance, the present case, that not 0.3 parts of substance were operated on, but 100 parts. This is one of the simplest of arithmetical operations. If 0.3 of substance yields 0.1565 of carbon, 100 of substance will yield 52.17 of carbon. If 0.3 of substance yields 0.0391 of hydrogen, 100 of substance will yield 13.03 of hydrogen. And if 0.3 of substance yields 0.1044 of oxygen, 100 of substance will yield 34.80 of oxygen.

(d) *Chemically Empirical Composition*.—But the chemist further desires to know, not so much what percentages or ordinary unit-weights of elements are contained in the compound, but what relative number of chemical unit-weights or atomic weights are present—how many parts of carbon each weighing twelve, how many parts of hydrogen each weighing one, how many parts of oxygen each

* If the reader is not already familiar with the metric system of weights and measures, he is referred to the section on that subject in the latter part of the Manual. (*Vide* Index, "Metric System.")

weighing sixteen, how many parts of nitrogen each weighing fourteen, etc. This too is one of the simplest of arithmetical operations. Divide the percentage of carbon by 12, of hydrogen by 1, of oxygen by 16, of nitrogen by 14, etc. Thus, in the present case, $52.17 \div 12 = 4.347$ atomic weights of carbon; $13.03 \div 1 = 13.03$ atomic weights of hydrogen; and $34.80 \div 16 = 2.171$ atomic weights of oxygen. Reducing these three fractional numbers of atomic weights to the lowest whole numbers (by assuming that the lowest of the three will represent 1 atomic weight—that is, by dividing the two higher of the three by the lowest), we find that the compound is composed of 2 atomic weights of carbon, 6 of hydrogen, and 1 of oxygen, thus: $4.347 \div 2.171 = 2$ of carbon; $13.03 \div 2.171 = 6$ of hydrogen; $2.171 \div 2.171 = 1$ of oxygen. The result is that of proportions of carbon each weighing 12, the substance contains 2; of proportions of hydrogen each weighing 1, the compound contains 6; and of proportions of oxygen each weighing 16, the compound contains 1. Finally, instead of the words "proportion of carbon weighing 12," the simple capital letter C may be used, which, as the reader now well knows, is not only the shorthand or symbol for the word carbon, but stands for 12 parts of carbon. Similarly, H may stand for 1 part of hydrogen, and O for 16 parts of oxygen; whence we arrive at C_2H_6O as the simplest chemically empirical expression, or *empirical formula*, of the substance under consideration.

(e) *Chemically Rational Composition*.—From the *empirical formula* of a substance we pass to a *two-volume formula*; that is to say, in accordance with the practice of chemists, the formula must, if possible, represent two volumes of the substance when in the state of vapor (see pp. 55 and 58). 2 parts of hydrogen gas, or 17 of ammonia gas, or 18 similar parts of water vapor, or 44 of carbonic acid gas, or $36\frac{1}{2}$ parts of hydrochloric acid gas, etc. occupy, if all are at the same temperature and under the same pressure, the same volume; whence we derive the formulæ H_2 , NH_3 , H_2O , HCl , and CO_2 as formulæ comparable with each other. If, now, the vessel which held these quantities of the respective substances be filled with the vapor of the substance supposed to be under examination at the same temperature and pressure, and the vessel and contents be weighed, the contents will be found to weigh 46 similar parts. C_2H_6O will be found, on adding up the atomic weights, to represent 46 parts by weight. Therefore C_2H_6O is the two-volume formula as well as the empirical formula, and thus the first step has been taken from the formula chiefly obtained by chemical art, an *empirical formula*, toward one largely obtained by, and that satisfies, the reason—a *rational formula* or *structural* or *constitutional formula*. Had the weight been found to be 92, the two-volume formula would have been $C_4H_{12}O_2$. The actual method of taking these weights of equal volumes of gases and vapors (specific gravity and vapor-density) will be described in the paragraphs on quantitative analysis.

(f) *Molecular Composition*.—Equal volumes of gases and vapors, being similarly affected by temperature and pressure, must be similarly constituted (Avogadro and Ampère's conclusion, pages 53 and 54). Whatever the number of molecules in such equal volumes

may be, it must be the same in each. Therefore the weights of equal bulks of gases and vapors under like conditions represent the relative weights of the respective molecules. Hence 2, 17, 18, 44, and 36.5 respectively represent the weight of one molecule of each of the substances hydrogen, ammonia gas, water vapor, carbonic acid gas, and hydrochloric acid gas, and 46 represents the molecular weight of the substance under consideration. C_2H_6O therefore represents the composition of a molecule of the substance. C_2H_6O is the *molecular formula* of the substance. (The substance is common alcohol.)

Acetic acid will serve as another simple illustration. Analysis and arithmetic yield the formula CH_3O . But metals only displace one-fourth of the hydrogen in any given weight of acetic acid. That fact cannot be shown by the formula CH_3O , but it can by the doubled formula $C_2H_4O_2$. This *chemical* evidence that the latter formula is, so far, correct is supported by the *physical* evidence of vapor-density as interpreted in the previous paragraph. (See also "Vapor-density" and "Raoult's Experiments" in Index.)

Check on Composition.—If the formula found is, even so far, the true formula of the substance, the centesimal composition found by experiment ought to be practically the same—the same within the limits of experimental error—as the centesimal composition obtained by calculation from the formula, thus:

	Calculated.	Found.
$C_2 = 24$	52.174	52.170
$H_6 = 6$	13.043	13.030
$O = 16$	34.783	34.800
46	100.000	100.000

From *composition* we now pass to *constitution*. What we know of the composition of a substance is reflected in its *empirical formula*. A step in advance is recorded in the *molecular formula*. What is afterward learnt about *constitution* is exhibited in the *structural formula*. If the letters in a formula—for instance, C_2H_6O —be regarded only as representing fixed weights, the formula expresses only facts; but we also regard the letters as representing *atoms* (Dalton's theory), and the whole formula as representing a *molecule* (Avogadro's theory), while in attempting structural formulæ we further depend on the idea of the valency of atoms. To this subject of *structure* or *constitution* we now pass.

CONSTITUTION OF ORGANIC COMPOUNDS.

In the molecule of an organic compound how are the atoms arranged? This is perhaps the greatest problem the chemist has to solve. Like the toy-puzzles of our youth, these chemical puzzles have to be attacked analytically and synthetically. How to sepa-


rate into its constituent bars of wood that apparently solid cube given to us by a friend in old days was not an uncongenial task, but, once accomplished, how to put these bars together again was a still more fascinating if more difficult labor. How to separate the groups of atoms or radicals in molecules of chemical substances, or at least how to find out the positions of those groups in a molecule, is a most difficult yet fascinating task for the skilled enthusiast in chemistry; and how to so marshal those groups (drawn perhaps from several different sources, and visible and tangible only in a state of combination and in mass) that he shall produce by art the compound originally only furnished by nature, is still more difficult, but also more fascinating—more fascinating, firstly, because it will furnish proof that his synthetical work was sound; secondly, because by artificially and perhaps cheaply producing a rare color, a rare perfume, a rare flavor, or a previously costly medicine, he may become a benefactor to his fellow-man; and, thirdly, because he may gain the honor of unveiling for all time one more of the truths of nature.

In practically attacking the problem of the constitution of a compound the chemist proceeds to note whether the substance is acid, alkaline, or neutral; to act on it with a base of known constitution if it is an acid, or with an acid of known constitution if it is a base, and to analyze the produced salts; to oxidize it; to deoxidize it; to chlorinize it; to remove or add hydroxyl (HO), carbonyl (CO), etc.; to substitute hydrogen by a compound radical, and *vice versâ*; to heat it; to electrolyze it; and, generally, to perform many such operations in the hope that the lines of chemical cleavage in the molecule will be detected, the essential groupings of atoms in the molecule be discovered, and even the positions of atoms or groups of atoms in relation to each other be reasonably inferred. Briefly, similarity in properties implies similarity in constitution or structure. *Per contra*, similarity in structure being reasonably implied, reference to properties shows whether or not the reason is on the right track toward truth in the matter of constitution or structure, the advance toward error being prevented and toward truth being maintained, whatever be the result of the reference, new truths not infrequently being unveiled. Thus, by the way in chemistry do fact and theory ever discharge their obligations to each other.

For example, urea, which was the first organic body produced artificially, was obtained by Wöhler in 1828 on heating solution of ammonium cyanate; H_4NCNO became $\text{OC}(\text{NH}_2)_2$, the mere change in the position of the constituent atoms—that is, in the *structure* of the molecules (indicated roughly, but to the best of our judgment, by the change in the relative position of the letters in the two formulæ just given)—accounting for the differences in the properties of the two substances, just as the differences in the relative position of a given number of stone blocks which at first were put together to form a bridge, but afterward were put together to form a house—that is, the differences in the *structure* of the edifices—account for the differences in their properties.


NOTATION OF ORGANIC COMPOUNDS.

In order that we may convey to one another our conclusions respecting the constitution of organic compounds, notation has to be carried somewhat farther in organic than has already been shown to be necessary in inorganic chemistry (see pp. 42 and 46). The relative position of atoms and groups of atoms in a molecule may be indicated by placing the symbolic letters above or beneath one another as well as on one line, and the quantivalence of atoms, as well as the directions in which we conclude they are joined in the molecule, may be indicated by lines ($—$ $=$ or $\equiv$) or dots ($\cdot$ $:$ $:$), either completely or only partially employed throughout the formula; each dot, and especially each line or "bond" or "link," representing such union between two neighboring atoms or radicals as would be represented by the extended arms of two persons shaking hands. For instance, the statement just made that ammonium cyanate (H_4NCNO) becomes urea [empirically $\text{CH}_4\text{N}_2\text{O}$, or rationally $\text{OC}(\text{NH}_2)_2$] might be represented by either of the following forms of equation:




Here, bars in the first equation and dots in the second show not only the quantivalence, but especially the distribution, of the chemical affinity expressed by the quantivalence of each atom. Thus the first four bars or dots not only indicate the univalence of each of the four hydrogen atoms on the one hand, and four-fifths of the quantivalence of the first nitrogen atom on the other hand, the next bar or dot showing the remaining fifth, but the five bars or dots also indicate that of the total affinity of the nitrogen atom four-fifths are engaged with a corresponding amount of attraction offered by four univalent hydrogen atoms, while the other fifth is engaged with one-fourth of the total attraction of the adjacent carbon atom. And so on with the quadrivalence of the carbon atom, the quinquivalence of the second nitrogen atom, and the bivalence of the oxygen atom.

But it is unnecessary, indeed undesirable, thus to indicate the quantivalence of *each* atom in a molecule, the closeness of union of groups of atoms (radicals) within a molecule being best indicated by putting the symbolic letters in a formula as close together as written characters or printer's types will allow; moreover, an atom, such as nitrogen, may often pass from one degree of activity to another during a reaction. The following, therefore, are better arrangements:



Indeed, after a time the chemical student will find that his own imagination will often best supply that which is intended to be indicated by the lines or dots in the formulæ of organic compounds, actual lines or dots only being employed where their use tends to promote clearness in a formula. For printed lines look like bars, and these, and even dots, are liable to suggest separation, whereas in a chemical formula they should suggest the union of the atoms and radicals in the molecule of which the formula is the crude picture. While suggesting links and bonds, however, and the affinity (quantivalence or valency) of the atoms, they must be regarded as indicating line of force rather than anything more substantial. Again, symbols and formulæ, written or printed, are necessarily exhibited on surfaces, whereas the conception of a molecule should be that of a sphere—that of grapes on a bunch of apples on a tree, rather than balls on a billiard-table; or, still better, that of moons round a planet and planets round a sun, all kept in their places by force rather than by anything material; our conceptions should be *stereochemical*, should be stereoscopic pictures rather than pictures of objects on a plane surface without perspective. (See also p. 393.)

Finally, bars, dots, or what not must only be placed in a formula where actual experiment warrants, unless the statement is distinctly made or understood that the suggested formula is only hypothetical. The use of such apparently highly complicated graphic *rational formulæ* or *constitutional* or *structural formulæ*—as, for example, the following formula for uric acid—is fully justified by a series of well-defined experiments:



Here not only is the univalence of each of the hydrogen atoms, the bivalence of oxygen atoms, the trivalent character of each of the nitrogen atoms, and the quadrivalent nature of each carbon atom shown, either directly by bars attached to the symbols or suggestively by the position of a symbol of recognized quantivalence next to another symbol of recognized quantivalence, but the positions which experiment warrants us in believing that the radicals occupy within the molecule are indicated in the formula by the position of the symbols for those radicals (HN, imidogen; CO, carbonyl) near central atoms of carbon.

To the student the great advantage of extended formulæ, whether ordinary or graphic, consists in the relationships which they clearly exhibit between compounds which otherwise are not readily shown to be related to one another. The *constitution* or *structure*, so far as can at present be inferred, of chemical compounds of interest to the medical or pharmaceutical student will be found to be given,

when desirable, in this Manual. Students who desire to pursue the subject more fully must seek other guides.

The structural formulæ characteristic of modern chemistry may be regarded as pictures of our idea of architecture in nature's molecules. The first sketches are seen in such formulæ as $\text{HC}_2\text{H}_3\text{O}_2$. Among the earlier mid-century artists were Gerhardt, Williamson, Frankland, and, especially from 1858 onward, Kekulé.


Caution.—Our conception of the structure or constitution of masses or *moles* of matter, or of particles or *molecules*, or of the *atoms* of which we conceive molecules are composed, or of the *valency* of those atoms, are nothing more than conceptions. All varieties of *chemical formulæ* are but concrete reflections or pictures of those conceptions. Why do we make those conceptions, and why do we thus picture those conceptions? We make them because, firstly, the enormous number of facts that chemistry unfolds creates, in the healthy human mind, a demand for classification; and, secondly, because the healthy human mind instinctively demands the reason *why* facts are as they are. And why do we adopt the existing chemical formulæ as concrete reflections of our abstract ideas? Because, firstly, intercommunication between minds can only be accomplished satisfactorily either by human utterances or by those written or printed equivalents, or signs, or symbols of speech termed letters, etc. (singly or combined to form words, etc.); and, secondly, because, as regards any one mind, the desire to avoid utter mental confusion demands the adoption of some method of concretely ticketing and labelling our ideas, so that we can set them aside or take them up at a moment's notice; and no better method than that offered by letters and similar signs has yet been devised, whether for the concrete expression of one's own thoughts for one's self or for written or printed intercommunication between mind and mind. In chemistry, however, we take care to use formulæ, letters, or symbols, (dots, bars, brackets, or what not) only to represent our conceptions respecting facts, except when we designedly, openly, and temporarily use them as a mode of giving rein to the imagination, hoping thereby to be led to inferences which experiments shall prove to be facts.

From the consideration of the *composition* and *constitution* of organic compounds we now pass to the subject of *classification*.

HYDROCARBONS: NEUTRAL OR NORMAL, AND BASYLOUS.

Neutral or Normal Hydrocarbons.—The simplest compounds of carbon are those with hydrogen; and as the atom of carbon is quadrivalent and the atom of hydrogen univalent, it follows that if a single atom of carbon be fully saturated with hydrogen, the formula of the resulting molecule must be CH_4 . But carbon is of all elements that which is peculiarly and specially liable to unite with itself (as

magnets attract each other), so far, at all events, as a portion of the attractive power of its atom is concerned, the other portions of its power attracting and being attracted by other atoms, the result being, possibly, molecules of great complexity. The following graphic formulæ will illustrate this point:




These formulæ represent well-known hydrocarbons, the first being common marsh-gas, or methane, one molecule of which is otherwise represented by the shorter formula, CH_4 ; the next represents ethane, C_2H_6 ; the third propane, C_3H_8 ; while C_4H_{10} is the formula of butane or tetrane. The first three members of the series are gases; those which immediately follow are liquids, C_5H_{12} , C_6H_{14} , etc.; while the highest members are solids, several of which form the mixture of hydrocarbons known as common paraffin; indeed, the whole series are distinguished as the *paraffin series* of hydrocarbons. It will be observed that the four units of affinity of the carbon atom are, in the molecule of each substance, fully saturated either by the affinities of adjacent hydrogen atoms or by that of another carbon atom. The substances are illustrations of *saturated hydrocarbons* (neutral or normal hydrocarbons). They differ in composition by CH_2 : add CH_2 to the first, and you obtain the second; add CH_2 to the second, and you obtain the third; and so on. The members of this series resemble each other in containing, to a given number of carbon atoms, twice that number, with two added, of hydrogen atoms. Representing "any number" by the letter n , the *general* formula for members of this neutral series of hydrocarbons will be $\text{C}_n\text{H}_{2n+2}$. Like neutral inorganic salts, their elements have saturated each other's affinities; hence the molecules refuse further to unite by direct or indirect addition with atoms having attractive powers. Potassium is powerfully basylous, chlorine powerfully acidulous—each has great affinity for the other; but the product, potassium chloride, KCl , is comparatively neutral or normal; saturated hydrocarbons are in the same case, for they do not unite with any other substances.


Basylous Hydrocarbons.—Many hydrocarbon groups, such as "methyl," CH_3 , and "ethyl," C_2H_5 , apparently have strong basylous affinities, because in compounds they appear to play the part which in inorganic compounds is performed by those basylous metals, etc. (K , Fe , NH_4 , *e. g.*) which are commonly called inorganic radicals. Indeed, such hydrocarbon groups are often termed *organic radicals*, and to hold the theory that they exist is convenient; but any attempt to isolate them results in the production of neutral hydrocarbons, C_2H_6 , C_4H_{10} , etc.

Some hydrocarbons, however, which can quite easily be isolated, are basylous, such as ethylene, C_2H_4 , and other bivalent radicals


having the general formula C_nH_{2n} ; and acetylene, C_2H_2 , and other quadrivalent radicals having the general formula C_nH_{2n-2} . Such radicals are sometimes termed *unsaturated hydrocarbons*:




their compounds with, for example, bromine, being thus formulated:



But the first is probably methane, in which one atom of hydrogen is substituted by one of bromine, other salts containing the supposed non-isolable radicals being normal hydrocarbons in which atoms of hydrogen are substituted by atoms of acidulous elements or acidulous radicals, the residual hydrocarbon being the so-called basylous radical. And as regards the basylous hydrocarbons which can be isolated, they too probably are neutral hydrocarbons in which the carbon atoms are united to the extent of half or even three-fourths of their affinities, thus:



Bring bromine into contact with these so-called free basylous radicals, and in the case of ethylene one pair of carbon "arms" may be considered to unclasp, each of the two free arms clasping a one-armed bromine atom; while, in the case of acetylene first one pair of arms unclasp and take in two bromine individuals, and then another pair unclasp and take in two more individuals of a bromine molecule.



Molecules.—Instead of the foregoing *plane-conceptions* of molecules, *stereo-conceptions* have been suggested (see p. 390). Thus a tetrahedron may represent a carbon atom, or the area of influence of the

(central) carbon atom, its corners or summits representing the foci of its affinities or valencies—foci at which atoms of hydrogen or of compound radicals will find place or in the vicinity of which they will oscillate. Two or more of such tetrahedra may be regarded as united by their corners, edges, or faces according to circumstances. This subject cannot be pursued here. Chemists seem to be gliding from statical notions of chemical structure to dynamical ideas, slowly discarding the principle of affinities between fixed particles for that of atoms conditioned by motion. (See the memoirs of Van't Hoff, Lebel, Mayer, Guye, and others.)

Series of Hydrocarbons.—Three distinct series of hydrocarbons have now incidentally been alluded to—namely, the paraffin series, C_nH_{2n+2} ; the olefine series, C_nH_{2n} ; and the acetylene series, C_nH_{2n-2} . Twelve or fourteen other series are known, as, the terpene series, C_nH_{2n-4} ; the benzene series, C_nH_{2n-6} ; the cinnamene series, C_nH_{2n-8} ; the anthracene series, C_nH_{2n-18} , etc. Each member of any such series obviously differs in composition from the preceding or succeeding member by CH_2 . Either series will therefore be an illustration of an *homologous* series (from *ὁμός*, *homos*, the same, and *λόγος* *logos*, proportion) of compounds. There will be similar *homology*, of course, between the members of the series of alcohols derived from these hydrocarbons, or between the haloid salts, the ethers, the aldehydes, the acids, etc. Homology is necessarily concurrent with step-by-step variation in the properties of members of a series.

Substitution.—The atoms of hydrogen in any member of either of the series of hydrocarbons may be substituted by radicals of all kinds—basylous and acidulous, elementary and compound; by chlorine or bromine, hydroxyl or sulphydroxyl, oxygen or sulphur, amidogen or imidogen, carboxyl, etc. Very large numbers of organic compounds have thus been obtained artificially; still larger numbers have been proved, analytically, to have distinct existence; while it is certain that still larger numbers exist of which we do not know the constitution and only partially know the composition.

Note.—The idea of substitution, in chemistry, involves or includes the conception of the unity of a molecule as opposed to the old-fashioned conception of duality; involves the unitary conception under which we picture a molecule of, say, anhydrous Epsom salt as $MgSO_4$ rather than MgO, SO_3 ; or, figuratively, involves that conception of oneness or wholeness in a building which allows of one kind of brick being substituted by another kind without change in the structure *quâ* structure. That the idea of substitution also involves, in chemistry, a somewhat unwieldy notation and an extremely unwieldy nomenclature appears at present to be inevitable.

Procedure as regards Further Study.

Several of these series of hydrocarbons and their substitutional derivatives will now be described, special notice being given to the compounds of medical and pharmaceutical interest. Some members of the paraffin, olefine, acetylene, terpene, benzene,

naphthalene, and anthracene series will be treated of, together with their haloid, nitrous, and acetic derivatives, etc; the alcohols or hydroxyl substitution-compounds will then be noticed as a class; and afterward the carbohydrates, amyloids, aldehydes, acids, glucosides, and alkaloids.

The series of chief interest to medical and pharmaceutical and—indeed, to all—students is the first, known by either of the four names *Paraffin*, *Fatty*, *Marsh-gas*, or *Methane* Series. The *Benzene* or *Aromatic* Series has great general chemical interest. The *Terpene* Series has considerable pharmaceutical interest.

A very large number of compounds of carbon will thus be brought under notice—far larger than that of any other element. The mere number, however, need not dismay the student. The relation of the derivatives of one hydrocarbon to that hydrocarbon will be found to obtain between the next set of derivatives studied with their hydrocarbon, and so on; hence as the student progresses he is soon looking for compounds which he already expects to exist, instead of finding his mind overburdened with what at first sight he might fear would be an intricate and endless subject.

The methods of examining morbid urine will afterward be experimentally considered. There will then remain to be studied by the medical and pharmaceutical pupil, but by aid of some other guide than the author's, certain galenical as distinguished from chemical substances, solid and liquid, which can only be fairly regarded from a pharmacist's rather than a chemist's point of view, and a still larger number, doubtless, not yet brought within the grasp of chemist or pharmacist, and of which, therefore, we must at present be content to remain in ignorance. An opportunity, however, will be afforded of noticing the effect of such indefinite organic matter as a vomit, or the contents of a stomach, in masking or preventing the reactions by which mineral and vegetable poisons are detected.


A section on quantitative analysis will complete the Manual.

QUESTIONS AND EXERCISES.

What do you understand by Organic Chemistry?—Give methods of ascertaining the presence of carbon, hydrogen, and nitrogen in organic compounds.—Give an outline of the methods by which the quantities of carbon, hydrogen, oxygen, and nitrogen are determined in organic compounds.—How would you convert centesimal into "atomic" composition?—Define *empirical*, *molecular*, and *rational* formulæ.—How is the constitution of an organic compound ascertained?—What do you understand by graphic chemical formulæ?—Define "stereo-chemical" formulæ.—Give graphic formulæ of two or three saturated hydrocarbons.—What do you mean by an organic radical?—Give illustrations.—Give the general formulæ of different series of hydrocarbons, with special illustrations.—Define *substitution* as understood in organic chemistry.





THE PARAFFIN SERIES OF HYDROCARBONS.

METHANE, *Marsh-gas*, *Light Carburetted Hydrogen*, *Methyl Hydride*, *Fire-damp*, CH_4 .—This gaseous hydrocarbon may be made from its elements by uniting the carbon with sulphur and the hydrogen with sulphur or oxygen, and passing these over red-hot copper. It occurs naturally in coal-mines and in the mud-volcanoes of the Crimea, is frequently associated with the crude petroleum that issues from the earth, and is constantly rising in bubbles to the surface of stagnant pools in marshy places. It is a non-luminous constituent of ordinary coal-gas. It is inodorous and colorless. It may be produced by acting on methyl iodide with zinc on which copper has been deposited, and in other theoretically interesting ways, but economically by heating a mixture of 2 parts of dry sodium acetate, 3 of lime, and 2 of caustic soda, or, better, potash.




Two Notes on the Notation of the Foregoing and Similar Formulæ, and on the Constitution of Salts.—(a) Soda, NaHO , contains bivalent oxygen, univalent sodium, and univalent hydrogen. The chemical valency of the oxygen atom is double that of either of the other atoms—a relationship perhaps better realized if the symbol for the oxygen be placed between those of hydrogen and sodium, NaOH , or HONa . So HOK , HOH , etc. The student must expect to find the symbols of a formula placed where apparently they will best reflect our knowledge of the structure of the molecule pictured. (b) Acetic acid, $\text{C}_2\text{H}_4\text{O}_2$, by action of chlorine (presented as PCl_3) loses hydroxyl, OH , and yields acetyl chloride, $\text{C}_2\text{H}_3\text{OCl}$. Hence acetic acid would seem in constitution to be acetyl hydrate, $\text{C}_2\text{H}_3\text{O}\cdot\text{OH}$, especially when we find that acetyl chloride by reaction with water, HOH , yields again acetic acid (and HCl). Sodium will only displace one atom of hydrogen from water, yielding HONa , and will only displace one atom of hydrogen from acetic acid, yielding sodium acetate, $\text{C}_2\text{H}_3\text{O}\cdot\text{ONa}$. Further, chlorine will not displace more than one portion or atom of hydroxyl, OH , from acetic acid. So that three atoms of the hydrogen in acetic acid apparently perform different functions to those of the fourth atom, and, apparently, the two atoms of oxygen perform different functions. Hence our necessity for separating in the formula the letters representing those atoms, $\text{C}_2\text{H}_3\text{O}\cdot\text{OH}$. Once more, acetates may be formed from two different methyl compounds—sodium acetate by the direct combination of sodium methide, CH_3Na , and carbonic acid gas, CO_2 , giving $\text{CH}_3\cdot\text{CO}\cdot\text{ONa}$; and ammonium acetate by the combination of methyl cyanide, CH_3CN , with water (2HOH), yielding $\text{CH}_3\cdot\text{CO}\cdot\text{ONH}_4$. From these and other facts and modes of reasoning arise our justification—from them, indeed, comes the necessity—for thus extending the formulæ for acetates. Less extended formulæ are of course correct, and even occasionally more useful: $\text{C}_2\text{H}_4\text{O}_2$, $\text{C}_2\text{H}_3\text{O}_2\text{H}$, $\text{C}_2\text{H}_3\text{O}\cdot\text{OH}$, $\text{CH}_3\cdot\text{CO}\cdot\text{OH}$ form an illustration of a set of formulæ for a substance, either member of which set may be used according to

circumstances. (See also pp. 286 and 299.) The following would be reasonable *graphic* formulæ, like those on p. 390 or 392:



ETHANE, C_2H_6 , *Dimethyl, Ethyl Hydride*.—This is one of the constituents of crude petroleum. It also results on heating ethyl iodide with granulated zinc or zinc covered with copper, and then adding water to the zinc iodide and zinc ethide first produced.





Ethane is sometimes regarded as dimethyl or methylmethane, CH_3CH_3 ; that is to say, as being derived from methane by the substitution of an atom of hydrogen in methane, CH_4 , by methyl, CH_3 : its properties, however, are not those of a radical. It is also considered to be ethyl hydride, $\text{C}_2\text{H}_5\text{H}$: its properties, however, are not those of such a substance. The other hydrocarbons of the paraffin series are also similarly regarded as containing radicals. Such views of constitution are useful as enabling composition to be remembered and relationships to be realized, especially if their hypothetical character be fully recognized; but these hydrocarbons are apparently single homogeneous substances, and, whatever other views of their constitution be held, this last should be dominant.

PROPANE, *Methyl Ethyl*, C_3H_8 .—This gas, like methane, occurs dissolved in the Pennsylvanian petroleum springs.

TETRANE OR BUTANE, C_4H_{10} .—Two varieties exist, *normal butane* or *diethyl*, $\text{C}_2\text{H}_5\cdot\text{C}_2\text{H}_5$, found in petroleum, and *isobutane* or *trimethyl methane*, $\text{CH}(\text{CH}_3)_3$, formed by artificial means.

Turning back to the highly extended formulæ for methane, ethane, propane, and butane given on p. 391, the reader will see why there should only be one ethane or propane, while two butanes (two methyl-propanes) are possible. We can but replace one of the atoms of hydrogen, H, in methane, CH_4 , to form ethane, $\text{CH}_3\cdot\text{CH}_3$, and it matters not which; hence only ethane (one ethane) can result. In ethane, $\text{CH}_3\cdot\text{CH}_3$, if an atom of hydrogen be displaced by methyl, CH_3 , it can but be a hydrogen atom of one of the two methyl groups ($\text{CH}_3\cdot\text{CH}_3$), and it matters not which, though two *di*-derivatives may exist. (See the respective ethylene and ethylidene chlorides, p. 411.) But in propane, $\text{CH}_3\cdot\text{CH}_2\cdot\text{CH}_3$, a CH_2 group exists, as well as CH_3 groups. Now, CH_2 is a different group to CH_3 ; hence if we

displace one of its two atoms of hydrogen (it matters not which) by methyl to get butane, we should expect to get a butane of different properties to the butane obtained by displacing one of the atoms of hydrogen in the methyl groups by methyl; and two butanes, and two only, do actually exist. Normal butane may be thus formulated, $\text{CH}_3 \cdot \text{CH}_2 \cdot \text{CH}_2 \cdot \text{CH}_3$, while isobutane would be either $\text{CH}_3 \cdot \text{CH}_3 \cdot \text{CH} \cdot \text{CH}_3$, or a practically identical formula, $\text{CH}_3 \cdot \text{CH} \cdot \text{CH}_3 \cdot \text{CH}_3$.



PENTANE, C_5H_{12} .—Three varieties are possible, and three only; three are known, and three only; the second, or *isoamylic hydride*, yielding the ordinary amylic alcohol and valerianic acid,

HEXANES, C_6H_{14} .—Five are possible, five are known.

HEPTANES, C_7H_{16} .—Nine are possible, five are known.

OCTANES, C_8H_{18} .—Eighteen isomers possible, three known.

NONANE, C_9H_{20} ; **DECANE**, $\text{C}_{10}\text{H}_{22}$; and every paraffin hydrocarbon up to $\text{C}_{24}\text{H}_{50}$, as well as some others and derivatives of far higher members of the paraffin series of hydrocarbons, are known.

Petroleum Spirit, Paraffin Oil, Paraffin.


Benzinum, U. S. P.; *Benzin* (*Petroleum Spirit*, B. P.) *Pentane*, C_5H_{12} , *Hexane*, C_6H_{14} , etc., known also as *Benzoline*, *Petroleum Benzin*, and *Petroleum Ether*, "is a colorless, very volatile, and highly inflammable liquid obtained from petroleum, and consisting of a mixture of the lower members of the paraffin or marsh-gas series of hydrocarbons. Boiling-point, 122° to 140° F. (50° to 60° C.). Specific gravity, about 0.670 to 0.700." (Benzine or benzol is quite a different fluid; *vide* Index.)

Paraffin Oil, the *Paraffinum Liquidum* of the German Pharmacopœia, is a mixture of the higher fluid members of the paraffin series of hydrocarbons, a clear oily liquid obtained from petroleum after distilling off the lower-boiling portions. Specific gravity, not below 0.840. Boiling-point, not below 360° C. (680° F.). Digested and agitated with warm sulphuric acid for a day or two, the oil is not colored and the acid only tinged brown; metallic sodium under similar conditions is not tarnished. *Petrolatum Liquidum*, U. S. P., is similar to the above, having a specific gravity between 0.875 and 0.945; is soluble in chloroform, ether, carbon bisulphide, benzol, benzin, and the fixed and volatile oils. Alcohol boiled with the oil should not become acid. *Soft Paraffin* (*Petrolatum Molle*, U. S. P. 1890, the *Petrolatum* of U. S. P. 1880), *Soft Petroleum Ointment*, officially termed *Paraffinum Molle* in England, *Unguentum Paraffini* in Germany, *Pétroléine* in France, and "known in commerce by various fanciful names," is a semi-solid mixture of paraffins, usually

obtained by purifying the less volatile portions of petroleum. It is "white or yellowish, translucent, soft, greasy, free from acidity, alkalinity, or any unpleasant odor or flavor even when warmed to 120° F. (48.9° C.). Specific gravity at the melting-point, from about 0.840 to 0.870. Melts at 95° to 105° F. (35° to 40.5° C.) or even somewhat higher; volatilizes without giving acrid vapors, and burns with a bright flame, leaving no residue; insoluble in water, slightly soluble in absolute alcohol, freely soluble in ether, chloroform, benzol, etc. It is not saponified by solutions of alkalies."—*B. P. Hard Paraffin* (*Petrolatum Spissum*, *Hard Petrolatum*, *Hard Petroleum Ointment*, U. S. P. 1890) (*Paraffin Durum*, B. P.), (*Petrolatum*, U. S. P., 1880), commonly termed *Paraffin Wax* or simply *Paraffin*, is "a mixture of several of the harder members of the paraffin series of hypocarbons; usually obtained by distillation from shale, separation of the liquid oils by refrigeration, and purification of the solid product. It is colorless, semi-transparent, crystalline, inodorous, and tasteless; slightly greasy to the touch. Specific gravity, 0.82 to 0.94. Insoluble in water, slightly soluble in absolute alcohol, freely soluble in ether. It melts at 110° to 145° F. (43.3° to 62.8° C.), and burns with a bright flame, leaving no residue."


Paraffin resists all ordinary reagents (hence the original name *Paraffin*, from *parum affinis*, without affinity), but may, by continued boiling with sulphuric acid and solution of potassium bichromate, be oxidized to *cerotic acid*, $C_{27}H_{54}O_2$, and by continued digestion with nitric and sulphuric acids yields acids of the acetic series and *paraffinic acid*, $C_{24}H_{48}O_2$ (Pouchet).

SUBSTITUTION-PRODUCTS OF METHANE.—The paraffins all form substitution-derivatives with the halogens, chlorine acting energetically, bromine less so, and iodine scarcely at all. In the preparation of chlorine and bromine substitution-products by acting on the hydrocarbons, the mono-derivatives are always mixed with the higher derivatives, even though the quantities are taken in relation to their combining proportions; thus, if methane and chlorine are mixed in the proportion of $CH_4 + Cl_2$, not only will *monochloromethane*, or *methyl chloride*, CH_3Cl , be formed, but *dichloromethane*, CH_2Cl_2 , and *trichloromethane*, $CHCl_3$, with free hydrogen. The best method of obtaining the mono-derivatives is to act on the alcohols by haloid acids or by phosphorus compounds:



Chloroform.


Trichloromethane, or *Chloroform*, $CHCl_3$, may be made by acting on methane with chlorine, as already indicated:




also as shown on p. 402, but on a larger scale by the official process, as follows:

Process.— $1\frac{1}{2}$ fluidounces of spirit and 24 of water are placed in a retort or flask of at least a quart capacity; 8 ounces of chlorinated lime and 4 of slaked lime are added, the vessel connected with a condenser, and the mixture heated until distillation commences, the source of heat then being withdrawn. The condensed liquid should fall into a small flask containing water, at the bottom of which about a drachm of chloroform will slowly collect.


Explanation of the Process.—Though there is some doubt as to the exact reaction, the following seems to be most probable. The calcium hypochlorite believed to be present in the chlorinated lime (see the remarks in connection with the latter, p. 114) readily yields up oxygen and chlorine to organic substances, the calcium being liberated as hydrate. The alcohol used in making chloroform is thus probably first converted into aldehyde:*




The action of chlorine on aldehyde then probably gives chloral (*chlor-aldehyde*):



The hydrochloric acid being at once neutralized by some of the liberated calcium hydrate to form calcium chloride and water, more freed calcium hydrate and chloral gives calcium formate and chloroform.





Or, irrespective of the chemistry of each step in the process, and regarding only the materials and the products, four molecules of alcohol and eight of calcium hypochlorite give two of chloroform, three of calcium formate, five of calcium chloride, and eight of water, thus:



The calcium hydrate placed in the generating vessel is not essential, but is useful in preventing secondary decompositions, the calcium hydrate obtainable from the reaction being insufficient for this purpose.

* The special formulæ for alcohol, aldehyde, and the formate used in the accompanying equations will be better understood when the constitution of alcohols and acids has been considered.

Chlorine converts chloroform into *tetrachloromethane* or *carbon tetrachloride*, CCl_4 , completing the chlorine substitution-products of methane.



Chloroform is purified by shaking it with water, and then with pure sulphuric acid (containing no trace of nitric acid), which chars and removes hydrocarbons, etc., but does not affect chloroform. It is freed from any trace of acid by agitation with lime, and from moisture by solid calcium chloride. It is finally rectified.

Properties.—The sp. gr. of pure chloroform is at least 1.500, perhaps higher. It is liable to slowly decompose when exposed to air and light: $4\text{CHCl}_3 + 3\text{O}_2 = 4\text{COCl}_2 + 2\text{H}_2\text{O} + 2\text{Cl}_2$. The resulting chlorine may be detected by zinc iodide and starch, and the carbon oxychloride by baryta-water: $2\text{COCl}_2 + 2\text{H}_2\text{O} = 2\text{CO}_2 + 4\text{HCl}$. To render it stable, a minute amount (one volume in one hundred or less) of absolute alcohol is necessary: hence the specific gravity of medicinal chloroform is about 1.497 (*Chloroformum*, U. S. P., the *Purified Chloroform* of the 1880 Pharmacopœia). It readily and entirely volatilizes at common temperatures, having, to the last drop, its pleasant characteristic odor. It has a sweetish taste, is limpid, colorless, soluble in alcohol (1 in 16 gives *Spiritus Chloroformi*, U. S. P.) and ether, and slightly in water. It may be so frozen at low temperatures that any impurities shall remain in the still fluid portion (Pictet). Boils at 142°F . It burns with a sluggish, green, smoky flame. It reduces Fehling's solution. It should be neutral to test-paper, indicating absence of acid; give no precipitate with solution of silver nitrate, indicating absence of ordinary chlorides; remain colorless when heated with potash, indicating absence of aldehyde; and give no more color than is producible by the absolute alcohol that is present to any sulphuric acid with which it may be shaken, even after the mixture has been set aside for half an hour, indicating absence of hydrocarbons, etc. Alcohol may be detected by the iodoform test (see Index), or by shaking with a little of the dye termed "Hofmann's violet," which gives the chloroform a purple tint if alcohol be present, but affords no color with pure chloroform. At the temperature of melting ice chloroform unites with water to form a crystalline compound, $\text{CHCl}_3 \cdot 18\text{H}_2\text{O}$.

Aqua Chloroformi, U. S. P., the official chloroform-water, is made by saturating a sufficient quantity of water with chloroform.

Iodoform.


Tri-iodomethane, or *Iodoform*, CHI_3 (*Iodoformum*, U. S. P.), analogous in constitution to chloroform, the iodine occupying

the place of the chlorine, is made by mixing together 1 part of alcohol, 2 parts of crystallized sodium carbonate, and 10 parts of water, the whole being heated to about 150° F., and 1 part of iodine gradually added in small portions. When the fluid becomes colorless the iodoform is allowed to settle. The iodoform is collected on a filter, washed thoroughly with water, and dried between filtering-paper. (This reaction forms a very delicate means of testing the presence of alcohol. *Vide* "Alcohol, Test for," in Index.)

Iodoform occurs as yellow, shining, six-sided scales. It is volatile at ordinary temperatures. Almost insoluble in water, soluble in alcohol or ether. Warmed with an alcoholic solution of potash, potassium formate and iodide are produced: $\text{CHI}_3 + 4\text{KOH} = \text{HCOOK} + 3\text{KI} + 2\text{H}_2\text{O}$; and the resulting fluid, heated with a little nitric acid, yields free iodine, recognized by its color or by giving a blue color with starch.

Chloroform, Iodoform, and Bromoform may also be obtained on passing a current of electricity through hot strong alcohol containing potassium chloride, iodide, or bromide, respectively, carbonic anhydride being simultaneously supplied; or, again, by the action of bleaching-powder, or a mixture of chlorinated soda with potassium bromide or iodide, on acetone. The latter, "*ketone chloroform*," is a commercial article.

SUBSTITUTION-PRODUCTS OF ETHANE.—Ethane, like methane, yields substitution-derivatives. *Monobromethane, ethyl bromide, ethylic bromide, or hydrobromic ether*, $\text{C}_2\text{H}_5\text{Br}$, may be prepared by gradually adding 6 parts of bromine to a mixture of 6 parts of ethylic alcohol and 1 of amorphous phosphorus contained in a flask fitted with an upright condenser, care being taken to keep the apparatus cool.



When all the bromine has been added, the mixture is poured into a retort and distilled over a water-bath, the resulting ethylic bromide freed from excess of bromine by washing with a small quantity of dilute soda or potash, then washed with water, and rectified over calcium chloride and redistilled.

For its preparation on a large scale De Vrij's method is preferable, $\text{C}_2\text{H}_5\text{HSO}_4 + \text{KBr} = \text{C}_2\text{H}_5\text{Br} + \text{KHSO}_4$ (see *Pharm. Journ.*, Feb. 15, 1879), or the same method as modified by Greene (*P. J.*, July 12, 1879), by Remington (*P. J.*, May 29, 1880), or by Wolff (*P. J.*, July 3, 1880).

Mon-iodo-ethane, ethyl iodide or ethylic iodide, $\text{C}_2\text{H}_5\text{I}$, may be made, like the bromide, by mixing 7 to 8 parts of amorphous phosphorus and 70 of absolute alcohol with 100 parts of iodine. The complete decomposition takes three or four hours, after

which it may be treated as above. It should be kept in a dark place, as light favors decomposition and liberation of iodine.

The paraffins give rise to many substitution-derivatives by displacement of their hydrogen by compound acidulous radicals. The following, chiefly from ethane and pentane, are of pharmaceutical interest:

Spirit of Nitrous Ether.

Ethyl Nitrite, Nitrous Ether, $C_2H_5NO_2$.—A “spirit,” probably containing nitrous ether, was one of the earliest known medicinal compounds, its discovery being generally ascribed to Raymond Lully.

Process.—To a third of a test-tubeful of rectified spirit add about a tenth of its bulk of sulphuric acid, rather more of nitric acid, and warm the mixture; as soon as ebullition commences the vapor of nitrous ether (with other substances) is evolved, recognized by its odor. A long bent tube, kept very cool, may be adapted by a perforated cork to the test-tube, and thus a little of the product be condensed and collected.

The above process, conducted on a larger scale, with definite quantities of materials and slight modifications, temperature regulated by a thermometer and a well-cooled condenser, etc. (see p. 127), is the official process for the preparation of a concentrated solution of nitrous ether, etc. in spirit; diluted with nearly three times its bulk of rectified spirit, it forms the official variety of the “spirit of nitrous ether” (*Spiritus Ætheris Nitrosi*, U. S. P.) of pharmacy, containing about 5 per cent. of the crude ether.


“Take of—

Sodium nitrite	770 grms.
Sulphuric acid	520 “
Sodium carbonate	10 “
Potassium carbonate (anhydrous)	30 “
Deodorized alcohol	sufficient.
Water	sufficient.”

Dissolve the sodium nitrite in 1000 cc. of water in a suitable flask connected with a condenser kept cold by ice-cold water; then add 550 cc. of deodorized alcohol, and mix well. Through a cork fitted into the mouth of the flask insert a funnel-tube dipping below the surface of the liquid. With the condenser connect a receiver, and keep this surrounded by a freezing mixture. Then gradually introduce into the flask, through the funnel-tube, the sulphuric acid, previously diluted with 1000 cc. of water. Distillation will usually commence before the whole of the acid has been added. When all the acid has been introduced, regulate the distillation by the application or withdrawal of a gentle heat until no more nitrous ether distils over. Wash the distillate, first, with 100 cc. of ice-cold water to remove any alcohol which may have passed over, and then remove any traces of acid by washing the ether with 100 cc. of ice-cold water in which the sodium carbonate had previously been dis-

solved. Carefully separate the ether from the aqueous liquid, and agitate it, in a well-stoppered vial, with the potassium carbonate to remove traces of water. Then filter it through a pellet of cotton, in a covered funnel, into a tared bottle containing 2000 cc. of deodorized alcohol. Ascertain the weight of the nitrous ether filtered into the alcohol by noting the increase of weight of the tared bottle and contents, and then add enough deodorized alcohol to make the mixture weigh 22 times the weight of the nitrous ether added. Lastly, transfer the product to small, dark amber-colored, well-stoppered vials, and keep them in a cool place, remote from lights or fire.

Disregarding the other products, the following equation represents the chief decomposition that occurs in the operation :



Properties.—*Spirit of Nitrous Ether*, U. S. P., is “a clear, mobile, volatile, and inflammable liquid of a pale-yellowish or faintly greenish-yellow tint, having a fragrant, ethereal, and pungent odor free from acidity, and a sharp, burning taste. Sp. gr. 0.836 to 0.842 at 15° C. When freshly prepared, or even after being kept for some time with but little exposure to light and air, it is neutral to litmus-paper. When long kept or after having been freely exposed to air and light, it acquires an acid reaction, but it should not effervesce when a crystal of potassium bicarbonate is dropped into it. If a test-tube be half filled with the ‘spirit,’ and put into a water-bath heated to 65° C. (149° F.) until it has acquired this temperature, the ‘spirit’ should boil distinctly upon the addition of a few small pieces of broken glass.” It should not effervesce, or only feebly so, when shaken with a little sodium bicarbonate (showing absence of appreciable quantities of free nitrous, acetic, or other acid). Any aldehyde, which is generally present, may be detected by the potash test (see “Aldehyde, Test for,” in Index). The great tendency of this aldehyde to become converted into acetic acid by the absorption of oxygen from the air renders spirit of nitrous ether unstable, and pharmacists are obliged to neutralize such acid, generally by potassium bicarbonate, before adding it to medicines containing iodides, etc.

A very old variety of spirit of nitrous ether, or rather “sweet spirit of nitre,” still largely sold in Great Britain, is made from spirit of wine and nitric acid, as ordered in the London Pharmacopœias, except that the distillation is continued until the product has a sp. gr. of 0.850. It may contain little or no nitrite, but is popular.


Test.—Any nitrous radical may be detected by adding ferrous sulphate and sulphuric acid to some of the spirit of nitrous ether, the usual black compound being produced.

Official (U. S. P.) Test of Strength.—“If 5 cc. of recently prepared spirit of nitrous ether be introduced into a nitrometer, and followed, first, by 10 cc. of potassium iodide, and then by 10 cc. of




sulphuric acid, the volume of nitrogen dioxide generated at the ordinary indoor temperature (assumed to be at or near 25° C. or 77° F.) should not be less than 55 cc. (corresponding to about 4 per cent. of pure ethyl nitrite)."

(For the detection of methyl alcohol in spirit of nitrous ether *vide* "Methylated Sweet Spirit of Nitre" in Index.)

Pure Ethyl Nitrite.—Dr. Leech states that the physiological and the therapeutic action of "spirit of nitrous ether" is similar to that of an alcoholic solution of ethyl nitrite of similar nitrous strength. The nitrite was prepared for Dr. Leech in the Research Laboratory of the Pharmaceutical Society of Great Britain by Hare's process of mixing sodium nitrite, sulphuric acid, and alcohol at a low temperature, but perhaps for the first time in a pure condition. The nitrite separates as a pale yellow layer. It may be washed rapidly with a little water and dried with anhydrous potassium carbonate. As it is decomposed by prolonged contact with water, Dr. Leech was supplied, according to suggestions and experiments by the late John Williams, with a pure solution of the ether in absolute alcohol containing 5 per cent. of glycerin as a preservative. It should be dispensed and used from small bottles to avoid loss by volatilization and to prevent absorption of moisture from the air.



Nitro-compounds.—There are two nitro-ethylic compounds having similar composition, but differing very much in properties—namely, *ethyl nitrite* ($\text{C}_2\text{H}_5\text{NO}_2$), which boils at 63.5° F. (17.5° C.) and has a sp. gr. of 0.900 (at 0° C.; water = 1; 0.917 to 0.920; Dunstan and Dymond), and *nitro-ethane* ($\text{C}_2\text{H}_5\text{NO}_2$), which boils at about 235° F. (nearly 113° C.) and has a sp. gr. of 1.058. There are also two nitro-methylic compounds—namely, methyl nitrite (CH_3NO_2) and nitro-methane (CH_3NO_2). The nitrites are easily decomposed; the nitro-compounds are stable. The official spirit of nitrous ether contains ethyl nitrite. Possibly the nitrites contain the nitrogen in the trivalent or unsaturated condition, while in the nitro-compounds it is in the quinquivalent or saturated state. Moreover, the reaction of the two sets of bodies warrants the conclusion that in the nitrites the methyl or ethyl is united to oxygen, in the nitro-compounds to the nitrogen. Each hydrocarbon furnishes only one mono-nitrite; each hydrocarbon furnishes only one mono-nitro compound. The nitrites are, the nitro-compounds are not, saponifiable; on reduction the nitrogen of the former does not, while that of the latter does, remain with the radicals, yielding amines. There are two similar nitro-amyl compounds.




Acetic Ether, or Ethyl Acetate.

Ethyl Acetate, or *Acetic Ether*, $\text{CH}_3\cdot\text{CO}\cdot\text{OC}_2\text{H}_5$ or $\text{C}_2\text{H}_5\text{C}_2\text{H}_3\text{O}_2$.—To a little dried sodium acetate, in a test-tube, add a small quantity of rectified spirit of wine and some sulphuric acid. Adapting a long bent tube in the usual manner, heat the test-tube and so distil over acetic ether, which may be collected in another test-tube, kept cool by partial immersion in cold water.


The official proportions (*Æther Aceticus*, B. P.) are: rectified spirit, $32\frac{1}{4}$ fluidounces; sulphuric acid, $32\frac{1}{2}$ fluidounces; sodium acetate, 40 ounces; potassium carbonate, freshly dried, 6 ounces. To the spirit slowly add the acid, keeping the fluid cool, and, the product being cold, add the acetate, mixing thoroughly. Distil 45 fluidounces. Digest the distillate with the potassium carbonate for three days in a stoppered bottle. Separate the ethereal fluid, and again distil until all but about 4 fluidounces have passed over. Preserve the resulting acetic ether in a well-closed bottle and in a cool place. It is a colorless liquid with an agreeable ethereal odor. *Æther Aceticus*, U. S. P., has the specific gravity 0.889 to 0.897. Boiling-point, about 76°C . (168°F). If a small portion of the ether be carefully poured upon some concentrated sulphuric acid, no dark ring should be developed at the point of contact of the two layers (absence of readily carbonizable organic impurities). Soluble in all proportions in rectified spirit and in ether.

“When 25 cc. of acetic ether are shaken, in a graduated tube, with 25 cc. of water just previously saturated with the ether, upon separation the ethereal layer should not measure less than 24.5 cc. (absence of an undue proportion of alcohol or water).”



Amyl Acetate, $\text{CH}_3\cdot\text{CO}\cdot\text{OC}_5\text{H}_{11}$ or $\text{C}_5\text{H}_{11}\text{C}_2\text{H}_3\text{O}_2$. (Fousel oil, or ordinary amylic alcohol, is a mixture of two or more alcohols derived from pentane, but the derivatives may be simply termed amyl compounds; *vide* Pentylic or Amylic Alcohol in Index.)

To a small quantity of amylic alcohol in a test-tube add some potassium acetate and a little sulphuric acid, and warm the mixture; the vapor of amyl acetate is evolved, recognized by its odor, which is that of the jargonelle pear. If a condensing-tube be attached, the essence may be distilled over, washed by agitation with water to free it from alcohol, and separated by a pipette.



Fruit-essences.—Amyl acetate, prepared with the proper equivalent proportions of constituents, as indicated by the above equation, is largely manufactured for use as a flavoring agent by confectioners. Amyl valerianate ($C_5H_{11}C_5H_9O_2$) is similarly used under the name of apple oil. Ethyl butyrate ($C_2H_5C_4H_7O_2$) closely resembles the odor and flavor of pine-apple; ethyl ceanthylate ($C_2H_5C_7H_{13}O_2$) recalls greengage; ethyl pelargonate ($C_2H_5C_9H_{17}O_2$), quince; ethyl suberate ($C_2H_5C_8H_{15}O_4$), mulberry; ethyl sebacate ($C_2H_5C_{10}H_{19}O_4$), melon. Salicylic aldehyde (the old salicylol or salicylofus acid), $C_6H_4 \cdot OH \cdot COH$, is the essential oil of meadow-sweet (*Spiræa ulmaria*), and may be prepared artificially by the oxidation of salicin. (*Vide Index*, "Salicin.") Acid methyl salicylate ($CH_3HC_7H_4O_3$), or gaultheric acid, forms the chief part of the essential oil of winter-green (*Gaultheria procumbens*), the fresh leaves of which yield about 0.4 per cent. of oil. Acid methyl salicylate, produced by synthesis, is official (*Methyl Salicylas*, U. S. P.). It is a colorless, optically inactive liquid, soluble in alcohol, glacial acetic acid, and carbon bisulphide. The white precipitate produced by adding sodium hydrate to a solution of methyl salicylate should dissolve at 100° after a few minutes; if this liquid be diluted and excess of hydrochloric acid added, the precipitate formed should, after being recrystallized, answer to all the tests of purity and identity for salicylic acid. Oil of sweet-birch (*Betula lenta*) is methyl salicylate, and is official—*Oleum Betulæ Volatile*, U. S. P. Salicylic aldehyde may also be prepared artificially by heating chloroform and sodium phenol. Salicylic acid ($C_6H_4 \cdot OH \cdot COOH$) can be obtained from methyl salicylate, but more cheaply from carbolic acid.

By mixing such *ethereal salts* (*alkyl salts or esters*; *vide Index*) with each other and with essential oils in various proportions the odor and flavor of nearly every fruit may be fairly imitated. (For a set of formulæ of fruit-essences see *Pharmaceutical Journal*, May 17, 1879.)

Amyl Nitrite, or Nitrite of Amyl.

Amyl Nitrite (*Amyl Nitris*, U. S. P.) ($C_5H_{11}NO_2$).—This may be prepared on the large scale by the direct action of nitric acid on amylic alcohol, the nitric acid being reduced to nitrous by a portion of the alcohol, and valerianic aldehyde with valerianic acid being produced. The heat must be very carefully regulated or the reaction may become extremely violent; indeed, with small quantities a violent explosion may occur. For experimental purposes it is preferable to pass nitrous gases, generated by the action of nitric acid on white arsenic or on starch, into the amylic alcohol (kept cool by standing the vessel in cold water) until the alcohol is saturated. The product is shaken with an aqueous solution of potassium hydrate or carbonate to remove free acids, and the oily liquid then separated and distilled. The portion distilling between 205° and 212° F. is the amyl nitrite.

The official amyl nitrite is a yellowish ethereal liquid; sp. gr. of liquid should be between 0.870 and 0.880, of vapor about 4.03; boiling-point, 96° C. (205° F.); soluble in spirit of wine, insoluble in

water; converted by fused caustic potash into potassium valerianate: exposed to the air, it yields amylic alcohol. If of good quality (for physiological purposes, although perhaps not chemically pure) about 80 per cent. (principally isoamyl nitrite) will distil between 194° and 212° F. (90° to 100° C.), the bulb of the thermometer being in the vapor and not touching the residual fluid.

The official and commercial varieties of "nitrite of amyl" are well known to be only ("chiefly," B. P.) real amyl nitrite. Since the British Pharmacopœia was published, the staff of the Research Laboratory of the Pharmaceutical Society have shown that the fluid may contain both alpha-amyl and beta-amyl nitrites, iso-butyl nitrite, and propyl nitrite, and have furnished specimens of these substances to Professor Cash, who finds that their physiological action is not primarily dependent on the amount of their nitrosyl. These nitrites are, of course, derived from the corresponding hydrates (see pp. 449 and 450) present in the amylic alcohol.

Nitropentane ($C_5H_{11}NO_2$) is similar to amyl nitrite in composition, but differing much in properties. It is obtained by reaction of amyl iodide on silver nitrite. It boils at 300° to 320° F. (For remarks made respecting the two similar derivatives of ethane see p. 405.)

QUESTIONS AND EXERCISES.

How would you prepare methane and ethane?—Give formulæ.—Give details of the production of chloroform from alcohol, tracing the various steps by equations.—Give the formulæ and state the constitution of the various chlorine derivatives of methane.—How is chloroform purified?—State the characters of pure chloroform.—Explain the official process for the preparation of nitrous ether.—Give the properties of nitrous ether as compared with nitro-ethane.—By what official method is the strength of spirit of nitrous ether to be estimated?—How is ethyl iodide made?—Mention the systematic names of several artificial fruit-essences.—What is the formula of amyl nitrite, and how is it prepared?


THE OLEFINE SERIES OF HYDROCARBONS.

The Olefine Series of Hydrocarbons consists of *unsaturated* hydrocarbons, having the general formula C_nH_{2n} . *Ethylene*, C_2H_4 ; *Propylene*, C_3H_6 ; *Butylene*, C_4H_8 ; *Amylene*, C_5H_{10} ; *Hexylene*, C_6H_{12} ; and *Heptylene*, C_7H_{14} , and many others are well known.


Ethylene, *Olefiant Gas*, or *Heavy Carburetted Hydrogen*, C_2H_4 , is the first of this series. It is formed in the destructive distillation of coal, and is the chief illuminating constituent of coal-gas. Coal-gas consists of 30 to 40 per cent. of methane, 40 to 50 per cent. of hydrogen, and from 5 to 7 per cent. of ethylene and its homologues. Hydrocarbons, normally fluid, but kept in the vaporous condition by the diluents, also contribute materially to the illuminating power of gas. The impurities are nitrogen, air, carbolic acid, carbon disulphide (CS_2), and some badly-smelling sulphur compounds.

Upward of one hundred and fifty distinct chemical substances have been obtained from the solid, liquid, and gaseous products of the destructive distillation of coal.

Preparation.—Ethylene may be prepared by dropping alcohol into a large retort or flask containing 10 ounces of sulphuric acid and 3 ounces of water heated to 160–165° C. The gas is washed in cold water and a solution of soda to free it from ether, alcohol, and sulphurous acid :



The product, when further heated, yields ethylene :




If the ethylene be passed into bromine under water until all the bromine disappears, *ethylene dibromide*, $\text{C}_2\text{H}_4\text{Br}_2$, or *dibromethane*, will be formed. (For constitution see p. 393.)

Properties.—A colorless and odorless gas, burning with a luminous flame.

Ethylene Sulphate, $\text{C}_2\text{H}_4\text{SO}_4$, is said to be contained in “Hoffman’s anodyne,” the *Spiritus Ætheris Compositus*, U. S. P., a solution of 25 cc. of ethereal oil, 325 cc. of ether, and 1650 cc. of alcohol. The so-called ethereal oil or heavy oil of wine is obtained by digesting spirit of wine and sulphuric acid together, then distilling, removing any acid from the distillate by washing with lime-water, and exposing the ethereal fluid to the air to facilitate escape of the more volatile fluids. The product is a mixture consisting probably of ethylene sulphate, ethyl sulphate, ether, dissolved ethylene, and other bodies.

Glycols.—The olefines form corresponding dihydric alcohols or glycols (named from glycol, the first member of the series), and these give two sets of aldehydes and acids. (See also p. 456.) Thus :



RELATION OF PARAFFINS TO OLEFINES.

1st. The paraffins may be converted into olefines either by acting on alcohols of the paraffin series by sulphuric acid or by acting on a monochloro-paraffin by caustic potash.







TABLE SHOWING RELATIONS BETWEEN PARAFFINS, OLEFINS, AND ACETYLENES.

	Paraffin, $C_n H_{2n+2}$.	Olefines, $C_n H_{2n}$.	Acetylenes, $C_n H_{2n-2}$.
Hydrocarbon	Ethane, $C_2 H_6$, or $CH_3 \cdot CH_3$	Ethylene, $C_2 H_4$	Acetylene, $C_2 H_2$.
Alkyl Salts Chloro-derivatives	Ethyl Chloride, or Monochloroethane, $C_2 H_5 Cl$	Ethylene Chloride, or Dichloroethane, $C_2 H_4 Cl_2$	Acetylene Chloride, or Tetrachloro-ethane, $C_2 H_2 Cl_4$.
Alcohol	Ethylic Alcohol, $C_2 H_5 OH$, or $CH_3 CH_2 OH$	Ethylene Glycol, $C_2 H_4 (OH)_2$, or $(CH_2 OH)_2$	
Acid	Acetic Acid, $CH_3 \cdot CO \cdot OH$	Oxalic Acid $(COOH)_2$	
Cyanide	Cyanethane, $CH_3 CH_2 CN$	Ethylene Cyanide, or Dicyanethane $C_2 H_4 (CN)_2$	

Inversely, the olefines may be converted into paraffins. By combining an olefine with hydrochloric acid a monochloro-paraffin results, which, when acted on by nascent hydrogen, yields a paraffin.



2d. The bromine, chlorine, and iodine additive derivatives of the olefines are either identical or isomeric with the substitution-derivatives of the paraffins. Thus $\text{C}_2\text{H}_4\text{Cl}_2$ is either dichlorethane or ethylene chloride; and the additive derivatives with the acids, such as hydrochloric, produce mono-substitution derivatives of the paraffins. In the case of the chloride, however, $\text{C}_2\text{H}_4\text{Cl}_2$, it has a different boiling-point and specific gravity according as it is prepared from ethylene and chlorine (*ethylene chloride*, alpha dichlorethane, or the old "Dutch liquid") or from monochlorethane (ethyl chloride) and chlorine (monochlorethyl chloride, beta-dichlorethane, or *ethylidene chloride*.) The former may be represented by the formula $\text{CH}_2\text{Cl}\cdot\text{CH}_2\text{Cl}$, and the latter as $\text{CH}_3\cdot\text{CHCl}_2$. It is the former also which yields glycol (by reaction of the chloride with silver acetate, and of the resulting ethylene acetate with an alkaline hydrate), hence the formula of the glycol also must be $\text{CH}_2\text{OH}\cdot\text{CH}_2\text{OH}$, and not $\text{CH}_3\cdot\text{CH}(\text{OH})_2$:



THE ACETYLENE SERIES OF HYDROCARBONS.

The members of the acetylene series, $\text{C}_n\text{H}_{2n-2}$, are characterized by forming metallic derivatives. *Acetylene* itself, C_2H_2 , is formed during the passage of electric sparks between carbon points in an atmosphere of hydrogen; it is the only member which can be formed directly from its elements. Other members of the series are *Allylene*, C_3H_4 ; *Crotonylene*, C_4H_6 ; etc. The hydroxyl derivative of allylene, known as *propargyl alcohol*, has the formula $\text{C}_3\text{H}_3\text{OH}$.

Preparation.—Acetylene may be obtained by heating ethylene bromide (dibromethane) with caustic potash, and passing the gas into a well-cooled ammoniacal solution of cuprous chloride, with which it reacts, forming a red precipitate, probably having the formula $\text{C}_2\text{H}_2\text{Cu}_2\text{O}$ (Blochmann), called cuprous acetylde. The pure acetylene gas may be obtained from the copper compound by heating with hydrochloric acid. Acetylene is yielded in a steady stream when water is dropped on to barium carbide. Acetylene is also formed by the incomplete combustion of coal-gas, as when an air-gas burner is lighted below. It has an unpleasant odor, well known in every chemical laboratory in which air-gas lamps are used.

QUESTIONS AND EXERCISES.

What are the properties of ethylene, and how is it prepared?—What alcohols are derived from the olefine series?—Mention the relations between the paraffins and olefines.—Give three methods of preparing acetylene.

THE TERPENE SERIES OF HYDROCARBONS.

The terpene series have the general formula C_nH_{2n-4} . Valylene, C_5H_6 , is the lowest, and Terebenthene or Pinene, $C_{10}H_{16}$ (pure oil of turpentine), the most common member of the series.

The hydrocarbons, called terpenes, $C_{10}H_{16}$, are very commonly met with in analyzing the volatile oils. Very few of these oils have been artificially produced. Their fragrance appears to be due to the non-terpenoid constituent (Wallach). They differ from one another in the power of deviating a ray of polarized light to the right or left. They may be divided into classes, of which several are interesting in pharmacy: (a) Terpenes or pinenes, boiling at about 156° C., and found in ordinary turpentine and other oils; (b) sylvestrene, found in Russian and Swedish turpentine; (c) phellandrene, lævo-rotatory from *Phellandrium aquaticum* and dextro-rotatory from eucalyptus oil (p. 417), boiling-point 170° C.; (d) citrenes (limonenes), boiling at about 175° C., and derived from the different species of Citrus; (e) di-pentene, found in some turpentines and oils of camphor and elemi; (f) terpinene, occurring in oil of cardamoms. Camphene, fenchene, and terpinolene are terpenes, but do not occur naturally in oils. The sesquiterpenes have the formula $C_{15}H_{24}$, and include canidene, found in oils of cubebs, savin, cade, betel, camphor, galbanum, patchouli, juniper, asafoetida, colo, and olibanum; caryophyllene, found in oil of cloves, and other isomers of these. They boil at a much higher temperature than the terpenes, but resemble them in other respects.

Oil of Turpentine (*Oleum Terebinthinæ*, U. S. P.).—Turpentine itself is really an oleo-resin of about the consistence of fresh honey. It flows naturally or by incision from the wood of most coniferous trees, larch (*Pinus Larix*) yielding Venice turpentine; *Abies balsamea* furnishing Canadian Turpentine or Canada Balsam (*Terebinthina Canadensis*, U. S. P.); the bark of *Pistachia terebinthus*, the variety termed Chian Turpentine (containing about 1 part of essential oil to 7 of resin), and the *Pinus australis* (*palustris*), *P. abies*, *P. pinaster* and *P. tæda*, affording the common American Turpentine (*Terebinthina*, U. S. P.). *Pinus maritima* gives the French or Bordeaux Turpentine, and *P. picea* the old fragrant Strasburg Turpentine. By distillation with steam this crude turpentine is separated into colophony, rosin or resin, which remains in the still, and essential oil of turpentine, often termed simply turpentine, spirit of turpentine, or "turps," which distils over. Mixed with alkali to saturate resinous acids and redistilled in a current of steam, oil of turpentine furnishes about 80 per cent. of rectified oil of turpentine. *Pinus sylvestris* and *P. Ledebourii* furnish Russian Turpentine, which, according to Tilden, consists of two terpenes and cymene, and also

(Wallach) a *lævogyre* limonene. This turpentine is probably a by-product in the preparation of common *wood tar* (*Pix liquida*, U. S. P.); its odor is very pleasant, quite different from that of ordinary turpentine. The leaves of the *Pinus sylvestris*, or Scotch fir, are in Germany broken down to a woolly condition, producing pine wool, fir wool, or wadding used in making vermin-repelling blankets; and this substance, or, still better, the fresh leaf, by distillation with water, yields *Fir-wool Oil* (*Oleum Pini Sylvestris*, B. P.), consisting, according to Tilden, of two terpenes, like those of Russian turpentine, and cymene. This oil, diffused through water by aid of magnesia, forms the *Vapor Olei Pini Sylvestris*, B. P. The terpene of Bordeaux turpentine (terebenthene) rotates a ray of polarized light more than, and in the opposite direction to, the terpene of American turpentine, *Oleum Terebinthinæ*, U. S. P.

Turpentine "commences to boil at about 320° F. (160° C.), and almost entirely distils below 356° F. (180° C.), little or no residue remaining," whereas petroleum spirit, with which turpentine might be mixed, covers much wider limits of temperature during its distillation. Petroleum spirit also, when the small round flame of the end of a piece of twine is brought near to some of the spirit in a cup, gives a momentary flash of flame at a much lower temperature than that at which turpentine flashes. Thus tested in the specially-arranged flashing apparatus of the Petroleum Act, Mr. Boverton Redwood found that the flashing-point of turpentine was lowered 18° by 1 per cent. of petroleum spirit. The specific gravity of oil of turpentine is from about 0.855 to 0.870.

Under the influence of heat and sulphuric acid or other chemical agents pure oil of turpentine ($C_{10}H_{16}$) yields many derivatives of considerable chemical interest. Amongst them are two optically inactive terpene isomers, named *terebene* and *colophene*, used for inhalation and as disinfectants and deodorizers. When acted on by gaseous hydrochloric acid the product is a white crystalline monohydrochloride, $C_{10}H_{16}HCl$. In the sunlight it slowly oxidizes, and hydrolyzes to a crystalline body, $C_{10}H_{18}O_2$. Bromine acts violently on turpentine and terpenes, resulting in dibromides, which yield cymene when heated, $C_{10}H_{16}Br_2 = C_{10}H_{14} + 2HBr$. Crystals of *terpin hydrate*, $C_{10}H_{18}(OH)_2 \cdot H_2O$, also *terpinol*, $(C_{10}H_{16})_2 \cdot H_2O$, are used therapeutically instead of terebene.

The official *Terebene*, $C_{10}H_{16}$ (*Terebenum*, U. S. P.), is a colorless, mobile liquid, having a somewhat aromatic odor and taste. Boils between 156° and 160° F. In its chemical properties it very much resembles oil of turpentine.

Volatile Oils.

The *Volatile* or *Essential Oils* exist in various parts of plants. They usually are mixtures of the liquid hydrocarbons or *elæoptens* (from *ἐλαίον*, *elaion*, oil, and *ὀπτωμαί*, *optomai*, to see) with oxidized hydrocarbons, which are commonly solid or camphor-like bodies termed *stearoptens* (from *στέαρ*, *stear*, suet), and which on cooling often crystallize out; or on distilling an oil the stearopten may remain in the retort, being less volatile than the elæopten. The

oils are also often associated with further oxidized bodies termed *resins*.

The process by which volatile oils are usually obtained from herbs, flowers, fruits, or seeds may be imitated on the small scale by placing the material (bruised cloves or caraways, for instance) in a tubulated retort, adapting the retort to a Liebig's condenser, and passing steam, from a Florence flask, through a glass tube to the bottom of the warmed retort. The steam in its passage through the substance will carry the particles of oil over the neck of the retort into the condenser, and thence, liquefied and cooled, into the receiving vessel, where the oil will be found floating on the water. It may be collected by running off the distillate through a glass funnel having a stopcock in the neck, or by letting the water from the condenser drop into an old test-tube which has a small hole in the bottom, or any similar tube placed in a larger vessel, the water and oil being subsequently run off separately from the tube as from a pipette. The water will in most cases be the ordinary official medicated water of the material operated on (*Aqua Anethi, Anisi, Aurantii Floris, Carui, Cinnamomi, Fœniculi, Menthæ Piperitæ, Menthæ Viridis, Pimentæ, Rosæ* (from *Rosæ Centifoliæ Petala*, B. P.), *Sambuci*). Volatile oils, like fixed oils, stain paper, but the stain of the former is not permanent, like that of the latter. Oils of lemon and orange are sometimes obtained by mere pressure of the rind of the fruit.

The following official (U. S. P.) waters may be made by distributing 2 parts of volatile oils over 4 parts of precipitated calcium phosphate, and percolating with 1000 parts of distilled water: *Aqua Anisi, Cinnamomi, Fœniculi, Menthæ Piperitæ, Menthæ Viridis*.

Aqua Aurantii Florum Fortior, U. S. P. (*Aqua Aurantii Florum*, U. S. P. 1880), is a saturated solution of volatile orange-flower oil, and the *Aqua Aurantii Florum*, U. S. P. 1890, is made by mixing equal volumes of the *stronger orange-flower water* and distilled water.

The presence of alcohol in an essential oil may be detected and its quantity estimated by shaking with an equal bulk of pure glycerin. The latter dissolves the alcohol, and is augmented in volume according to the amount of alcohol present (Boettger). (For tests for alcohol see Index, "Alcohol.")

A large number of volatile oils are employed in medicine, either in the pure state, in the form of saturated aqueous solution (medicated waters), solution in spirit of wine, 1 in 5 (*Essentia Anisi* and *Essentia Menthæ Piperitæ*, B. P.), and 1 in 50 (*Spiritus Cajuputi, Cinnamomi, Juniperi, Lavandulæ, Menthæ Piperitæ, Myristicæ, Rosmarini*), or as leading constituents in various barks, roots, leaves, etc. The strength of *Spiritus Anisi*, U. S. P., and *Sp. Cinnamomi*, U. S. P., is 10 of oil to 90 of deodorized alcohol. *Sp. Menthæ*

Piperitæ, U. S. P., and *Sp. Menth. Viridis*, U. S. P., are of similar strength, but also contain whatever may be extracted by the 100 parts of the spirit from 1 part of the dried herb. *Spiritus Aurantii*, U. S. P., contains 5 of oil and 95 of alcohol; *Sp. Gaultheriæ*, U. S. P., *Sp. Juniperi*, U. S. P., *Sp. Lavandulæ*, U. S. P., and *Sp. Myristicæ*, U. S. P., contain 5 of oil and 95 of alcohol; *Spiritus Limonis*, U. S. P., is made with 10 of oil, 10 of fresh-grated lemon-peel, and deodorized alcohol sufficient to produce 1000 of filtered product. *Perfumes* ("scents" or "essences," including "lavender-water" and "eau de Cologne") are for the most part solutions of essential oils in spirit of wine or spirituous infusions of materials containing essential oils. The following oils are, directly or indirectly, official in the Pharmacopœias:* 1. Volatile oil of *Bitter Almond* (*vide* Index). 2. Oil of the fruits of *Ajwain* or *Omum*, *Carum Ajowan*, or *Ptychotis Ajowan* (*Fructus Ptychotis*, P. I.) contains *cymol* or *cymene* ($C_{10}H_{14}$) and a stearopten (*Ajwainka-phul*, flowers of ajwain) identical with *thymol*, $C_{10}H_{14}O$. 3. Oil of *Dill* (*Oleum Anethi*, B. P.), a pale, yellow, pungent, acrid liquid distilled from dill-fruit; it contains a hydrocarbon, *anethene* ($C_{10}H_{16}$), and an oxidized oil ($C_{10}H_{14}O$) identical with the carvol of oil of caraway (Gladstone). 4. Oil of *Aniseed* (*Oleum Anisi*, U. S. P.), a colorless or pale-yellow liquid, of sweetish warm flavor, distilled in Europe from the anise-fruit (*Pimpinella Anisum*) (*Anisum*, U. S. P.), but chiefly, in China, from the fruit of star-anise (*Illicium Anisatum*) (*Illicium*, U. S. P.); it is a mixture of a hydrocarbon isomeric with oil of turpentine and *anethol*, a stearopten ($C_{10}H_{12}O$) which crystallizes out at low temperatures. The melting-point of anethol is between 74° and 75° F. (Flückiger). The congealing-point of the natural oils appears to be dependent on the proportion of the fluid to the solid constituent, a very small quantity of the former lowering the congealing- and melting-points very considerably. 5. Oil of *Chamomile* (*Oleum Anthemidis*, B. P.), a bluish or, when old, yellow oil of characteristic odor and taste, distilled from chamomile-flowers (*Anthemis*, U. S. P.). The official variety (*Anthemis nobilis*) yields about 0.2 per cent. of an oil composed of a hydrocarbon ($C_{10}H_{16}$) and an oxidized portion ($C_{10}H_{16}O_2$), which, heated with potash, gives *potassium angelate* ($KC_5H_7O_2$), whence is obtained *angelic acid* ($HC_5H_7O_2$). According to Demareay, Kopp, and Köbig, the oil is a mixture of the angelates of butyl and amyl and similar bodies. Naudin has also obtained from chamomile *anethen*, a hydrocarbon crystallizing in needles. The flowers of another variety (*Matricaria Chamomilla*) (*Matricaria*, U. S. P.) contain a stearopten ($C_{10}H_{16}O$) having the composition of laurel-camphor. 6. Oil of *Horse-radish* root (*Armoracæ Radix*, B. P.) is, according to Hofmann, butyl or tetryl sulphocyanate (C_4H_9CNS): it is the chief active ingredient of *Spiritus Armoracæ Compositus*, B. P. 7. Oil of *Sweet-Orange* peel (*Aurantii Dulcis Cortex*, U. S. P.) and oil of *Bitter-Orange* rind (*Aurantii Amari Cortex*, U. S. P.; *Oleum Aurantii Corticis*, U. S. P.), the

* The student is not expected to remember, but to understand, all that follows respecting the volatile oils.

former the flavoring constituent of the official syrup of the *peel* (*Syrupus Aurantii*, U. S. P.) and the oils of various species of *Citrus*—namely, 8, *lemon* (*Oleum Limonis*, U. S. P.), from lemon-peel (*Limonis Cortex*, U. S. P.); 9, *lime*, 10, *bergamot* (*Oleum Bergamottæ*, U. S. P.), 11, *citron* and a variety of citron termed *cedra*—resemble each other in composition, all containing *hesperidene*, a hydrocarbon ($C_{15}H_{34}$), and a small quantity of oxidized hydrocarbons [$C_{10}H_{10}O_5$, $C_{15}H_{16}O$, and (Wright and Piesse) $C_{20}H_{30}O_3$], etc. Tilden states that lemon oil distilled from the fresh peel consists chiefly of a terpene, $C_{10}H_{16}$, boiling at $176^\circ C.$, with small quantities of a terpene boiling below 160° , and a hydrous terpene, the odor of the oil being due to the mixture. Oil of bergamot appears to owe its fragrance to 40 or 50 per cent. of *linalool acetate*, $C_{10}H_{17}C_2H_3O_2$. It also contains a stearopten, *bercapten*, $C_{12}H_8O_4$. Expressed lime essence contains a soft resin. 12. *Oleum Aurantii Florum*, U. S. P., oil of *Neroli* or *Orange-flower*, the aqueous solution of which is official in the forms of water (*Aqua Aurantii Florum Fortior*, U. S. P.) and syrup (*Syrupus Aurantii Florum*, U. S. P.), contains a fragrant hydrocarbon ($C_{10}H_{16}$), colorless when fresh, but becoming red on exposure to light, and an inodorous oxidized hydrocarbon. Strong acids, especially nitric, attack the oil in orange-flower water, coloring the fluid of a rose tint. 13. Oil of *Petit Grain*, distilled from the leaves and shoots of the orange tree, consists chiefly of *aurantiol acetate*, $C_{10}H_{17}C_2H_3O_2$. 14. The leaves of *Boldo* (*Peumus Boldus*), a Chilean shrub (tonic and hepatic), yield 2 per cent. of essential oil (and, according to Bourgon and Verne, an alkaloid, *boldine*). 15. Oil of *Buchu-leaves* (*Buchu*, U. S. P.) consists chiefly of a fluid oil, $C_{10}H_{18}O$, holding in solution a crystalline stearopten, diosphenol, ($C_{14}H_{22}O_3$, Flückiger; $C_{10}H_{16}O_2$, Spica, Shimoyana also). 16. Oil of *Cannabis indica*, see p. 423. 17. Oil of (the lesser) *Cardamoms*, from the seeds of the capsules (*Cardamomum*, U. S. P.), is chiefly a hydrocarbon ($C_{10}H_{16}$) isomeric with oil of turpentine (terpene and probably limonene) and a camphor resembling turpentine-camphor ($C_{10}H_{16}^3H_2O$). 18. Oil of *Cajuput* (*Oleum Cajuputi*, U. S. P.) is a mobile bluish liquid, consisting of *hydrous cajuputene* or *cajuputol* ($C_{10}H_{16}H_2O$); *cajuputene*, $C_{10}H_{16}$; *cineol*, $C_{10}H_{17}OH$; a sesquiterpene, $C_{15}H_{24}$; as well as butyric, valeric, and benzoic aldehydes. The latter, repeatedly distilled from phosphoric anhydride, yields cajuputene itself ($C_{10}H_{16}$), which has the odor of hyacinths. Fresh cajuput oil has a green hue, which is perhaps transient, for the color of the oil of trade is due to copper (Guibourt and Histed): certainly the green coloring-matter of pure cajuput oil is organic and either oily or chlorophylloid. 19. Oil of *Caraway-fruit* (*Carum*, U. S. P., *Oleum Carui*, B. P., *Oleum Cari*, U. S. P.) is a mixture of *carvene* ($C_{15}H_{24}$) and *carvol* ($C_{10}H_{14}O$). 20. Oil of *Cloves* (*Caryophyllus*, U. S. P., *Oleum Caryophylli*, U. S. P.) and of *Pimento* or *Pimenta*, U. S. P., or *Allspice* (*Oleum Pimentæ*, U. S. P.), both heavier than water, contain a liquid hydrocarbon ($C_{15}H_{24}$), *eugenol* ($C_{10}H_{12}O_2$), a solid body, *eugenin*, isomeric with the eugenic acid, a second crystalline substance, *caryophyllin* ($C_{10}H_{16}O$), isomeric with

common camphor, and a salicylic compound. 21. Oil of *Cascarilla*-bark (*Cascarilla*, U. S. P.) has not been fully examined. 22. Oil of *Cinnamon*-bark (*Cinnamomum Cassia*, U. S. P.) and of *Cassia*-bark is mostly cinnamic aldehyde (C_8H_7COH). Boiled with nitric acid, it furnishes benzoic aldehyde (C_6H_5COH) and benzoic acid (C_6H_5COOH); with chloride of lime it yields calcium benzoate ($(C_6H_5COO)_2Ca$), and with caustic potash gives potassium cinnamate (C_8H_7COOK). The specific gravity of oil of Ceylon cinnamon is about 1.040, and of Chinese cinnamon (oil of cassia) about 1.060. Both are termed *Oleum Cinnamomi* in U. S. P. 23. Oil of *Citronella*, a grass oil, from *Andropogon nardos*, is chiefly composed of *citronellol* ($C_{10}H_{16}O$ and $C_{10}H_{18}O$, Wright), probably isomeric with the *absinthol* from the *Artemisia Absinthium* or wormwood (*Absinthium*, U. S. P.) (Gladstone). Kremer also obtains heptoic aldehyde ($C_7H_{14}O$), a terpene ($C_{10}H_{16}$), etc. 24. Oil of *Copaiva* (*Oleum Copaibæ*, U. S. P.), and, 25, of *Cubebs* (*Oleum Cubebæ*, U. S. P.), are hydrocarbons having the formula $C_{15}H_{24}$. This *cubebene* is sometimes associated with a camphor, *hydrous cubebene* ($C_{15}H_{24}, H_2O$). Oil of cubebs also contains a small quantity of a terpene ($C_{10}H_{16}$). 26. Oil of *Coriander* (*Oleum Coriandrum*, U. S. P.) seems to have the composition of hydrous oil of turpentine ($C_{10}H_{16}, H_2O$). 27. The fruits of *Cumin*, or *Cummin* (*Cuminum Cyminum*), an ingredient of many curry-powders, contain about 3 per cent., and those of *Water Hemlock* or *Cowbane* (*Cicuta virosa*) about $1\frac{1}{2}$ per cent., of an essential oil composed of *cymol* or *cymene* ($C_{10}H_{14}$) and *cumic aldehyde* ($C_9H_{11}COH$). The latter is an aldehyde readily uniting with alkaline bisulphites and by oxidation yielding *cuminic acid* ($C_9H_{11}COOH$). *Cymol* also occurs in *Garden Thyme* (*Thymus vulgaris*). 27a. The fresh flowering herb of *Erigeron canadense*, or Canadian fleabane, yields an essential oil (*Oleum Erigerontis*, U. S. P.). 28. The fresh leaves of various species of *Eucalyptus* (*E. globulus*, *E. oleosa* or *cneorifolia*, *E. dumosa*, and other "mallee"—that is, "scrub" or shrub-like eucalypts) furnish about 1 per cent. of the essential *Oil of Eucalyptus* (*Oleum Eucalypti*, U. S. P.). The more volatile portion of this oil consists partly of *cymene* and three terpenes—namely, *pinene*, *limonene*, and, especially in *E. amygdalina*, *phellandrene* (p. 412). The latter is more readily alterable than other terpenes, and is characterized by yielding a crystalline mass with nitrous anhydride. Oxidized bodies also are present, $C_{10}H_{14}O$, $C_{10}H_{16}O$, and 40 to 50 per cent. of an oil (or, when cooled, a camphor) having the same composition as *cajuputol*, and as the chief constituent of *wormseed oil*—namely, *eucalyptol*, $C_{10}H_{18}O$ or $C_{10}H_{16}, H_2O$; boiling at about $174^\circ C.$, freezing at about $0^\circ C.$, and having a specific gravity of about 0.927. It is not yet known satisfactorily to which of the constituents of eucalyptus oil its medicinal efficacy is due. The sp. gr. of the oils varies greatly—namely, 0.030 to 0.040 above or below 0.900. The official density is 0.915 to 0.925. Different species of eucalyptus may yield oils differing in specific gravity, flavor, and odor. *E. maculata*, var. *citriodora*, contains an aldehyde or ketone similar to that of citronella. Like the turpentines, the eucalyptus oils are good solvents of resins. Voiry states that eucalyptol is

present also in the oil of *Lavandula spica*, oil of *spike* or "foreign" oil of lavender. *Red gum* (*Eucalypti Gummi*, B. P.) is from the *E. rostrata* and other species, and is used solely for its astringent properties. 29. *Elecampane-root* (*Inula Helenium*) (*Inula*, U. S. P.) by distillation with water yields solid volatile *helenin* (C_6H_8O), a camphor-oil or *inulol* ($C_{10}H_{16}O$), and *inulic anhydride* or *lactone* ($C_{15}H_{20}O_2$), as well as, according to Marpmann, crystals of *alantic acid* ($C_{15}H_{22}O_3$) and fluid *alantol* ($C_{20}H_{32}O$), each more powerfully antiseptic than helenin. 30. *Oleum Fœniculi*, U. S. P., oil of *Fennel-fruit* (*Fœniculum*, U. S. P.), differs in odor, but contains the same proximate constituents as oil of anise. 31. Oil of *Geranium*, or *Ginger Grass* oil, from *Andropogon schœnanthus* and various species of *Pelargonium*, contains *geraniol* ($C_{10}H_{18}O$). 32. *Grains of Paradise* (*Amomum melegueta*), Guinea Grains or Melegueta Pepper, *Semina Cardamomi Majoris*, contain essential oil ($C_{10}H_{16}$ and $C_{10}H_{16}O$) and a highly pungent principle, termed by Thresh *paradol*, $C_9H_{14}O_2$, isomeric with capsaicin of the same chemist. 32a. Oil of *American Pennyroyal* (*Hedeoma*, U. S. P.), "the leaves and tops of *Hedeoma pulegioides*," contains *hedeomol* ($C_{10}H_{18}O$), and yields *isoheptoic acid* ($C_7H_{14}O_2$) and other substances (Kremer). 33. Oil of *Juniper* (*Oleum Juniperi*, U. S. P.), the active constituent of juniper tops and berries, contains a hydrocarbon ($C_{10}H_{16}$) which by contact with water yields a white crystalline hydrous compound ($C_{10}H_{16}, H_2O$) and a polymeric hydrocarbon ($C_{20}H_{32}$). 34. Oil of *Lavender* (*Oleum Lavandulae*, B. P.) is distilled from the flowers of *Lavandula vera*, and is the same as *Oil of Lavender-flowers* (*Oleum Lavandulae Florum*, U. S. P.); it has not been satisfactorily examined. 34a. Oil of *Myrcia* (*Oleum Myrcia*, U. S. P.), oil of bay or bayberry oil (sp. gr. about 1.040), is obtained from the leaves of *Myrcia acris*. 35. Oil or butter or camphor of *Orris* (*Iris Florentina*) is a soft solid lighter than water. Flückiger and Hanbury found it to be chiefly myristic acid associated with a little essential oil. 36. Oil of *Peppermint* (*Oleum Menthæ Piperitæ*, U. S. P.) consists of a hydrocarbon, *menthene* ($C_{10}H_{18}$), different from that of most volatile oils, depositing crystalline peppermint camphor, known as menthol, $C_{10}H_{18}, H_2O$, when exposed to low temperatures. It is official (*Menthol*, U. S. P., also *Emplastrum Menthol*, B. P.), and has the following characters: "In colorless acicular crystals, usually more or less moist from adhering oil, or in fused crystalline masses. Its melting-point should not exceed $110^\circ F.$ ($43.3^\circ C.$). It has the odor and flavor of peppermint, producing warmth on the tongue, or, if air be inhaled, a sensation of coolness. It is sparingly soluble in water and readily soluble in rectified spirit, the solutions having a neutral reaction. Boiled with sulphuric acid diluted with half its volume of water, menthol acquires an indigo-blue or ultramarine color, the acid becoming brown. It should be entirely dissipated by the heat of a water-bath."—B. P. Its alcoholic solution rotates a ray of polarized light to the left. "If a few crystals of menthol be dissolved in 1 cc. of glacial acetic acid, and then 3 drops of sulphuric acid and 1 drop of nitric acid added, no green color should be produced (absence of thymol)."—U. S. P. It is also yielded by

the oil of *Mentha arvensis* (vars. *piperascens* and *glabrata*). 37. Oil of *Spearmint* (*Oleum Menthae Viridis*, U. S. P.), the common mint of the kitchen garden, contains a liquid having the formula $C_{10}H_{20}O$ or $C_{10}H_{18}, H_2O$; also, according to Gladstone, an oil ($C_{10}H_{14}O$) isomeric with carvol. 38. Oil of *Pennyroyal* (*Mentha Pulegium*) contains, according to Kane, $C_{10}H_{16}O$ (*pulezone*; Plaisner). 38a. The leaves and tops of *Melissa officinalis* or *Balm* (*Melissa*, U. S. P.) yield a volatile oil containing a camphor. 39. Oil of *Nutmeg* (*Oleum Myristicæ*, B. P. and U. S. P.) is composed of a hydrocarbon, *myristicene* ($C_{10}H_{16}$), and *myristicol* ($C_{10}H_{16}O$), and *cymene* ($C_{10}H_{14}$) (Gladstone). *Mace* (*Macis*, U. S. P.), the arillus or net-like envelope of the nutmeg, appears to yield similar bodies, and also *myristicin*, $C_{12}H_{14}O_3$ (Semmler). 40. Oil or *Otto* or *Attar* of *Cabbage-rose Petals* (*Rosæ Centifoliæ*, U. S. P., *Oleum Rosæ*, U. S. P.) gives the fragrance to rose-water (*Aqua Rosæ* and *Aqua Rosæ Fortior*, U. S. P.). It resembles most other volatile oils in being composed of a hydrocarbon and an oxidized portion, but differs from all in this respect, that the hydrocarbon is solid and is destitute of odor, while the oxygenated constituent is liquid and the source of the perfume. According to Flückiger, the solid hydrocarbon ($C_{18}H_{16}$) yields succinic acid as the chief product of its oxidation by nitric acid, and in other respects affords evidence of belonging to the paraffin series of fats. 41. Oil of *Rosemary-tops* (*Oleum Rosmarini*, U. S. P.) exists in the plant to the extent of from $1\frac{1}{2}$ to 3 parts per 1000. It chiefly consists of a hydrocarbon ($C_{10}H_{16}$) resembling that from myrtle, *Myrtus communis*, but also contains camphor ($C_{10}H_{16}O$) and borneol ($C_{13}H_{18}O$) in variable proportions. 42. Oil of *Rue* (*Oleum Rutæ*, B. P.) contains a small quantity of hydrocarbon ($C_{10}H_{16}$) with some rutic aldehyde ($C_{10}H_{20}O$), and methyl-nonyl ketone, $C_{11}H_{22}O$ or $CH_3-CO-C_9H_{19}$. Gorup-Besanez and Grimm have obtained oil of rue ($C_{11}H_{22}O$) artificially as one of the products of the destructive distillation of calcium acetate and caprate. (See Ketones.) According to Greville Williams, it is chiefly euodic aldehyde ($C_{11}H_{22}O$), some lauric aldehyde ($C_{12}H_{24}O$) also being present. 43. Oil of *Sage* (*Salvia*, U. S. P.) contains about 40 per cent. of salviol, $C_{10}H_{16}O$; about 20 per cent. of two $C_{10}H_{16}$ hydrocarbons, boiling at 156° and 167° C. respectively; about 10 per cent. of a camphor, $C_{10}H_{16}O$; and about 10 per cent. of cedrene, $C_{15}H_{24}$ (Muir). 44. Oil of *Savin* (*Oleum Sabinæ*, U. S. P.) obtained from the tops of *Juniperus Sabina* or *Savina* (*Sabina*, U. S. P.), contains several hydrocarbons, but none isomeric with oil of turpentine (Tilden). 45. Oil of *Elder-flowers* (*Sambucus*, U. S. P.) occurs in very small quantity; it has a butyraceous consistence; it contains a hydrocarbon, *sambucene* ($C_{10}H_{16}$), and probably a camphor. 46. Oil of *Sandal-wood* (*Oleum Santali*, U. S. P.) or *Oil of Santal* is composed (Chapoteaut) of two bodies: mostly of a substance having a formula $C_{15}H_{24}O$ (boiling at 572° F.), and a small quantity of a substance having the formula $C_{15}H_{26}O$ (boiling at 600° F.). It occurs to the extent of about $2\frac{1}{2}$ per cent. in the fragrant white or yellow sandal-wood of India, *Santalum album*, a small tree of the natural order Santalaceæ, and not to be confounded with the *Pterocarpus santalinus*, a tree of the natural order Legu-

minosæ and furnishing the inodorous Red Sandal-wood or Red Sander's Wood or *Barwood* of the dyer. 47. Oil of *Sassafras-root* (*Oleum Sassafras*, U. S. P.) (*Sassafras*, U. S. P.), sp. gr. 1.094, contains nine-tenths of its weight of *safrol* or *sassafrol*, $C_{10}H_{10}O_2$, also *eugenol* and a small quantity of a terpene. *Sassafras* camphor, $C_{10}H_{10}O_2$, is deposited when the oil is exposed to a low temperature. 48. Oil of *Mustard* (*Oleum Sinapis Volatile*, U. S. P.) is allyl sulphocyanate. (See Index.) If contaminated with alcohol, its sp. gr. is below 1.015. 49. Oil of *Sweet Flag* (*Acorus calamus*) contains the hydrocarbon $C_{10}H_{16}$. The rhizome (*Calamus*, U. S. P.) also contains *acorin*, $C_{36}H_{60}O_6$, a bitter glucoside, and an alkaloid, *calamine*. 50. Oil of common garden *Thyme* (*Thymus vulgaris*, *Oleum Thymi*, U. S. P.) is composed of *cymene* or *cymol* ($C_{10}H_{14}$), *thymene* ($C_{10}H_{16}$), and *thymol* ($C_{10}H_{14}O$). *Thymol*, U. S. P., crystallizes out when oil of thyme or of *ptychotis*, etc. is kept at a low temperature for a day or two. It may also be obtained by shaking the oils with caustic alkali, and treating the separated alkaline liquid with an acid. It may be purified by distillation or by crystallization from alcohol. It would seem that as an antiseptic thymol is far stronger than carbolic acid. 51. Oil of *Turmeric* (*Curcuma longa*) is said by Jackson and Menke to be chiefly an alcohol having the formula $C_{19}H_{27}OH$. They name it *turmerol*. It is a light yellow volatile oil, having the sp. gr. 0.902. It is to this oil that turmeric (hence curry-powder partly) owes its flavor and odor. 52. Oil of *Valerian-root* (*Valerianæ*, U. S. P.) is a mixture of a hydrocarbon, *valerene* or *borneene* ($C_{10}H_{16}$), *valerol* ($C_6H_{10}O$), and (Gersek) 10 per cent. of borneol valerianate. Valerol slowly oxidizes to valerianic acid, known by its smell. A similar change occurs at once if oil of valerian be allowed to fall, drop by drop, on heated caustic potash: $C_6H_{10}O + 3KHO + H_2O = K_2CO_3 + C_4H_9COOK + 3H_2$. By the action of sulphuric acid on the potassium valerianate thus produced valerianic acid is obtained. 53. Oil of *Verbena*, *Lemon-grass* Oil, or *Indian Melissa* Oil, is obtained from *Andropogon citratus* (*Oleum Andropogi Citrati*, P. I.). 54. Oil of *Ginger* (*Zingiber*, B. P.) is, according to Thresh, a complex mixture of hydrocarbons and their oxidation-products; *cymene* ($C_{10}H_{14}$) is present, a terpene, aldehydes, and ethereal salts. (For an analysis of ginger by Thresh, and for papers on "Soluble Essence of Ginger," see the *Pharmaceutical Journal* for August 30 and September 6, 1879, and March 4, 1882.) 55. American Wormseed (*Chenopodium*, U. S. P.) contains a volatile oil (*Oleum Chenopodii*, U. S. P.; p. 417).

Caoutchouc (or *India-rubber*) and *Gutta Percha*.

Caoutchouc is the hardened juice of *Dichopsis Gutta*, *Hevea* (*Siphonia*) *Brasilensis*, *Castilloa elastica*, *Urceola elastica*, *Ficus elastica*, and other plants. Heated moderately with sulphur, it takes up 2 or 3 per cent., and forms *vulcanized india-rubber*; at a higher temperature a hard horny product, termed *ebonite* or *vulcanite*, results. *Gutta Percha* (*Gutta Percha*, B. P.) is the concrete drop or juice of the *percha* (Malay) tree, the *Isonandra gutta*, and of other

Sapotaceous plants. White gutta percha is obtained by precipitating a solution of ordinary gutta percha in chloroform by alcohol, washing the precipitate with alcohol, and finally boiling in water and moulding into the desired form while still hot. The official solution of gutta percha (*Liquor Gutta Percha*, B. P.) is made by digesting *thin* slices of gutta percha in 12 parts by weight of chloroform, and then "fining" by shaking with 1 part of lead carbonate and setting aside till the fluid is clear.

These two elastic substances, in the pure state, are hydrocarbons ($\alpha\text{C}_5\text{H}_4$), usually slightly oxidized. When caoutchouc is distilled a terpene, $\text{C}_{10}\text{H}_{16}$, called cautchin, is obtained.

Official india-rubber (*Elastica*, U. S. P.) is "the prepared milk-juice of various species of *Hevea* (nat. ord. *Euphorbiaceæ*), known in commerce as Para rubber." It melts at about 125°C .

Camphors.

In addition to the stearoptens or camphors already mentioned as being contained in or formed from volatile oils, there is one that is a common article of trade. It is obtained from the wood of *Camphora officinarum*, or camphor-laurel, in Japan (termed, in Europe, Dutch camphor, because imported by the Dutch, and in China known as Formosa camphor), by a rough process of distillation with water, and is purified by resublimation (*Camphora*, U. S. P.). The formula of laurel-camphor is $\text{C}_{10}\text{H}_{16}\text{O}$. Sp. gr. 0.990 to 0.995; melting-point, 175°C .; boiling-point, 205°C . Bromine heated with camphor gives *monobrom-camphor* ($\text{C}_{10}\text{H}_{15}\text{BrO}$) and hydrobromic acid. Monobrom-camphor is camphor in each molecule of which an atom of hydrogen has been displaced by one of bromine. Recrystallized, it occurs in white prisms. The essential oil, from which doubtless camphor is derived by oxidation, is easily obtained from the wood, and is occasionally met with in commerce under the name of *liquid camphor* or *camphor oil*. It contains hydrocarbons resembling terebinthene and citrene, and hydrous camphor ($\text{C}_{10}\text{H}_{16}\text{O}, \text{H}_2\text{O}$) as well as camphor. By exposure to air it becomes oxidized and deposits common camphor. Camphor distilled with phosphoric anhydride yields cymene, $\text{C}_{10}\text{H}_{14}$. There is another kind of camphor, *borneol*, in European markets, less common than laurel-camphor, but highly esteemed by the Chinese; it is obtained from the *Dryobalanops aromatica*, and denominated Sumatra or Borneo camphor. It differs slightly from laurel-camphor in containing more hydrogen, its formula being $\text{C}_{10}\text{H}_{18}\text{O}$. It may be obtained by acting on camphor with hydrogen, the camphor being dissolved in some inert liquid, such as toluene, and sodium added; the sodium forms a compound, $\text{C}_{10}\text{H}_{15}\text{ONa}$, while the hydrogen thus liberated acts on another portion of the camphor, forming borneol, $\text{C}_{10}\text{H}_{17}(\text{OH})$ —a better result being obtained if absolute alcohol is used instead of toluene (Jackson and Menke). It is accompanied in the tree by a volatile oil ($\text{C}_{10}\text{H}_{16}$) isomeric with oil of turpentine. This oil, *borneene*, is also occasionally met with in trade under the name of *liquid camphor* or *camphor oil*, but differs from laurel-camphor oil in not depositing crystals on exposure to air.

The constitution of the camphors is still somewhat doubtful. Camphor is soluble to a slight extent in water (40 grains per gallon, Pooley). The official camphor-water (*Aquæ Camphoræ*, U. S. P.), or *camphor mixture*, is such a solution.

Common camphor and many other of the camphors, oily hydrocarbons, and oxidized hydrocarbons yield *camphoric acid*, $C_8H_{14}(COOH)_2$, and camphoronic acid, $C_7H_8(OH)(COOH)_2$, when attacked by oxidizing agents. Such reactions indicate natural relationships. Camphoric acid is a good antiseptic.

Cantharidin ($C_{10}H_{12}O_4$), the active blistering principle of cantharides (*Cantharis*, U. S. P.) and other vesicating insects (such as *Mylabris cichorii* or *Telini Fly*, P. I., common in India), has most of the properties of a camphor or a stearopten. It slowly crystallizes, from an alcoholic tincture of the beetles, in fusible, volatile, micaceous plates. The following process for the extraction of cantharidin is by Fumouze: Powdered cantharides are macerated with chloroform for twenty-four hours; and this treatment is repeated twice with fresh quantities of solvent, the residue having been well squeezed each time. The collected solutions are then distilled, and the dark-green residue treated with carbon disulphide, which dissolves fatty, resinous, and other matters and precipitates the cantharidin. The precipitate is placed on a filter, washed with carbon disulphide, and recrystallized from chloroform. The same process, omitting the final recrystallization, may be used for the quantitative estimation of cantharidin in cantharides. The average quantity found is from 4 to 5, or occasionally 10 or even 12, parts in 1000. Cantharidin is readily soluble in warm glacial acetic acid (Tichborne), and still more readily in acetic ether or chloroform. Cantharides from which the fat has been removed by petroleum ether yield their cantharidin with great facility.

Massing and Dragendorff consider cantharidin to be an anhydride, and that with the elements of water it forms *cantharidic acid* ($H_2C_{10}H_{12}O_5$). Piccard gives the vapor-density of cantharidin as about 6.5, and its formula $C_{10}H_{12}O_4$. Homolka assigns to it the formula $C_8H_{13}O_2 \cdot CO \cdot COOH$.

Resins, Oleo-Resins, Gum-Resins.

Resins seem to be the oxidized products of terpenes and the allied hydrocarbons; they occur in plants, generally in association with volatile oils. They closely resemble camphors and stearoptens, but are not volatile, and differ from oils and fats mainly in being solid and brittle. For convenience they are classified as resins, oleo-resins, and gum-resins, the distinctions being founded as much on physical as on chemical properties.

Oleo-resins are mixtures of a resin and a volatile oil.

Gum-resins are mixtures of a resin or oleo-resin and gum.

Balsams are commonly described as resins or oleo-resins which yield benzoic and cinnamic acids; they are benzoin (*Benzoinum*, U. S. P.), balsam of Peru (*Balsamum Peruvianum*, U. S. P.), balsam of Tolu (*Balsamum Tolutanum*, U. S. P.), and storax, and are treated of under the respective acids.

Some oleo-resins, containing neither of the above acids, are often termed balsams (e. g. balsam of copaiva and Canada balsam); these will be treated under the head of Oleo-resins.

A physico-chemical method for the identification of the chief resins, gum-resins, and balsams will be found in the *Pharmaceutical Journal* for November 17, 1877.

Resins appear to be somewhat antiseptic. Beer is said never to turn sour in casks lined with Burgundy pitch. The resin of hops has perhaps a similar effect in retarding oxidation of alcohol.

RESINS.*—1. *Resin, rosin, or colophony* (*Resina*, U. S. P.) is the type of this class. Its source is the oleo-resin or true turpentine of the conifers, a body which by distillation yields spirit of turpentine and a residuum of rosin. "Brown" and "white" rosin are met with in trade. The former is the residue of American, the latter of Bordeaux, turpentine (from *Pinus Abies*, etc. and *Pinus maritima* respectively). The chief constituents of brown resin are *pinic acid* ($\text{HC}_{20}\text{H}_{29}\text{O}_2$) and *sylvic acid*, identical in composition, but differing in properties (*vide* Isomerism), the former being soluble and the latter insoluble in cold spirit of wine. White resin or "galipot" is chiefly *pimaric acid*, also isomeric with pinic acid. Pinic acid, cautiously heated, yields *colophononic* or *colopholic acid*. Rosin by destructive distillation yields *resin oil*, the first portion being "pale," the next "blue," and the third "green resin oil." Mixed with other oils, they are used for lubricating purposes and in the manufacture of printing ink. Among the products of the destructive distillation of resin Tichborne has found "*colophononic hydrate*" ($\text{C}_{10}\text{H}_{22}\text{O}_3, \text{H}_2\text{O}$), a white inodorous crystalline substance, and by depriving this of water has obtained white crystalline *colophonine* ($\text{C}_{10}\text{H}_{22}\text{O}_3$). Resin is soluble in oil of turpentine. Contact with sulphuric acid immediately colors it strongly red. It is a constituent of four of the fourteen plasters (*Emplastra*) of the U. S. Pharmacopœia. 2. *Arnica* ($\text{C}_{20}\text{H}_{30}\text{O}_4$), the chief acrid and one of the active principles of *arnica* (*Arnica Flores*, U. S. P.; *Arnica Rhizoma*, B. P.), is a resin, and, probably, a glucoside. 3. *Cannabin*, said to be the active principle of *Indian Hemp* (*Cannabis Indica*, U. S. P.), was obtained in 1846 by T. and H. Smith, and is a resin. Personne in 1857 isolated a volatile oil, also said to possess much medicinal activity, consisting of *cannabene* ($\text{C}_{18}\text{H}_{20}$) and a solid crystalline, "*cannabene hydride*" ($\text{C}_{18}\text{H}_{22}$). A sesquiterpene ($\text{C}_{15}\text{H}_{24}$) is also said to be present. Preobraschensky has stated, and since reasserted, that the active principle is nicotine. Kennedy searched for nicotine by two methods, but found none. Hay found an alkaloid, *tetano-cannabine*; Siebold and Bradbury, also H. F. Smith, an alkaloid termed *cannabinine*. Warden and Waddell, after careful investigations, consider that the active principle of the plant has yet to be isolated. Jahns finds choline present. The native names of Indian hemp—that is, of the cultivated "dried flowering or fruiting tops of the female plants of *Cannabis sativa*"—are *ganga* and

* The student is not expected to remember, but to understand, all that follows respecting the resins.

gunjah. It is chiefly grown in Bengal. *Guaza* is the name of the Bombay product, which includes the wild plant. Both are used for smoking, and form the equivalent of the tobacco luxury of Western nations. *Bhang*, or *sidee*, consists of the dried leaves, fruit, and twigs of the wild plant. Its infusion is drunk, as tea is in Europe and elsewhere. *Hashish*, made from bhang, corresponds to our *Extractum Cannabis Indicae*, B. P. *Charas* or *churras* is a resinous exudation of the plant, and is also used for smoking. All these preparations are stimulating and narcotic. 4. *Capsicum-fruit* contains a resin (p. 425). 5. *Castorin*, a resinous matter, is the name given to the chief constituent of *Castor* (*Castoreum*, B. P.), the dried preputial follicles and included secretion of the beaver (*Castor Fibre*). 6. *Copal*.—The best copal is the exuded resin of trees of extinct forests, and is found beneath the surface of the ground in the neighborhood of existing trees. It appears to be a mixture of acids, but its character is still obscure. 6a. *Doundaké-bark*, an African febrifuge, from *Sarcocephalus esculentus*, owes its activity to resinoid substances, according to Heckel and Schlagdenhauffen. 7. *Dragon's Blood*, a crimson-red resin found as an exudation on the mature fruits of a rotang or rattan palm (*Calamus draco*). It consists of resins having the probable formula $C_{20}H_{24}O_4$ and $C_{20}H_{20}O_4$ (Johnston). 8. *Ergotin* is a very active resinoid constituent of *Ergot* (*Ergota*, U. S. P.), or the sclerotium (compact mycelium or spawn) of *Claviceps purpurea*, produced within the pales and replacing the grain of the common rye, *Secale cereale*. According to Wenzell, ergot contains two alkaloids, *ecboline* and *ergotine*, to the former of which, he says, the activity of ergot is due. Blumberg considers these alkaloids to be identical. Tanret states that an unstable alkaloid termed *ergotinine* occurs in ergot to the extent of 1 per 1000, and that it is accompanied by a camphor; also *ergosterin*, $C_{26}H_{40}O, H_2O$, resembling cholesterin. Dragendorff and Podwissotzki assert that ergot owes most of its activity to *sclerotic* or *sclerotinic acid*, present to the extent of about 4 per cent. Recent investigations seem to show that *cornutine* is an active alkaloid of ergot, associated with ergotinic and sphacelinic acids, picrosclerotine and ergotinine. The activity really seems to be due to a combination of alkaloids and acids, and not to any one constituent, as no principle representing the full activity of ergot has been extracted. The same may be said of a similar therapeutical agent, the root-bark of *Gossypium herbaceum* (*Gossypii Radicis Cortex*, U. S. P.), the activity of which appears to reside in a red resin. Ergot also contains *choline*, which by decomposition may yield trimethylamine. "*Ergotin*" (*Ergotinum*, B. P.) is an alcoholic extract of an aqueous extract of ergot. 9. *Guaiacum Resin* is a mixture of substances. (See Index.) 10. *Jalap Resin*. (See Index.) 11. *Kousso* (*Cusso*, U. S. P., formerly *Brayera*) yields yellow crystals of a resinoid body readily soluble in alkaline liquids, *kosin* or *koussin* ($C_{31}H_{38}O_{10}$). It is, perhaps, an anhydride. 12. *Mastic* (*Mastiche*, U. S. P.) is a resinous exudation obtained by incision from the stem of the mastic or lentisk tree. Nearly nine-tenths of mastic is *mastichic acid* ($C_{20}H_{32}O_3$), a resin soluble in alcohol; the remainder consists of

masticin ($C_{20}H_{32}O$), a tenacious, elastic resin, and a terpene having the formula $C_{10}H_{16}$. 13. *Mezereon* (*Mezereum*, U. S. P.), the dried bark of *Daphne Mezereum*, mezereon, and *Daphne laureola*, spurge laurel, owes its acidity to a resin. 14. *Pepper* contains resin. (See Index.) 15. *Burgundy Pitch* (*Pix Burgundica*, U. S. P.) is the melted and strained exudation from the stem of the spruce fir, *Abies excelsa*. The term Burgundy is a misnomer, the resin never having been collected at or near Burgundy—Finland, and to a smaller extent Baden and Austria, being the countries whence it is derived. Its constituents closely resemble those of common resin. It is often adulterated and imitated by a mixture of resin with palm oil, water, etc., from which it may be readily distinguished by its duller yellow color, highly aromatic odor, greater solubility in alcohol, and almost complete solubility in twice its weight of glacial acetic acid (Hanbury). 16. *Podophyllum Resin*.—In preparing the resin of podophyllum or may-apple (*Resina Podophylli*, U. S. P.), an alcoholic extract of the rhizome and rootlets (*Podophyllum*, U. S. P.) is poured into cold water; the resin is then deposited. This resin is the chief active principle of podophyllum-root. According to Guareschi, podophyllin contains a glucoside resembling convolvulin. Podwissotzki has extracted from podophyllum a little crystalline coloring matter, fat, a bitter crystalline acid, a bitter crystalline neutral principle, and an amorphous acid resin. Kürsten states that the latter yields a crystalline active substance, *podophyllotoxin*, $C_{23}H_{24}O_9$. 17. *Pyrethrin* is the name of the acrid resinous active principle of the root of *Anacyclus pyrethrum*, or *Pellitory-root* (*Pyrethrum*, U. S. P.). According to Buckheim, the action of alkalies breaks it up into piperidine and pyrethric acid. The crystalline poisonous principle obtained by Bellesme from *Pyrethrum carneum*, the powder of which (and *P. roseum*, and especially *P. cinerariæfolium*, or Dalmatian insect powder) is the well-known "insecticide," has not yet been analyzed. 18. The resins of rhubarb have already been alluded to in connection with Chrysophanic Acid. 19. *Rottlerin*, $C_{11}H_{10}O_3$ (*mallotoxin*, $C_{11}H_{10}O_3$, Perkin), is the name given by Anderson to a crystalline resin from kamala (*Kamala*, U. S. P.), the minute glands that cover the capsules of *Rottlera tinctoria*: to this and, apparently, allied resins (*isorottlerin*, A. G. Perkin) kamala owes its activity as an anthelmintic.

OLEO-RESINS.—1. "*Capsicin*," a term suggestive of a definite chemical substance, is a name somewhat unhappily accorded to an indefinite substance, an oleo-resin, obtained by digesting the alcoholic extract of capsicum-fruit (*Capsicum*, U. S. P.) in ether and evaporating the clear ethereal fluid to dryness. Besides volatile oil and resin, capsicum-fruits contain much fatty matter, which Thresh states is chiefly free palmitic acid. (See also *Capsicine* and *Capsaicin*, in Index.) 2. *Copaiva* (*Copaiba*, U. S. P.) is a mixture of essential oil ($C_{15}H_{24}$), *copaivaöl*, $C_{20}H_{32}$ (Strauss), with 2 or more per cent. of brown soft resin, and 30 to 60 of a yellow dark crystalline resin consisting mostly of *copaivic acid* ($C_{20}H_{32}O_2$) with *oxycopaivic acid*, $C_{20}H_{28}O_3$ (Fuhling), and *metacopaivic acid*, $C_{22}H_{34}O_4$

(Strauss). Copaiva, containing about equal parts of this acid and of the oil, heated with a fourth of its weight of the official magnesium carbonate, yields a transparent fluid, owing to the formation of magnesium copaivate and solution of this soap in the essential oil. With an equal weight of the carbonate enough soap is produced to take up the whole of the essential oil and form a mass capable of being rolled into pills. A much smaller quantity of calcined magnesia, as might be expected, effects the same result, but more time, often several days, is required before complete reaction is effected. The *Massa Copaibæ*, U. S. P., is formed from 6 parts of magnesia and 94 of copaiba, and a little water. Quicklime has a similar effect. Perhaps carbonate reacts more quickly because of its fine state of division and admixture of hydrate, in which case calcium and magnesium hydrates may be expected to act better than the calcined preparations, and in much smaller quantity than magnesium carbonate. Copaiva, unlike, 3, *Wood Oil*, or *Gurjun Balsam* (*Dipterocarpi Balsamum*, P. I.), a similar oleo-resin from the *Dipterocarpus turbinatus* (*D. Lævis*, P. I.) is almost entirely soluble in absolute alcohol and in petroleum spirit. Copaiva is often slightly fluorescent; Gurjun balsam is highly fluorescent. The stated analogy of Gurjun balsam to copaiva is borne out by its chemical composition; for by distillation it yields about 40 per cent. of an essential oil identical in composition with oil of copaiva, the non-volatile portion being resinous. The adulteration of copaiva with fixed oil is best detected by heating 20 or 30 drops in a capsule until all essential oil has evaporated. (Turpentine is betrayed by its odor during this evaporation.) The residue, copaiva resin, is brittle if pure, and more or less sticky or soft if fixed oil is present. The limit of brittleness is stated, by Siebold, to be reached when 1 per cent. of oil has been added to the copaiva, that amount preventing the residue being reduced to a fine powder. "On adding 1 drop of copaiba to 19 drops of carbon disulphide, and shaking the mixture with 1 drop of a cold mixture of equal parts of sulphuric and nitric acids, it should not acquire a purplish-red or violet color (absence of Gurjun balsam)."—U. S. P. *Resina Copaibæ*, U. S. P., is the residue left after distilling off the volatile oil from copaiba.

4. Oleo-resin of cubebs (*Oleo-resina Cubebæ*, B. P.) is an ethereal extract of cubebs decanted from waxy matter. (See *Piperine and Oil of Cubebs*.) 5. *Elemi* (*Elemi*, B. P.) is an exudation from a tree growing in the Philippine Islands. It consists of volatile oil ($C_{10}H_{16}$) with 80 or more per cent. of two resins, the one ($C_{20}H_{32}O_2$) soluble in cold alcohol, the other, *Amyrin*, (C_5H_8)₅H₂O, almost insoluble, associated with *Amyric acid* ($(C_5H_8)_4O_4$ (Buri). There is an α and a β amyryl, each having the formula $C_{30}H_{49}OH$ (Vesterberg). It also contains small quantities of two crystalline bodies soluble in water, *Bryoidin*, $(C_5H_8)_3H_2O$, and *Breidin* (Flückiger). The *icacin* of Stenhouse and Groves is either identical with amyryl or perhaps has the formula $(C_6H_8)_9H_2O$. All these bodies are probably hydrous terpenes. 6. *Wood-tar* (*Pix Liquida*, U. S. P.) is mixture of several resinoid and oily bodies (amongst others creasote; see Index) obtained by destructive distillation from the wood of *Pinus sylvestris*

and other pines. When heated, it yields a terebinthinate oil (*Oleum Picis Liquidæ*, U. S. P.) and a residue of *pitch*. (*Earth Pitch*, or *Asphalte*, appears to be a partially oxidized petroleum.) *Oleum Cadinum*, U. S. P., "*Huile de Cade*," or *Juniper Tar Oil*, is the product of the similar destructive distillation of the *Juniperus Oxycedrus*. 7. *Turpentine*.—These oleo-resins have been mentioned in connection with *oil of turpentine*, their volatile, and *resin*, their fixed constituent. 8. *Common Frankincense Thus Americanum*, B. P.) is the concrete turpentine of *Pinus tæda*. 9. *Canada Balsam* (*Terebinthina Canadensis*, B. P.) is largely gathered in the province of Quebec, and is the turpentine or oleo-resin of the balm of Gilead fir (*Abies balsamea*). 10. *Sumbul-root*, from *Ferula Sumbul* (*Sumbul*, U. S. P.), contains 9 per cent. of resin, to which probably it owes its stimulating properties. The resin consists of two parts, one soluble in ether and the other in alcohol, together with valerianic, sumbulic, and sumbuolic acids. By dry distillation it yields a blue oil. 11. *Oleo-resin of Lupulin* (U. S. P.) is an ethereal extract of the yellow glandular powder (*Lupulinum*, B. P.) attached to the small nuts at the base of the scales which form the aggregate fruit of the *Humulus Lupulus*, or hop (*Humulus*, U. S. P.). It contains essential oil of hop (valerol, $C_6H_{10}O$), a terpene, $C_{10}H_{16}$, oxidized oil or resin, bitter extract containing the hop-bitter, lupulinic acid ($C_{32}H_{50}O_7$), and tannic acid. It generally contains a good deal of earthy dust, but should not yield more than 15 per cent. of ash and not more than 30 or 40 per cent. of matter insoluble in ether. *Oleo-resinæ Aspidii, Capsici, Cubeæ, Lupulini, Piperis*, and *Zingiberis* are official in the United States Pharmacopœia. 12. *Pix Canadensis* (*Canada Pitch* or *Hemlock Pitch*) is the concrete juice of *Abies Canadensis*.

GUM-RESINS.—1. *Ammoniacum* (*Ammoniacum*, U. S. P.) is an exudation from the *Dorema Ammoniacum*. It contains nearly 20 per cent. of gum, a little volatile oil, and about 70 of resin ($C_{40}H_{50}O_9$, —Johnston). 2. *Asafoetida* (*Asafoetida*, U. S. P.) formerly spelt *assafoetida*, is a gum-resin obtained by incision from the living root of *Ferula Narthex*. It contains from 50 to 70 per cent. of a resin which is partly *ferulaic acid* ($C_{10}H_{10}O_4$), 25 to 30 per cent. of gum (about two-thirds arabin, one-third bassorin, p. 116), a little vanillin, and 3 to 5 per cent. of volatile oil, which (Semmler) contains two sulphur compounds, $C_{17}H_{14}S$ and $C_{11}H_{20}S_2$, two terpenes, $C_{10}H_{16}$, and a sesquiterpene, $C_{15}H_{24}$. 3. *Euphorbium*, an old drug which is an emetic and purgative gum-resin, contains an amorphous active resin ($C_{20}H_{32}O_4$), crystalline *euphorbon* ($C_{26}H_{44}O_2$) and mucilage (Flückiger). 4. The ordinary or Siam *Gamboge* (*Gambogia*, U. S. P.) of European trade is obtained from the *Garcinia Hanburii*; the gamboge of India (*Cambogia Indica vel Mysoriensis*, P. I.) from *G. pictoria*. When of best quality it contains about 20 per cent. of a gum and 80 to 75 per cent. of a yellow resin termed *gambogic acid* ($C_{20}H_{24}O_4$). 5. *Galbanum* (*Galbanum*, B. P.) contains from 20 to 25 per cent. of gum, about 65 per cent. of resin ($C_{40}H_{54}O_7$), and 3 or 4 per cent. of volatile oil. Moistened with alcohol, and then with hydrochloric acid, galbanum yields a purple color, due, probably, to the produc-

tion and oxidation of resorcin. Galbanum heated for some time to 212° F. with hydrochloric acid, the liquid separated and shaken with ether or chloroform, and the latter evaporated, yields somewhat less than 1 per cent. of colorless acicular crystals of *umbelliferone* ($C_9H_6O_3$). "Umbelliferone is soluble in water; its solution exhibits, especially on addition of an alkali, a brilliant blue fluorescence which is destroyed by an acid. If a small fragment of galbanum is immersed in water, no fluorescence is observed, but it is immediately produced by a drop of ammonia. The same phenomenon takes place with asafoetida, and in a slight degree with ammoniacum; it is probably due to traces of umbelliferone pre-existing in those drugs. Umbelliferone is also produced from many other aromatic umbelliferous plants, as *Angelica*, *Levisticum*, and *Meum*, when their respective resins are submitted to dry distillation; also from the resin of *Daphne mezereum*. The fluorescence of umbelliferone may be beautifully shown by dipping some bibulous paper into water which has stood for an hour or two on lumps of galbanum, and drying it. A strip of this paper, placed in a test-tube of water with a drop of ammonia, will give a superb blue solution, instantly losing its color on the addition of a drop of hydrochloric acid" (Flückiger and Hanbury). 6. *Myrrh* (*Myrrha*, U. S. P.), an exudation from the stem of *Balsamodendron myrrha*, contains about half its weight of soluble arabinoid gum, 10 per cent. of insoluble gum (probably bassorin), $2\frac{1}{2}$ of volatile oil isomeric with thymol and carvol (Köhler), and about 25 per cent. of resin (myrrhic acid). (For a note by R. H. Parker on the spurious gums imported with myrrh see the *Pharmaceutical Journal* for July 17, 1880.) 7. *Olibanum* (P. I.). *Thus masculum* or *Arabian Frankincense* (from various species of *Boswellia*) is about one-third gum and nearly two-thirds resin ($C_{40}H_{30}O_6$), with a little hydrocarbon ($C_{10}H_{16}$) and oxidized hydrocarbon volatile oils. It has always been an important ingredient of *incense*—myrrh, storax, benzoin, and such fragrant combustible resinous substances being other constituents. 8. *Scammony*. (See Index.)

Gum-resins need only to be finely powdered and rubbed in a mortar with water to yield a medicinal *emulsion* in which the fine particles of resin are held in suspension by the aqueous solution of gum.

QUESTIONS AND EXERCISES.

What are the general chemical characters of volatile oils?—How do volatile oils usually differ chemically from fixed oils?—Describe the usual process by which volatile oils are obtained.—How does natural turpentine differ from turpentine of trade?—With what object is commercial turpentine rectified?—What is the chemical nature of india-rubber and gutta percha?—How is india-rubber *vulcanized* and converted into *ebonite* or *vulcanite*?—Mention the difference in composition between the volatile oils of *Anthemis nobilis* and *Matricaria Chamomilla*.—Give the systematic name for oil of horseradish.—State the general composition of the oil of lemon, lime, bergamot, citron, and cedra.—Name the constituents of oil of cloves.—In what respect does otto of roses differ from other oils?—What class of substances forms the chief part of oil of rue?—How is camphor oil related to camphor?—In what respects do Borneo or Sumatra

camphor and camphor oil differ from the corresponding products of Japan and China?—How may borneol be artificially prepared?—How do resins occur in nature?—Distinguish between resins and camphors.—Mention the points of difference of resins, oleo-resins, gum-resins, and balsams.—Name the sources of common resin or rosin.—Enumerate some official articles of which the active constituents are resins.—Give the distinguishing characters of Burgundy pitch.—What is the average proportion of oil and of resin in the so-called balsam of copaiva?—Explain the effect of magnesium carbonate, magnesia, and lime on copaiva.—Why do ammoniacum, asafoetida, gamboge, galbanum, myrrh, and similar substances give an emulsion by mere trituration with water?

THE BENZENE SERIES OF HYDROCARBONS.

The Benzene or Aromatic Series, C_nH_{2n-6} .—This series is of great general interest. It yields, like other families of hydrocarbons, alcohols, haloid derivatives, aldehydes, acids, etc., obtained, however, as a rule by special rather than general methods. Just as the consecutive members of the paraffin series of hydrocarbons may be regarded as derived by the displacement of a hydrogen atom of the previous member by the methyl (CH_3) group, or of a hydrogen atom in methane by a paraffin radical, so the consecutive members of the benzene series may for convenience of study be viewed as obtained by the displacement of hydrogen atoms in *benzene* by paraffin radicals, as in the following examples:

Benzene or Phenöene, C_6H_6 .

Toluene, Benzöene, or Methylphenöene, C_7H_8 or $C_6H_5 \cdot CH_3$.

Xylene or Dimethylphenöene, C_8H_{10} or $C_6H_4 \cdot (CH_3)_2$.

Mesitylene or Trimethylphenöene, C_9H_{12} or $C_6H_3 \cdot (CH_3)_3$.

Cymene or Methylpropylphenöene, $C_{10}H_{14}$ or $C_6H_4 \cdot CH_3 \cdot C_3H_7$.

It is, perhaps, desirable, as suggested by Odling, to designate the first member of this series by the name *phenöene* rather than *benzene*, as its hydrate is termed *phenol*, and its derivatives *phenyls*; e. g. phenylamine. Toluene (first obtained from balsam of Tolu, hence the name) then becomes benzöene; from it benzoic acid is derived.

The members of the benzene series are *unsaturated* hydrocarbons. A molecule of benzene itself readily absorbs two, four, or six atoms of chlorine, these being added on to the benzene, forming what are termed *additive* compounds, as distinguished from the true substitutional compounds, in which the hydrogen atoms in benzene are actually substituted by chlorine, bromine, etc. The derivatives of benzene may more or less readily be reconverted into benzene, a fact supporting the close structural or constitutional relationship between the many benzenoid bodies.

Bodies having an aromatic odor are somewhat characteristic of the benzene series, hence the latter was originally termed the *aromatic series* of organic compounds.

Benzene, or Benzol.

Benzene or Phenöene, C_6H_6 (commercially known as *Benzol*).*—

* Care must be taken to distinguish between benzene, C_6H_6 , and benzin, 19 *

Commercially it is obtained from the portion of coal-tar boiling below 100°C . It is partially purified by shaking successively with sulphuric acid, water, and caustic soda, and then redistilling, the product still containing large quantities of toluene and other impurities. If pure benzol is required, the liquid must be subjected to a freezing mixture, when the benzol crystallizes out, leaving some impurities in solution; the crystals are well drained. Bromine is then added to the liquid resulting from the melting of the crystals until a permanent coloration results. The liquid is again washed with caustic soda, and distilled. Benzene boils at 81°C . It may be artificially produced by heating benzoic acid with lime or by passing acetylene through red-hot tubes. It is a colorless, limpid, refractive liquid, having a specific gravity of 0.899 at 0°C . It is a powerful solvent of grease, and under the name of "Benzene Collas" was introduced by M. Collas in 1848 for cleansing purposes.

Benzene, when acted on by chlorine and bromine in the presence of a little iodine, forms all derivatives from monochloro- and monobromo-benzene ($\text{C}_6\text{H}_5\text{Cl}$ and $\text{C}_6\text{H}_5\text{Br}$) to hexachloro- and hexabromobenzene (C_6Cl_6 and C_6Br_6). It also forms iodine and fluorine derivatives, nitro-derivatives, etc.

Nitrobenzene (nitrobenzol, artificial oil of bitter almonds, or essence of mirbane), $\text{C}_6\text{H}_5\text{NO}_2$, is obtained by mixing fuming nitric acid or a mixture of nitric and sulphuric acids with benzene, the vessel being kept cool by immersion in water. It is a yellow liquid, heavier than water, having a strong odor of oil of bitter almonds, though of very different nature. (*Vide* "Oil of Bitter Almonds" in Index.) When acted on by nascent hydrogen it yields aniline.

Aniline or *Phenylamine* or *Amidobenzene*, $\text{C}_6\text{H}_5\text{NH}_2$.*—Mix 13 parts of iron filings, 7 or 8 of the ordinary acetic acid, and 13 of nitrobenzene in a large flask (with an upright condenser) placed in a water-bath, and set the whole aside for some time. After the mixture has digested for several hours the supernatant liquid is poured off from the deposit of iron filings and distilled in a current of steam. By this method the nitrobenzene yields, first, aniline, distilled over as a yellow oil, and afterward a red oil, which is a mixture of azobenzene ($\text{C}_6\text{H}_5 - \text{N} = \text{N} - \text{C}_6\text{H}_5$), hydrazobenzene ($\text{C}_6\text{H}_5 - \text{NH} - \text{NH} - \text{C}_6\text{H}_5$), and azoxybenzene ($\text{C}_6\text{H}_5 - \text{N} \diagdown \text{O} \diagup \text{N} - \text{C}_6\text{H}_5$).

petroleum ether, benzolin, etc. (*Petroleum Spirit*, B. P.), which are mixtures of paraffin hydrocarbons of lower boiling-points. *Benzin* (U. S. P.), $\text{C}_5\text{H}_{12} : \text{C}_6\text{H}_{14}$, and other hydrocarbons of the paraffin series having the boiling-point of 122° – 140°F ., require six times the bulk of alcohol for solution, whereas benzene, C_6H_6 , dissolves in less than its own bulk. Specific gravity of benzene, about 0.850; of benzin, about 0.700.

* Aniline may be obtained from indigo, hence its name, *anil* being Portuguese for indigo.


Aniline, $C_6H_5NH_2$ (mixed with toluidine, $C_7H_7NH_2$), when acted on by arsenic acid or chlorinated lime produces *rosaniline*, $C_{20}H_{19}N_3$, whose salts and derivatives form most of the well-known aniline colors.

Constitution of Amines.—Amines are usually viewed as derivatives of ammonia, one, two, or three atoms of hydrogen being replaced by one, two, or three univalent organic radicals or equivalents of radicals of higher quantivalence. The products were formerly known as amidogen (NH_2) bases, imidogen (NH) bases, and nitrile (N) bases, but are now termed primary, secondary, and tertiary amines. The class includes certain alkaloids.


Amides result when NH_2 displaces OH in acids. Acetic acid = $CH_3 \cdot CO \cdot OH$; hence *acetamide* = $CH_3 \cdot CO \cdot NH_2$. Aniline boiled with strong acetic acid yields *phenyl-acetamide*, or *acetanilid* (*Acetanilidum*, U. S. P.) or "*antifebrin*," a febrifuge, $C_6H_5 \cdot NH \cdot C_2H_3O$ (a rival of "*antipyrine*," $C_{11}H_{12}N_2O$, *phenyl dimethyl-pyrazolon*, $C_6H_5 \cdot (CH_3)_2C_3HN_2O$ (*Phenazonum*, B. P.), from phenylhydrazine or aniline and aceto-acetic ether). *Monobrom-acetanilide*, $C_6H_4Br \cdot NH \cdot C_2H_3O$, is a sedative and febrifuge. *Phenacetin* (*Phenacetinum*, B. P.), or *para-acetphenetidin*, $C_6H_4 \cdot OC_2H_5 \cdot NH \cdot C_2H_3O$, is another febrifuge. *Phenecoll*, *amido-acet-phenetidin*, $C_6H_4 \cdot OC_2H_5 \cdot NHCOCH_2NH_2$, *Para-phenetol-carbamide*, or *dulcin*, is a body having a very powerful sweet taste, and proposed for use, like saccharin, in place of sugar.

Acetanilid occurs in "white, shining, micaceous, crystalline laminae or a crystalline powder, odorless, having a faintly burning taste, permanent in the air. Soluble at $15^\circ C.$ ($59^\circ F.$), in 194 parts of water and in 5 parts of alcohol in 18 parts of boiling water and in 0.4 part of boiling alcohol; also soluble in 18 parts of ether, and easily soluble in chloroform. Heated to $113^\circ C.$ ($235.4^\circ F.$), it melts. It is neutral to litmus-paper. On heating about 0.1 gram. with a few cc. of concentrated solution (1 in 4) of potassium or sodium hydrate the characteristic odor of aniline becomes noticeable. On now adding chloroform and again heating, the disagreeable odor of isonitril (which is poisonous) is evolved. On boiling 0.1 gram. for several minutes with 2 cc. of hydrochloric acid, a clear solution results, which, when mixed with 3 cc. of a 5 per cent. aqueous solution of carbolic acid, and afterward with 5 cc. of a filtered, saturated solution of chlorinated lime (*Calx chlorata*), acquires a brownish-red color, becoming blue upon supersaturation with ammonia. A cold saturated aqueous solution, added to ferric chloride, should not affect the color of the latter (absence of aniline salts and various allied substances)." — *U. S. P.*

Toluene, *Benzöene*, *Methyl-phenöene*, or *Methyl-benzene* (commercially known as *Toluol*), $C_6H_5CH_3$, forms the principal portion of coal-tar, boiling between 100° and $120^\circ C.$; it may be synthetically made by acting on monochlorobenzene and iodomethane by sodium.




It is also obtained by the dry distillation of Tolu balsam. It is an inflammable, refractive liquid, boiling at $111^\circ C.$ It may be directly oxidized to benzoic acid.



Having both a phenyl (C_6H_5) and a methyl (CH_3) group in its molecule, it forms two sets of isomeric derivatives—one (*a*) in which, by acting on toluene in the cold, the atoms of hydrogen are displaced in the phenyl group, and the other (*b*) by acting on boiling toluene, in which the atoms of hydrogen in the methyl group are displaced.*

- a* { Toly chloride, or methylmonochlorobenzene, $C_6H_4Cl \cdot CH_3$.
 Toly dichloride, or methyldichlorobenzene, $C_6H_3Cl_2 \cdot CH_3$.
 Toly trichloride, or methyltrichlorobenzene, $C_6H_2Cl_3 \cdot CH_3$.
b { Monochloromethylbenzene, or benzylchloride, $C_6H_5 \cdot CH_2Cl$.
 Dichloromethylbenzene, or benzyl dichloride, $C_6H_5 \cdot CHCl_2$.
 Trichloromethylbenzene, or benzyl trichloride, $C_6H_5 \cdot CCl_3$.


Dichloromethylbenzene, when acted on by glacial acetic acid and zinc chloride and water, produces benzoic aldehyde, C_6H_5COH (Jacobson). By acting on trichloromethylbenzene by water in sealed tubes benzoic acid results.




Cymene, $C_{10}H_{14}$.—Propylmethylbenzene, $C_6H_4(CH_3)(C_3H_7)$, occurs in several volatile oils, and is readily obtained by the removal of hydrogen from the terpenes ($C_{10}H_{16}$) of those oils. Many of the members of the benzene series have an aromatic odor, hence the synonym *aromatic series*.

CONSTITUTION OF THE BENZENE SERIES.

The fact that benzene forms three additive derivatives with chlorine, $C_6H_6Cl_2$, $C_6H_6Cl_4$, and $C_6H_6Cl_6$, one molecule uniting with not more than six atoms of chlorine, and that it affords no isomeric monosubstitution derivatives (but only one toluene, $C_6H_5CH_3$, one benzoic acid, C_6H_5COOH , and one only of all such derivatives, led Kekulé to represent benzene by the following figure (*a*), in which each atom of carbon is assumed to be "linked" to adjacent atoms of carbon by three-fourths of its affinity, the remaining fourth of its attraction being exerted toward the equivalent attraction of another atom, thus (Fig. *a*):

Fig. *a*.

Benzene.

Fig. *d*.Fig. *b*.Fig. *e*.Fig. *c*.

Hexachlorobenzene.

* "Benzyl" is the name given to the derivatives of benzene when substitution takes place in the methane nucleus, "tolyl" when in the phenylene nucleus.

In a *monosubstitution-derivative* such as chlorobenzene, C_6H_5Cl , no matter where the atom of chlorine be placed, it bears the same relation to the atoms of hydrogen; hence there can be *only one variety* of such a derivative. The experimental evidence of the truth of this inference is as follows: Displace H in benzene by another radical X, and obtain C_6H_5X . In the latter displace H by Y, and obtain C_6H_4XY . Now displace X by H, and obtain C_6H_5Y . Lastly displace Y by X, and obtain C_6H_5X . The first C_6H_5X and the last C_6H_5X are identical in properties, yet manifestly the latter X is in a different position to the first X: whence we conclude that actual position matters nothing if relative position is unchanged. Such hydrocarbons are *symmetrical*. Such mono-X compounds are *unsymmetrical*. Further displacements of H by X in C_6H_5X result in *more than one variety* of C_6H_4XX . In dichlorobenzene, $C_6H_4Cl_2$, the atoms of chlorine may (representing, for the moment, benzene by a hexagonal figure (*b*), and assuming that the carbon atoms are at the angles) be placed either at 1 and 2, 1 and 3, or 1 and 4, the chlorine atoms being either near to each other, separated by one carbon atom or by two carbon atoms. So with other *di-derivatives*. In trichlorobenzene, $C_6H_3Cl_3$, the atoms of chlorine may be placed at 1, 2, and 3, 1, 2, and 4, or 1, 3, and 5; 1, 2, and 4 being the same as 1, 3, and 4; 1, 2, and 3 the same as 1, 6, and 5, etc.; that is to say, the chlorine atoms must all three be near to each other, or two near to each other and one be separated, or all three be separated as far from each other as possible in the molecule. So with other *tri-derivatives*. Hence, theoretically, there can only be three isomeric di- and trichlorobenzenes; which has been verified by experiment. (For other illustrations, and for nomenclature, see p. 445.)

In the *additive compounds* a second quarter of the affinity of the carbon atoms for each other is freed, so to say, for exertion toward the added chlorine atoms (Fig. *c*). The mind may be aided in forming an idea of the constitution of molecules like benzene by other figures, such as the prism (*d*) or wheel-like hexagon (*e*)—devices all of which embody the idea of the interdependence of each atom of the molecule on every other atom, the idea conveyed by the linkages, not only of a series of links of an open chain, as pictured on previous pages, but of a chain without terminal links, a continuous or closed or endless chain—the open chain with its ends linked together, and even sometimes also having cross linkages.

OTHER SERIES OF HYDROCARBONS.

The Naphthalene Series, C_nH_{2n-12} . 

Naphthalene ($C_{10}H_8$; synonyms, *Naphtalin*, *Naphtalene*) is the chief member of the series. It is in condensed benzene nuclei (Erlenmeyer and Graebe, 1866). It is a volatile white crystalline body, in shining laminae, having a strong characteristic odor, insoluble in water, very soluble in boiling alcohol, ether, chloroform, carbon bisulphide, and the fixed and volatile oils. Naphtalin is now

official (*Naphtalinum*, U. S. P.), and is much used for increasing the luminosity of coal-gas. Sulphuric acid should not be discolored by it, and if heated on a water-bath for five minutes should not acquire more than a pale-reddish tint (absence of contamination derived from coal-tar).

Naphthalene by oxidation yields phthalic acid, $C_6H_4(COOH)_2$, the anhydride of which, $C_6H_4(CO)_2O$, fused with phenol, forms *phenolphthalein* (U. S. P.), an alkalimetric indicator. With other phenols various colored bodies are produced; for example, with resorcinol *fluorescin*, which, treated with bromine, gives *eosin*. Such bodies are termed *phthaleins*. Of the two *naphthyl-alcohols*, or *naphthols*, or *monoxynaphthalenes*, $C_{10}H_7(OH)$, α and β , the latter is the well-known powerful antiseptic.

Beta-naphthol (*Naphtol*, U. S. P.) crystallizes in colorless volatile leaflets, very soluble in boiling water and alcohol, and in cold chloroform, ether, and in solutions of caustic alkalies. Ferric chloride imparts a greenish color to the solution of β -naphthols, dinaphthol, $C_{20}H_{12}(OH)_2$, separating out.

"Naphthol should dissolve in 50 parts of ammonia-water without leaving a residue (absence of naphthalin), and the solution should not have a deeper tint than pale yellow (absence of various other organic impurities). If 0.1 gm. be mixed, in a test-tube, with 1 drop of syrup and 5 cc. of water, and about 3 cc. of concentrated sulphuric acid be then poured into the tube held in a slanting position, so that the liquids may form separate layers, a yellowish-brown color will appear at the zone of contact, which becomes darker on standing (absence of, and distinction from, alpha-naphthol, which produces at once a crimson color, turning deep blue in the upper part of the zone on standing)." — U. S. P.

The Anthracene Series, C_nH_{2n-18} .



Anthracene, $C_{14}H_{10}$, is the only noteworthy member of this series, its importance being due to the fact that *artificial madder*, or *alizarin*, is formed from it by the following reactions: Anthracene is first oxidized to anthraquinone by the influence of nascent oxygen. By acting on anthraquinone with fuming sulphuric acid it is easily converted into a derivative, which yields potassium alizarate when fused with caustic potash.

Chrysophanic Acid ($CH_3C_{14}H_5(OH)_2O_2$) and the *aloins* are related to anthraquinone, chrysophanic acid being a dihydroxy-derivative of methylanthraquinone, and the *aloins* ($C_{16}H_{18}O_7$) yielding on oxidation aloxanthin or tetrahydroxy-methylanthraquinone.

Aloins.

Aloins.—The aloes of pharmacy (*Aloe Barbadosensis*, U. S. P., and *Aloe Socotrina*, U. S. P.) is an evaporated juice, doubtless much altered by the temperature to which it is subjected.

Aloe Purificata, U. S. P., is the evaporated alcoholic extract of Socotrine aloes. It contains a yellow crystalline substance, *Aloinum*,

(U. S. P.), slightly varying chemically, but not medicinally, as derived from the respective species of aloes. Aloin is slightly soluble in cold water or spirit, but readily soluble in the hot fluids. Dissolved in alkalis, it rapidly decomposes, absorbing oxygen, but it is only slowly altered in neutral or acidified solutions.

Aloin may readily be obtained from either kind of aloes by warming with three or four times its weight of amyl alcohol, pouring off the solution, allowing it to stand for a few hours to crystallize, and washing the deposited aloin with ether or carbon bisulphide to remove resinoid matters. It forms in "minute acicular crystals or a microcrystalline powder, varying in color from yellow to yellowish-brown, odorless or possessing a slight odor of aloes, of a characteristic bitter taste, and permanent in the air."—*U. S. P.*

Tests.—Ferric chloride turns an aqueous solution of aloin greenish-black. Basic lead acetate slowly precipitates aloin.

Barbaloin, U. S. P.—This substance, first obtained by T. and H. Smith, occurs in minute crystals in Barbadoes aloes. It yields, by the action of bromine and chlorine, substitution-compounds. Nitric acid dropped upon it produces a red color which soon fades. Boiled for some time with strong nitric acid, barbaloin gives, together with oxalic and picric acids, a yellow substance, chrysammic acid, which furnishes beautiful red salts (Tilden). Anthracene ($C_{14}H_{10}$) has been obtained by deoxidation of barbaloin.

Nataloin.—This body was discovered by Flückiger in Natal aloes. It crystallizes readily in rectangular plates, either from spirit or from water. No bromine or chlorine substitution-derivatives have yet been found, but an acetyl compound has been analyzed (Tilden). Nataloin moistened with nitric acid gives a red coloration which does not fade. When boiled with nitric acid it yields no chrysammic acid, but only oxalic and picric acids.

Socaloin, U. S. P., or *Zanaloïn*.—Histed and Flückiger have shown that Socotrine or Zanzibar aloes yields an aloin distinct from those just described. It forms tufted acicular prisms. Nitric acid scarcely alters the color of socaloin (difference from barbaloin). Neither socaloin nor barbaloin affords any color when vapor from a glass rod moistened with nitric acid is brought near to a drop of oil of vitriol containing a minute fragment of the aloin, while nataloin gives rise to a blue coloration.

Analysis.—To aloin or powdered aloes on a white plate add strong nitric acid. *No color* = socaloin. *Crimson color* = nataloin or barbaloin. To another portion add strong sulphuric acid and vapor of nitric acid. *A blue color* = nataloin. *No blue color* = barbaloin.

(For other methods of distinguishing the aloins and other allied bodies see a paper on Aloins by Cripps and Dymond, *Pharmaceutical Journal*, February 7, 1885.)

E. von Sommaruga and Egger ("Pharmacographia") arrived at the conclusion that the aloins form an homologous series, and that they have the composition indicated in the following formulæ: Socaloin, $C_{15}H_{16}O_7$; nataloin, $C_{16}H_{18}O_7$; barbaloin, $C_{17}H_{20}O_7$. Tilden's subsequent experiments indicate, however, that barbaloin ($C_{16}H_{18}O_7$)

and socaloin ($C_{16}H_{18}O_7$) are isomeric in the anhydrous state, but that socaloin and its derivatives in the hydrous condition contain more water of crystallization than barbaloin. Nataloin ($C_{16}H_{18}O_7$) seems to be isomeric with the others, but is less soluble, and does not yield either chrysammic acid or chloro- or bromo-derivatives ($C_{16}H_{15}Cl_3O_7$; $C_{16}H_{15}Br_3O_7$). The acetyl-derivatives appear to have the formula $C_{16}H_{15}(C_2H_3O)_3O_7$. According to Grænewold, the formula for the aloin from the aloes of Barbadoes and Curaçoa is $C_{16}H_{16}O_7$, and that from Natal aloes $C_{24}H_{26}O_{10}$.

QUESTIONS AND EXERCISES.

What is the formula of benzene?—How is it artificially and commercially prepared?—Draw out an equation explanatory of the production of aniline.—What is the relation between toluene and benzoic acid?—Give the formulæ of naphthalene and anthracene.—Explain by equations the production of alizarin or artificial madder.—Give tests for distinguishing the aloins.

The student is referred to the accompanying table for a general view of the relations of four series of hydrocarbons (the paraffin, benzene, naphthalene, and anthracene series) to each other. Three members of the paraffin series are shown, two of the benzene series, and one each of the naphthalene and anthracene series. Beneath each hydrocarbon are given its chief derivatives. A glance along the table shows the relations of the derivatives to each other.

ALCOHOLS.

Alcohols are those bodies in which one or more hydrogen atoms of the hydrocarbons are displaced by one or more hydroxyl (OH) groups, forming (a) monhydroxyl derivatives, (b) dihydroxyl derivatives, etc.; they are, in fact, hydrates of unsaturated or radical hydrocarbons, just as caustic potash and slaked lime are potassium and calcium hydrates, thus:

C_2H_5OH Ethyl hydrate.	KHO, or KOH Potassium hydrate.
$C_2H_4(OH)_2$ Ethylene hydrate, or glycol.	$Ca(HO)_2$ Calcium hydrate.
$C_3H_5(OH)_3$ Glyceric hydrate, or glycerin.	$Bi(OH)_3$ Bismuth hydrate.

a. Monhydroxyl Derivatives of Paraffins.

The Ethylic Series of Alcohols, $C_nH_{2n+1}OH$.—The alcohols, or *carbinols* (Kolbe), are *primary*, *secondary*, or *tertiary* according as one, two, or three atoms of hydrogen in the first (or hydrogen) member of the series (methyl alcohol or *carbinol* itself, CH_3OH) are

To

Hy

Al

As

M

D

T

To face page 437.

SYNOPTICAL TABLE SHOWING THE RELATIONS BETWEEN THE PRINCIPAL MEMBERS OF THE PARAFFIN, BENZENE, NAPHTHALENE, AND ANTHRACENE SERIES OF HYDROCARBONS, AND BETWEEN THEIR DERIVATIVES.


HYDROCARBON ...	CH_4	Methane.	C_2H_6 or CH_3CH_3	Ethane or Methyl-methane	C_3H_8 or $\text{C}_2\text{H}_5\text{CH}_3$	Propane or Ethyl-methane	C_6H_6	Phenol or Benzene	C_7H_8 or $\text{C}_6\text{H}_5\text{CH}_3$	Benzoene, Toluene or Methylphenol	C_{10}H_8	Naphthalene	$\text{C}_{14}\text{H}_{10}$	Anthracene
ALKYL SALT, OR ETHERAL SALT, ETC.	CH_3Cl	Methyl chloride or Chloromethane	$\text{C}_2\text{H}_5\text{Cl}$ or $\text{CH}_3\text{CH}_2\text{Cl}$	Ethyl chloride or Chloroethane	$\text{C}_3\text{H}_7\text{Cl}$ or $\text{C}_2\text{H}_5\text{CH}_2\text{Cl}$	Propyl chloride or Chloropropane	$\text{C}_6\text{H}_5\text{Cl}$	Phenyl chloride or Chlorobenzene	$\text{C}_7\text{H}_7\text{Cl}$ or $\text{C}_6\text{H}_5\text{CH}_2\text{Cl}$	Chlorotoluenes	$\text{C}_{10}\text{H}_7\text{Cl}$	Naphthalene chloride or Chloronaphthalene	$\text{C}_{14}\text{H}_9\text{Cl}$	Anthracene chloride or Chloroanthracene
AMINE ...	CH_3NH_2	Methylamine	$\text{C}_2\text{H}_5\text{NH}_2$ or $\text{CH}_3\text{CH}_2\text{NH}_2$	Ethylamine	$\text{C}_3\text{H}_7\text{NH}_2$ or $\text{C}_2\text{H}_5\text{CH}_2\text{NH}_2$	Propylamine	$\text{C}_6\text{H}_5\text{NH}_2$	Phenylamine or Aniline	$\text{C}_7\text{H}_7\text{NH}_2$ or $\text{C}_6\text{H}_5\text{CH}_2\text{NH}_2$	Tolylamine or Benzylamine	$\text{C}_{10}\text{H}_7\text{NH}_2$	Naphthalamine		
MONOHYDRIC ALCOHOL	CH_3OH or HCH_2OH	Methyl alcohol or Carbinol	$\text{C}_2\text{H}_5\text{OH}$ or $\text{CH}_3\text{CH}_2\text{OH}$	Ethyl alcohol or Methylcarbinol	$\text{C}_3\text{H}_7\text{OH}$ or $\text{C}_2\text{H}_5\text{CH}_2\text{OH}$ or $(\text{CH}_3)_2\text{CH}_2\text{OH}$	Primary Propyl Alcohol or Ethyl carbinol or Secondary Propyl Alcohol or Dimethyl carbinol	$\text{C}_6\text{H}_5\text{OH}$	Phenyl alcohol or Phenol	$\text{C}_7\text{H}_7\text{OH}$ or $\text{C}_6\text{H}_5\text{CH}_2\text{OH}$	Tolyl alcohol or Cresol or Benzyl alcohol or Phenyl carbinol	$\text{C}_{10}\text{H}_7\text{OH}$	Naphthyl alcohol or Naphthol	$\text{C}_{14}\text{H}_9\text{OH}$	Anthracene alcohol or Anthrol
DIHYDRIC ALCOHOL			$\text{C}_2\text{H}_4(\text{OH})_2$ or $\text{CH}_2\text{OH}\cdot\text{CH}_2\text{OH}$	Ethylene glycol or Glycol	$\text{C}_3\text{H}_6(\text{OH})_2$ or $\text{C}_2\text{H}_4\text{OH}\cdot\text{CH}_2\text{OH}$	Propylene glycol	$\text{C}_6\text{H}_4(\text{OH})_2$	Pyrocatechin Resorcin Hydroquinone	$\text{C}_7\text{H}_6(\text{OH})_2$ or $\text{C}_6\text{H}_4\text{OH}\cdot\text{CH}_2\text{OH}$	Salicyl alcohol, Saligenol or Hydroxybenzyl alcohol	$\text{C}_{10}\text{H}_6(\text{OH})_2$	Naphthaquinol or Dihydroxynaphthalene	$\text{C}_{14}\text{H}_8(\text{OH})_2$	Anthraquinol or Dihydroxyanthracene
TRIHYDRIC ALCOHOL					$\text{C}_3\text{H}_5(\text{OH})_3$ or $\text{C}_2\text{H}_3(\text{OH})_2\text{CH}_2\text{OH}$	Glycerol or Glycerin	$\text{C}_6\text{H}_3(\text{OH})_3$	Pyrogallol or Pyrogallie acid			$\text{C}_{10}\text{H}_5(\text{OH})_3$	Trihydroxynaphthalene		
ALDEHYDE	CH_2O or HCO_2H	Formic aldehyde	$\text{C}_2\text{H}_4\text{O}$ or $\text{CH}_3\text{CO}_2\text{H}$	Acetic aldehyde	$\text{C}_3\text{H}_6\text{O}$ or $\text{C}_2\text{H}_5\text{CO}_2\text{H}$	Propionic aldehyde			$\text{C}_7\text{H}_6\text{O}$ or $\text{C}_6\text{H}_5\text{CO}_2\text{H}$	Benzoic aldehyde				
KETONE					$\text{C}_3\text{H}_6\text{O}$ or $\text{CH}_3\text{CO}_2\text{CH}_3$	Acetone or Dimethyl ketone								
NITRILE	HCN	Acetonitrile or Methyl cyanide	$\text{C}_2\text{H}_3\text{CN}$ or $\text{CH}_3\text{CH}_2\text{CN}$	Propionitrile or Ethyl cyanide	$\text{C}_3\text{H}_5\text{CN}$ or $\text{C}_2\text{H}_5\text{CH}_2\text{CN}$	Butyronitrile or Propyl cyanide (primary)	$\text{C}_6\text{H}_5\text{CN}$	Benzonitrile or Phenyl cyanide	$\text{C}_7\text{H}_7\text{CN}$ or $\text{C}_6\text{H}_5\text{CH}_2\text{CN}$	Benzyl cyanide	$\text{C}_{10}\text{H}_7\text{CN}$	Naphthomitrile or Naphthyl cyanide		
MONOBASIC ACID (Monohydric)	HCO_2OH	Formic acid	$\text{C}_2\text{H}_4\text{O}_2$ or $\text{CH}_3\text{CO}_2\text{OH}$	Acetic acid or Methylformic acid	$\text{C}_3\text{H}_6\text{O}_2$ or $\text{C}_2\text{H}_5\text{CO}_2\text{OH}$	Propionic acid or Ethylformic acid			$\text{C}_7\text{H}_6\text{O}_2$ or $\text{C}_6\text{H}_5\text{CO}_2\text{OH}$	Benzoic acid or Phenylformic acid				
MONOBASIC ACID (Dihydric)	$\text{OH}\cdot\text{CO}_2\text{OH}$	Hydroxyformic acid or Carbonic acid	$\text{C}_2\text{H}_4\text{O}_3$ or $\text{CH}_2\text{OH}\cdot\text{CO}_2\text{OH}$	Glycollic acid or Hydroxyacetic acid	$\text{C}_3\text{H}_6\text{O}_3$ or $\text{C}_2\text{H}_4(\text{OH})_2\text{CO}_2\text{OH}$	Lactic acid or Hydroxypropionic acid			$\text{C}_7\text{H}_6\text{O}_3$ or $\text{C}_6\text{H}_4(\text{OH})_2\text{CO}_2\text{OH}$	Salicylic acid or Hydroxybenzoic acid				
MONOBASIC ACID (Trihydric)					$\text{C}_3\text{H}_6\text{O}_4$ or $\text{C}_2\text{H}_2(\text{OH})_3\text{CO}_2\text{OH}$	Glyceric or Dihydroxypropionic acid			$\text{C}_7\text{H}_6\text{O}_4$ or $\text{C}_6\text{H}_3(\text{OH})_3\text{CO}_2\text{OH}$	Dihydroxybenzoic acid				
DIBASIC ACID			$\text{C}_2\text{O}_4\text{H}_2$ or $\text{COOH}\cdot\text{COOH}$	Oxalic acid	$(\text{CH}_2\text{CO}_2\text{OH})_2$	Malonic acid†								
ETHER	$\text{CH}_3\cdot\text{O}$	Methyl ether	$(\text{C}_2\text{H}_5)_2\text{O}$	Ethyl ether	$(\text{C}_3\text{H}_7)_2\text{O}$	Propyl ether	$(\text{C}_6\text{H}_5)_2\text{O}$	Phenyl ether						

The nitrile of Formic acid or Formonitrile is HCN (Hydrocyanic acid).


† The existence of $\text{OH}\cdot\text{CO}_2\text{OH}$ is not definitively proved. The association of two OH groups with one atom of Carbon is unusual. (See page 312.)

‡ This cannot be obtained by direct oxidation of Propylene glycol.


Note.—This Table is a slight extension of one compiled by Dunstan.




displaced by one, two, or three atoms of any radical having the general formula C_nH_{2n+1} . Thus :




Carbinol.



Typical primary carbinol.




Typical secondary carbinol.



Typical tertiary carbinol.

The primary oxidize their CH_2OH group to aldehydes (bodies containing the aldehydic group, COH) and acids (bodies having the group $COOH$ or *carboxyl*) ; the secondary oxidize their $CHOH$ group to form a ketone (a body having *carbonyl*, CO'' , as a group, as acetone $CH_3 - CO - CH_3$), and by further oxidation break up, forming bodies with less carbon units than the original alcohol ; while the tertiary yield a ketone and an acid. The primary alcohols alone are of practical interest to medical and pharmaceutical students. The tertiary alcohols are said to be depressants instead of stimulants. (For examples of primary, secondary, and tertiary alcohols see p. 449.)

General Method of Preparing Primary Alcohols.—By acting on the monochloro-derivatives of a paraffin by potassium or silver acetate an ethereal salt (acetate) is produced, which when saponified with caustic potash yields the alcohol. For instance—



If the chloro-derivatives were directly acted upon by the potassium hydrate, hydrocarbons of the olefine or acetylene series would result.

The chief primary alcohols are, however, otherwise obtained.

Note on Nomenclature.—The student will perceive that such names as those in the following set are used indifferently : ethylic acetate, ethyl acetate, acetate of ethyl, acetic ether.

Methylic Alcohol.

Methyl (*μέθν, methu*, wine, and *ῥλη, ule*, wood) Alcohol or Carbinol, CH_3OH or HCH_2OH (*Pyroxylic Spirit* or *Wood Naphtha*), is a product of the destructive distillation of wood, and is now obtained in large quantities as a by-product in the manufacture of beet-sugar in France. By oxidation it yields formic acid. (See p. 339.)

Methylated Spirit.—Spirit of wine, containing 10 per cent. of wood spirit, constitutes ordinary *methylated spirit*, a spirit issued duty free for the use of manufacturers, the methylic alcohol, etc. not interfering with technical applications. From its nauseous taste and odor, however (not due to methyl alcohol, but to tarry

bodies), it cannot take the place of gin, brandy, or other spirit; hence, while industry is benefited, intemperance is discouraged and the revenue not injured.

Detection of Methylic Alcohol in Presence of Ethylic Alcohol.—Three or four methods have been proposed for the detection of methylated spirit in various liquids; that most used by pharmacists is by J. T. Miller. For the application of the test to tinctures and similar spirituous mixtures some of the spirit is first separated by distilling off a drachm or so from about half an ounce of the liquid placed in a small flask or test-tube having a long bent tube attached. Into a similar apparatus put 30 grains of powdered red potassium chromate, $\frac{1}{2}$ an ounce of water, 25 minims of strong sulphuric acid, and 30 or 40 minims of the spirit to be tested. Set the mixture aside for a quarter of an hour, and then distil nearly half a fluidounce. Place the distillate in a small dish, add a very slight excess of sodium carbonate, boil down to about a quarter of an ounce, add enough acetic acid to impart a distinct but feeble acid reaction; pour the liquid into a test-tube, add a grain of silver nitrate dissolved in about 30 drops of water, and heat gently for a couple of minutes. If the liquid then merely darkens a little, but continues quite translucent, the spirit is free from methylic alcohol; but if a copious precipitate of dark-brown or black metallic silver separates, and the tube, after being rinsed out and filled with clean water, has a distinct film of silver, which appears brown by transmitted light (best seen by holding it against white paper), the spirit is methylated. The experiments are best performed by daylight.

Explanation.—This test depends for its action on the reducing powers of formic acid. In the above operation the ethylic alcohol becomes oxidized to acetic acid (the natural acid of the ethyl series), which does not reduce silver salts, a minute quantity only of formic acid being produced, while the methylic alcohol yields formic acid (the natural acid of the methyl series) in a comparatively large quantity. Aldehyde, which is also a reducing agent, is simultaneously produced, but removed in the subsequent ebullition with carbonate of sodium.


Methylated Sweet Spirit of Nitre.—The preparation of spirit of nitrous ether from methylated spirit is illegal in Great Britain, and, probably, is very rarely practised. For the detection of methylic alcohol in this liquid Mr. Miller suggests the following modification of the foregoing process:

Shake about an ounce of the sample with 20 or 30 grains of anhydrous potassium carbonate, and, if needful, add fresh portions of the salt until it ceases to be dissolved, then pour off

the supernatant spirit. This serves to neutralize acid and to remove water, of which an abnormal quantity may be present. Introduce $\frac{1}{2}$ a fluidounce of the spirit into a small flask; add 150 grains of anhydrous calcium chloride in powder, and stir well together; then, having connected the flask with a condenser, place it in a bath of boiling water, and distil $1\frac{1}{2}$ fluidrachms, or continue the distillation until scarcely anything more comes over. The operation is rather slow, but needs little attention, and should be done thoroughly. The distillate contains nearly the whole of the nitrous ether and other interfering substances, while in the retort there remains a non-volatile compound of calcium chloride and methylic alcohol, if the latter be present. Now add to the contents of the flask a fluidrachm of water, which decomposes the compound just referred to, and draw over the half-drachm of spirit required for testing. Then proceed as described in the foregoing paragraphs.

Ethylic Alcohol.

Ethyl Alcohol, or Methyl Carbinol, commonly called simply *Alcohol* (C_2H_5OH or CH_3CH_2OH).—It is a colorless liquid, having a boiling-point of 173.6° F. and sp. gr. 0.7935. Ethyl alcohol may be obtained by passing ethylene into strong sulphuric acid. The product, ethylhydrogen sulphate, when distilled with water, yields alcohol:



On the large scale alcohol is produced by fermentation. All fermented bread retains a little alcohol, sometimes as much as 1 in 400.

Formation of Alcohol.—Ferment two or three grains of sugar by dissolving it in a test-tube full of water, adding a little yeast (*Cerevisiæ Fermentum*, B. P.) or a piece of the so-called German or dried yeast, and setting the whole aside for several hours in a warm place at a temperature of 75° to 85° F.; carbonic acid gas is evolved, and, if the tube be inverted in a small dish containing water, may be collected in the upper part of the tube and subsequently tested: the solution contains alcohol. If the experiment be made on large quantities (4 ounces of sugar, 1 of yeast, and 1 pint of water), the fermented liquid should be distilled, one-half being collected, shaken with a little lime, soda, or potash to neutralize any acetic acid and decompose ethereal salts, and again distilled till one-half has passed over; the product is dilute spirit of wine. It may be still further concentrated or rectified by repeating this process of *fractional distillation*, the separate fractions being redistilled,

then fractions having fairly near boiling-points being mixed and again distilled, and so on until the ethylic alcohol, the other alcohols (chiefly amylic alcohol), and the water, having had the cohesive tendencies of their molecules thus overcome, are separated from each other. The heads of stills can be so adapted, especially on the large scale, as to condense and deliver at once the substances having different boiling-points.

Fermentation.—The act of fermentation is commonly the result of, or rather accompaniment of, some vital action. *Alcoholic fermentation* would appear to be always attended by, or to attend development of, life and free multiplication of cellular structure. It follows the development of the fungus already referred to as constituting the chief active part of yeast, the *Saccharomyces cerevisiae*. In the presence of this fungus, with small quantities of phosphates and albumenoid matter, *glucose* is converted into alcohol and carbonic acid gas, together with small portions of glycerin, succinic acid, and other substances. Yeast also contains a soluble ferment, *invertase*, analogous to diastase, which is capable of converting sucrose into glucose. Therefore if yeast be used, *sucrose*, or cane-sugar, may be converted into carbonic acid gas and alcohol, the soluble ferment first converting the sucrose, not itself fermentable by the saccharomyces, into glucose.



Not more than 20 per cent. by weight of alcohol can be obtained in a fermenting fluid, for more than this proportion prevents fermentation.

Other kinds of fermentation, arising from the action of special ferments which have not received in all cases distinctive names, are the following: *Viscous* or *Mannitic fermentation*, which occurs when beer or saccharine juices, such as that of the beet-root, become "ropy." Gum, mannite, and carbonic acid gas are produced. (For *Lactic* and *Butyric fermentations* see "Lactic Acid.") *Putrefactive fermentation* occurs when a liquid containing albumenoid matter is exposed to the air. Infusoria appear in the liquid, using up the dissolved oxygen, and the ferments of the genus *vibrio* are developed. These are protected from oxygen, which is fatal to them, by a thin surface layer crowded with bacteria—small rod-like organisms having powers of locomotion. The vibrionic action, or putrefaction, proceeds with evolution of sulphuretted hydrogen, together with other gases having unpleasant odors and of complex chemical constitution. (For *Acetic fermentation* see "Acetic Acid;" for *Ammoniacal fermentation* see "Urine.")

Fermentation by Certain Soluble Albumenoids.—(For the conversion of starch into sugar by diastase see "Starch;" of amygdalin into benzoic aldehyde, hydrocyanic acid, and glucose by emulsin see "Amygdalin;" of salicin into saligenin and glucose see "Salicin;" of potassium myronate into allyl sulphocyanate, etc. by myrosin, see "Mustard;" of cane-sugar into grape-sugar by the soluble fer-

ment in yeast see the foregoing paragraphs.) Many soluble ferments or *hydrolysts* occur in the germinating seeds and other parts of plants, and play an important part in nutrition.

The nomenclature of *ferments* and *fermentation* is slowly emerging from unavoidable confusion caused by growth of knowledge on the subject. The word *fermentation* originally described the action that goes on in the preparation of alcoholic fluids or of dough, for it is derived from the Latin *ferves*, I boil or seethe, in allusion to the production of gas. But discoveries of ferments so multiplied as to force classification, resulting in the names *organized ferments* (yeast, for example) and *unorganized ferments* (diastase, for example), also termed *insoluble ferments* and *soluble ferments*. Moreover, the word *fermentation* itself became scarcely applicable to many of these actions, which, otherwise strictly analogous, afforded no gas, no boiling or seething. Hence the word *zymosis* (from ζύμη, zume, leaven) for the action of organized ferments (or *zymes*?), while the soluble or unorganized ferments are termed *enzymes*, and their action one of *enzymosis*. All enzymes promote *hydrolysis* (*vide* Index), hence are *hydrolysts*.

Alcoholic Fermentation.—The chief reaction results, as already stated, in the formation of alcohol and carbonic acid gas, though 3 per cent. of glycerin, 0.5 of succinic acid, and traces of several other substances are simultaneously produced. (*Vide* "Fousel Oil," in Index.) By this reaction is formed the spirit of the various kinds of wine, beer, and liqueurs, such as Orange Wine (*Vinum Aurantii*, B. P.), made "by the fermentation of a saccharine solution to which the fresh peel of the bitter orange has been added;" Sherry Wine (*Vinum Xericum*, B. P.), the fermented juice of the grape; Whiskey (*Spiritus Frumenti*, U. S. P.), containing from 44 to 50 per cent. of pure alcohol; Bay Rum, or Spirit of Myrcia (*Spiritus Myrciæ*, U. S. P.), by distilling rum with leaves of *Myrcia acris*, etc., or by dissolving the oils in alcohol; and others.

Alcoholic drinks vary much in strength. Cider or apple wine, perry or pear wine, and good beer (ale and porter or stout) contain 4 to 6 per cent. of real alcohol; good light wines, both "red" and "white," and natural sherry also, 10 to 12 per cent.; strong sherry and port, which are commonly "fortified"—that is, contain added spirit—16 to 18 per cent.; while "spirits" (gin, rum, brandy, whiskey, etc.) and "liqueurs" (ratafia, almond-flavored; maraschino, cherry-flavored; curaçoa, orange-flavored; chartreuse, a composite-flavored liqueur, etc.), are "under proof" or "over proof," terms explained in a following paragraph. For British excise purposes "beer" is any such liquid or substitute which contains more than 2 per cent. of proof spirit. The well-known effects of these spirituous fluids on the animal system would appear to be due primarily to alcohol, and secondarily to ethereal derivatives of alcohols. Some owe a part of their effect to non-volatile substances, for beer from which all alcohol, etc. has been removed by ebullition is said to have considerable effect on the human economy.

The official (U. S. P.) wines are all made with "white wine" (*Vinum Album*, U. S. P.), a kind of natural sherry, containing not

less than 10 or more than 14 per cent. of absolute alcohol. *Vinum Rubrum*, U. S. P., is of similar strength—a kind of natural port wine.

Varieties of Alcohol.—The weak spirit, concentrated by distillation till it contains 84 per cent. by weight of pure alcohol, is an ordinary article of trade; its specific gravity at 60° F. is 0.8382. This is common *Spirit of Wine*, the *Spiritus Rectificatus* of the British Pharmacopœia. The official *Proof Spirit** (*Spiritus Tenuior*, B. P.) contains 49½ per cent. by weight, 57 by volume, of alcohol, and is made by diluting 100 volumes of “rectified spirit” with water until the well-stirred product measures 156 volumes. Sixty volumes of water will be required for this purpose, the liquids occupying less bulk after than before admixture. In the language of the Excise authorities, the rectified spirit of the Pharmacopœia would be described as “56 degrees over proof” (56 O. P.); that is, 100 volumes contain as much alcohol as is present in 156 volumes of proof spirit. Obviously, proof spirit may be made by diluting with water rectified spirit of any other strength than that mentioned above. Thus 100 fluidounces of a spirit of “70 over proof” may be diluted to 170, or the same quantity of a spirit of “50 over proof” may be diluted to 150, and so on. The specific gravity of proof spirit at 60° F. is 0.920. Spirit 10 per cent. “under proof” contains as much alcohol as would be present in spirit formed of 90 volumes of proof spirit mixed with sufficient water to form 100 volumes. According to British law, gin is not “adulterated” with water if it is not weaker than 35 degrees under proof; nor brandy, whisky, or rum if they are not weaker than 25 degrees under proof.

Alcohol, U. S. P., contains 99 per cent. by weight; *Alcohol Dilutum*, U. S. P., 41 per cent. by weight (48.6 by volume), of real alcohol, the remainder being water. The former has a sp. gr. not higher than 0.797, the latter about 0.937 at 15.6 C., or 0.789 and 0.936 respectively at 25° C. The stronger boils at 78° C.

Real Alcohol (C_2H_5HO) may be prepared from spirit of wine by removing the water which the latter contains. This is accomplished, partially, by anhydrous potassium carbonate, and finally and entirely by recently fused calcium chloride. In operating on, say, 1 pint, 2 ounces of dried potassium carbonate is placed in a bottle that can be well closed, and frequently shaken during two days with the spirit. Meanwhile put rather more than a pound of calcium chloride into a covered crucible, and subject it to a red heat for half an hour; then pour the fused salt on to a clean stone slab, cover it quickly with an inverted porcelain dish, and when it has congealed break it up into small fragments and enclose it in a dry stoppered bottle. Put 1 pound of this fused calcium chloride into a flask,

* *Proof spirit* is so termed from the fact that in olden times a proof of its strength was supposed to be afforded by moistening a small quantity of gunpowder and setting light to the spirit; if it fired the powder, it was said to be “over-proof;” if not, “under proof.” The weakest spirit that would stand this test was what we should now describe as of sp. gr. 0.920.

pour over it the spirit decanted from the potassium carbonate, and, closing the mouth of the flask with a cork, shake them together, and allow them to stand for twenty-four hours with repeated agitation. Then, attaching a dry condenser closely connected with a receiver from which free access of air is excluded, and applying the flame of a lamp to the flask, distil about 2 fluidounces, which should be returned to the flask, after which the distillation is to be continued until 15 fluidounces have been recovered. The foregoing details are those of the British Pharmacopœia. The official *Ethyl Alcohol*, U. S. P., should not contain more than 1 per cent. by weight of water, and should have a specific gravity of 0.797 to 0.789. The product should be colorless, and of a characteristic but agreeable odor. It is entirely volatilized by heat, is not rendered turbid when mixed with water, does not cause anhydrous copper sulphate to assume a blue color even after the two have been well shaken together, and should leave no smell on a piece of blotting-paper when allowed to evaporate spontaneously on it. What little water remains may, if necessary, be removed by the cautious addition of a little metallic sodium. If 5 per cent. of sodium be used, solution of sodium ethylate or caustic alcohol results (*Liquor Sodii Ethylatis*, B. P.) by replacement of the hydroxyl hydrogen by sodium: $\text{Na}_2 + 2\text{C}_2\text{H}_5\text{OH} = 2\text{C}_2\text{H}_5\text{ONa} + \text{H}_2$. The solution contains 19 per cent. of sodium ethylate.

There is another strength of alcohol which is official, containing 92.5 per cent. by weight of pure ethyl alcohol and about 7.5 per cent. of water. This is known as *Alcohol Deodoratum*, U. S. P., or *Deodorized Alcohol*. It has a sp. gr. of about 0.816 to 0.808.

Spirit of French Wine (*Spiritus Vini Gallici*, U. S. P.), or *Brandy*, is a colored and flavored variety of alcohol distilled from French wine. Its color is that of light sherry, and is derived from the cask in which it has been kept, but it is commonly deepened by the addition of burnt sugar. Its taste is due to the volatile flavoring constituent of the wine, often increased by the addition of artificial essences.

"A pale, amber-colored liquid, having a distinctive odor and taste, and a slightly acid reaction; specific gravity should not be more than 0.941, nor less than 0.925, corresponding, approximately, to an alcoholic strength of 39 to 47 per cent. by weight or 46 to 55 per cent. by volume. If 100 cc. of brandy be very slowly evaporated in a tared capsule on a water-bath, the last portions volatilized should have an agreeable odor, free from harshness (absence of fusel oil from grain or potato spirit), and the residue, when dried at 100° C. (212° F.), should not weigh more than 1.5 gm. This residue should have no sweet or distinctly spicy taste (absence of added sugar, glycerin, or aromatic substances). It should almost completely dissolve in 10 cc. of cold water, forming a solution which is colored not deeper than light green by a few drops of dilute ferric chloride, made by mixing the latter with 10 volumes of water (absence of more than traces of oak tannin from casks). To render 100 cc. of brandy distinctly alkaline to litmus should require not more than 1 cc. of potassium hydrate (limit of free acid)."—U. S. P.

The foregoing words are also used in describing *Whiskey* (*Spiritus Frumenti*) in the United States Pharmacopœia, except that the sp. gr. is to be "not above 0.930 nor below 0.917, corresponding approximately with an alcoholic strength of 44 to 50 per cent. by weight or 50 to 58 per cent. by volume," and that the acidity is not to be greater in 100 cc. than 1.2 cc. of soda solution will neutralize.

Tests.—There are no specific tests for alcohol when mixed with complex matters. It is, however, easily isolated and concentrated by fractional distillation, and is then recognizable by conjoint physical and chemical characters. Thus its odor and taste are characteristic; it is lighter than water, volatile, colorless, and when tolerably strong inflammable, burning with an almost non-luminous flame; it readily yields aldehyde (see below) and acetic ether (*vide* Index), each of which has a characteristic odor, and in presence of hot acid alcohol reduces red potassium chromate to a green chromium salt.

According to Lieben, 1 of alcohol in 2000 of water can be detected by adding to some of the warmed liquid a little iodine, a few drops of solution of soda, again warming gently, and setting aside for a time; a yellowish crystalline deposit of *iodoform* (CHI_3) is obtained. Under the microscope the latter presents the appearance of hexagonal plates or six-rayed and other varieties of stellate crystals.



Other alcohols, aldehydes, gum, turpentine, sugar, and several other substances give a similar reaction.

Tests of Purity.—Oil or resin is precipitated on diluting spirit of wine with distilled water, giving an opalescent appearance to the mixture. The sp. gr. of rectified spirit is 0.838. Fousel oil, aldehyde, and such impurities are detected by silver nitrate. (*Vide* Index, "Alcohol, Test for Purity of.") Water in absolute alcohol may be detected by adding to a small quantity a little highly-dried copper sulphate, which becomes blue ($\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$) if water be present, but retains its yellowish-white anhydrous character (CuSO_4) if water be absent.


Note.—Most ethyl derivatives are formed from alcohol, such as ethyl nitrite in spirit of nitrous ether, iodo-ethane, etc. These have been treated under "Ethane." Aldehyde and acetic acid are obtained from alcohol by oxidation.

Alcohols and Ethers.—As already stated, just as such elementary radicals as potassium (K) form hydrates (as KOH) and oxides (as K_2O), so do such compound radicals as ethyl (C_2H_5) form hydrates, as common alcohol, $\text{C}_2\text{H}_5\text{OH}$, and other alcohols, p. 436, and oxides as common ether, $(\text{C}_2\text{H}_5)_2\text{O}$, and other ethers, p. 447.


But *Sulphur Alcohols*, or *Thio-alcohols*, CH_3SH , $\text{C}_2\text{H}_5\text{SH}$, etc., analogous to sulphydrates, KHS, etc., are known. They were

originally termed *mercaptans* (*mercurius captans*), from the readiness with which they took mercury captive, $(\text{C}_2\text{H}_5\text{S})_2\text{Hg}$. *Sulphur Ethers*, or *Thio-ethers*, also are known, $(\text{CH}_3)_2\text{S}$, $(\text{C}_2\text{H}_5)_2\text{S}$, etc. The vapors of such sulphur compounds have an extremely unpleasant smell.

Sulphonic Acids are products of the oxidation of the sulphur alcohols just mentioned. For example:




They also may be formed by acting on hydrocarbons with sulphuric acid. Examples:




Sulphonic acids are isomeric with acid sulphites, the characteristic sulphonic group or radical being SO_3H , but the acid sulphites of the organic radicals are extremely unstable, the corresponding sulphonic acids very stable. *Orthophenolsulphonic acid*, $\text{C}_6\text{H}_4\text{OH}\cdot\text{SO}_2\cdot\text{OH}$, *sozolic acid*, or *aseptol*, is a non-poisonous, non-irritating antiseptic. The *di-iodoparaphenolsulphonic acid*, or *sozoiodol*, $\text{C}_6\text{H}_2\text{I}_2\text{OH}\cdot\text{SO}_2\cdot\text{OH}$, has similar properties, and is used instead of iodoform.

Saccharin (*Glusidum*, B. P.; synonym, *Glucosimide*), which is a harmless, non-alimentary, purely sweetening agent two or three hundred times as sweet as sugar, is *benzoyl-sulphonic imide*. Fahlberg obtains it by converting the toluene, $\text{C}_6\text{H}_5\text{CH}_3$, of coal-tar into toluene-sulphonic acid (above); this into a calcium salt, then into a sodium salt, and the latter into toluene-sulphonic chloride, by action of phosphorus trichloride and chlorine; the liquid orthochloride into amide by ammonium carbonate; the amide is then oxidized by potassium permanganate to sulphamidobenzoate and water; hydrochloric acid then precipitating benzoyl-sulphonic imide or "saccharin" with elimination of water. "Soluble saccharin" is saccharin in which hydrogen is displaced by sodium. The following formulæ illustrate the stages of manufacture:



Sulphonal (B. P.), a new hypnotic, is a crystalline, colorless, inodorous, tasteless substance, a product of the action of permanganate solution on mercaptol—a liquid resulting from the reaction

of hydrochloric acid, mercaptan, and acetone. Its descriptive name is diethylsulphon-dimethylmethane, and the following is its descriptive formula:




ETHERS.

Ethyllic (or Ordinary) Ether.—Into a capacious test-tube put a small quantity of spirit of wine and about half its bulk of sulphuric acid; mix and gently warm; the vapor of ether, recognized by its odor, is evolved. Adapt a cork and long bent tube to the test-tube, and slowly distil over the ether into another test-tube. Half the original quantity of alcohol now placed in the generating-tube will again give ether; and this operation may be repeated many times.

On the larger scale, and according to the following official process (*Æther*, B. P.), the addition of alcohol, instead of being intermit-

Fig. 41.



Preparation of Ether.

ting, is continuous, a tube conveying alcohol from a reservoir into the generating-vessel. Mix 10 fluidounces of sulphuric acid with 12 fluidounces of rectified spirit in a glass retort or flask capable of containing at least 2 pints, and, not allowing the mixture to cool, connect the retort or flask, by means of a bent glass tube, with a Liebig's condenser, and distil with a heat sufficient to maintain the liquid in brisk ebullition. (If a thermometer also be inserted in the tubulure of the retort or through the cork of the flask, the temperature may be still


more carefully regulated—between 284° and 290° F.) As soon as the ethereal fluid begins to pass over, supply fresh spirit in a continuous stream, and in such quantity as to about equal the volume of the fluid which distils. For this purpose use a tube furnished with a stopcock to regulate the supply, as shown in Fig. 41, connecting one end of the tube with a vessel containing the spirit supported above the level of the retort or flask, and passing the other end through the cork of the retort or flask into the liquid. When a total of 50 fluidounces of spirit has been added, and 42 fluidounces of ether have distilled over, the process may be stopped. This is *Æther*, U. S. P.

To *partially purify* the liquid dissolve 10 ounces of calcium chloride in 13 ounces of water, add $\frac{1}{2}$ an ounce of lime, and agitate the mixture in a bottle with the impure ether. Leave the mixture at rest for ten minutes, pour off the light supernatant fluid, and distil it with a gentle heat until a glass bead of the specific gravity 0.735 placed in the receiver begins to float. The ether and spirit retained by the calcium chloride and by the residue of each rectification may be recovered by distillation and used in a subsequent operation.

Explanation of Process.—On the addition of sulphuric acid to alcohol in equal volumes, one molecule of each reacts and gives a molecule of ethylhydrogen sulphate and one of water :



More alcohol then gives ether and sulphuric acid by the reaction of one molecule of the alcohol on one of ethylhydrogen sulphate (sometimes termed ethylsulphuric acid or sulphethylic acid or sulphovinic acid) :



The water of the first reaction and the ether of the second distil over, while the sulphuric acid liberated is attacked by alcohol and reconverted into ethylhydrogen sulphate. So that the sulphuric acid originally employed finally remains in the retort in the form of ethylhydrogen sulphate. The effect, however, of a small quantity of sulphuric acid in thus converting a large quantity of alcohol into ether is limited, secondary reactions occurring to some extent after a time.

Mixed Ethers.—That $\text{C}_2\text{H}_5-\text{O}-\text{C}_2\text{H}_5$ represents the constitution of ether is indicated by the result of the reaction of, say, methyl-alcohol on ethylsulphuric acid, a single definite substance, methyl-ethyl ether, $\text{CH}_3-\text{O}-\text{C}_2\text{H}_5$, resulting.

Ethers of various radicals, $\text{R}-\text{O}-\text{R}$, and several *mixed ethers*, $\text{R}_a-\text{O}-\text{R}_b$, and *sulphur-ethers*, or *thio-ethers*, $\text{R}-\text{S}-\text{R}$, are known.

Properties.—Pure ethylic ether is gaseous at temperatures above 95°F .; hence the condensing-tubes employed in its distillation must be kept as cool as possible. At all ordinary temperatures it rapidly volatilizes, absorbing much heat from the surface on which it is placed. A few drops evaporated consecutively from the back of the hand produce great cold, and if blown in the form of spray the cooling effect is so rapid and intense as to produce local anæsthesia. Evaporated by aid of a current of air from the outside of a thin narrow test-tube containing water, the latter is solidified to ice. Its vapor is very heavy, more than twice and a half as heavy as air, and nearly forty times as heavy as hydrogen ($\text{H}_2=2$; $\text{C}_4\text{H}_{10}\text{O}=74$; or as 1 to 37). In a still atmosphere it will flow a considerable distance along a table or floor before complete diffusion occurs. The

vapor is highly inflammable; hence the importance of keeping candle and other flames at a distance during manipulations with ether. Exposed to the action of air and light, ether becomes charged with a little hydrogen peroxide.

Purification.—To imitate the process of partial purification above described, add to the small quantity of ether obtained in the foregoing operation a strong solution of calcium chloride and a little slaked lime; the latter absorbs any sulphurous acid that may have been produced by secondary decompositions, while the former absorbs water; on shaking the mixture and then setting aside for a minute or two, the ether will be found floating on the surface of the solution of calcium chloride.

This ether, redistilled until the distillate has a sp. gr. not higher than 0.735 and boiling-point not higher than 105° F., is the ether of the British Pharmacopœia. It still contains about 8 per cent. of alcohol. The latter may be removed by well shaking the ether with half of its bulk of water, setting aside, separating the floating ether, and again shaking it with water; alcohol is thus washed out. This washed ether containing water (for water and ether are to some extent soluble the one in the other; 50 measures of pure ether agitated with an equal volume of water are reduced to 45 measures) is placed in a retort with solid calcium chloride and a little caustic lime, and once more distilled; pure dry ether (*Æther Purus*, B. P.) results. Sp. gr. not exceeding 0.720, indicating about 99 per cent. of real ether. Shaken with about a fourth of its bulk of solution of potassium iodide and a little starch mucilage, little or no blue color is produced, indicating absence of an impurity. The exact nature of the latter is not known, but inasmuch as it develops hydrogen peroxide, Dunstan and Dymond suggest that the test for the latter should always be applied. Poleck and Thummel state that the impurity in ether which causes potash to yield a brown color is vinyl alcohol. On shaking pure ether with half its bulk of a dilute solution of potassium bichromate acidified by sulphuric acid, and setting aside, the supernatant ether has no blue color.

Æther, U. S. P., contains about 96 per cent. of real ether, and 4 per cent. of alcohol containing a little water; sp. gr. 0.725 to 0.728 at 15° C.; boiling-point, 37° C.

Upon evaporation, ether should leave no residue. If 10 cc. of it be poured, in portions, upon clean, odorless blotting-paper and allowed to evaporate spontaneously, no foreign odor should become perceptible when the last traces of ether leave the paper. When 20 cc. of ether are shaken in a graduated tube, with 20 cc. of water just previously saturated with ether, the ethereal layer, upon separation, should not measure less than 19.8 cc. (absence of an undue amount of alcohol or water). If 10 cc. of ether be shaken occasionally, within one hour, with 1 cc. of potassium iodide, no color should be developed in either liquid (absence of aldehyde, etc.).—*U. S. P.*


Spiritus Ætheris, U. S. P., is a mixture of 325 cc. of ether with

675 cc. of alcohol. *Spiritus Ætheris Compositus*, U. S. P., contains 325 cc. of stronger ether, 650 cc. of alcohol, and 25 of ethereal oil. It is the old "Hoffmann's Anodyne."

ALCOHOLS—continued.

Propylic and Butylic Alcohols.


The primary and secondary propyl alcohols— $\text{C}_2\text{H}_5\text{CH}_2\text{OH}$ and $(\text{CH}_3)_2\text{CHOH}$ —and the four isomeric butyl alcohols ($\text{C}_4\text{H}_9\text{OH}$; see below) are of little pharmaceutical interest.




Amylic Alcohol.

Pentyl or *Amylic Alcohol* (*Fousel Oil*) (*Alcohol Amylicum*, B. P.) ($\text{C}_5\text{H}_{11}\text{HO}$ or $\text{C}_4\text{H}_9\text{CH}_2\text{OH}$), is a constant accompaniment of ethylic or common alcohol ($\text{C}_2\text{H}_5\text{OH}$), especially when the latter is prepared from sugar which has been derived from starch; hence the name, from *amylum*, starch. The sugar of potato starch yields a considerable quantity; hence the alcohol is often called *potato oil*. It is also termed *fousel oil* or *fusel oil* (from *φύω*, *phuō*, to produce), in allusion to the circumstance that the supposed oil is not simply educed from a substance already containing it, as is usually the case with oils, but is actually produced during the operation. It was described as oil probably because it resembled oil in not readily mixing with water, but it is soluble to some extent in water, and is a true spirit, homologous with spirit of wine. It often contains variable proportions of propylic, butylic, and caprylic alcohols. (See also Valerianic Acid.) When used for medicinal purposes, "it should be redistilled, and the product, passing over at 262° to 270° F. (or about 128° to 132° C.), alone be collected for use."


Amylic alcohol is a "colorless liquid with a penetrating and oppressive odor and a burning taste. When pure, its specific gravity is .818. Sparingly soluble in water, but soluble in all proportions in alcohol, ether, and essential oils. Exposed to the air in contact with platinum-black, it is slowly oxidized, yielding valerianic acid" ($\text{C}_4\text{H}_9\text{COOH}$). Two allotropic varieties of amylic alcohol exist—one, α , having no action on the other, β , lævo-rotating a polarized ray. The amylic alcohol of trade probably contains both varieties.



Primary
α, or inactive
amylic alcohol.



Primary
β, or active
amylic alcohol.



Tertiary
amylic alcohol,
"amylylene hydrate."

The constitution of the variety of amylic alcohol ($\text{C}_5\text{H}_{11}\text{OH}$) termed *tertiary amylic alcohol*, or *dimethyl-ethylcarbinol*, is shown in the above graphic formula. It is used in medicine in place of chloral hydrate, and is known as amylylene hydrate, for it contains the elements of amylylene (C_5H_{10}) and water (H_2O).

The pentylic salts of pharmaceutical interest are all derived from amylic alcohol.

Other Monhydroxyl Alcohols.


Among the higher alcohols are the following:

Cetylic Alcohol ($\text{C}_{16}\text{H}_{33}\text{OH}$), or *Cetyl Hydrate*, formerly termed *ethal*, obtained by saponifying spermaceti (*Cetaceum*, U. S. P.), which consists of cetyl palmitate ($\text{C}_{16}\text{H}_{33}\text{C}_{16}\text{H}_{31}\text{O}_2$), or *cetine*. Spermaceti is the solid crystalline fat accompanying sperm oil in the head of the spermaceti whale.

Cerylic Alcohol ($\text{C}_{27}\text{H}_{55}\text{OH}$) is obtained in a similar manner from Chinese wax (ceryl cerotate).

Melissic Alcohol ($\text{C}_{30}\text{H}_{61}\text{OH}$) is obtained in a similar manner from melissic palmitate, the portion of beeswax soluble in hot alcohol. *Yellow Wax* (*Cera Flava*, U. S. P.) and the same bleached by exposure to moisture, air, and sunlight, or *White Wax* (*Cera Alba*, U. S. P.), is the prepared honeycomb of the hive-bee. According to Brodie, it is in the main a compound of the melissic ($\text{C}_{30}\text{H}_{61}$) and cerotic ($\text{C}_{26}\text{H}_{53}\text{COO}$) radicals with about 5 per cent. of *ceroleine*, the body to which the color, odor, and tenacity of wax are due. Amongst the possible adulterants of wax are paraffin and *ceresine*. The latter is the purified native *ozokerite* of Galicia, a solid hydrocarbon, largely used as a substitute for beeswax, especially in Russia. Both paraffin and *ceresine* reduce the melting-point of wax, which should not be lower than 146°F . (63.3°C .) when taken in the manner described in connection with the quantitative determination of temperature. (*Vide Index*.) The amount is obtained by destroying the wax with warm oil of vitriol, and afterward with fuming sulphuric acid, which scarcely affects paraffin and *ceresine*. Pure wax will not yield more than about 3 per cent. to cold rectified spirit, whereas rosin, etc. would be extracted by the spirit. Solution of soda extracts nothing from pure wax, but dissolves fat acids, fat, rosin, Japan wax, etc. and the alkaline fluid then yields a precipitate of acids on the addition of hydrochloric acid. Soap would be dissolved from wax on boiling the sample with water, and the aqueous fluid would yield oily acid on adding hydrochloric acid. Flour or any starch would be detected in the cooled aqueous fluid by iodine.

The *Allylic Series of Alcohols* ($C_nH_{2n-1}OH$) (monhydric alcohols).—*Allylic alcohol* (C_3H_5OH) may be obtained by heating 4 parts of glycerin with 1 of oxalic acid, the receiver being changed at $195^\circ C.$, and the liquid collected till the temperature rises to $260^\circ C.$ The first product is formic acid, which reacts on glycerin, forming monoformin:




This, on further heating, yields allylic alcohol:




By the action of the haloid acids it produces iodine, bromine, and chlorine derivatives by replacing the OH by I, Br, or Cl; these derivatives, when digested with potassium sulphocyanate, yield allyl sulphocyanate or artificial *Oil of Mustard* (identical in composition with the chief constituent of the natural oil), allyl sulphocyanate being the body to which mustard owes its power of inducing inflammatory action on the skin ("Mustard Poultice" and *Charta Sinapis*, U. S. P.).

Mustard (*Sinapis*, U. S. P.) is a powdered mixture of black or, rather, reddish-brown, mustard-seeds (*Sinapis Nigræ Semina*, B. P.) from the *Brassica nigra*, and white mustard-seeds (*Sinapis Alba Semina*, B. P.) from the *Brassica alba*. The white mustard-seed contains *sinalbin* ($C_{30}H_{44}N_2S_2O_{16}$), a glucoside which, in contact with the myrosin in an aqueous extract of mustard, yields the sulphocyanate of the radical *acrinyl*, a body which forms part of the essential oil of mustard paste.



The black contains the albumenoid ferment *myrosin*, resembling the *emulsin* of almonds, and also potassium myronate, or *sinigrin*. The latter is the body which, under the influence of the former, yields the chief part of the pungent oil of mustard paste. The amount of myrosin in black mustard is scarcely sufficient to decompose the whole of the sinigrin, while in white mustard the amount is more than sufficient to decompose the sinalbin. Hence the most effective mustard is a mixture of white and black.


The ferments act most effectively—hence the maximum amount of pungency is produced—in mustard paste at temperatures not exceeding $100^\circ F.$ This temperature is contemplated in preparing *Cataplasma Sinapis*, B. P.



Crude oil of mustard often contains *allyl cyanide*, C_3H_5CN .

In the Pharmacopœia of India the seed of *Brassica juncea*, *Rai*, or *Indian Mustard Plant*, is official in addition to that of *B. alba* and *B. nigra*. It is the common mustard of warm countries. It does

not differ chemically from other mustard. Allyl compounds are also met with in several other cruciferous and liliaceous plants. *Oil of Garlic* (*Allium*, U. S. P.) owes its odor to an allyl sulphide ($C_3H_5)_2S$, some say to a "disulphide." The former may be artificially obtained by acting on allyl iodide by potassium sulphide;



Decylene Alcohol, $C_{10}H_{19}OH$, belongs to this series. *Menthol* (*Menthol*, B. P.), obtained from oil of peppermint, is said by some to consist wholly of this alcohol.

QUESTIONS AND EXERCISES.


Give an outline of the relations between alcohols and acids.—Give a general method of preparing the primary alcohols of the ethylic series.—Name the source of methylic alcohol.—What is "methylated spirit"?—Describe the mode of detecting methylated spirit in a tincture.—How can artificial ethylic alcohol be prepared?—Write a few sentences on the formation, purification, and concentration of alcohol, and explain the difference between rectified spirit, proof spirit, and absolute alcohol.—What quantity of water must be added to 1 gallon of spirit of wine 56 degrees over proof to convert it into proof spirit?—How far must 5 pints of spirit of wine of 53 degrees over proof be diluted to become proof spirit?—*Ans.* 7 pints 13 ounces.—State the specific gravity of proof spirit. State the proportion of alcohol commonly present in malt liquors, light wines, port and sherry, and "spirit," and state the extent to which spirits may be diluted without "adulteration."—Enumerate the characters of alcohol.—Whence is brandy obtained? and to what are due its color and flavor?—Describe the official process for the preparation of ether, giving equations.—Offer a physical explanation of the mode of producing local anaesthesia.—How is commercial ether purified?—Is "amylic alcohol" a simple or a complex body?—How is allylic alcohol prepared?—In what relation does allylic alcohol stand to oil of mustard and oil of garlic?

Alcohols of the $C_nH_{2n-7}OH$ Series.—*Phenols and Benzylic Alcohols.*—These are alcohols only in the sense of being hydroxyl derivatives of hydrocarbons. Unlike the paraffin alcohols, they do not yield aldehydes, oxidation acids, or ketones.

Carbolic Acid.

*Phenol, Phenic Alcohol, Phenic Acid, or Carbolic Acid** (C_6H_5OH) may be artificially obtained by heating benzene with sulphuric acid, which forms benzene-sulphonic acid (p. 445), ($C_6H_5HSO_4$) this, when heated with potash, yielding potassium phenate or carbolate, and this, with acids, the phenol:

* Ordinary carbolic acid is a mixture of phenol, cresol, and other homologues.



Commercially, carbolic acid is obtained from that part of coal-tar boiling between 180° and 190° C. When purified, it is a colorless* crystalline body (*Acidum Carbolicum*, U. S. P.). A crystalline, so-called hydrous acid ($\text{C}_6\text{H}_5\text{OH}, \text{H}_2\text{O}$) may also be obtained.

At temperatures above 95° F. ordinary carbolic acid is an oily liquid. It is only slightly soluble in water, but readily dissolved by alcohol, ether, and glycerin (*Glyceritum Acidi Carbolici*, U. S. P., contains about 20 per cent. of carbolic acid dissolved in glycerin). At 60° F. (15.5° C.) 100 parts of the acid are liquefied by the addition of 5 to 10 parts of water (100 of acid and 10 of water added forming the *Acidum Carbolicum Liquefactum*, B. P.); dissolve 30 to 40 of water, and are dissolved by 1800 to 1200 of water, the former and latter of these numbers being respectively characteristic of the acicular and pulverulent varieties of the acid. In odor, taste, and solubility (and in appearance when liquefied by heat or by the addition of 5 per cent. of water) it resembles creasote, a wood-tar product for which carbolic acid has been substituted. Besides phenol ($\text{C}_6\text{H}_5\text{OH}$), coal-tar oil contains *cresol*, cresylic acid ($\text{C}_7\text{H}_7\text{OH}$) or ($\text{C}_6\text{H}_4\text{CH}_3\text{OH}$), the alcohol of toluene, while wood-tar oil furnishes *guaiacol* ($\text{C}_7\text{H}_8\text{O}_2$)—also a product of the destructive distillation of guaiacum resin, boiling-point 200° C.—and *creasol* ($\text{C}_8\text{H}_{10}\text{O}_2$) or creasote. Certain coloring-matters may be obtained by the oxidation of carbolic acid: ammonia, or, still better, phenyl-ammonia (aniline or phenylamine) mixed with it, and then a small quantity of solution of a hypochlorite, gives a blue liquid. No very satisfactory chemical method can be found for distinguishing creasote from carbolic acid, as creasote contains phenol, the chief difference consisting in the fact that the former boils only at 370° F., while the latter readily dries up at 212°. Some other physical differences exist: thus carbolic acid does not affect a ray of polarized light; creasote twists it slightly to the right. Carbolic acid is either solid or may be solidified by cooling; creasote is not solidified by the cold produced by a mixture of hydrochloric acid and sodium sulphate. Creasote from coal (impure or crude carbolic acid) gives a jelly when shaken with albumen or with collodion; creasote from wood (*Creosotum*, U. S. P.) is scarcely affected, especially if quite free from even all natural traces of carbolic acid. Coal-creasote is soluble in solution of potash and in the strongest solution of ammonia (Read), wood-creasote scarcely soluble. The coal-product is soluble in twenty volumes of water, and a neutral solution of ferric chloride strikes a more or less permanent green or blue color with the liquid; wood-creasote is less soluble (*Aqua Creosoti*, U. S. P.; *Creasote-water* is said to contain 1 in 129) and not permanently colored blue by ferric chloride. An alcoholic solution of the coal-creasote is colored

* Phenol soon assumes a pink color, owing (Fabrini) to the action of hydrogen peroxide and ammonia in presence of traces of copper, iron, or lead.

brown by ferric chloride, a similar solution of true creasote-green. A dilute solution of creasote, such as creasote-water, is not affected by agitation with spirit of nitrous ether, while a similar solution of phenol becomes red. A few drops of the spirit of nitrous ether are placed in a test-tube, then about a drachm of the aqueous fluid, and an equal volume of sulphuric acid is poured down the sides of the tube. A pink or red color results if phenol be present, especially after standing aside a short time (Eykmán ; MacEwan). A solution of carbohc acid gives, with excess of bromine-water, an insoluble white precipitate of tribromophenol, $C_6H_2Br_3OH$. This reaction is useful in quantitative estimations of carbohc acid. The extent of absorption of iodine by alkaline solutions of this and other phenols (thymol, naphthol, etc.) serves also for quantitative purposes. According to Morson, pure creasote is unaffected when mixed with an equal volume of commercial glycerin, while carbohc acid is miscible in all proportions, and will carry into solution even a considerable quantity of creasote.


Carbohc acid and alkalis yield *carbulates* or *phenylates*, as C_6H_5OK , C_6H_5ONa . Alcoholic solutions of the latter and of mercuric chloride yield yellow crystalline mercuric phenylate or *phenol-mercury* (C_6H_5O)₂Hg.

Carbohc acid is a powerful *antiseptic* (*ἀντί, anti*, against, and *σῆπω, sēpō*, to putrefy.) In large doses it is poisonous, *antidotes* being a mixture of olive oil and castor oil, freely administered, or a mixture of slaked lime with about three times its weight of sugar rubbed together with a little water. Carbohc acid is soluble in sulphuric acid, *sulphocarbohc acid*, *phenol-sulphonic acid* ($C_6H_4(OH)SO_3H$), or *sulphophenic acid* being formed. On diluting and mixing with oxides, hydrates, or carbonates *sulphocarbohcates* are formed. The formula of *sodium sulphocarbohcate* is $NaC_6H_5SO_4 \cdot 2H_2O$ or $C_6H_4(OH)SO_3Na \cdot 2H_2O$. It is obtained by saturating sulphocarbohc acid by barium carbonate, and decomposing the resulting soluble barium sulphocarbohcate, $(C_6H_4OHSO_3)_2Ba$, by sodium carbonate until a precipitate of barium carbonate ceases to form. The filtrate on evaporation yields colorless, neutral, prismatic crystals of the salt (*Sodii Sulphocarbohcates*, or *Sodium Sulphocarbohcate*, U. S. P.), the old sulphocarbohcate of soda. *Sulphocarbohcate of Zinc*, $(C_6H_4OHSO_3)_2Zn \cdot H_2O$ (*Zinci Sulphocarbohcates*, B. P.), may be obtained by saturating sulphocarbohc acid with zinc oxide.

Trinitro-phenol ($C_6H_2(NO_2)_3OH$) is formed on slowly dropping carbohc acid into fuming nitric acid ; it is the yellow dye known as *carbazotic acid* or *picric acid* ; most of the picrates are explosive by percussion.

Both carbohc acid and benzene are secondary products obtained in the manufacture of coal-gas ; hence, indeed, the word *phenic* and thence *phenyl* (from *φαῖνω, phainō*, I light, an allusion to the use of coal-gas).


By heating phenol with zinc dust benzene results :




Salicylic acid is made from phenol. (*Vide Salicylic Acid.*)

The official test for phenols is a brownish or violet color on the addition of a drop of ferric chloride.


Constitution of Phenol.—Phenol (C_6H_5OH) may be regarded as benzene (C_6H_6) in which one atom of hydrogen (H) is displaced by hydroxyl (OH). When two atoms of hydrogen in benzene are displaced by two of hydroxyl, *resorcin* ($C_6H_4_2OH$) results, a colorless, crystalline antiseptic having many advantages over carbolic acid in surgical operations. Its name was given in allusion to its original source, resin, and to certain similarities with orcin. It occurs in white flat prisms readily soluble in most liquids. It may be made by passing benzol vapor into hot sulphuric acid and heating the product (benzenedisulphonic acid, $C_6H_4(SO_2\cdot OH)_2$) with excess of soda.



Ortho-, Meta-, and Para-aromatic Compounds.—Resorcin is one of a group of three metameric dihydroxyl-benzenes. Their chemical relationships warrant the conclusion (on the atomic theory) that the cause of their differences in properties is a difference of position of the two atoms of hydroxyl in the molecule, these being, respectively, next to each other, separated by one atom (of CH), and by two atoms (of CH) (see Constitution of Benzene, p. 432), thus:



The foregoing formulæ may conveniently be shortened as follows:



In these formulæ the letters *o*, *m*, or *p* indicate the position of the atoms of hydroxyl (OH) in relation to each other: *o* signifying the *ortho position* of one atom in relation (next, or 1, 2) to the other; *m* signifying the *meta position* of one atom in relation (next but one, or 1, 3) to the other; and *p* signifying the *para position* of one atom in relation (next but two, or 1, 4) to the other. Among the benzene or aromatic compounds there are many such metameric trios (three xylenes, three phthalic acids, etc.), their occurrence strongly supporting "the benzene-ring" hypothesis of Kekulé as to the constitution of benzene compounds.



Cresol, or *Tolyl Alcohol*, $C_6H_4OH\cdot CH_3$, one of the alcohols of toluene, $C_6H_5CH_3$, is always found with crude phenol; artificially it may be made in the same manner as phenol, by acting on toluene with sulphuric acid and heating the resulting sulphonic acid (C_6H_4-

(SO₃H)CH₃) with potash. With ferric chloride it gives a brown coloration. The three forms, ortho-, meta-, para-, are known.



Benzylic Alcohol, Phenylcarbinol, C₆H₅CH₂OH, is isomeric with cresol, but has the hydroxyl group replaced in the methane nucleus, and not in the benzene nucleus of toluene. Having the CH₂OH group, on oxidation it yields benzoic aldehyde, C₆H₅COH (oil of bitter almonds), and benzoic acid, C₆H₅COOH.

b. Dihydroxyl Derivatives of Hydrocarbons.

Dihydric or Dihydroxylic or Diacid Alcohols.—Glycols, C_nH_{2n}-(OH)₂ series (see also p. 409).—Glycols may be viewed as dihydroxyl derivatives of the paraffins, the alcohols of the ethylic series being mono-derivatives:



They are prepared by acting on di-iodo-derivatives of the paraffins by silver acetate, and then treating with potash.



The glycols yield very interesting results on oxidation, forming two sets of acids, the lactic and the succinic series.

Aromatic Glycols, C_nH_{2n-8}(OH)₂, and *Saligenin Alcohols*.—Resorcin, pyrocatechin, and hydroquinone are dihydric alcohols of benzenes. (For their constitution see *Phenol*.)

Resorcin.

Synonyms.—Resorcinol; Metadioxybenzol.

It is obtained from various resins (such as *galbanum* and *asafoetida*), and on a large scale, in the arts, from crude benzene-sulphonic acid; it is purified by sublimation, and finally crystallized from benzene. *Resorcinum*, U. S. P., is a colorless substance which crystallizes in rhombic needles or plates, having a peculiar faint odor and an unpleasant taste. It is very soluble in water and alcohol, also in glycerin and ether, almost insoluble in chloroform. Ferric chloride colors the aqueous solution of resorcin a dark violet. If a small quantity of tartaric acid be gently heated with a trace of resorcin and a few drops of concentrated sulphuric acid, a crimson color will be produced, which will become a pale yellow on dilution with water. When resorcin is gently heated it should not emit a smell of phenol.

Toluene Dihydric Alcohols—*Orcin*, C₆H₃(OH)₂CH₃. This is found in lichens. *Hydroxybenzylic alcohol*, salicylic alcohol, saligenol, saligenin, C₆H₄OH·CH₂OH. This is obtained from the salicin of willow-bark. Having the hydroxyl group in the methane nucleus

as well as in the benzene nucleus, salicylic aldehyde ($C_6H_4OH \cdot COH$) and salicylic acid ($C_6H_4OH \cdot COOH$) are formed on oxidation.

c. Trihydroxyl Derivatives of Hydrocarbons.

Trihydric Alcohols.— $C_nH_{2n-1}(OH)_3$ series. Glycerols.


Glycerin.

Synonyms.—Glycerol;* Propenyl Alcohol; Glycerin, $C_3H_5(OH)_3$.

The propenyl (glycyl or glyceryl) of glycerin, in combination with many of the acidulous radicals of the acids, oleic, palmitic, stearic, etc., forms most of the solid fats and oils. When these latter substances are heated with metallic hydrates (even with water—hydrogen hydrate—at a temperature of 500° to 600° F.), double decomposition occurs, oleate, palmitate, or stearate of the metal is formed, and glycerol (propenyl hydrate) is set free. Hence glycerin is a by-product in the manufacture of soap, hard candles, and lead plaster. (*Vide Index.*)

Properties.—Glycerin is viscid when pure, specific gravity 1.28 (not below 1.25, U. S. P.), has a sweet taste, and is soluble in water or alcohol in all proportions. It has remarkable powers as a solvent, is a valuable *antiseptic* even when diluted with 10 parts of water, and useful as an emollient. In vacuo it may be distilled unchanged, but under ordinary atmospheric pressure it is decomposed by heat, especially if distillation be attempted in a flask or retort. In a shallow open vessel heat readily vaporizes it if a little water be present. From damp air glycerin absorbs moisture slowly, but in considerable proportions. Perfectly pure and anhydrous glycerin, at a few degrees below the freezing-point of water, sometimes solidifies to a mass of crystals.

Tests.—Heat 1 or 2 drops of glycerin in a test-tube, alone or with strong sulphuric acid, acid potassium sulphate, or other salt powerfully absorbent of water; vapors of acrolein, acrylic aldehyde (from *acer*, sharp, and *oleum*, oil), are evolved—



recognized by their powerfully irritating effects on the eyes and respiratory passages. If the glycerin be in solution, the latter must be evaporated as low as possible before the test is applied.

To a little weak solution of borax, reddened by the addition of phenol-phthalein, add a few drops of the solution (neutralized, if necessary) suspected to contain glycerin; if any is

* It will be noticed that all the alcohols have the termination *-ol*—carbinol, glycol, glycerol, saligenol, pyrogallol.

present, the color will be discharged, owing to the liberation of free boric acid, but will reappear on heating the solution; this reaction is also given by other polyhydric alcohols, such as mannite or glucose.

Add a few drops of the fluid suspected to contain the glycerin to a little powdered borax; stir well together; dip the looped end of a platinum wire into the mixture, and expose to an air-gas flame; a deep-green color is produced (Senier and Lowe).

If a very dilute solution of glycerin be mixed with zinc carbonate and then dried at 100°C ., and the dried mass extracted with absolute alcohol, on evaporation a sweetish residue is left behind. This is the official method of detecting glycerin.

The glycerin liberates boric acid, which colors the flame. (See p. 335.) Ammoniacal salts, which similarly affect borax, must first be got rid of by boiling with solution of sodium carbonate. Acids must also be neutralized. Liquids containing much indefinite organic matter must sometimes be evaporated to dryness, the residue extracted by alcohol, and the latter tested for the glycerin. To detect traces, liquids must be concentrated.

Glycerin, by action of very strong nitric acid, yields *nitro-glycerin*, or *glyceryl nitrate* ($\text{C}_3\text{H}_5\text{NO}_3$). It is highly explosive, a very small quantity being liable to explode during preparation and with great violence. 75 parts of nitroglycerin, absorbed by 25 of porous silica, yield a pasty mass more convenient to handle than nitroglycerin itself; it is used for blasting under the name of *dynamite*. Tablets of chocolate, weighing $2\frac{1}{2}$ grains and containing $\frac{1}{100}$ grain of nitroglycerin, constitute the *Tabellæ Nitroglycerini*, B. P. A "1 per cent." solution in rectified spirit (that is, 1 part by weight in 100 by volume) is the *Spiritus Glonoini*, or *Spirit of Nitroglycerin*, U. S. P.; *Liquor Trinitrini*, B. P.; *Liquor Nitroglycerini*, B. P.; or *Liquor Glonoini*. The latter solution has a specific gravity of 0.844, whereas the former solution has a range between 0.826 and 0.832 at 15°C .

"If the specific gravity of the spirit be higher than 0.840, or if 10 cc. of it be rendered turbid by less than 10 cc. of water, the spirit should be rejected."

Besides glycerin itself (*Glycerinum*, U. S. P.), solutions or mixtures of starch and of yolk of egg and glycerin (*Glyceritum Amyli*, U. S. P., *Glyceritum Vitelli* or *Glyconin*, U. S. P.) are official. Also *Glyceritum Acidi Carbolici*, U. S. P.; *Glyceritum Acidi Tannici*, U. S. P.; *Glyceritum Boroglycerini*, U. S. P.; and *Glyceritum Hydrastis*, U. S. P.

FATTY BODIES.

Processes of Extraction.—Fixed oils and fats are extracted from animal and vegetable substances by pressure or straining, with or without the aid of heat, or digestion in solvents, as ether, etc., and evaporation of the solvent.

Constitution and General Relations.—Fixed oils and fats are, apparently, almost as simple in constitution as ordinary inorganic salts. Just as potassium acetate ($\text{KC}_2\text{H}_3\text{O}_2$) is regarded as a compound of potassium (K) with the characteristic elements of all acetates ($\text{C}_2\text{H}_3\text{O}_2$), so soft soap is considered to be a compound of potassium (K) with the elements characteristic of all oleates ($\text{C}_{18}\text{H}_{33}\text{O}_2$), and hence is chemically termed potassium oleate ($\text{KC}_{18}\text{H}_{33}\text{O}_2$). Olive oil (*Oleum Olivæ*, U. S. P.), from which soap is officially prepared, is mainly oleate of the trivalent radical *glyceryl* (C_3H_5), the formula of such a fluid oil being $\text{C}_3\text{H}_5\text{C}_{18}\text{H}_{33}\text{O}_2$, and its name *oleine*. The formation of a soap therefore, on bringing together oil and a moist oxide or hydrate, is a simple case of double decomposition (or, rather, metathesis), as seen already in connection with lead plaster (p. 213), or in the following equation relating to the formation of common hard soap:



Berthelot has succeeded in preparing oil artificially from hydrogen oleate or oleic acid, $\text{HC}_{18}\text{H}_{33}\text{O}_2$, and glycerin, and it is said to be identical with the pure oleine of olive oil and of other fixed oils.

Olive oil is liable to contain cotton-seed oil. The admixture may be detected by Bechi's test: Take 1 grain of crystallized silver nitrate and dissolve it in the smallest possible quantity of water (about 1 cc.), and add 200 cc. of alcohol (96°). The addition also of 20 cc. of sulphuric ether is a good one, in that it makes the reagent better miscible with the oil to be examined, but it is not necessary. On the other hand, prepare a solution composed of 85 parts of amyl alcohol and 15 parts of oil of rape-seed. These reagents should be made as needed, and not kept on hand for any length of time. To apply the test, take 10 cc. of the oil to be examined, add 1 cc. of the alcoholic solution of silver nitrate, and then from 8 to 10 cc. of the mixture of amyl alcohol and oil of rape, agitating strongly and then heating on a water-bath for five or ten minutes. In the case of pure oils the color remains the same as it was after the addition of the reagents. If cotton-seed oil be present, there will be produced a brownish color or turbidity of a varying grade, from a very light brown to a deep maroon or black, according to the quantity of cotton oil present.

Hard fats chiefly consist of *stearine*—that is, of tristearate of glyceryl ($\text{C}_3\text{H}_5\text{C}_{18}\text{H}_{35}\text{O}_2$). Mr. Wilson, of Price's Candle Company, obtains stearic and oleic acids and glycerin by simply passing steam, heated to 500° or 600° F., through melted fat. Both the glycerin and fat-acids distil over in the current of steam, the glycerin dissolving in the condensed water, the fat-acids floating on the aqueous liquid. From glyceryl oleate and hydrogen hydrate there result hydrogen oleate and glyceryl hydrate.* The oleic acid (*Acidum*

* Any such decomposition of water and fixation of its elements, whether direct as above, or indirect through the intermediate agency of saponification, is termed *hydrolysis* (ὑδωρ, *hudor*, water, λύω, *luo*, to decom-

Oleicum, U. S. P.) is separated by cooling and pressing the mixture. It is "a straw-colored liquid, nearly odorless and tasteless, and with not more than a very faint acid reaction. Unduly exposed to air, it becomes brown and decidedly acid. Specific gravity, 0.860 to 0.890. It is insoluble in water, but readily soluble in alcohol, chloroform, and ether. At 40° to 41° F. (4.5° to 5° C.) it becomes semi-solid, melting again at 56° to 60° F. (13.3° to 15.5° C.). It should be completely saponified when warmed with potassium carbonate; and an aqueous solution of this salt, neutralized by acetic acid and treated with lead acetate, should yield a precipitate which, after washing with boiling water, is almost entirely soluble in ether," showing the absence of any important quantity of stearic and palmitic acids, the lead stearate and palmitate being insoluble in ether.

In a mixture of oils or fats and free fatty acids the latter may be estimated by taking advantage of their solubility in spirit of wine, and the formation of a neutral soap on shaking the spirituous solution with caustic soda, phenolphthalein being used as indicator. (See the section on the use of the caustic-soda solution in Volumetric Analysis.)

The author found (*Pharmaceutical Journal*, March, 1863) that oleic acid readily combines with alkaloids and most of the metallic oxides or hydrates, forming *oleates* which are soluble in fats. In this way active medicines may be administered internally in conjunction with oils or externally in the form of ointments (*Oleatum Hydrargyri*, U. S. P., and *Oleatum Zinci*, U. S. P.; *Oleatum Veratrinæ*, U. S. P.). Tichborne considers the formula of mercuric oleate to be $\text{Hg}(\text{C}_{18}\text{H}_{33}\text{O}_2)_2\text{H}_2\text{O}$.

Some fats, such as "suint" from sheep's wool and the unctuous matter from bristles, feathers, horn, and hair generally, yield by saponification, etc. fatty acids, and, instead of glycerin, *cholesterin*, an alcoholoid crystalline substance. The "lanolin" of pharmacy is cholesterin fat which has absorbed a large volume of water.

Adeps Lanæ, or Wool Fat, B. P., is "the purified cholesterin fat of sheep's wool." It is "a yellowish, tenacious, unctuous substance; almost inodorous, with a melting-point varying from 100° F. (37.8° C.) to 112° F. (44.4° C.); readily soluble in ether and in chloroform, sparingly soluble in rectified spirit. 10 grains should dissolve almost completely in 14 fluidrachms of boiling ethylic alcohol, the greater part separating in flocks on cooling. Ignited with free access of air, it burns, leaving but a trace of ash. 50 grains dissolved in 4 fluidrachms of ether and 2 drops of tincture of phenolphthalein added, should not require more than 2 grain-measures of volumetric solution of soda to produce a permanent red coloration. The solution in chloroform poured gently over the surface of sulphuric acid acquires a purple-red color. Heated with solution of soda, no ammoniacal odor should be evolved."

Adeps Lanæ Hydrosus, or *Hydrous Wool Fat*, U. S. P., is "the

pose). The fixation of water without such actual separation of its elements from each other is termed *hydration*.

purified fat of the wool of sheep, mixed with not more than 30 per cent. of water." It is commonly known as "lanoline." It is a "yellowish-white or nearly white ointment-like mass, having a faint, peculiar odor;" has a melting-point about 104° F. (40° C.).

Hydrous wool fat is insoluble in water, but mixes with twice its weight of it, dissolves with turbidity in ether and chloroform to form a neutral solution. When heated on a water-bath it leaves a residue of not less than 70 per cent., which should be completely soluble in ether and chloroform, and only partially so in alcohol; should give the same reactions as wool fat. "If 10 grm. of hydrous wool-fat be heated, together with 50 cc. of water, on a water-bath until the fat is melted, there should result an upper, translucent, and light-yellow, fatty layer, and a lower, clear, aqueous layer, which latter should not yield glycerin upon evaporation."—*U. S. P.*

Soaps.

Olive oil boiled with solution of potash yields potassium soap, or *soft soap* (*Sapo Mollis*, *U. S. P.*, or *Green Soap*); with soda, sodium soap, or *hard soap* (*Sapo*, *U. S. P.*), or *white Castile soap*, as distinguished from the variety of hard Castile or Marseilles soap, which is "mottled" by iron; soap mixed with ammonia, an ammonium soap (*Linimentum Ammoniacæ*, *U. S. P.*); and with lime-water, calcium soap (*Linimentum Calcis*, *U. S. P.*),—all oleates, chiefly of the respective basylous radicals. Their mode of formation is indicated in the equation on p. 459. The alkali soaps are soluble in alcohol, the others insoluble. A green soap, much used on the continent of Europe, and indeed official in Germany (formerly as *Sapo Viridis*, now as *Sapo Kalinus Venalis*), is made by adding indigo to ordinary soft soap, the yellow color of the soap yielding with the indigo a greenish compound. The official characters of hard soap are—"grayish-white, dry, inodorous; horny and pulverizable when kept in dry warm air; easily moulded when heated; soluble in rectified spirit; not imparting an oily stain to paper; incinerated, it yields an ash which does not deliquesce;" and of soft soap—"yellowish-green, inodorous, of a gelatinous consistence; soluble in rectified spirit; not imparting an oily stain to paper; incinerated, it yields an ash which is very deliquescent." Curd soap (*Sapo Animalis*, *B. P.*) is "a soap made with soda and a purified animal fat, consisting principally of stearine." It will, of course, chiefly contain sodium stearate. In pharmacy it is often advantageously employed instead of the "hard soap."

The hard soap met with in trade is made from all varieties of oil, the commoner kinds being simply the product of the evaporated mixture of oil and alkali, while the better sorts have been separated from alkaline impurities and the glycerin by the addition of common salt or excess of lyè to the liquors, which causes the precipitation of the pure soap as a curd. Potash soap is not so readily precipitable by salt; moreover, some soda soap results. Saponification on the small scale is much facilitated by first well mixing the oil with 5 per cent. of sulphuric acid, and letting this mixture stand for twenty-four hours. The dark product is then readily soluble

when boiled with soda, and the clear fluid yields a crust of white soap on cooling. If required quite free from alkali, the resulting soap is boiled with water until dissolved, salt added, and the whole cooled. A cake of pure soap results.

Yellow soap is a common, cheap soap, containing a good deal of resin soap, resin consisting chiefly of acids—pinic, sylvic, pimaric, etc.—which readily unite with alkalies to form true soaps.

Saponification.—This term is now extended in chemistry so as to include any process analogous to the foregoing—any reaction in which an alkali decomposes any ethereal salt or alkyl salt.

Solid Fats.

1. *Lard* (*Adeps*, U. S. P.) is the purified internal fat of the abdomen of the hog—the perfectly fresh *omentum* or *flare*, freely exposed to the air to dissipate animal odor, rubbed to break up the membranous vesicles, melted at about 130° F. (54.4° C.), and filtered through paper or flannel. *Lard Oil* (*Oleum Adipis*, U. S. P.), which is chiefly olein, is a “fixed oil expressed from lard at a low temperature.” Sp. gr. 0.90 to 0.920 at 15° C. 2. *Benzoinated Lard* (*Adeps Benzoinatus*, U. S. P.) is prepared lard heated over a water-bath with benzoin (1 part to 50), which communicates an agreeable odor and prevents or retards rancidity. Purified lard is a mixture of oleine (*lard oil*, removable by pressure) and stearine. *Margarine*, formerly supposed to be a constituent of lard and other soft fats, is now regarded as a mere mixture of *palmitine* (the chief fat of palm oil) and stearine. 3. *Suet*, the internal fat of the abdomen of the sheep, purified by melting and straining, forms the official *Sevum*, U. S. P.; it is almost exclusively composed of stearine ($C_3H_5 \cdot 3C_{18}H_{35}O_2$). 4. *Expressed Oil of Nutmeg* (*Oleum Myristicæ Expressum*, B. P.), commonly but erroneously termed *Oil of Mace*, is a mixture of a little volatile oil with much yellow and white fat; the latter is *myristin* or myristate of glyceryl ($C_3H_5 \cdot 3C_{14}H_{27}O_2$). 5. *Oil of Theobroma*, or *Cacao Butter* (*Oleum Theobromatis*, U. S. P.), chiefly stearine, but with one higher and some lower homologues (Heintz), is a solid product of the roasted and crushed seeds or *cocoa-nibs* of *Theobroma cacao*. They contain from one-half to two-thirds of this fat. [Cocoa is too rich for use as food, hence is diluted with farina (*cheap cocoa*) or sugar (*chocolate*) or has a portion of its fat extracted, while its solubility is, in one brand, usefully increased by a slight addition to its potassium.] 6. *Cocoa-nut Oil*, or butter, a soft fat contained in the edible portion of the nut of *Cocos nucifera*, or *cocconut* of the shops, is a body containing glyceryl united with six acidulous radicals—namely, the caproic ($C_6H_{11}O_2$), caprylic ($C_8H_{15}O_2$), rutic ($C_{10}H_{19}O_2$), lauric ($C_{12}H_{23}O_2$), myristic ($C_{14}H_{27}O_2$), and palmitic ($C_{16}H_{31}O_2$)—radicals which, like some from resin, when united with sodium, form a soap differing from ordinary hard soap (sodium oleate) by being tolerably soluble in a solution of sodium chloride; hence the use of cocoa-nut oil and resin in making *marine soap*, a soap which, for the reason just indicated, readily yields a lather in sea-water. 7. *Kokum Butter*, *Garcinia Oil*, or *Concrete Oil* of

Mangosteen, a whitish or yellowish-white fat obtained from the seeds of *Garcinia Indica* or *G. purpurea*, is composed of stearine, myristicine, and oleine. It is recognized officially in the Pharmacopœia of India (*Garciniæ Purpureæ Oleum*).

Butter commonly yields $87\frac{1}{2}$ per cent. of insoluble fat acids by saponification and decomposition of the soap by acid. Other animal fats, with which butter is likely to be adulterated, yield about $95\frac{1}{2}$. Hence the percentage of fat acids, and, especially, volatile acids, insoluble acids, and soluble acids, yielded by a suspected sample of butter, indicates purity or the opposite. Occasionally, however, a sample of genuine butter may not conform to the figures, hence they cannot be relied on to show the exact extent of sophistication.

Fixed Oils.

Fixed and volatile oils are naturally distinguished by their behavior when heated; they also generally differ in chemical constitution—a fixed oil being, apparently, a combination of a basylous with an acidulous radical, as already stated, while a volatile oil is more commonly a neutral or normal hydrocarbon, mixed with a comparatively small proportion of a body—containing oxygen as well as carbon and hydrogen—to which the odor of the oil is generally due. The latter bodies are now articles of trade under the name of “concentrated essential oils.”

Drying and Non-drying Oils.—Among fixed oils, most of which are oleate with a little palmitate and stearate of glyceryl, a few, such as—1, *Linseed Oil* (*Oleum Lini*, U. S. P., contained in *linseed* or *flaxseed*, the ground residue of which “should yield, when extracted with disulphide of carbon, not less than 25 per cent. of fixed oil,” is *linseed meal*), and, 2, *Cod-liver Oil* (*Oleum Morrhue*, U. S. P.), and, to some extent, *castor* and *croton*, are known as *drying oils*, from the readiness with which they absorb oxygen and become hardened to a resin. Linseed commonly contains 37 or 38 per cent. of oil; 25 to 27 per cent. is obtained by submitting the ground seeds to hydraulic pressure, 10 to 12 per cent. remaining in the residual *oil-cake*. *Boiled oil* is linseed oil which has been boiled with lead oxide. This treatment increases the already great tendency of linseed oil to resinify, forming *linoxyn* ($C_{32}H_{54}O_{11}$) on exposure to air. The drying oils appear to contain linoleine, an oily body distinct from oleine. Cod-liver oil contains an unimportant trace of iodine, 1 in 1,000,000 or 2,000,000 parts, according to Stanford; a little choline also and other bases, Gautier and Mourgues having recently isolated *aselline*, $C_{25}H_{32}N_4$, and *morrhaine*, $C_{19}H_{27}N_3$, besides butyl-, amyl-, and hexylamines and dihydro-lutidine. Among the *non-drying oils* are the following: 3. *Almond Oil* (*Oleum Amygdalæ Expressum*, U. S. P.), indifferently yielded by the bitter (*Amygdalæ Amara*, U. S. P.) or sweet seed (*Amygdalæ Dulcis*, U. S. P.) to the extent of 45 and 50 per cent. respectively. (3a) *Cotton-seed Oil* (*Oleum Gossypii Seminis*, U. S. P.) contains oleine and some palmitine. Sp. gr. 0.920 to 0.930. It should not be permanently colored dirty-yellow by sulphuric acid. 4. *Croton Oil* (*Oleum Crotonis*, B. P., and *Oleum Tiglii*, U. S. P.). Geuther states that no such acid as crotonic is

obtainable from croton oil, but acetic, butyric, valerianic, and higher members of the oleic series, together with *tiglic acid*, $\text{HC}_5\text{H}_9\text{O}_2$. H. Senior states that alcohol separates croton oil into a soluble oil containing the powerful vesicating principle of croton oil and an insoluble non-vesicating but powerful purgative principle. Kobert states that free crotonoleic acid is both the vesicant and the purgative.

5. *Lycopodium* (U. S. P.), a yellow powder composed of the spores of the common club-moss (*Lycopodium clavatum*), contains a large proportion of a very fluid fixed oil; also an alkaloid (Bödeker), $\text{C}_{32}\text{H}_{52}\text{N}_2\text{O}_3$.

6. *Olive Oil* (*Oleum Olivæ*, U. S. P.), already noticed (p. 459). "If 1 grm. of olive oil be agitated in a test-tube with 2 grm. of cold mixture prepared from equal volumes of strong sulphuric acid and of nitric acid of sp. gr. 1.185, and the mixture be set aside for half an hour, the supernatant oily layer should not have a darker tint than yellowish, nor should a green or red layer separate on standing if 1 grm. of the oil be shaken for a few seconds with 1 grm. of a cold mixture of sulphuric acid (sp. gr. 1.830) and nitric acid (sp. gr. 1.250), and 1 grm. of carbon disulphide; and if 5 drops of the oil are let fall upon a thin layer of sulphuric acid in a flat-bottomed capsule, no brown-red or dark-brown zone should be developed within three minutes at the line of contact of the two liquids (absence of appreciable quantities of other fixed oils of similar physical properties)." — U. S. P.

7. *Castor Oil* (*Oleum Ricini*, U. S. P.), chiefly a *ricinoleate of glyceryl* ($\text{C}_3\text{H}_5\text{C}_{18}\text{H}_{33}\text{O}_3$) or *ricinoleine*, a slightly oxidized oleine, soluble, unlike most fixed oils, in alcohol and in glacial acetic acid. Castor-oil seeds were stated, by Tuson, to contain an alkaloid, *ricinine*. Beck has recently confirmed Tuson, giving as the formula $\text{C}_{24}\text{H}_{32}\text{N}_7\text{O}_3$. It possesses no purgative property. They also contain an albumose, *ricin*, resembling, physiologically, but not quite chemically, the *abrin* of jequirity.

8. *Oil of Male Fern* (*Filix Mas*, B. P.), a vermifuge obtained by exhausting the rhizome (*Aspidium*, U. S. P.) with ether and removing the ether by evaporation—a dark-colored oil containing a little volatile oil and resin, and officially termed an oleoresin (*Oleoresina Aspidii*, U. S. P.). Its chief active constituent appears to be *filicic acid*, $\text{C}_{35}\text{H}_{42}\text{O}_{13}$.

9. *Fixed Oil of Mustard*, a bland, inodorous, yellow or amber oil yielding, by saponification and action of sulphuric acid, glycerin, oleic acid, and *erucic acid* ($\text{HC}_{22}\text{H}_{41}\text{O}_2$) (Darby).

10. *Arachis Oil* (*Oleum Arachis*, P. I.) is found to the extent of 40 or 50 per cent. in the seeds of the *Arachis hypogæa* (P. I.), the ground-nut or earth-nut (so called because the pod of the herb in the growth of its stalk downward is forced beneath the surface of the ground and there ripens). It is chiefly oleine, but contains hypogæine, palmitine, and arachine. The oil is largely used in India in the place of olive oil, and is becoming much employed in Europe, especially for soap-making.

11. *Sesamé Oil*, or *Oil of Sesamum* (*Oleum Sesami*, U. S. P.) (Gingelly, Teal, or Benné Oil), from the seeds of *Sesamum indicum*, is also largely used in Europe. It has most of the characters of the best olive oil. It may be detected in olive oil by shaking the sample with a solution of pyrogallic acid in strong hydrochloric acid, and separating and boiling the acid fluid, a purplish color resulting if sesamé be

present. 12. *Shark-liver Oil*, from *Squalus carcharias* (*Oleum Squalæ*, P. I.), is used as a substitute for cod-liver oil in India.


Trihydric Alcohols of the $C_nH_{2n-9}(OH)_3$ series.

Pyrogallol or *Pyrogallic Acid*.—Trihydroxybenzene, $C_6H_3(OH)_3$. (*Vide Index*.)

d. Other Polyhydroxyl Derivatives of Hydrocarbons.

Only one *tetrahydric alcohol* is known—namely *Erythrite*, or *Lichen Sugar*, $C_4H_6(OH)_4$, found in *Protococcus vulgaris*, *Roccella tinctoria*, and *R. fuciformis*. *Quercite*, the sugar of acorns, is *penthydric*: *Mannite* is *hexahydric*. *Sorbite* occurs in the fruits of the order *Rosaceæ*.

Hexahydric Alcohols.—*Mannite*, $C_6H_8(OH)_6$.—Boil manna with 15 or 16 parts of alcohol, filter, and set aside; mannite separates in colorless shining crystals or acicular masses to the extent of from 60 to 80 per cent. of the manna. It is closely related to the ordinary sugars, glucose becoming mannite by action of nascent hydrogen:



Mannite does not undergo fermentation in contact with yeast. With nitric acid it forms explosive *nitromannite*, $C_6H_8(NO_3)_6$.

Manna, U. S. P., is a concrete saccharine exudation obtained by making transverse incisions in the stems of cultivated trees of *Fraxinus Ornus*. It occurs in stalactitic pieces, varying in length and thickness, flattened or somewhat concave on their inner surface, and of a pale yellowish-brown color, and nearly white externally. This manna, which is known as flake manna, is crisp, brittle, porous, crystalline in structure, and readily soluble in about 6 parts of water. Odor faint, resembling honey; taste sweet and honey-like, combined with a slight acidity and bitterness. It contains about 10 per cent. of moisture. Mannite is also met with in celery, onions, asparagus, certain fungi and sea-weeds, occurs in the exudations of apple and pear trees, and is produced during the viscous fermentation of sugar.

Dulcite, isomeric with mannite, is formed by the action of nascent hydrogen on inverted milk-sugar. It differs from mannite by oxidizing to mucic acid, $C_6H_{10}O_8$, when treated with nitric acid.

QUESTIONS AND EXERCISES.

How is phenol artificially and commercially prepared?—How would you distinguish carbolic acid from creasote?—Give the formulæ and


systematic names for picric acid, sodium carbolate, and resorcin.—Give names for the bodies having the formulæ $C_6H_4OH \cdot CH_3$ and $C_6H_5CH_2OH$.—What are glycols? how prepared?—Give formula and mention the chief properties of glycerin.—What is the specific gravity of glycerin?—By what test is glycerin recognized?—Enumerate some official preparations in which glycerin is employed. Give a sketch of the general chemistry of fixed oils, fats, and soaps.—What is the difference between hard and soft soap?—Which soaps are official?—Name the source of lard.—How is "Prepared Lard" obtained?—Mention the chief constituent of suet.—Whence is cacao-butter obtained?—Why is *marine soap* so called? and from what fatty matter is it almost exclusively prepared?—What do you understand by *drying* and *non-drying* oils?—In what respect does castor oil differ from other oils?—How is oil of male fern (*Ex. Filicis Liquidum*) prepared?—Classify pyrogallol (*pyrogallie acid*), erythrite, mannite, and duleite.—Describe the source and characters of manna.

CARBOHYDRATES.

Under the name *carbohydrates* have been grouped a large number of compounds containing carbon with hydrogen and oxygen in the same proportion as in water, and whose constitution has only recently been understood. They include sugars, dextrin, starch, cellulose, etc. The molecules of some of the latter are very complex, but are resolved by hydrolysis into sugars such as glucose.

The most commonly occurring carbohydrates contain six carbon atoms or even multiples of that number, but analogous bodies with three, five, seven, or nine carbon atoms in the molecule have lately been discovered.

The sugars are the simplest of the carbohydrates in constitution, and a large number of them, some identical with previously known natural sugars, some previously unknown, have been synthesized. They are partially oxidized polyhydric alcohols, having one of their alcohol groupings oxidized into an aldehyde or ketone group. For example, the trihydric alcohol glycerin, on gentle oxidation with bromine, yields a body, *glycerose*, having all the characters of a sugar.



This, however, is not stable, but spontaneously condenses into a glucose, $C_6H_{12}O_6$.

Erythrose, $C_4H_8O_4$, is an example of a sugar with four, and *arabinose*, $C_5H_{10}O_5$, of one with five, atoms of carbon. Most of the natural sugars are glucoses ($C_6H_{12}O_6$) or compounds of two or three molecules of glucoses minus water (*bioses* or *trioses*).

Sugars with seven, eight, and nine atoms of carbon have been constructed by treating glucoses with hydrocyanic acid, which combines with the aldehyde or ketone group to form the nitrite of an acid containing one more carbon atom. This on hydrolysis gives the acid, the lactone of which may be reduced to the corresponding

sugar by the action of nascent hydrogen. This process is then repeated to get an eight-carbon sugar, and so on. One of these seven-carbon sugars was found to be identical with a natural sugar, *perseite*, but most of them have not yet been found occurring naturally.

Glucoses, $C_6H_{12}O_6$.

Glucoses, $C_6H_{12}O_6$.—There are two chief types of these six-carbon sugars, differing from each other in the position of the alcohol grouping that has undergone oxidation, and classed accordingly as aldehyde and ketone sugars—*aldose* and *ketose*. *Dextrose* is an example of the first, and *lævulose* or *fructose* of the second class. Each of these classes contains a very large number of physical isomers, differing from each other in their action on polarized light and in some other respects; these may be most readily distinguished from one another by means of the physical characters of the compounds they form with phenyl-hydrazine. (See Index.) The large number of these isomers is accounted for, on the stereo-chemical theory, by the circumstance of there being no less than four asymmetrical carbon atoms in each molecule. Thus there are three dextroses—dextro-rotatory, lævo-rotatory, and inactive; three analogous mannoses; three fructoses or lævuloses, etc. Ordinary lævulose, or, better, fructose, is not, therefore, the mere optical isomer of dextrose, each of them having dextro-, lævo-, and inactive forms.

All the glucoses above mentioned have been obtained artificially, the starting-point being an artificial glucose (or *acrose*, $C_6H_{12}O_6$) obtained by the condensation of formic aldehyde, $CH_2O(6CH_2O = C_6H_{12}O_6)$; it is probably in a similar way that natural sugars are produced by plants.

Dextrose, or *Grape-sugar*, or *Glucose* (from $\gamma\lambda\upsilon\kappa\acute{\iota}\varsigma$, *glucūs*, sweet), is often seen in the crystallized state in dried grapes or raisins and other fruits; it is also the variety of sugar met with in diabetic urine. Its crystalline character is quite distinct from that of cane-sugar, the latter forming large four- or six-sided rhomboidal prisms, while grape-sugar occurs in masses of small cubes or square plates. Grape-sugar is also less soluble in water, but more soluble in alcohol, than cane-sugar.

According to Fresenius, the percentage proportion of saccharine matter in the dried fig is 60 to 70; grape, 10 to 20; cherry, 11; mulberry, 9; currant, 6; whortleberry, 6; strawberry, 6; raspberry (*Rubus Idæus*, U. S. P.), 4.

Fructose or *lævulose* is lævogyrate, while sucrose and glucose possess right-handed rotation; the latter twist a ray of polarized light from left to right to an extent dependent on the amount of sugar present—a fact easy of application in estimating the amount of sugar in syrups or in diabetic urine.

Fructose or *lævulose* is the uncrystallizable or very difficultly crystallizable constituent of inverted cane-sugar. It is found in the grape, fig (*Ficus*, U. S. P.), cherry, gooseberry, strawberry, peach, plum, and other fruits, often with dextrose or with cane-sugar. Fruit-sugar reduces cupric salts and silver ammonio-nitrate.

Artificial Formation of Grape-sugar from Cane-sugar—Tests for Sugar.—Dissolve a grain or two of common cane-sugar in water. To a portion of this solution placed in a test-tube add more water, two or three drops of solution of copper sulphate, a considerable quantity of solution of potash or soda (enough to turn the color of the liquid from a light to a dark blue), and heat the mixture to the boiling-point; no obvious immediate change occurs. To another portion of the syrup add a drop of sulphuric acid, and boil for ten or twenty minutes; then add the copper solution and alkali, and heat as before; a yellowish-red precipitate of cuprous oxide (Cu_2O) falls. This test is exceedingly delicate.

The above reaction is due to the conversion of the *cane-sugar* ($\text{C}_{12}\text{H}_{22}\text{O}_{11}$) into *inverted sugar*—or *lævulose*, $\text{C}_6\text{H}_{12}\text{O}_6$ (so called because its solution causes left-handed rotation of a ray of polarized light, cane-sugar having an opposite effect) and *grape-sugar*, $\text{C}_6\text{H}_{12}\text{O}_6, \text{H}_2\text{O}$ —by the influence of the sulphuric acid, and to the reducing action of the lævulose and grape-sugar on the cupric solution. The formation of a precipitate immediately, without the action of acid, shows the presence of the latter sugars—its formation only after ebullition with acid indicating, in the absence of starch or dextrin, cane-sugar. In this reduction-process the sugar is oxidized and broken up into several substances, but the exact nature of the reaction has not been ascertained.

Dextrin also reduces the copper salt to suboxide, unless its solution is cold and very dilute. It does not, however, so act on a solution of cupric acetate acidified with acetic acid, while glucose produces with this liquid the usual red cuprous precipitate (Barfoed).

Sugar from Starch.—Boil starch with a little water and a drop of sulphuric acid as for dextrin, but continue the ebullition for several minutes; on testing a portion of the cooled liquid with iodine and another portion with the heated alkaline solution of a copper salt, as described above, it will be found that the starch has nearly all become converted into a sugar—dextrose. Maltose is also formed, at first, but by the continued action of the acid is changed to dextrose. When made on a large scale a warm (131°F.) mixture of starch and water of the consistence of cream is slowly poured into a boiling solution of 1 part of sulphuric acid in 100 of water, the whole boiled for some time, the acid neutralized by chalk, the mixture filtered, the liquid evaporated to a thick syrup and set aside; in a few days it crystallizes to a granular mass resembling honey. In this operation a small quantity of dextrin remains with the glucose; but if the process be conducted under pressure, conversion, according to Manbré, is complete. Sugar made from the starch of rice, maize, etc. is largely used

for table syrups, confectioneries, bee-food, and as a partial substitute for malt in brewing. It is known as *patent sugar*, *saccharine*, *maltose*, etc.

In the United States the dealers term the syrups "glucose," and the further evaporated solid product "grape-sugar." The former contain one-third or more of dextrose, about one-fifth of maltose, one-fourth or more of dextrin, and about one-sixth or one-fifth of water; the latter often contain about three-fourths of dextrose, from none up to one-third of maltose, and one-seventh or one-sixth of water.

Galactose (from milk-sugar), *Sorbinose* (from mountain-ash berries), *Inosite* (from muscles), *Mannose*, (from mannite), *Gulose*, *Formose*, β -*acrose*, *Rhamnose*, *Dambose* (from a caoutchouc), and *Scyllite* (from many fish), are other glucoses.

Saccharoses, or Bioses, $C_{12}H_{22}O_{11}$.

Cane-sugar, or *Sucrose* (*Saccharum Purificatum*, U. S. P.), is a frequent constituent of vegetable juices. Thus it forms the chief portion of cassia-pulp (*Cassia Pulpa*, U. S. P.), is contained in the carrot and turnip, but is most plentiful in the sugar-cane; much, however, is now obtained from the sugar-maple and beet-root. On evaporation of the juice common *brown* or *moist sugar* crystallizes out; this by re-solution, filtration through animal charcoal, evaporation to a strong syrup, and crystallization in moulds, yields the compact crystalline conical loaves known in trade as *lump sugar*. From a slightly less strong syrup, slowly cooled, the crystals termed *sugar-candy* are deposited, white or colored, according to the color of the syrup. The official syrup (*Syrupus*, U. S. P.) is an aqueous solution, consisting of 85 grammes of sugar dissolved in water and made up to 100 cc.

The sugar in fresh fruits is mainly cane-sugar, but by the action of the acid, or possibly of a ferment in the juice, it is gradually converted into *inverted sugar*, a variety differing from cane-sugar in being uncrystallizable, and in having an inverted or opposite influence on polarized light, twisting the ray from right to left (*lævo-gyrate*, having *lævo-rotation*). Ripe hips (*Rosæ Caninæ Fructus*, B. P.) contains 30 per cent. of such sugar, besides gum and acid malates and citrates. Fruit-sugar, as gathered in the form of syrup by bees, is probably a mixture of these two varieties. It is gradually altered to a crystalline or granular mass of grape-sugar, as seen in dried fruits, such as raisins (*Uvæ*, B. P.) and the prune (*Prunum*, U. S. P.), and in solidified honey (*Mel*, B. P., U. S. P.). This, the common form of grape-sugar, is *dextrogyrate*, and hence is sometimes termed *dextrose*, to distinguish it from *lævulose*. Diluted with twice its weight of water, it yields a liquid having the sp. gr. 1.101 to 1.150. Honey often contains pollen, hairs, spores, the dust and dirt from the flowers, and various flocculent matters which cause it to ferment and yield mannite, alcohol, and acetic acid; hence for use in medicine it is directed (*Mel Despumatum*, U. S. P.) to be clarified by melting and straining while hot through flannel previously

moistened with warm water. A mixture of clarified honey 80 per cent., acetic acid 10 per cent., and water 10 per cent. is official under the name of *Oxymel* (from ὄξις, *oxus*, acid, and μέλι, *meli*, honey). A similar mixture of honey with acetic acid containing the soluble portions of squill-bulbs (*Scilla*, B. P.) is known as *Oxymel of Squill* (*Oxymel Scillæ*, B. P.). Honey and cane-sugar are the bases of the official *Confections*.

Maltose, $C_{12}H_{22}O_{11}$.—This crystallizable sugar is formed, together with dextrin, when diastase or dilute acids act upon starch. In the case of diastase it is the ultimate product, but the dilute acids may convert it into dextrose. It differs also from dextrose in its optical activity.

Cane-sugar, maltose, and grape-sugar yield alcohol and carbonic acid gas by fermentation, the cane-sugar nearly always passing into grape-sugar before the production of alcohol commences.



In *bread-making* some of the starch is converted into dextrin, and this into sugar by the ferment. The above action then goes on, the liberation of gas producing the *rising* or swelling of the mixture of flour, water, and yeast (dough), the temperature to which the mass is subjected in the oven causing escape of most of the alcohol and further expansion of the bubbles of carbonic acid gas in every part of the now spongy loaf. The carbonic acid gas gradually evolved when flour is worked up for bread with a mixture of dry sodium bicarbonate and tartaric acid (best preserved by previous admixture with dried flour and a little magnesium carbonate)—*baking-powder*—exerts similar influence. The least objectionable method of introducing carbonic acid gas, however, is that of Daughlish, whose patent aerated bread is made from flour by admixture with carbonic acid water under pressure by the aid of machinery. On removal from the cylinder the resulting dough expands by the natural elasticity of the imprisoned carbonic acid gas, and the bake-oven completes the process. The crumb of bread is official (*Mica Panis*, B. P.). All fermented bread retains, obviously, a little alcohol, sometimes 0.25 per cent.

Action of Alkali on Sugar.—To a little solution of grape-sugar add solution of potash or soda or solution of potassium carbonate, and warm the mixture; the liquid is darkened in color from amber to brown according to the amount of sugar present. A trace of picric acid greatly intensifies the color.

Tests.—The copper reaction, the fermentation process, and the effect of alkalies form three good tests of the presence of grape-sugar and, indirectly, of cane-sugar. A piece of merino or other woollen material, previously dipped in a solution of stannic chloride and dried, becomes of a brown or black color when dipped in a solution of glucose and heated to about 300° F. by holding before a fire.

Melitose or *Melitriose* (from eucalyptus) is a triose, giving on hydrolysis galactose, dextrose, and fructose. *Meletizose* (from the larch), *Trehalose* (from Turkish manna), and *Maltose* (from starch), belong to the saccharoses.

"*Honey-dew*" is a viscid saccharine matter occasionally met with on the leaves of the lime, maple, black alder, rose, and other trees, being a sweet principle exuded from aphides. Sometimes it is sufficiently abundant to dry and fall on the ground, forming a veritable "shower of manna." It is a mixture of cane-sugar, inverted sugar, and dextrin.

Barley-sugar is made by simply heating cane-sugar till it fuses, a change from the crystalline to the uncrystallizable condition occurring. *Treacle* (*Theriaca*, B. P.), *Molasses* or *Melasses* (from *Mel*, honey), or *Golden Syrup* chiefly results from the application of too much heat in evaporating the syrups of the sugar-cane; it is a mixture of cane-sugar with uncrystallizable sugar and more or less coloring matter. Liquorice-root (*Glycyrrhizæ Radix*, B. P.) contains a considerable quantity of uncrystallizable sugar.

Caramel.—Heat a grain or two of sugar in a test-tube until it blackens and froths; the product is *caramel* or *burnt sugar* (the *Saccharum Ustum* of pharmacy). It is used as a coloring agent for gravies, confectioneries, spirits, vinegar, and similar materials. It is a mixture of caramels.

Milk-sugar, or *Lactose* ($C_{12}H_{22}O_{11}H_2O$) (*Saccharum Lactis*, U. S. P.), the sweet principle of milk of animals, is not susceptible of alcoholic or vinous fermentation by ordinary yeast; certain varieties of the fungus, however, convert it into alcohol. It resembles grape-sugar in reducing an alkaline solution of copper with precipitation of suboxide. It is obtained from milk by adding a few drops of acid, stirring, setting aside for the *curds* to separate, filtering, evaporating the *whey* to a small bulk, filtering again if necessary, and allowing to cool and crystallize. The deposited crude "*sugar-sand*" is afterward refined and recrystallized. It usually occurs in trade "in cylindrical masses two inches in diameter, with a cord or stick in the axis, or in fragments of cakes—grayish-white, crystalline on the surface and in its texture, translucent, hard, scentless, faintly sweet, gritty when chewed." Thus obtained, milk-sugar has the formula above given, but if deposited during evaporation the crystals are anhydrous, $C_{12}H_{22}O_{11}$. It is soluble in 6 parts of cold and 3 of boiling water; slightly soluble in alcohol, insoluble in ether. Powdered milk-sugar is used in pharmacy as a vehicle for potent solid medicines. Milk-sugar is convertible, by the action of dilute acids, into galactose and dextrose; these may be reunited to form milk-sugar.

"If about 1 gram. of powdered sugar of milk be sprinkled upon about 5 cc. of cold sulphuric acid contained in a flat-bottomed

capsule, the acid may acquire a greenish or reddish, but no brown or brownish-black, color within half an hour (absence of cane-sugar).'
—*U. S. P.*

Saccharic Acid, $\text{H}_2\text{C}_6\text{H}_8\text{O}_8$ or $\text{C}_4\text{H}_4(\text{OH})_4(\text{COOH})_2$, is the result of oxidizing sucrose, dextrose, starch, gum, and lignin by nitric acid. *Mucic Acid*, isomeric with saccharic acid, may be obtained in the same way by acting on lactose, gum, and dulcite.

QUESTIONS AND EXERCISES.

Into what three classes may the carbohydrates be divided?—How is grape-sugar obtained from cane-sugar?—How are cane-sugar and grape-sugar analytically distinguished?—How is dextrose obtained from starch?—Mention the chief sources of cane-sugar.—Give chemical explanations of the processes of bread-making.—What is the difference between fruit-sugar and honey?—What is oxymel?—Describe the effect of heat on cane-sugar.—How is milk-sugar obtained?—How does it differ from other sugars?—Whence are mucic and saccharic acids obtained?

Amyloses, or Amyloids, $n\text{C}_6\text{H}_{10}\text{O}_5$.

Starch ($n\text{C}_6\text{H}_{10}\text{O}_5$) is contained in large or small quantities in nearly every plant. It forms about 60–70 per cent. of wheat and from 20 to 30 per cent. of potatoes. The starch officially recognized in the United States Pharmacopœia (*Amylum*) is that of maize (*Zea Mays*).

Processes.—Rasp or grate or scrape with a knife a portion of a clean raw potato, letting the pulp fall on to a piece of muslin placéd over a small dish or test-glass, and then pour a slow stream of water over the pulp; minute particules or *granules* of starch pass through the muslin and sink to the bottom of the vessel, fibrous matter remaining on the sieve. This is potato starch. Even diseased potatoes furnish good starch by this method. Wheat starch may be obtained by tying up some flour in a piece of calico and kneading the bag in a slow stream of water flowing from a tap, the washings running into a deep vessel, at the bottom of which the white starch collects; the sticky matter remaining in the bag is *gluten*.

The *blue starch* of the shops is artificially colored with smalt or indigo to neutralize the yellow tint of recently washed linen; it should not be used for medicinal purposes. Starch dried in mass splits up into curious columnar masses, resembling the basaltic pillars of Fingal's Cave in Staffa or those of the Giant's Causeway in the north of Ireland. The cause of the phenomenon, which may also be seen in grain tin, is not conclusively known.

Gluten is the body which gives tenacity to dough and bread. It seems to be a mixture of vegetable fibrin, vegetable casein, and an albuminous matter termed gluten. These substances and gluten itself are closely allied; each contains about 16 per cent. of nitrogen. *Wheaten Flour* (*Farina Tritici*, B. P.) contains about 72 per cent. of starch and 11 of gluten, as well as sugar, gum, fine bran, water, and ash. The compactness of barley, well seen in husked or pearl barley (*Hordeum decortiatum*, B. P.), is said to be due to the large amount of vegetable fibrin present. During germination the fibrin is destroyed; hence, probably, the cretaceous character of malt. *Oatmeal* (*Avenæ Farina*, U. S. P.), popular as "porridge," is rich in albumenoids or flesh-forming constituents, containing nearly 16 per cent. *Sago* is granulated starch from the sago palm; *tapioca* from the bitter cassava; each has less than 1 per cent. of albumenoids. The white translucent rice grains are the husked seeds of *Oryza sativa*. Rice (*Oryza*) and its flour, or ground rice (*Oryzæ Farina*), are official in the Pharmacopœia of India. Rice is a staple article of food in tropical countries. Ground rice resembles flour of wheat in composition, but contains from 85 to 90 per cent. of starch.

Mucilage of Starch.—Mix two or three grains of starch with first a little and then more water, and heat to the boiling-point; starch mucilage (*Mucilago Amyli*, B. P.) results. 1 part of starch to 200 of water gives "*Starch Test-solution*," U. S. P.

This mucilage or paste is not a true solution; by long boiling, however, a portion of the starch becomes dissolved. In the latter case the starch probably becomes somewhat altered.

Chemical Test.—To some of the mucilage add a very little free iodine; a deep-blue color is produced.

This reaction is a very delicate test of the presence of either iodine or starch. The starch must be in the state of mucilage; hence in testing for starch the substance supposed to contain it must be first boiled with water. The solutions used in the reaction should also be cold, or nearly so, as the blue color disappears on heating, though it is partially restored on cooling. The iodine reagent may be iodine-water or tincture of iodine. In testing for iodine its occurrence in the free state must be ensured by the addition of a drop, or even less, of chlorine-water. Excess of chlorine must be avoided, or iodine chloride will be formed, which does not color starch.

The so-called *starch iodide* scarcely merits the name of a chemical compound, the state of union of its constituents being so feeble as to be decomposed at 100° F. Substances that attack free iodine remove that element from starch iodide. The alkalies, hydrosulphuric acid, sulphurous acid, and other reducing agents, destroy the blue color. With nitric acid starch yields an explosive compound (*Xyloidin*), $C_{12}H_{16}(NO_2)_4O_{10}$ or $C_{12}H_{16}O_6(NO_2)_4$.

Composition of Starch-granules.—Starch-granules consist mainly of *granulose*, soluble in cold water and giving an indigo color with

iodine, and *starch cellulose*, insoluble in water, and giving with iodine a dirty yellow color, with, possibly, other carbohydrates. The starch cellulose forms an external coating upon the granule, and also exists, mixed with the granulose, inside the granule. If this coating be broken by mechanical means, the continued application of cold water will remove all the granulose, leaving the cellulose insoluble. By the action of diastase, ptyalin, and other ferments, and by other means, the granulose may be converted into sugar and dextrin, leaving the starch cellulose unacted upon.

Microscopical Examination of Starches.


All kinds of starch afford the blue color with iodine, showing their chemical similarity. Physically, however, the granules of different starches differ from each other; hence a careful microscopical examination of any starch, or of any powder or vegetable tissue containing starch, enables the observer to state, with a high degree of probability, the source of the starch, either at once if he has much experience, or after comparing the granules in question with authentic specimens. A glance at the accompanying eight engravings* (Figs. 42 to 49) of common starches will show to what extent different starch-granules naturally differ in size, shape, general appearance, distinctness and character of the rugæ, and position of the more or less central point or hilum. While from different starches individual granules may be picked out which much resemble each other, the appearance of each starch as a whole is fairly characteristic; that is to say, each *group* of granules differs in one or more characters from similar *groups* of granules of other starches.

A quarter-inch object-glass will commonly suffice for the microscopical observation of starch. A very little of the starch is mixed on a glass slide with a drop of water, a piece of thin covering-glass placed on the drop and gently pressed, so as to provide a very thin layer for observation. Instead of water, diluted spirit of wine, diluted glycerin, turpentine or other essential oil, Canada balsam, and other fluids may be used in cases where the markings or other appearances are not well defined. The illumination also of the granules may be varied, the light being reflected or transmitted, concentrated or diffused, white or colored, polarized or plain. Polarized light is especially valuable in developing differences and in intensifying the effects of obscure markings. By polarized light the granules of potato starch appear as if traversed by a black cross; wheat starch-granules and many others also peculiarly and characteristically influence polarized light. Distinctive characters will sometimes present themselves only when the granules are made to roll over in the fluid in which they have been temporarily mounted or when the slide is gently warmed. Starches which have already been subjected to the influence of heat—partly, as in sago or tapioca, or almost entirely, as in bread—will of course differ in appearance from

*By permission of Messrs. Longmans & Co. these engravings have been copied, with very few modifications, from the plates in two of the three volumes of the original edition of Pereira's *Materia Medica*.

STARCHES
(Magnified 250 diameters).

Figs. 42 to 49.



granules of the same starch before being dried, cooked, or torrefied. The characters of a starch will also somewhat vary according to the age and condition of the plant yielding it.

The description of the microscopical characters of the official varieties of starch is as follows: 1. Wheat starch: A mixture of large and small granules, which are lenticular in form, and marked with faint concentric striæ surrounding a nearly central hilum. 2. Maize starch: Granules more uniform in size, frequently polygonal, somewhat smaller than the large granules of wheat starch, and having a very distinct hilum, but without evident concentric striæ. 3. Rice starch: Granules extremely minute, nearly uniform in size, polygonal, hilum small and without striæ.

(For plates and descriptions of the characters of other starches occurring in plants used for medicinal purposes the reader is referred to works on *Materia Medica*, and to the indexes of Journals of Pharmacy, as well as to general works and magazines on microscopy. For engravings of starch-granules *in situ*, vide Berg's "Anatomischer Atlas," published by Gaertner, Berlin.)

The student may place fair confidence in the accompanying lithographs and in most of the published engravings of starch-granules; but in microscopical analyses of importance the worker should, if possible, himself obtain actual specimens of starches for comparison from the respective seeds, fruits, and other tissues.

Inulin, $(C_6H_{10}O_5)_n \cdot H_2O$ (Kiliani), is associated with similar bodies, *pseudo-inulin*, $(C_6H_{10}O_5)_{16} \cdot H_2O$, and *inulenin*, $(C_6H_{10}O_5)_{10} \cdot 2H_2O$ (Tanret). It is a white powder, apparently occupying the place of starch in the roots of many plants, especially those of the natural order *Compositæ*. 20 to 45 per cent. has been obtained from elecampane (*Inula helenium*). It is also contained in the dahlia, colchicum, arnica, dandelion, chicory, artichoke, etc. It is soluble in boiling water, nearly all being redeposited on cooling. Iodine turns it yellow. Long ebullition converts it into a kind of gum. Like starch, inulin is convertible into sugar, but by its own special ferment, the existence of which, in the Jerusalem artichoke, has been demonstrated by Professor J. R. Green. This ferment differs from diastase in being without the power of converting starch into sugar.

Lichenin ($nC_6H_{10}O_5$) is a white starch-like powder largely contained in many lichens—Iceland "moss," *Cetraria Islandica*, and many others. It is soluble in boiling water, and the fluid gelatinizes on cooling. It may be precipitated from its aqueous solution by alcohol. With iodine it gives a reddish-blue color.


Glycogen, or *animal starch*, is the name given to the solid matter stored in the liver and resulting from the dehydration of the digested hydrated food which has been carried to the liver by the portal vein.

Dextrin ($nC_6H_{10}O_5$).—Mix a grain or two of starch with half a test-tubeful of cold water and a drop or two of sulphuric acid, and boil the mixture for a few minutes; no mucilage is formed, and the liquid, if sufficiently boiled, yields no blue color with iodine; the starch has become converted into *dextrin* and some sugar. Dextrin is also produced if starch is

maintained at a temperature of about 320° F. for a short time. Dextrin is now largely manufactured in the latter way, and a paste of it is used by calico-printers as a vehicle for colors; it is termed *British gum*. The change may also be effected by *diastase*, a peculiar ferment existing in malt. Mix two equal quantities of starch with equal amounts of water, adding to one a little ground malt, then heat both slowly to the boiling-point: the mixture without malt thickens to a paste or pudding; that with malt remains thin, its starch having become converted into dextrin and a sugar termed maltose.

Diastase is probably a mixture, but possibly an oxidation-product, of the coagulable albumenoids. It is so named from *διάσπαισις* (*diastasis*), *separation*, in allusion to the separation, or rather alteration, it effects among the constituent atoms of the molecule of starch. This function is shared by the saliva, pancreatic juice, bile, and the intestinal and other juices. The function is completely destroyed when the albumenoids are coagulated by a temperature of from 176° to 178° F.

The Action of Diastase upon Starch.—Diastase has scarcely any action upon unbroken starch-granules. The granules must be ruptured by gelatinization with heat and moisture or in some other way. When a solution containing diastase, such as a cold-water infusion of malt, is allowed to act upon gelatinous starch or starch-paste at 140° to 160° F., liquefaction occurs. It is possible to operate so that when liquefaction has taken place the solution shall give no reaction for sugar or dextrin. If this solution be concentrated and allowed to cool, a glistening white precipitate of soluble starch falls. Soluble starch is probably the result of the partial decomposition of the more complex molecule of granulose or gelatinous starch. The next step in the action of diastase upon gelatinous starch is the breaking down of the soluble starch-molecule into dextrin and a sugar called maltose. At least ten dextrans are successively produced, each simpler than the one preceding it, the proportion of maltose being correspondingly increased. The dextrans first produced give a red or brown color with iodine, while those last produced, and having a simpler molecule, give no color with iodine. The final reaction may be expressed thus:




The dextrans are distinguished by their rotatory power, their reducing action on cupric salts, and in other ways.

Starch heated with glycerin is converted into the soluble variety. The latter may be precipitated from an aqueous solution by strong alcohol. A strong solution in water gradually gelatinizes, owing to reconversion into insoluble starch (Zulkowsky).

The Action of Dilute Acids upon Starch.—Dilute acids act upon gelatinous starch in the same way as diastase, except that the final product is glucose.

Malt (the word malt is said to be derived from the Welsh *mall*, soft or "rotten") is simply barley which has been softened by steeping in water, allowed to germinate slightly, and further change then arrested by the application of heat in a kiln. During germination the gluten breaks up and yields a glutinous substance termed vegetable gelatin, diastase, and other matters. To the vegetable gelatin is due much of the "body" of well-malted and slightly hopped beer; it is precipitated by tannic acid; hence the thinness of ale (pale or bitter) brewed with a large proportion of hop or other materials containing tannic acid. A portion of the diastase, reacting on the starch of the barley, converts it into dextrin, and, indeed, carries conversion to the further stage of maltose, as will be explained immediately. The temperature to which the malt is heated is made to vary, so that the sugar of the malt may or may not be partially altered to a dark-brown coloring material: if the temperature is high, the malt is said to be *high-dried* and is used in porter-brewing; if low, the product is of lighter color and is used for ale. The diastase remaining in malt is still capable of converting a large quantity of starch into dextrin and sugar (maltose); hence the makers or distillers of the various spirits operate on a mixture of malted and unmalted grain in preparing liquors for fermentation.

Extract of Malt is an evaporated infusion of malt. Taken with food, its diastase aids in the conversion of starch into a variety of sugar termed maltose, and dextrin, and, *pro tanto*, assists enfeebled digestive powers.



As diastase begins to lose its power at temperatures above 150° F., that degree should not be exceeded in evaporating the infusion; indeed, if the dissolved albumenoid matters are to be retained, the evaporation should be conducted at 120° F.

The following method serves for the estimation of the diastatic power of malt extract: 1.5 gram of the extract is dissolved in 15 cc. of water and mixed with a mucilage of .1 gram of starch in 100 cc. of water. The mixture is raised to 140° F. in temperature, and tested from time to time by adding two drops of iodine solution to 5 cc. of it, and comparing with 5 cc. of a similar mixture to which no starch has been added. No difference of tint between the two solutions indicates completion of the reaction. Good malt extract will accomplish this within half an hour; some samples will take less time, but many commercial extracts will require three hours or more.

Gum is a frequent constituent of vegetable juices, existing in large quantity in several species of *Acacia*. The nature of gums is very little known, though most probably they all belong to the carbohydrates. According to Fremy, gum is a calcium salt, sometimes partially a potassium salt, of the *gummic* or *arabic* radical, though consisting mostly of arabin or arabic acid alone. The formula of

gummie acid is said to be $\text{H}_2\text{C}_{12}\text{H}_{18}\text{O}_{10}, \text{H}_2\text{O}$, but, from the important researches of O'Sullivan, it would seem to be far more complex, a multiple of the empirical formula, $\text{C}_6\text{H}_{10}\text{O}_5$ —Raoult ($\text{C}_5\text{H}_{10}\text{O}_5$)₇. Gum differs from dextrin in yielding mucic acid when oxidized by nitric acid. Good adhesive mucilages may be made from such gum-arabic substitutes as "ghatti," "amrad," etc. *Cerasin*, or cherry-tree gum, is a calcium metagummate, an insoluble modification of acacia gum. *Bassorin*, *traganthin*, or *adraganthin*, ($\text{C}_{12}\text{H}_{20}\text{O}_{10}$) is a form of gum which is insoluble in water, but absorbs large quantities of that liquid and forms a gelatinoid mass: it occurs largely in tragacanth, combined, like arabin, with calcium. *Pectin*, or vegetable jelly ($\text{C}_{32}\text{H}_{40}\text{O}_{28}, 4\text{H}_2\text{O}$), is the body which gives to expressed vegetable juices the property of gelatinizing: it forms the chief portion of Irish or carrageen "moss" (*Chondrus crispus*). Ceylon "moss" (*Gracillaria lichenoides* and *G. confervoides*, P. I.) contains from one-third to three-fourths of vegetable jelly.

The mucilage of marshmallow-root (*Althea officinalis*) and of linseed or common flaxseed (*Linum usitatissimum*) is a gum-like substance containing much mineral matter. It is the basis of the infusions termed *mallow tea* and *linseed tea*. Somewhat similar mucilage occurs in infusion of bael: it is also largely yielded by the seeds of the quince (*Pyrus Cydonia*), as well as by the bark of the red or slippery elm (*Ulmus*, U. S. P.). *Salep*, the powdered dried tubers of many species of *Orchis*, contains a large quantity of such matter. *Squill* also. The Indian Okra (*Hibisci Capsula*, P. I., from *Hibiscus esculentus*) and *Ispaghul* or *Spogel* seed (*Ispaghula Semina*, P. I., from *Plantago ispaghula*) also appear to contain a considerable quantity. In *Sassafras-pith* (*Sassafras Medullæ*, U. S. P.) starch and mucilage occur.

Cellulin, or *Cellulose*, $\text{C}_6\text{H}_{10}\text{O}_5$, the woody fibre of plants, familiar, in the nearly pure state, under the forms of "cotton-wool" (*Gossypium Purificatum*, U. S. P., hairs of the seed of various species of *Gossypium*), paper, linen, and pith, is another substance isomeric, probably polymeric, with starch. Lignin is a closely-allied body lining the interior of woody cells and vessels. By the action of nitric acid of various strengths on cellulul, mono-, di-, or tri-nitro-cellulins are readily formed: $\text{C}_6\text{H}_9\text{O}_4\text{NO}_3$, $\text{C}_6\text{H}_8\text{O}_3(\text{NO}_3)_2$, $\text{C}_6\text{H}_7\text{O}_2(\text{NO}_3)_3$. Trinitrocellulin is highly-explosive gun-cotton; dinitrocellulin is not sufficiently explosive for use instead of gunpowder; mononitrocellulin is scarcely at all explosive. The heat of a water-bath may explode trinitrocellulin, but not dinitrocellulin if pure. The three displaceable atoms of hydroxyl in cellulul may be displaced by bodies other than the nitric radical.

Dinitrocellulin (*Pyroxylinum*, U. S. P.).—Mix 22 parts of sulphuric acid and 14 of nitric in an earthenware mortar. When cooled to about 32° C. (90° F.), immerse 1 part of cotton-wool in the mixture, and stir it with a glass rod, so that it is thoroughly and uniformly wetted by the acids. Macerate until a sample, washed with water and then with alcohol, is soluble in a mixture of one volume of alcohol and three of stronger ether. Transfer the cotton to a vessel

containing a considerable volume of water, stir it rapidly and well with a glass rod, decant the liquid, pour more water upon the mass, agitate again, and repeat the affusion, agitation, and decantation until the washing ceases to give a precipitate with barium chloride or to taste acid. Drain the product on filtering-paper and dry on a water-bath.

Pyroxylin may also be made by soaking 7 parts of white filtering-paper, which has been washed in hydrochloric acid and dried, in a mixture of 140 parts of sulphuric acid (sp. gr. 1.82) and 70 of nitric acid (1.37) for three hours, and well washing the product (Guichard).

Mononitrocellulin and *trinitrocellulin* are insoluble in a mixture of alcohol and ether; *dinitrocellulin*, or *pyroxylin*, is soluble, the solution forming ordinary collodion (*Collodium*, U. S. P.). The official proportions are 3 grm. of pyroxylin dissolved in a mixture of 75 cc. of ether and 25 of alcohol. After digesting for a few days, decant the liquid from any sediment and preserve it in a well-corked bottle. It dries rapidly upon exposure to the air, and leaves a thin, transparent film, insoluble in water or spirit. *Flexible Collodion* (*Collodium Flexile*, U. S. P.) is a mixture of collodion (92 parts), Canada balsam (5 parts), and castor oil (3 parts). *Blistering Collodion* (*Collodium Vesicans*, B. P.), *Collodium Cantharidatum*, U. S. P., or *Cantharidal Collodion*, is a solution of pyroxylin containing the active blistering principle of cantharides. A *Styptic Collodion* (*Collodium Stypticum*, U. S. P.), containing tannic acid, is also official. Many articles of utility and beauty are now made of pyroxylin variously colored and sold under the name of *xylonite* (ξύλον, *xulon*, wood) or *celluloid* (cellulin-like).

Tunicin, or animal cellulose, exists. It is extracted from ascidians and cynthians.

Isomerism.—Allotrophy.—Polymorphism.

The composition of dextrin is represented by the same formula as that of starch—namely, $C_6H_{10}O_5$; for it has the same percentage composition as starch. Inulin (p. 476) and cellulose (p. 479) have also a similar formula. There are many other bodies similar in centesimal composition, but dissimilar in properties; such substances are termed *isomeric* (from *ἴσος*, *isos*, equal, and *μέρος*, *meros*, part); and their condition is spoken of as one of *isomerism*. There is sometimes good reason for doubling or otherwise multiplying the formula of one of two *isomers*, *isomerides*, or isomeric bodies. Thus a molecule of ethylene (olefiant gas), the chief illuminating constituent of coal-gas, is represented by the formula C_2H_4 , while a molecule of amylene, an anæsthetic liquid hydrocarbon, obtained from amylic alcohol, though having the same percentage composition as olefiant gas, is represented by the formula C_5H_{10} ; for amylene, when gaseous, is about twice and a half as heavy as ethylene, and must contain, therefore, in each molecule, twice and a half as many atoms, for (Avogadro) these equal volumes must contain equal numbers of molecules; its formula is, consequently, constructed to represent these proportions. (Read again pages 36 to 60.) This variety of isomerism is termed *polymerism* (from *πολύς*, *polūs*, many

or much, and μέρος, part). Formic aldehyde, CH_2O , acetic acid, $\text{C}_2\text{H}_4\text{O}_2$, and lactic acid, $\text{C}_3\text{H}_6\text{O}_3$, are, obviously, *polymers*. Metastannic acid (*vide* p. 243) is a polymeric variety, or *polymeride*, of stannic acid. An illustration of a second variety of isomerism is seen in the case of ammonium cyanate and urea, bodies already alluded to in connection with cyanic acid. These and several other pairs of chemical substances have dissimilar properties, yet are similar in elementary composition and in the centesimal proportion of the elements, and we cannot escape the conclusion that each molecule possesses the same number of atoms. But the reactions of these bodies indicate the probable nature of their construction; and this is shown in their formulæ by the disposition of the symbols. Thus ammonium cyanate is represented by the formula NH_4CNO , urea by $\text{CO}(\text{NH}_2)_2$. Such bodies are termed *metameric* (from μετά, *meta*, a preposition denoting change, and μέρος), and their condition spoken of as one of *metamerism*. (For two metameric nitro-methylic and also two nitro-ethylic bodies see p. 405.) Ethyl acetate (p. 406) is metameric with butyric acid (p. 488), for they have the same percentage composition and their vapors have the same specific gravity, and each therefore might be represented by the formula $\text{C}_4\text{H}_8\text{O}_2$; but their properties warrant us in assuming that their atoms occupy different positions in the two molecules—justify us in giving $\text{CH}_3\text{CO}\cdot\text{OC}_2\text{H}_5$ as a picture of a molecule of ethyl acetate, and $\text{C}_3\text{H}_7\text{COOH}$ as a picture of a molecule of butyric acid. Methyl acetate ($\text{CH}_3\text{COO}\cdot\text{CH}_3$), propionic acid ($\text{C}_2\text{H}_5\text{COOH}$),* and ethyl formate ($\text{H}\cdot\text{CO}\cdot\text{OC}_2\text{H}_5$) are isomers of the metameric variety, or *metamers* or *metamerides*; also quinine and quinidine, cinchonine and cinchonidine, and many of the volatile oils, etc. The isomerism of starch and dextrin may be of a polymeric or of a metameric character; but we do not yet know which, and must therefore at present give them identical formulæ, though it is most probable that many of the carbohydrates are multiples of the mere empirical formulæ, since dextrin ($x\text{C}_6\text{H}_{10}\text{O}_5$) by hydration produces maltose, $\text{C}_{12}\text{H}_{22}\text{O}_{11}$, which would point to the formula of dextrin as being at least $(\text{C}_6\text{H}_{10}\text{O}_5)_2$, while the extent to which it lowers the freezing-point of a solvent points to $(\text{C}_6\text{H}_{10}\text{O}_5)_7$. Patient accumulation of facts and the aid of the theory of valency will, doubtless, sooner or later, fully explain all cases of isomerism.

Substances similar in composition and constitution, yet differing in properties, are termed *allotropic* (ἄλλος, *allos*, another, τροπός, *tropos*, condition). Thus ordinary phosphorus, kept at a temperature of about 450° F. in an atmosphere from which air is excluded, becomes red, opaque, insoluble in liquids in which ordinary phosphorus is soluble, oxidizes extremely slowly, and only ignites when heated to near 500° F. (red or amorphous phosphorus). A black allotropic variety of phosphorus is known. There are also three *allotropes* of carbon which are respectively crystalline, graphitic, and amorphous. Sulphur may be obtained in the viscous as well as in the hard, brit-

* For explanation of formulæ see section on Aldehydes and Acids, p. 482.


tle condition. Another illustration of *allotropy* is seen in the varieties of tartaric acid which have different optical properties, but otherwise are identical; they are in neither of the above-mentioned states of isomerism, but are allotropic modifications of the same substance. The constitution of such bodies is perhaps best conceived by the aid of stereochemical hypotheses. Occasionally one and the same substance crystallizes in two distinct forms; its state is then described as one of *polymorphism* (*πολvs, polus*, many; *μορφή, morphē*, form). Sulphur is polymorphous. It crystallizes by slow cooling in (1) prismatic crystals of sp. gr. 1.98, while in nature it occurs in (2) octahedra of sp. gr. 2.07. Melted and poured into water, sulphur takes up (3) the form of caoutchouc of sp. gr. 1.96. These differences warrant the statement that sulphur occurs in three distinct allotropic conditions. Possibly such conditions result from the association of different numbers of atoms in the molecule of the element; that is, allotropic bodies may simply be *physically polymeric*, or in some other way be mere *physical isomerides*.

QUESTIONS AND EXERCISES.

How is wheat starch or potato starch isolated?—Define gluten and gluten.—Enumerate the proximate principles of wheaten flour.—Is starch soluble in water?—Which is the best chemical test for starch?—Distinguish physically between the varieties of starch.—Into what compound is starch converted by heat?—What occurs when a mixture of starch and water is allowed to flow into hot diluted sulphuric acid?—If two equal amounts of starch with water be heated, one containing a small quantity of ground malt, what effects ensue?—Write a short article on the chemistry of malting.—What is the nature of gum arabic? and how is it distinguished from "British gum"?—Mention the properties of the products of the action of nitric acid of various strengths on cellulose.—How is pyroxylin prepared?—Explain *isomerism*, giving several illustrations.—Give examples of polymeric bodies.—State the formula of a body metameric with urea.—Define allotropy and polymorphism, giving illustrations.


ALDEHYDES AND ACIDS.

General Formation.—The aldehydes and acids may be artificially formed by oxidation of the primary alcohols, glycols, etc. Monhydric alcohols, having only one hydroxyl (OH) group, form monobasic acids; dihydric alcohols (glycols) having two hydroxyl groups, yield monobasic and dibasic acids; and so on. Thus:




It will be seen that the groups COH and COOH denote respectively an aldehyde and an acid, the H in the COOH group being replaceable by a metal, such as $\text{CH}_3\cdot\text{CO}\cdot\text{ONa}$ (sodium acetate). (See also pp. 409 and 456.)

Acids may also be obtained by acting on the *nitriles* or cyanides of the hydrocarbon radicals with hydrochloric acid and water. Thus:




Many aldehydes and acids occur in nature; for example, oil of meadow-sweet (salicylic aldehyde), oil of bitter almonds (benzoic aldehyde), citric acid in lemons.

General Reactions.—Aldehydes all form crystalline compounds with acid potassium sulphite, by oxidation they yield an acid, and by the action of nascent hydrogen they yield an alcohol, while acids, by nascent hydrogen, yield an aldehyde, and then an alcohol. With oxides, hydrates, carbonates, and sometimes with metals, acids form metallic derivatives. With the alcohols, acids yield alkyl† or ethereal salts, as; for instance, acetic ether. By the action of phosphorus, chloride, iodide, or bromide their hydroxyl group is replaced by chlorine, iodine, or bromine:



Like inorganic acids, they form anhydrides by the elimination of water:



The important aldehydes and acids will now be mentioned.

*The reactions of *nitriles* indicate that the radical present is united to the carbon of the cyanogen, while the reactions of *isonitriles* (or *carbamines*) indicate that the radical is united to the nitrogen. Hence such formulæ as CH_3CN and CH_3NC .

†*Alkyl Salts.* *Alkyl*, from the Arabic article *al*, the, as in alkali, alcohol, etc., and the termination common to the names of such radicals as ethyl, amyl, and phenyl, and as seen in methyl, the prototype of such names. (See p. 437.) In Germany the word *ester*, a mere variation of the word *ether*, is similarly employed. In the scientific chemistry of both countries it is thus sought to restrict the name *ethers* to the oxides of radicals, as common ether, $(\text{C}_2\text{H}_5)_2\text{O}$ (*Æther*, U. S. P.).

The Acetic Series.

Acids of the Acetic Series, $C_nH_{2n+1}CO\cdot OH$ (monobasic), formed by the two general methods given—namely, from primary alcohols of the ethylic series and from cyanides of the paraffin hydrocarbon radicals.

Formic Acid, $H\cdot COOH$. Formic aldehyde, $H\cdot COH$. (See p. 339.)


Acetic Acid, CH_3COOH (Methylformic acid). Obtained by the oxidation of alcohol and in other ways. (See p. 297.)

Aldehyde, or *Acetic Aldehyde*, C_2H_4O or CH_3COH .


Preparation.—Place together in a capacious test-tube or flask about 4 parts of spirit of wine, 6 of black manganese oxide, 6 of sulphuric acid, and 4 of water, and gently warm the mixture; aldehyde (*alcohol dehydrogenatum*), a highly volatile liquid, is immediately formed, and its vapor evolved, recognized by its peculiar somewhat fragrant odor. Adapt a cork and rather long bent tube to the test-tube, and let some of the aldehyde slowly distil over into another test-tube, the condensing-tube being kept as cool as possible. Set the distillate aside for a day or two; the aldehyde will have nearly all disappeared and acetic acid be found in the tube. Test the exposed liquid by litmus-paper; it will be found to have an acid reaction: make it slightly alkaline by a drop or two of solution of sodium carbonate, then boil to remove any alcohol and aldehyde present, add sulphuric acid, and notice the characteristic odor of the acetic acid evolved.

These experiments will enable the process of acetification described in connection with acetic acid to be more fully understood. Pure diluted alcohol is not oxidized by exposure to air alone; but in presence of a ferment, *bacterium aceti*, it is oxidized first to aldehyde and then to acetic acid.


In the above process the black manganese oxide and sulphuric acid furnish nascent oxygen:



The nascent oxygen then acts on the alcohol, just as the oxygen of the air acts on the alcohol in fermented infusion of malt, beer, or wine, giving aldehyde:



The aldehyde rapidly, even when pure (more rapidly when impure), absorbs oxygen and yields acetic acid:



Tests.—Aldehyde heated with solution of potash gives a brownish-yellow resinous mass of peculiar odor. Its aqueous solution reduces salts of silver, giving a mirror-like coating to the cleaned sides of a test-tube. When acted on by phenol dissolved in sulphuric acid, it gives a red color. Aldehyde on keeping or in contact with sulphuric acid, zinc chloride, etc. yields two polymerides—metaldehyde ($\alpha\text{C}_2\text{H}_4\text{O}$) and paraldehyde, $\text{C}_6\text{H}_{12}\text{O}_3$, the latter having a characteristic odor. The official paraldehyde (*Paraldehydum*, U. S. P.) boils at 123° – 125° C., dissolves in water, spirit, or ether, is neutral, and if it contains no ordinary aldehyde will not be colored on standing for two hours with solution of potash or soda. Add a little silver ammonium nitrate to a strong solution of paraldehyde, and on warming and allowing to stand a silver mirror will be formed. A mixture of 8 cc. of paraldehyde and an equal amount of alcohol, with 1 drop of phenolphthalein, should acquire a pink color, with, at the most, 0.5 cc. of normal potassium hydrate (limit of free acid).


CHLORAL.

Chloral, or *Trichloraldehyde*, CCl_3COH , is a chlorine substitution-derivative of aldehyde, though it cannot directly be obtained by acting on aldehyde by chlorine, because condensation-products are formed.


Process.—Pass a rapid stream of dry chlorine into pure absolute alcohol so long as absorption occurs. During the first hour or two the alcohol must be kept cool, afterward gradually warmed till ultimately the boiling-point is reached. The preparation of a considerable quantity occupies several days. The crude product is mixed with three times its volume of sulphuric acid and distilled, again mixed with a similar quantity of sulphuric acid and again distilled, and finally rectified from quicklime.

The formation of chloral would at first sight seem to be due to the production from the alcohol ($\text{CH}_3\text{CH}_2\text{OH}$) of aldehyde (CH_3COH), through the removal of hydrogen by the chlorine, and the substitution of chlorine for hydrogen in the aldehyde (CH_3COH), with formation of *chlor-aldehyde* or *chloral* (CCl_3COH). But the reactions are far more complicated, being as follows:

Aldehyde and hydrochloric acid are first formed:




The nascent aldehyde unites with alcohol, forming acetal:




Acetal* by further chlorination yields trichloroacetal:



Trichloroacetal, when acted on by hydrochloric acid, yields ethyl chloride and chloral alcoholate:




From the alcoholate chloral is liberated by treatment with sulphuric acid.



Properties.—It is a colorless liquid, of oily consistence. Sp. gr. 1.502. Boiling-point, 201.2° F. Its vapor has a penetrating smell and is somewhat irritating to the eyes. Mixed with water, heat is disengaged, and solid white, crystallizable *chloral hydrate*, $\text{CCl}_3\text{CH}(\text{OH})_2$, or *Hydrate of Chloral* (*Chloral Hydras*, B. P.), results. "Chloral hydrate," termed in the United States Pharmacopœia "Chloral," with "Chloral Hydrate" as a synonym, is a true glycol, the water not being simply water of crystallization, but of combination, the systematic name being trichlorethylidene glycol:




Chloral hydrate fuses when heated, solidifies at about 120° F.; boils at from 202° to 206° F. It sublimes as a white crystalline powder. Both chloral and chloral hydrate are soluble in water, alcohol, ether, and oils. Oils and fats are also soluble in chloral hydrate. The aqueous solution should be neutral and give no reaction with silver nitrate. Chloral, especially if it contains a trace of acid, may undergo a spontaneous change into an opaque white isomeric modification, *metachloral*, insoluble in water, alcohol, or ether, but convertible by prolonged contact with water or by distillation into the ordinary condition. By action of weak alkalis chloral yields formate of the alkali-metal and chloroform:



Chloral, or rather strong aqueous solution of chloral hydrate (3 in 4), injected beneath the skin yields chloroform and produces narcotic effects (Liebreich, Personne). Chloroform itself admits of similar hypodermic use (Richardson). If administered by the stomach, 30 to 80 grains of solid hydrate are required. The final products of the reaction of the chloroform and blood are sodium formate and chloride. A strong spirituous solution of potash effects the same transformation: $\text{CHCl}_3 + 4\text{KOH} = \text{H}\cdot\text{COOK} + 3\text{KCl} + 2\text{H}_2\text{O}$.

* *Methylal*, $\text{CH}_2(\text{OCH}_3)_2$, the lowest term of the series, is occasionally used as a soporific.

Solution of ammonia and moist calcium hydrate, as well as weak solutions of fixed alkalies, convert chloral hydrate into formate of the metal and chloroform. The reaction with the slaked lime being especially definite and complete (Wood), it may be employed in ascertaining the richness of a sample of commercial chloral hydrate in chemically pure chloral hydrate.



From the foregoing equation and molecular weights it is obvious that 100 grains of chloral hydrate, if quite dry, will yield by distillation, with 30 grains of slaked lime and an ounce of distilled water (in a small flask and a long bent tube kept cool by moistened paper), 72.2 grains of chloroform by weight or (the sp. gr. of chloroform being taken at 1.497) 47.56 grains by measure, or about 52 minims. 100 grains of the official chloral hydrate "should yield not less than 70 grains of chloroform." (Any such definite quantity of chloroform, on account of its volatile nature, is perhaps best measured, the weight being obtained by multiplying the volume by 1.5.)

Small quantities of chloral hydrate in dilute solutions may be estimated by converting its chlorine into hydrochloric acid by nascent hydrogen, and titrating with volumetric solution of silver nitrate (Short). A quantity of solution containing not more than .05 of a gram is placed in a small flask with granulated zinc and acetic acid, and allowed to stand twenty-four hours: the solution is then poured off and the zinc washed two or three times with distilled water; a little yellow potassium chromate is added, and it is then titrated with decinormal silver nitrate solution in the usual way, the acetic acid and zinc acetate not interfering with the indications. 1000 cc. of the silver solution indicate 5.52, nearly, of chloral hydrate.

Pure Chloral Hydrate.—Liebreich, who first proposed the use of chloral hydrate, gives the following as the characteristics of a pure article: Colorless, transparent crystals. Does not decompose by the action of the atmosphere, does not leave oily spots when pressed between blotting-paper, affects neither cork nor paper. Smells agreeably aromatic, but a little pungent when heated. Tastes bitter, astringent, slightly caustic. Seems to melt on rubbing between the fingers. Dissolves in water like candy, without first forming oily drops, and the solution is neutral or faintly acid to test-paper. Dissolves in carbon bisulphide, petroleum, ether, water, alcohol, oil of turpentine, etc. Its solution in chloroform gives no color when shaken with sulphuric acid. Boiling-point, 203° to 205° F. It volatilizes without residue. Distilled with sulphuric acid, the chloral should pass over at 205° to 207° F. Melting-point, 133° to 136° F., again solidifying at about 120°. Gives no chlorine reaction on treat-

ing the solution in water (acidulated by nitric acid) with silver nitrate.

Impure Chloral Hydrate.—Yellowish, cloudy. Decomposes; leaves spots by pressing between blotting-paper; decomposes corks and paper of the packing. Smells pungent and irritating; on opening the bottle is sticky and often emits fumes. Taste strongly caustic. With water forms oily drops or is partially insoluble. Boils at a higher temperature. On treating it with sulphuric acid turns brown, with formation of hydrochloric acid. Gives chlorine reaction on treating the solution in water (acidulated by nitric acid) with silver nitrate.

Alcoholates of chloral are obtained on combining alcohols with chloral. Chloral alcoholate or trichlorethylidene ethyl ether, $\text{CCl}_3\text{CH} < \begin{smallmatrix} \text{OH} \\ \text{OC}_2\text{H}_5 \end{smallmatrix}$, is obtained by mixing alcohol with chloral; it is in fact "chloral hydrate" with one hydroxyl group replaced by (OC_2H_5) .

Bromal, CBr_3COH , *Bromal Hydrate*, $\text{CBr}_3\text{CH}(\text{OH})_2$, and *Bromal Alcoholates* are produced when bromine instead of chlorine attacks alcohol. *Iodal*, CI_3COH , also exists.

BUTYL CHLORAL, $\text{C}_3\text{H}_4\text{Cl}_3\text{COH}$, originally, but erroneously, termed *croton chloral*, is a product of the action of dry chlorine on cold aldehyde. Its name expresses its constitution; it is *chlorinated butyric aldehyde*, ordinary chloral being chlorinated acet-aldehyde. *Butyl-chloral hydrate*, *croton-chloral hydrate*, wrongly so called, *hydrate of butyl-chloral* (*Butyl-chloral Hydras*, B. P.), *hydrous butyl-chloral*, $\text{C}_3\text{H}_4\text{Cl}_3\text{CH}(\text{OH})_2$ (*trichlorobutylidene glycol*), occurs "in pearly white crystalline scales, having a pungent but not acid odor, resembling that of hydrous chloral, and an acid nauseous taste. It fuses at about 172°F . (77.8°C .) to a transparent liquid, which in cooling commences to solidify at about 160°F . (71.1°C .). Soluble in about 50 parts of water, in its own weight of glycerin and of rectified spirit, and nearly insoluble in chloroform. The aqueous solution is neutral or but slightly acid to litmus-paper. It does not yield chloroform when heated with solutions of potash or soda or with milk of lime."

The Acetic Series of Acids—continued.

Propylic or *Propionic Acid* (ethyl-formic acid), $\text{C}_2\text{H}_5\text{COOH}$, is produced by oxidation of propylic alcohol.

Butyric or *Tetrylic Acid* (propyl-formic acid), $\text{C}_3\text{H}_7\text{COOH}$, is formed by general methods; also during the fermentation of cheese. It is found as a glyceric salt in butter (whence its name).

Pentylic, *Valerianic*, or *Valeric Acid*, $\text{C}_4\text{H}_9\text{COOH}$.—There are several varieties of this acid, the valerianic acid from valerian and angelica-root, and that artificially formed from amylic alcohol (*vide* p. 363), being the iso-primary valerianic acid, or iso-propylacetic acid, $\text{CH}(\text{CH}_3)_2\text{CH}_2\text{COOH}$, the normal having the constitution of $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{COOH}$.

Palmitic Acid, $\text{C}_{15}\text{H}_{31}\text{COOH}$, from soft fats; *Stearic Acid*,

$C_{17}H_{35}COOH$, from suet, tallow, and the hard fats; *Cerotic Acid*, $C_{26}H_{53}COOH$, from beeswax; and *Melissic Acid*, $C_{29}H_{59}COOH$, from beeswax and from canauba wax (from the leaves of *Copernicia cerifera*, a Brazilian palm), belong to the acetic series.

Stearic Acid, $HC_{18}H_{35}O_2$, is official (*Acidum Stearicum*, U. S. P.). It is a hard, bright, white solid without odor or taste, soluble in alcohol or ether. Melts at $69.2^\circ C$. "If 1 grm. of the acid and 1 of sodium carbonate be boiled with 30 cc. of water in a capacious flask, the resulting solution, while hot, should not be more than opalescent (limit of undecomposed fat)."

The Lactic Series.

Acids of the Lactic Series, $C_nH_{2n}(OH)COOH$.—This series is formed of hydroxy-derivatives of the acetic series, one atom of hydrogen being replaced by the hydroxyl group.




Acetic acid.



Hydroxyacetic or glycollic acid.

Though they possess only one carboxyl ($COOH$) group, yet, having an alcoholic hydroxyl group, they may sometimes form di-substitution derivatives with the metals.

They are best formed by hydrolysis of the nitriles produced on combining hydrocyanic acid with aldehydes or ketones; also by partial oxidation of glycols by diluted nitric acid, and by acting on monochloro-derivatives of the acids of the acetic series by moist silver oxide:



Monochloroacetic acid.

Glycollic acid.

Carbonic Acid or *Hydroxyformic Acid*, $OH \cdot CO \cdot OH$, the first of this series, has been studied already. *Carbamide* or *Urea*, $NH_2 \cdot CO \cdot NH_2$, the normal amide of carbonic acid, is interesting historically as being the first organic body synthetically produced from inorganic sources. (See Index, "Urea, Artificial Production of.") The acid amide of carbonic acid, *carbamic acid*, $NH_2 \cdot CO \cdot OH$, occurs as an ammonium salt, $NH_2 \cdot CO \cdot ONH_4$, in the ammonium carbonate of pharmacy. Ethyl carbamate, or *Urethane*, $NH_2 \cdot CO \cdot OC_2H_5$, is a mild hypnotic.

Glycollic Acid (Hydroxyacetic Acid), $CH_2OH \cdot COOH$, is found in the leaves of the Virginia creeper; artificially it may be obtained by carefully oxidizing glycol and by the action of silver oxide on dextrose and lævulose.

Lactic Acid (Hydroxypropionic Acid), $C_2H_4(OH)COOH$. At least three isomeric lactic acids are known, the fermentative lactic acid (ethylidene* lactic acid), $CH_3CH(OH)COOH$ (see p. 347), and sarcolactic acid, from flesh, being those of importance.

* Bodies having the CH_3CH group are called ethylidene compounds. Compare chloral hydrate, trichlorethylidene glycol, $CCl_3CH(OH)_2$.

The Acrylic Series.

Acids of the Acrylic Series, $C_nH_{2n-1}COOH$.

Acrylic Acid, C_2H_3COOH or $CH(CH_2)COOH$, is formed by oxidizing acrolein (acrylic aldehyde; see Glycerin) by silver oxide.

Crotonic Acid, or *Methacrylic Acid*, C_3H_5COOH or $CH(CH_3)COOH$, formerly supposed to be a constituent of croton oil, may be formed by oxidizing crotonic aldehyde, and by acting on allyl cyanide, C_3H_5CN , by water and hydrochloric acid: $C_3H_5CN + 2H_2O + HCl = C_3H_5CO\cdot OH + NH_4Cl$.

Oleic Acid, $C_{18}H_{34}O_2$, or $CH(C_{16}H_{32})COOH$, is found as a glyceric salt in many fats and oils.


Preparation.—Olive oil is saponified with caustic potash, and the resulting soap decomposed by tartaric acid, which liberates oleic and stearic acids. The oleic and stearic acids are heated with lead oxide, forming lead oleate and stearate, the former being dissolved out from the latter by ether. The ether is evaporated and the lead oleate treated with hydrochloric acid, which liberates the oleic acid.

Elaidic Acid (isomeric with oleic acid) is formed by passing nitrogen peroxide into oleic acid; it is more stable than oleic acid, distilling unchanged.

The Benzoic or Aromatic Series.

Acids of the Benzoic or Aromatic Series, $C_nH_{2n-7}COOH$.—The acids of this series are formed by oxidizing hydrocarbons, by oxidation of alcohols of the benzylic series, and by acting on the cyanides of the members of the benzene series. All the acids of this series, with the exception of benzoic acid, possess many isomers.

Benzoic Acid, C_6H_5COOH , occurs naturally in gum benzoin (gum benjamin), which contains from 12 to 15 per cent., the rest being mainly composed of two resins having the formulæ $C_{40}H_{46}O_9$ and $C_{30}H_{40}O_5$. Benzoic acid may be obtained by oxidizing benzoic aldehyde, C_6H_5COH , which may be prepared from trichloromethylbenzene. (See Toluene, p. 431.) Benzöene (toluene), $C_6H_5CH_3$, may be directly oxidized into benzoic aldehyde, the methyl group (CH_3) being resolved into $COOH$, evidence that benzoic acid is really a benzène derivative, not a phenène derivative. (For other modes of obtaining benzoic acid artificially, see p. 335.) It may also be produced from hippuric acid (benzamidacetic acid), p. 340. Benzoic acid heated with lime yields benzene:




Benzoic Aldehyde, or *Benzaldehyde*, C_6H_5COH , forms the greater part of oil of bitter almonds. (See Amygdalin, p. 499.) It is a colorless liquid, soluble in 30 parts of water and in all proportions in ether and alcohol. With acid potassium sulphite it, like other aldehydes, forms a crystalline compound, $C_6H_5\cdot COH\cdot NaHSO_3$.

Benzoyl Chloride, C_7H_5OCl , results from the action of chlorine on benzaldehyde (formerly termed *benzoyl hydride*, C_7H_5OH), or from the action of phosphorus pentachloride on benzoic acid (*benzoyl hydrate*, C_7H_5OOH). Benzaldehyde also results from the oxidation of the benzyl alcohol (C_7H_7OH) of balsam of Peru.




The other acids of this series are not very important.

The Hydroxybenzoic Series.

Acids of the Hydroxybenzoic Series, $C_nH_{2n-8}OH \cdot COOH$.—Just as the acids of the lactic series are related to the acetic series, so are the acids of the hydroxybenzoic (or salicylic) series related to the benzoic series.



Salicylic or Hydroxybenzoic Acid, $C_7H_6O_3$ or $C_6H_4OH \cdot COOH$ (*Acidum Salicylicum*, U. S. P.). Natural and artificial salicylates of methyl are described on p. 407. It also occurs in several species of violet (Mandelin). Salicylic acid may be made by the oxidation of salicylic aldehyde (*vide infra*), or by the action of carbonic acid on phenol or carbolic acid (Kolbe). To accomplish this, the phenol is mixed with caustic soda, forming sodium-phenol or sodium carbolate, C_6H_5ONa . The sodium-phenol is then saturated with carbonic acid at the ordinary temperature, by which sodium phenyl-carbonate is produced. The latter on being heated in closed vessels is transformed into sodium salicylate, from which salicylic acid may be obtained by the action of hydrochloric acid, and purification by recrystallization from alcohol. It appears to be identical with the natural acid.



Phenyl Salicylate.

Synonyms.—Salicylate of Phenyl; Salicylic Phenol.

Salol, $C_6H_4OH \cdot CO \cdot OC_6H_5$, is a new antiseptic, antipyretic, anti-rheumatic remedy. It is a white crystalline powder with a slight aromatic odor, soluble in alcohol, ether, chloroform, and the fixed and volatile oils, insoluble in water. When heated on platinum-foil it burns completely away.

Salol, U. S. P., should conform to the following tests: On warming a small portion of the substance with enough sodium hydrate

to dissolve it, and then supersaturating the liquid with hydrochloric acid, salicylic acid will separate, and the odor of phenol will become perceptible. In an alcoholic solution bromine-water, added in excess, produces a white precipitate. On adding a few drops of very dilute ferric chloride (1 in 200) to 10 cc. of an alcoholic solution (1 in 50) of salol, the liquid will acquire a violet tint. If, however, a few drops of the alcoholic solution be added to 10 cc. of the ordinary diluted ferric chloride, a whitish cloudiness, but no color, will be produced on agitation. On shaking 1 grm. of salol with 50 cc. of water the filtrate should not be affected by very dilute ferric chloride (absence of uncombined carbolic or salicylic acid), nor by barium chloride (absence of sulphate or phosphate), nor by silver nitrate (absence of chloride).

Table showing the Relations between the Benzoic and Hydroxybenzoic Acids.

Benzoic acid	$C_6H_5 \cdot CO \cdot OH$.
Hydroxybenzoic or salicylic acid .	$C_6H_4OH \cdot CO \cdot OH$.
Dihydroxybenzoic acid	$C_6H_3(OH)_2 \cdot CO \cdot OH$.
Trihydroxybenzoic or gallic acid .	$C_6H_2(OH)_3 \cdot CO \cdot OH$.

Salicylic acid, like carbolic acid, is a powerful antiseptic, but is free from the taste and smell of carbolic acid. It is only slightly soluble in cold water, but readily soluble in hot water, alcohol, ether, and in aqueous solutions of such alkali-metal salts as borax, sodium phosphate, or potassium citrate, which it converts into acid salts with formation of a salicylate. A similar antiseptic, *cresotic acid* (hydroxytoluic acid, $C_6H_3OH \cdot CH_3 \cdot COOH$), is similarly obtained from cresol or cresylic acid, $C_6H_4OH \cdot CH_3$. Ferric chloride strikes a violet coloration with both salicylic and cresotic acids. Both acids have antipyretic powers. The true salicylates of the alkali-metals, and probably therefore the cresotates, are very feeble antiseptics. Sodium salicylate (*Sodii Salicylas*, U. S. P.), $NaC_7H_5O_3$, the old salicylate of soda, made by neutralizing salicylic acid with sodium hydrate or carbonate, forms small, nearly colorless, lamellar crystals, soluble in alcohol and readily soluble in water. Carbolic acid often containing cresylic acid, commercial salicylic acid may contain cresotic acid. An alcoholic solution of salicylic acid allowed to evaporate spontaneously, exposure to dust being avoided, should leave a white residue free from color even at the points of the crystals. Salicylic acid yields colored substances on being nitrated and etherified, etc. *Iodosalicylic acid* and *di-iodosalicylic acid*, $C_7H_3IO_3$ and $C_7H_4I_2O_3$, are used in medicine.

Salicylic Aldehyde, or Hydroxybenzoic Aldehyde, $C_6H_4OH \cdot COH$ (Salicylous Acid, Hydride of Salicyl).—Found in the essential oil of meadow-sweet (*Spirea ulmaria*); also obtained by the oxidation of saligenin. (See p. 507.) It may be artificially formed by the action of chloroform on sodium-phenol.

Preparation.—Mix 10 parts of phenol with 20 parts of sodium hydrate dissolved in 30 parts of water in a flask having

1. *Balsam of Peru* (*Balsamum Peruvianum*, U. S. P.), an exudation from the trunk of *Myroxylon Pereiræ*, is a mixture of oily matter with about one-quarter or one-third resinous matter and 6 per cent. of cinnamic acid. The oil, by fractional distillation in an atmosphere of carbonic acid gas and under diminished pressure, furnishes *benzyl-hydrate*, or *benzylic alcohol* ($C_6H_5CH_2OH$), *benzyl benzoate* ($C_6H_5CO\cdot OC_7H_7$), and *benzyl cinnamate* ($C_8H_7CO\cdot OC_7H_7$), or *cinnamein* (Kraut). By action of alcoholic solution of potash it yields potassium benzoate and cinnamate and benzylic alcohol; also *cinnamic alcohol* ($C_8H_7CH_2OH$), otherwise known as *peruvine* or *styrone*; it also often holds in solution *metacinnamein* or *styracin* ($C_{18}H_{16}O_2$), isomeric with cinnamic aldehyde (C_8H_7COH). The resin of balsam of Peru seems to result from the action of moisture on the oil. Any admixture of resin, oil, storax, benzoin, or copaiva with balsam of Peru is detected by mixing 6 grains of slaked lime with 10 drops of the balsam, when a soft product results if the specimen be pure, but hard if impure; further, the mixture, on being warmed until volatile matter is expelled and charring commences, gives no fatty odor. 2. *Balsam of Tolu* (*Balsamum Tolutanum*, U. S. P.) is an exudation from the trunk of *Myroxylon Toluifera*; in composition it closely resembles balsam of Peru, but is more susceptible of resinification. It contains benzyl benzoate and cinnamate, cinnamic acid, a little benzoic acid (Busse), and about 1 per cent. of a volatile hydrocarbon, *tolene*, $C_{10}H_{16}$. The cinnamic-acid crystals may be seen with a lens when a little of the balsam is pressed between two warmed pieces of glass. Old hard balsam of Tolu is a convenient source of cinnamic acid, which may be extracted by the same process as that by which benzoic acid is obtained from benzoin—namely, ebullition with alkali, filtration, and precipitation by hydrochloric acid. 3. *Storax* (*Styrax*, U. S. P.) is an oleo-resin obtained from the *Liquidambar orientale*. It contains a volatile oil termed *styrol*, *cinnamene*, or *cinnamol* (C_8H_8)—which possibly (Berthelot) is condensed acetylene, $4C_2H_2$ —cinnamic acid, styracin, or cinnamyl cinnamate ($C_8H_7CO\cdot OC_9H_9$), and a soft and a hard resin. Styrol differs from similar hydrocarbons in being converted into a polymeric solid, termed *metastyrol* or *draconyl*, on heating to about 400° F. For medicinal use storax (*Styrax Præparatus*, B. P.) is purified by solution in alcohol, filtration, and removal of the alcohol by distillation. By oxidation with red potassium chromate and sulphuric acid it yields an odor resembling that of essential oil of bitter almonds.


DIBASIC ACIDS.

Dibasic Acids are acids having two carboxyl ($COOH$) groups in the molecule.


The Succinic Series.

Acids of the Succinic Series, $C_nH_{2n}(COOH)_2$.—These acids may be formed by the oxidation of glycols or by the action of water and hydrochloric acids on the cyanides of the olefines, obtained by acting on the olefine dibromo-additive derivatives by potassium cyanide.

Oxalic Acid, $C_2O_4H_2$ or $(COOH)_2$, is the first of this series. It may be obtained by oxidizing glycol, $C_2H_4(OH)_2$:



Also by the action of carbonic anhydride on metallic sodium:



(For other methods see *Oxalic Acid*, p. 316.)

Oxamide, $C_2O_2(NH_2)_2$, the analogue of urea—carbamide, $CO(NH_2)_2$ —is formed on mixing ethyl oxalate with ammonia or by passing cyanogen into aqueous hydrochloric acid, $C_2N_2 + 2H_2O = (CONH_2)_2$.

Succinic Acid, $C_2H_4(COOH)_2$.—(See p. 356.)

The Malic Series.

Acids of the Malic Series, $C_nH_{2n-1}OH(COOH)_2$.—*Malic* or *hydroxysuccinic acid*, $C_2H_3(OH)(COOH)_2$, is obtained artificially by acting on bromosuccinic acid, $C_2H_3Br(COOH)_2$, with moist silver oxide, the bromine being replaced by hydroxyl. It is contained in unripe mountain-ash berries, morello cherries, etc. (See p. 348.)

Asparagin (amidosuccinamic acid), $C_2H_3NH_2 < \begin{array}{c} CONH_2 \\ COOH \end{array}$. (See p. 349.)

The Tartaric Series.

Acids of the Tartaric Series, $C_nH_{2n-2}(OH)_2(COOH)_2$.—*Tartaric Acid* (dihydroxysuccinic acid), $C_2H_2(OH)_2(COOH)_2$, may be obtained by oxidizing erythrite, $C_2H_2(OH)_2(CH_2OH)_2$. (See p. 465. For other modes of formation see p. 318.) There are four isomeric tartaric acids, differing by their action on a ray of polarized light.

The Phthalic Series.

Acids of the Phthalic Series, $C_nH_{2n-8}(COOH)_2$.—*Phthalic Acid*, $C_6H_4(COOH)_2$, is obtained by the oxidation of naphthalene and naphthalene tetrachloride, or a mixture of benzene and benzoic acid. By distillation it forms phthalic anhydride, $C_8H_4O_3$, and this when heated with phenol and sulphuric acid yields *phenolphthalein* (B. P.), a light-yellow crystalline powder, which when dissolved in alcohol is used in alkalimetry for its property of turning brilliant red with the slightest excess of alkali. There are three phthalic acids: phthalic acid or orthophthalic acid, $C_6H_4 \cdot COOH \cdot COOH_{(o)}$; isophthalic acid or metaphthalic acid, $C_6H_4 \cdot COOH \cdot COOH_{(m)}$; and terephthalic acid or paraphthalic acid, $C_6H_4 \cdot COOH \cdot COOH_{(p)}$. (See p. 455.)

TRIBASIC ACIDS.

Tribasic Acids, having three carboxyl $(COOH)$ groups in the molecule.—*Tricarballic Acid*, or propane-tricarboxylic acid, C_3H_5-

TABLE SHOWING THE RELATIONS BETWEEN THE ACIDS OF THE ACETIC SERIES AND THE DIBASIC ACIDS.

Acids of the Acetic Series, $C_nH_{2n+1}COOH$.	Acids of the Succinic Series, $C_nH_{2n}(COOH)_2$	Acids of the Hydroxysuccinic or Malic Series, $C_nH_{2n-1}(OH)(COOH)_2$	Acids of the Dihydroxysuccinic or Tartaric Series, $C_nH_{2n-2}(OH)_2(COOH)_2$
Formic, $H.CO.OH$	Oxalic, $COOH.CO.OH$.	—	—
Acetic, $CH_3.CO.OH$.	Malonic, $CH_2(CO.OH)_2$	Tartaric (oxymalonic), $CH.OH.(COOH)_2$	Mesoxalic, $C(OH)_2(COOH)_2$
Propylic, $C_2H_5.CO.OH$.	Succinic, $C_2H_4(CO.OH)_2$	Malic (oxysuccinic), $C_2H_3.OH.(COOH)_2$	Tartaric (or dioxysuccinic), $C_2H_2(OH)_2(COOH)_2$
Tetrylic, $C_3H_7.CO.OH$	Pyrotartaric, $C_3H_6(CO.OH)_2$	Glutanic, $C_3H_5.OH.(COOH)_2$	Homotartaric, $C_3H_4(OH)_2(COOH)_2$

$(\text{COOH})_3$, is the first of these series; its hydroxy-derivative is *citric acid*, $\text{C}_3\text{H}_4(\text{OH})(\text{COOH})_3$, (hydroxy-propene-tricarboxylic acid), found in fruits. It has already been described. (See p. 323.)

OTHER POLYBASIC ACIDS.


Tetrabasic acids, as pyromellitic acid, $\text{C}_6\text{H}_2(\text{COOH})_4$, and *hexabasic* acids, as mellitic acid, $\text{C}_6(\text{COOH})_6$, are known.

QUESTIONS AND EXERCISES.

Give general methods for the formation of aldehydes and acids.—How is acetaldehyde prepared?—Describe the reactions that occur in the manufacture of chloral and chloral hydrate.—What is the nature of the action of alkalis on chloral hydrate?—Mention the characters of pure and impure chloral hydrate.—What relation has valerianic acid to amylic alcohol?—Give the relations between the acetic and lactic series of acids.—To what series do the following acids belong: Oleic, butyric, oxalic, and citric?—How is benzoic acid prepared?—Give the differences between balsams of Peru and Tolu and gum benzoin.—How is oil of bitter almonds prepared? and how can it be distinguished from so-called artificial oil of bitter almonds?—Give artificial methods of preparing salicylic aldehyde and acid.—Give systematic names of tartaric, succinic, carbonic, salicylic, and citric acids.

KETONES.

Just as primary alcohols on losing hydrogen yield aldehydes, so secondary alcohols (see p. 437) on losing hydrogen yield ketones:



Like aldehydes, ketones are converted by nascent hydrogen into the corresponding alcohols. Like aldehydes, ketones form crystalline compounds with acid sulphites. While, however, aldehydes by oxidation yield corresponding acids, ketones break up and yield acids whose molecules have a smaller number of carbon atoms.

Acetone, $\text{C}_3\text{H}_6\text{O}$, or *Dimethyl-ketone*, $(\text{CH}_3)_2\text{CO}$ or $\text{CH}_3\cdot\text{CO}\cdot\text{CH}_3$, the original and best known of the class, may be obtained by strongly heating calcium acetate, carbonate remaining. The calcium salts of other fatty radicals split up in a similar manner (hence perhaps the name, from *kéō*, *keō*, I split, and the original *acetone*), yielding other ketones, as propione, butyrone, valerone, etc. The mixed calcium salts give corresponding ketones. Thus acetate and caprate yield methyl-nonyl ketone, $\text{CH}_3-\text{CO}-\text{C}_9\text{H}_{19}$, the chief natural constituent of oil of rue. *Acetophenone*, or *phenylmethyl ketone*, $\text{C}_6\text{H}_5\cdot\text{CO}\cdot\text{CH}_3$, is known as *hyponone*.



Note.—There are many organic substances the composition of which has been established and the characters of which are definite, whether basic, acid, or neutral, but whose constitution is still so questionable that they cannot yet be classified with the hydrocarbons and derivatives of hydrocarbons. These are the glucosides, alkaloids, albumenoids, certain coloring matters, etc. They are described in the following pages.

THE GLUCOSIDES.

Source.—The glucosides are certain proximate vegetable principles which, by ebullition with dilute acid or other method of decomposition, take up the elements of water and yield glucose, accompanied by a second substance, which differs in each case according to the body operated on. Several of the glucosides which are of pharmaceutical interest will now be considered. Tannin has also been said to be a glucoside; it has been described among the Acids.

There are indications that all glucosides may be regenerated from the bodies into which they are thus converted.

Note on Nomenclature.—The first syllable of the names of glucosides and neutral principles generally is commonly given in allusion to origin; the last syllable is *in*, which sufficiently distinguishes them as a class.


ABSINTHIN ($C_{15}H_{10}O_4$), the bitter principle of *Artemisia Absinthium*, or *wormwood*, yields, when boiled with acids, glucose, volatile oil, and a resin of the aromatic series. (The liqueur termed *absinthe* is spirit of wine flavored with natural oil of wormwood, colored by chlorophyll, and slightly sweetened.)

AMYGDALIN ($C_{20}H_{27}NO_{11} \cdot 3H_2O$).—This body, obtained by Robiquet and Boutron-Charlard in 1830, was the first-discovered glucoside (Liebig and Wöhler, 1837). It is a white crystalline substance existing in the bitter (*Amygdala Amara*, U. S. P.), but not in the sweet almond (*Amygdala Dulcis*, U. S. P.). About 2 per cent. is readily extracted by strong alcohol from the cake left when the fixed oil has been expressed from bitter almonds. From the concentrated alcoholic solution ether precipitates the amygdalin.

Make an emulsion of two or three sweet almonds by bruising and rubbing them with water, and notice that it has no odor of essential oil of bitter almonds; add a grain or two of amygdalin: an odor of essential oil of bitter almonds is at once developed. Bruise two or three bitter almonds and rub with water: the volatile oil is again developed (*Oleum Amygdalæ Amaræ*, U. S. P.). Sp. gr. 1.060 to 1.070.

Bitter-almond Water (*Aqua Amygdalæ Amaræ*, U. S. P.) is made by filtering a mixture of 1 part of the oil with 999 parts of distilled water. The source of the benzaldehyde, or essential oil of bitter

almonds, in these reactions is the amygdalin, which, under the influence of *synaptase* or *emulsin* (a nitrogenous, casein-like ferment existing in both bitter and sweet almonds), splits up into the essential oil, hydrocyanic acid, and glucose:



As each molecule of amygdalin yields one of hydrocyanic acid, a simple calculation shows that 17 grains (mixed with emulsion of sweet almonds) will be required to form 1 grain of real hydrocyanic acid, a quantity equivalent to 50 minims of the diluted hydrocyanic acid of the British Pharmacopoeia. The hydrocyanic acid is probably in chemical combination with the oil to the extent of about 5 per cent. According to Linde, the occurrence of the benzaldehyde is preceded by the formation of benzaldehydecyanhydrin ($\text{C}_6\text{H}_5\text{CH}(\text{OH})\text{CN}$). The emulsin and amygdalin occur in different parts of the bitter almond.

Test.—The reaction between synaptase and amygdalin is applicable as a test of the presence of one by the addition of the other, even when mixed with much organic matter.

Jacobsen obtains true benzaldehyde artificially from benzodichloride (dichloromethylbenzene, $\text{C}_6\text{H}_5\text{CHCl}_2$), one of the dichlorotoluenes, by heating with glacial acetic acid and zinc chloride with a little water.

Cherry-laurel Water (*Aqua Laurocerasi*, B. P., by distillation with water from *Laurocerasi Folia*, B. P.) contains hydrocyanic acid, derived from a reaction similar to, indeed probably identical with, that described above, for bitter almond oil is simultaneously produced. But the proportion of amygdalin or analogous body in cherry-laurel leaves is most variable; hence normally the strength of the water is highly uncertain. The British Pharmacopoeia, however, directs that it shall contain 0.1 per cent. of real hydrocyanic acid, it being strengthened by the addition of hydrocyanic acid, or, if necessary, diluted by the addition of distilled water, until it has the prescribed strength.

Linseed yields a glucoside, *linamarin*, related to amygdalin; for it yields glucose and hydrocyanic acid on hydrolysis.

Cortex Pruni Serotinae.—The recently dried wild-cherry bark also furnishes by distillation an essential oil and hydrocyanic acid. Quince-seeds also (*Pyrus Cydonia*). The wild black cherry bark (*Prunus Virginiana*, U. S. P.), collected in autumn, contains amygdalin.

Caution.—Essential oil of almonds is of course highly poisonous. The purified oil or benzaldehyde is almost innocuous; it is obtained on distilling the crude oil with milk of lime and ferrous chloride and drying the product by shaking with fused calcium chloride. The so-called "artificial oil of bitter almonds" or "nitrobenzol" [$\text{C}_6\text{H}_5\text{NO}_2$], when taken in quantity, has been known to produce death. The presence of nitrobenzol in oil of bitter almonds is detected by adding a little of the oil to a mixture of zinc and diluted sulphuric acid, shaking well, setting aside for an hour or two, filter-

ing off the clear liquid, and adding a little potassium chlorate; a violet color (actual mauve) is produced. The reaction is due to the formation of phenylamine or aniline. (See p. 431.) Or the specimen may be shaken with sodium bisulphite to fix the benzaldehyde (for all such aldehydes form a compound with sodium bisulphite), and then with ether, which dissolves out, and on evaporation will yield, the nitrobenzol.

ARBUTIN ($C_{12}H_{16}O_7$) and *Methyl-arbutin* ($C_{13}H_{18}O_7$) are contained in the leaves of *Arctostaphylos Uva Ursi*, *Chimaphila umbellata* (*Chimaphila*, U. S. P., or *Pipsissewa*), and many ericaceous plants. They are bitter neutral bodies occurring in acicular crystals, and resolvable by acids into *hydroquinone* ($C_6H_6O_2$) and glucose, and by gentle oxidation into *quinone* ($C_6H_4O_2$) and formic acid. *Eriocolin* ($C_{34}H_{56}O_{21}$) is another bitter glucoside in bearberry-leaves.

BRYONIN ($C_{48}H_{80}O_{19}$, Walz).—The colorless, bitter, indistinctly crystalline principle of bryony (*Bryonia*, U. S. P., the root of *Bryonia alba* and *Bryonia dioica*).

CATHARTIC ACID.—“The glucoside acid that now is known to confer on the senna of Alexandria (from *Cassia acutifolia*) and of India (from *Cassia elongata*) (*Senna*, U. S. P.) its purgative property has been named by its discoverers (Dragendorff and Kubly) cathartic acid. Its formula has been stated as $C_{180}H_{192}N_4SO_{82}$ (but A. Gensz more recently states it to be $C_{30}H_{36}NO_{15}$). It is insoluble (?) in water, strong alcohol, and ether, but enters readily into either solution when combined with alkaline and earthy bases, in which state it exists in senna. Its ammonium salts give brownish flocculent precipitates with salts of silver, tin, mercury, copper, and lead. Antimonial salts, tannin, and yellow and red prussiates have no effect upon it. Alkalies, aided by heat, act destructively upon it. Boiled with a mineral acid, it splits into a peculiar kind of glucose and an acid that has been named cathartogenic; its formula is said to be $C_{132}H_{116}N_4SO_{44}$. The natural cathartate occurring in senna is prepared by partially precipitating by strong spirit a watery infusion of senna, concentrated to a syrupy state by evaporation *in vacuo*. The filtrate is now treated with a much larger bulk of absolute alcohol, and the precipitate thus obtained is purified by repeated solution in water and precipitation by alcohol. To obtain the pure acid advantage is taken of its colloidal properties; the crude cathartate is dissolved in moderately strong hydrochloric acid, and subjected to dialysis on a diaphragm of parchment paper. The minimum dose of this pure acid was found to be about $1\frac{1}{2}$ grains, which caused several stools with decided griping.”

“The cathartic combinations that I have made are—the cathartate of ammonium, prepared from cathartate of lead by my original process, and the mixed cathartates, prepared according to Dragendorff’s method as modified by myself. Of the former nearly pure salt I have found $3\frac{3}{4}$ grains to purge fairly as to amount, but slowly as to time and with considerable griping. Of the latter, $7\frac{1}{2}$ grains purged violently with much griping and sickness, which continued through the greater part of the day. It obviously would be improper to combine senna with any of its metallic precipitants should such be

desired, which is not likely. It is here satisfactory to observe that the cathartate of magnesium is soluble, and that the old-fashioned black draught agrees with new-fashioned science" (Groves).

Buckthorn Juice (Rhamni Succus, B. P. 1867) owes its cathartic properties to a substance apparently identical with cathartic acid. Possibly the purgative properties of the bark of the *Rhamnus Frangula* (*Frangula*, U. S. P.), black alder, buckthorn, also are due to cathartic acid.


COLOCYNTHIN ($C_{56}H_{84}O_{23}$?).—This substance is the active bitter and purgative principle of colocynth-fruit (*Colocynthis Pulpa*, U. S. P.): it is soluble in water and alcohol, but not in ether. By ebullition with acids it furnishes glucose and a resinoid body.

CONVOLVULIN. (See JALAPIN.)

COTOIN ($C_{22}H_{18}O_6$) appears to be the chief active principle of coto-bark, a Bolivian remedy for diarrhoea. A similar bark, *false coto* or *paracoto*, contains *paracotoin*, $C_{19}H_{12}O_6$, and *hydrocotoin*, $C_{15}H_{14}O_4$.

DAPHNIN ($C_{31}H_{34}O_{19}$) is the crystalline glucoside of the bark of *Daphne Mezereum* (*Mezerei Cortex*, B. P.). Boiled with dilute acids, it yields *daphnetin* ($C_{19}H_{14}O_9$) and glucose. The acrid principle of mezereon is resinoid.

DIGITALIN ($C_{27}H_{45}O_{15}$, Kosmann; $C_5H_8O_2$, Schmiedeberg).—This is an active principle of the foxglove, *Digitalis purpurea* (*Digitalis*, U. S. P.). On boiling a grain of digitalin with diluted sulphuric acid for some time, flocks of *digitaliretin* ($C_{15}H_{25}O_5$) separate, and glucose may be detected in the liquid.



Properties.—Digitalin occurs "in porous mammillated masses or small scales, white, inodorous, and intensely bitter, readily soluble in spirit, but almost insoluble in water and in pure ether; dissolves in acids, but does not form with them neutral compounds; its solution in hydrochloric acid is of a faint yellow color, but rapidly becomes green. It leaves no residue when burned with free access of air. It powerfully irritates the nostrils and is an active poison."—B. P. 1867. According to Pettenkofer, "an intense red color is produced if a trace of digitalin dissolved in water is mixed with a weak aqueous solution of inspissated bile and sufficient oil of vitriol added to raise the temperature to 158° F." Moistened with sulphuric acid and the liquid exposed to the vapor of bromine, a violet color is produced.

Process.—The process for the preparation of digitalin consists in dissolving the glucoside out of the digitalis-leaf (*Digitalis Folia*, B. P.) by alcohol, recovering the alcohol by distillation, dissolving the residue in water by the help of a small quantity of acetic acid, removing much of the color from the solution by animal charcoal, neutralizing most of the acetic acid by ammonia, precipitating the digitalin by tannic acid (with which it forms an insoluble compound), washing the precipitate, rubbing and heating it with spirit and lead oxide (which removes the acid in the form of insoluble lead tannate),

again decolorizing by animal charcoal, evaporating to dryness, washing out impurities still remaining by ether, and drying the residual digitalin. In this form digitalin is uncrystallizable and somewhat indefinite.

Pure Digitalin (?).—On treating commercial digitalin with chloroform an inert substance remains undissolved. The solution yields pure digitalin on evaporation; it may be crystallized from spirit in radiating needles (Nativelle). The therapeutic effect of the pure substance is identical with that of the preparations of digitalis, but more constant in its action, and, of course, intensely powerful.

Digitoxin ($C_{31}H_{53}O_7$) ($C_{31}H_{52}O_7$, Dragendorff) is a highly poisonous substance extracted from foxglove by Schmiedeberg. The same chemist regards commercial digitalin from foxglove-seeds as composed of varying proportions of three glucosides—namely, pure active *digitalin* ($C_5H_8O_2$), *digitonin* ($C_{31}H_{52}O_{17}$) (closely allied to saponin), and *digitalein*, with inactive digitalin or *digitin*. Kiliani says that Schmiedeberg's digitalein is a mixture, and that his "digitalin," which is a pure but uncrystallizable glucoside, $C_5H_8O_2$, is the true active principle. *Digitonin*, $C_{27}H_{46}O_{14}$ (Kiliani), yields on hydrolysis *digitogenin*, $C_{15}H_{24}O_3$, galactose, $C_6H_{12}O_6$, and dextrose, $C_6H_{12}O_6$.

ELATERIN ($C_{20}H_{28}O_5$).—Boil elaterium (*Elaterium*, B. P.), the dried sediment from the juice of the squirting cucumber fruit, *Ecballium Elaterium* (*Ecballii Fructus*, B. P.), with chloroform, filter, evaporate, wash with ether the precipitated elaterin, recrystallize it from chloroform, and again wash the crystals with ether. The product, the official *Elaterinum*, U. S. P., occurs in small hexagonal plates or prisms. A trituration of elaterin is official (*Trituratio Elaterini*, U. S. P.). It is a mixture of 1 part of elaterin with 9 of sugar of milk.

Elaterin is probably not a true glucoside. It does not always respond to the test for glucose after boiling with acids, and when it does the reaction is possibly due to *prophetin*, a glucoside stated by Walz to be present in elaterium.

Elaterin is the active principle of the so-called elaterium. Elaterium occurs in light friable greenish-gray cakes. Good specimens of this drug should yield not less than 20 per cent. of elaterin by the above process.

Test.—A little is placed in a watch-glass with a drop of liquefied carbolic acid, and then two drops of strong sulphuric acid: a carmine color is developed (Lindo).

FRANGULIN, $C_{21}H_{20}O_9$, is a glucoside found in the bark of *Rhamnus frangula*. It is decomposed on hydrolysis into *emodin*, $C_{15}H_{10}O_5$, and an unfermentable sugar, *rhamnose*, $C_6H_{12}O_5$.

GENTIOPICRIN, or *GENTIAN BITTER* ($C_{20}H_{30}O_{12}$), the neutral crystalline principle of the root of *Gentiana lutea* (*Gentiane Radix*, B. P.). It is soluble in water and weak spirit. Alkalies decompose it. Dilute acids convert it into *gentiogenin* and glucose. Gentian-root also contains a variety of tannin and a crystalline acid

($\text{HC}_{14}\text{H}_9\text{O}_5$) termed *gentianic* or *gentisic acid* or *gentisin*. Fused potash, etc. gives with the latter an acid ($\text{C}_7\text{H}_6\text{O}_4$), which has also, unfortunately, been called *gentisic acid*.

GLYCYRRHIZIN ($\text{C}_{24}\text{H}_{36}\text{O}_9$, Gorup-Besanez), or *Glycyrrhizic Acid* ($\text{C}_{44}\text{H}_{63}\text{NO}_{18}$, Habermann).—Liquorice-root (*Glycyrrhiza*, U. S. P.), in addition to uncrystallizable sugar, contains 3 or 4 per cent. of a sweet substance, *glycyrrhizin*, which, when boiled with hydrochloric acid or diluted sulphuric acid, yields a resinoid bitter body, *glycyrrhizin*, and an uncrystallizable sugar resembling glucose. *Glycyrrhizin* is only slightly soluble in cold water, but is taken up by diluted alcohol containing a little ammonia (*Extractum Glycyrrhizæ Fluidum*, U. S. P.) or by ammoniacal water. An infusion of the latter, evaporated to a pilular consistence, forms *Extractum Glycyrrhizæ Purum*, U. S. P. It is present in considerable quantity in the evaporated decoction (*Stick Liquorice*, *Spanish Liquorice*, or *Solazzi Juice*). The tropical substitute for liquorice is the root of *Abrus precatorius*, or *Indian Liquorice* (*Abri Radix*, P. I.), the *kunch* or *gunj* of Bengal, the *ratti* of Hindostan, and the *jequirity* or *jequerity* of Brazil, which also contains glucose and *glycyrrhizin*. The seeds yield by maceration a substance which acts as a poison when injected into the blood, but not when swallowed. Warden and Waddell regard the active principle as an albumenoid, and term it *abrin*. Bruylants and Venneman consider it to be a product of germination, and call it *jequeritin*. Bechamp and Dujardin regard the latter as a mixture of legumin and *jequerityzyme*. *Glycyrrhizin* has considerable power of disguising nauseous flavors. Roussin refers the sweet taste of liquorice not to pure *glycyrrhizin*, but to a combination of *glycyrrhizin* with alkalies, and states that ammoniacal *glycyrrhizin* has exactly the sweetness of liquorice-root. The formula of this *ammonium glycyrrhizate* is said by Habermann to be $(\text{NH}_4)_3\text{C}_{44}\text{H}_{60}\text{NO}_{18}$. Sestini finds that the *glycyrrhizin* of liquorice-root is chiefly calcium *glycyrrhizate*.

An ammoniated *glycyrrhizin* (*Glycyrrhizinum Ammoniatum*, U. S. P.) is directed to be prepared by precipitating a dilute ammoniacal percolate with sulphuric acid, washing, redissolving in ammoniacal water, reprecipitating, again washing, dissolving in solution of ammonia, and spreading on glass plates to dry until reddish-brown scales are obtained.


GUAIACIN.—Resin of *guaiacum* (*Guaiaci Resina*, U. S. P.), an exudation from the wood (*Guaiaci Lignum*, U. S. P.) of *Guaiacum officinale*, is probably a mixture of several substances, among which are *guaiaretic* or *guaiaretinic acid* ($\text{C}_{26}\text{H}_{26}\text{O}_4$, Hlasiwetz), *guaiaconic acid* ($\text{C}_{38}\text{H}_{40}\text{O}_{10}$, Hadelich), and *guaiacin*, a glucoside. On boiling *guaiacum* resin with diluted sulphuric acid for some time, glucose is found in the liquid, a green resinous substance (*guaiaretin*) remaining insoluble (Kosmann). Most oxidizing agents, and even atmospheric air, especially under the influence of certain organic substances, produce a blue, then green, and finally a brown color when brought into contact with an alcoholic solution of *guaiacum* resin.

These effects are said to be due to three stages of oxidation

(Jonas). They may be observed on adding the solution to the inner surface of a paring of a raw potato.

HELLEBORIN ($C_{36}H_{42}O_6$) and HELLEBOREIN ($C_{26}H_{44}O_{15}$) are crystalline glucosides occurring in the roots of black hellebore (*Helleborus niger*), or Christmas rose, and green hellebore (*H. viridis*), ranunculaceous herbs.

JALAPIN ($C_{31}H_{50}O_{16}$) and CONVULVULIN ($C_{34}H_{56}O_{16}$).—According to Keyser and Meyer, jalap resin contains two distinct substances—convolvulin, chiefly obtained from Mexican male jalap (*Ipomœa Orizabensis*), and jalapin, most largely contained in the true jalap (*Ipomœa purga*); the former is soluble in ether, the latter insoluble. Boil jalap resin with diluted sulphuric acid for some time, and filter; a substance which is probably a mixture of *jalapinol* ($C_{13}H_{24}O_3$) and *convolvulinol* ($C_{16}H_{30}O_3$), separates; glucose may be detected in the clear liquid. (It is to be regretted that the authors transpose these names, terming the old well-known jalapin convolvulin.)



Jalapic Acid.—This is contained in the portion of jalap resin soluble in ether. It may also be obtained from jalapin by ebullition with alkalis:



Jalap resin (*Resina Jalapæ*, U. S. P.) is obtained by digesting and percolating jalap-tubercles (*Jalapa*, U. S. P.) with spirit of wine, adding a little water, distilling off the spirit, pouring away the aqueous portion, which contains much saccharine matter, and washing and drying the residual resin. Jalap thoroughly exhausted by this process should furnish, according to the United States Pharmacopœia, not less than 12 per cent. of resin, of which resin (*Resina Jalapæ*, U. S. P.) not more than one-tenth should be soluble in ether—a test which excludes the so-called “Tampico” jalap, the resin of which is soluble in ether. The tincture of jalap is sometimes decolorized by animal charcoal and the evaporated product sold as “jalapin.”

Jalap resin is insoluble in oil of turpentine; common resin or rosin, soluble. If the presence of the latter is suspected, the specimen should be powdered, digested in turpentine, the mixture filtered, and the filtrate evaporated; no residue, or not more than yielded by the turpentine itself, should be obtained.

Tampico jalap, from *Ipomœa simulans*, yields a resin which apparently is chiefly convolvulin, but sometimes contains jalapin, for a sample obtained by Hanbury was entirely soluble in ether, and another extracted by Umney was almost wholly soluble, while Evans purified some, half only of which was soluble.

The *Kaladana resin* or *pharbitisin* of India (from *Pharbitis Nil*, P. I.) is a cathartic analogous to, if not identical with, resin of jalap.

LOGANIN, $C_{25}H_{34}O_{14}$, is a glucoside obtained from the pulp of the fruit and from the seeds of *Strychnos Nux Vomica*, Loganiaceæ, by Dunstan and Short. Boiled with dilute sulphuric acid, it yields glucose and *loganetin*.

OUABAIN ($C_{30}H_{46}O_{12}$, Arnaud, or $C_{30}H_{52}O_{14}$) is a very poisonous glucoside resembling strophanthin, found in arrow-poisons prepared from the wood of an *Acokanthera* and in the wood itself.

PICROTOXIN (*Picrotoxinum*, U. S. P.) is a crystalline bitter poisonous principle ($\pi\kappa\rho\acute{o}\varsigma$, *picros*, bitter, and $\tau\omicron\xi\kappa\acute{o}\nu$, *toxicon*, poison) occurring in *Cocculus Indicus*, the dried fruit of *Anamirta paniculata*. Ludwig regarded it as a glucoside, but its constitution is not yet satisfactorily ascertained. Barth and Kretschy state that the so-called picrotoxin may be separated into *picrotoxin* proper ($C_{15}H_{16}O_6 \cdot H_2O$), which is bitter and poisonous; *picrotin* ($C_{25}H_{30}O_{12} + nH_2O$), which is bitter, but not poisonous; and *anamirtin* ($C_{19}H_{24}O_{10}$), which is neither bitter nor poisonous. Schmidt asserts that the original picrotoxin is definite, and has the formula $C_{30}H_{34}O_{13}$, but that some solvents decompose it into *picrotoxinin*, $C_{15}H_{16}O_6$, which is poisonous, and *picrotin*, $C_{15}H_{16}O_7$, which is not poisonous.


QUASSIN ($C_{10}H_{12}O_8$, Wiggers, or $C_{31}H_{42}O_9$, Christensen), obtained from *Quassia Lignum*, B. P., is said to be a glucoside, but Oliveri and Denaro question the statement, and find quassin to have the formula $C_{32}H_{44}O_{10}$.

SAFFRANIN. (*Vide Index*.)

SALICIN ($C_{13}H_{18}O_7$).—This substance (*Salicinum*, U. S. P.) is contained in, and easily extracted from, the bark of willow and of other species of *Salix* and of *Populus*. It occurs in white, shining, bitter crystals, soluble in 28 parts of water or 65 of spirits at common temperatures.

Tests.—1. To a small portion of salicin placed on a white plate or dish add a drop of strong sulphuric acid; a deep-red color is produced.


2. Boil salicin with diluted sulphuric acid for some time; it is converted into *saligenin* or *saligenol* ($C_7H_8O_2$) and glucose. Test for the latter by the copper test.



3. To another portion of the liquid, carefully neutralized, add a ferric salt: a purplish-blue color is sometimes produced, due to the reaction of the saligenin and the ferric salt. The saligenin is, however, so rapidly decomposed by acids into *saliretin* (C_7H_6O) and water that this reaction is almost valueless as a test. The saligenin may, however, be obtained by action of synaptase on salicin.

4. Heat a mixture of about 1 part of salicin, 1 of red potassium chromate, $1\frac{1}{2}$ of sulphuric acid, and 20 of water in a test-tube; a fragrant characteristic odor is evolved, due to the

formation of salicylic aldehyde ($C_6H_4OH \cdot COH$), an essential oil identical with that existing in meadow-sweet (*Spiræa Ulmaria*) and in heliotrope.



SANTONIN ($C_{15}H_{18}O_3$).—This substance, used in *Trochisci Santonini*, U. S. P., is, apparently, the anhydride or, rather, the lactone,* of a weak acid (Hesse) insoluble in ammonia, but forming a soluble calcium salt. Indeed, by boiling santonin for twelve hours with baryta-water Cannizarro has obtained a salt from which hydrochloric acid separates *santoninic acid* ($C_{15}H_{20}O_4$). From a solution of calcium santonate the santonin is precipitated by acids. Boiled for some time with diluted sulphuric acid, it yields 87 per cent. of an insoluble resinous substance (*santonivretin*) and glucose (Kosmann). *Santonin* (*Santoninum*, U. S. P.) is official; it is soluble in an aqueous solution of twice its weight of sodium carbonate. Possibly (Berthelot) santonin resembles carbonic acid; in other words, is a phenol, $C_{15}H_{15}O_3H$. Its glucosidic character is questionable.

Process.—The process for its preparation consists in boiling *santonica* (*Santonica*, U. S. P.), the dried unexpanded flower-heads or capitula of *Artemisia pauciflora* with milk of lime (whereby calcium santonate is formed), straining, precipitating the santonin or santoninic acid by hydrochloric acid, washing with ammonia to remove resin, dissolving in spirit, and digesting with animal charcoal to get rid of coloring-matter, setting the spirituous solution aside to deposit crystals of santonin, and purifying by recrystallization from spirit (Mialhe).

Test.—To highly-diluted solution of ferric chloride add an equal bulk of concentrated sulphuric acid. To this reagent add the santonin, or powder or substance suspected to be santonin, and cautiously apply heat. A red, purple, and finally violet, color is produced (Lindo). *Santonin* added to warm alcoholic solution of potash yields a violet-red color.

Tanacetie Acid, from the leaves and tops of *Tanacetum vulgare*, or *Tansy* (*Tanacetum*, U. S. P.), is a yellow crystalline acid having the medicinal properties of santonin.

SAPONIN ($C_{32}H_{52}O_{17}, H_2O$) is a peculiar glucoside occurring in soapwort, the root of the common pink, and many other plants; its solution in water, even though very dilute, froths like a solution of soap. Heated with dilute acids, it yields sugar and *saponetin*, $C_{40}H_{66}O_{15}$, or *saponegol*, $C_{14}H_{22}O_2$ (Hesse). Pereira considered *smilacin* (*salseparin* or *parallin*) one of the principles of the supposed activity of the root of *Smilax officinalis*, or sarsaparilla (*Sarsaparilla*, U. S. P.), to be

*The hydroxyacids losing water furnish lactones. Aromatic compounds containing NH_2 in the ortho position, and losing water by the oxidation and removal of one or two atoms of that hydrogen, furnish bodies which may be distinguished as *lactams* and *lactims*.

closely allied to, if not identical with, saponin. According to Klunge ("Pharmacographia"), parallin by action of acids yields *parigenin*. The aqueous solutions of parallin froth when shaken. Von Schultz states that sarsaparilla contains three homologous glucosides analogous to saponin—namely, sarsaparill-saponin ($C_{20}H_{32}O_{10}$), sarsa-saponin ($C_{22}H_{36}O_{10}$), and parallin ($C_{26}H_{44}O_{10}$).

Saponin is also met with in the root of *Polygala Senega* (*Senegæ*, U. S. P.), though the active principle of senega is said to reside in *polygalic acid*, probably a glucoside derivative of saponin.

Saponin is readily obtained from the bark of *Quillaja saponaria*, or *soap-bark* (*Quillaja*, U. S. P.), by boiling the aqueous extract in alcohol and filtering while hot. Flocks of saponin separate on cooling. It is a white, non-crystalline, friable powder.

The alleged toxic properties of commercial saponin are said by Kobert to be due to *sapotoxin* and *quillaic acid*.

SCAMMONIN ($C_{34}H_{56}O_{16}$).—Boil resin of scammony (*Resina Scammonii*, U. S. P.) with diluted sulphuric acid for some time; glucose may then be detected in the liquid, a resinous acid termed *scammoniol* ($C_{14}H_{13}O_3$?) being produced at the same time.

Natural scammony (*Scammonium*, U. S. P.) is an exudation from incisions in the living root (*Scammoniae Radix*, B. P.) of *Convolvulus Scammonia*. It contains from 10 to 20 per cent. of gum, and therefore, when rubbed up with water, gives an emulsion. "Ether removes about 75 per cent. of resin" (B. P.). The official resin of scammony contains no gum, and therefore gives no emulsion when rubbed up with water. It is made by digesting the root in spirit, adding water, distilling off the alcohol, and washing the residual resin with hot water till free from gum. There seems to be little or no chemical difference between the extracted resin and the resin of the exuded scammony.

Resin of scammony is soluble in all proportions of ether. Spirigatis states that it is identical with the resin of Mexican male jalap, which also is soluble in ether. Sulphuric acid slowly redens it. It is said to be liable to adulteration with resin of true jalap, guaiacum resin, and common rosin. Resin of true jalap is insoluble in ether; guaiacum resin is distinguished by the color-tests mentioned under Guaiacin, and rosin by the action of sulphuric acid.

SCILLITIN.—Schroff, and afterward Riche and Remont, believed the bitter principle of the squill-bulb (*Scilla*, B. P.) to be a glucoside. Merck has extracted substances which he has termed *scillipicrin*, *scillitoxin*, and *scillin*. Schmiedeberg has given the name of *sines-trin* to a squill principle. But no definite crystalline principle has yet been obtained. Squill contains a large quantity of mucilage.

The bulbous root of *Crinum Asiaticum* is official in the Pharmacopœia of India (*Crini Radix*, P. I.) as a substitute for squill. It has not been chemically investigated.

STROPHANTHIN ($C_{20}H_{34}O_{10}$; Arnaud, $C_{31}H_{48}O_{12}$).—According to Fraser, this is the active principle of strophanthus-seed (*Strophanthus hispidus*, var. *Kombé*), and is a glucoside. He obtained it in crystals. Acids convert it into glucose and crystalline *strophan-*

thidin. Phosphomolybdic acid produces in solutions of strophanthin a bright bluish-green color. Helbing states that its aqueous solution yields, with a trace of solution of ferric chloride and a little strong sulphuric acid, a reddish-brown precipitate which after an hour or two turns green. Sulphuric acid colors strophanthin dark green, changing to reddish brown. Recent researches indicate that strophanthin is only one of the active principles of the different species of *Strophanthus*. *Strophanthus* is official, *Strophanthus*, B. P. and U. S. P., yielding *Tinctura Strophanthi*, B. P.

QUESTIONS AND EXERCISES.

Define glucosides, and mention those of pharmaceutical interest.—Draw out an equation illustrative of the development of oil of bitter almonds.—How much pure amygdalin will yield 1 grain of real hydrocyanic acid?—To what does cherry-laurel water owe its activity?—Mention the active principle of senna.—By what process is the glucoside of the purple foxglove prepared?—State the circumstances under which guaiacum resin and jalap resin yield glucose.—Mention a test for guaiacum resin.—How may the adulteration of jalap resin by rosin be detected?—Enumerate the tests for salicin.—How is santonin officially prepared?—Name the sources of saponin.—What is the difference between scammony and resin of scammony?—How would you detect resins of turpentine, guaiacum, or jalap in resin of scammony?

BITTER OR TONIC SUBSTANCES, ETC.

The following articles, employed medicinally in such forms as decoction, extract, infusion, tincture, etc., contain active principles which have not yet been thoroughly examined. Some of these principles have been isolated, and a few have been obtained in the crystalline condition; but their constitution has not been sufficiently well made out to admit of the classification of the bodies either among alkaloids, glucosides, acids, or other well-marked principles:

<i>Andrographis Caules et Radix</i> , P. I., from <i>Andrographis paniculata</i> , Kariyat.	<i>Buchu folia</i> .
<i>Anthemidis flores</i> .	<i>Calendula officinalis</i> . Marigold. (Calendulin, Stoltze.)
<i>Apocynum</i> . Canadian hemp.	<i>Calotropis Cortex</i> , P. I., from <i>Calotropis procera</i> and <i>C. gigantea</i> . Mudar.
<i>Asclepias Tuberosa</i> . Pleurisy-root (Asclepedin).	<i>Canellæ Cortex</i> . (Cascarillin, $C_{12}H_{11}O_4$.)
<i>Aurantii cortex</i> . (Hesperidin.)	<i>Caulophyllum</i> <i>Thalictroides</i> . Blue cohosh. Alkaloid?
<i>Azadirachtæ Cortex et Folia</i> , P. I., from <i>Azadirachta Indica</i> , Nim or Margosa. (A resin, $C_{36}H_{50}O_{11}$, Broughton.)	<i>Cimicifuga</i> (<i>Actæa</i>) <i>racemosa</i> . (Cimicifugin; said by Conard to be neutral, and by Falek alkaloidal.) Black snake-root. (<i>Cimicifugæ Rhizoma</i> , B. P.)
<i>Bonducellæ Semina</i> , P. I., from <i>Cæsalpina</i> (<i>Guilandina</i>) <i>Bon- ducella</i> . Bonduc-seeds or <i>Nickar-nuts</i> .	

- Cypripedium pubescens* (Cypripedin?). Ladies' Slipper.
Euonymus atropurpureus. Wahoo-bark. (Euonymin?)
Euonymi Cortex, B. P., is "the dried root-bark," the source of *Extractum Euonymi Siccum*, B. P.
Eupatorium perfoliatum. Thoroughwort or Boneset.
Gaulanthea (Tinospora) Radix et Caulis, P. I.).
Gynocardia semina, from *Gynocardia odorata*. (*Chaulmugra*, P. I.)
Hamamelis Virginica. Witch-hazel.
 The official portions are *Hamamelidis Cortex*, B. P., the source of *Tinctura Hamamelidis*, B. P., and *Hamamelidis Folia*, B. P., the source of *Extractum Hamamelidis Liquidum*, B. P.
Hydrocotyles Folia, P. I., from *Hydrocotyle Asiatica*. Indian pennywort.
Iris versicolor. Blue flag. (Iridin or Irisin?)
Lactuca. (Lactucin, etc.) The milk-juice, dried, yields *Lactucarium*. U. S. P.
Lappa, U. S. P., *Arctium Lappa*, *Lappa officinalis*. Burdock.
Lupulus.
Magnolia. Swamp sassafras, or beaver tree.
Marrubium. Horehound. Marrubein, a crystalline bitter substance (Mein).
Matico Folia. Matico.
Melia Azedarach. (Resin, Jacobs.)
Pepo. The seed of *Cucurbita Pepo*. A remedy for tapeworm.
Phytolacca Fructus et Radix. Poke fruit and root. Phytolaccin, a crystalline substance (Claassen).
Scutellaria. Skullcap.
Serpentaria. Virginia Snakeroot.
Soymida Cortex, P. I. Rohunbark, from *Soymida febrifuga*.
Taraxaci Radix. (Taraxacin.)
Toddalia Radix, P. I.
Triticum repens. Rhizome of couch-grass.
Veronica Virginica, roots and rhizome. Culvers root; *Leptandra*, U. S. P. (Leptandrin?)
Viburnum prunifolium. Black haw. (*Viburnin*.)

ALKALOIDS.

Constitution of Alkaloids, or Organic Bases.


Natural Alkaloids.—The natural organic bases, alkaloids, or alkali-like bodies (*είδος*, *eidos*, likeness), have many analogies with ammonia. Their constitution as a class is not yet satisfactorily known, but some are possibly direct derivatives of a single molecule of ammonia (NH_3) or of double, triple, or quadruple molecules (N_2H_6 , N_3H_9 , N_4H_{12}); others of ammonias in which the ammoniacal structure is largely merged in or conditioned by a benzenoid or aromatic structure, or, *vice versâ*, in which the benzenoid character is conditioned by the ammoniacal; while others again certainly appear to be benzenoid, but of a more or less nitrogen-benzene (pyridinoid) rather than a completely carbon-benzene character—benzene in which CH''' is displaced by N''' (p. 513).

Numerous artificial organic bases, having a simple ammoniacal constitution, have already been formed. These are sometimes termed *amido*, *imido*, and *nitrile bases*, or *amines*, and are primary,

secondary, and tertiary according as one, two, or three atoms of hydrogen in ammonia have been displaced by radicals, as seen in the following general formulæ (R = any univalent radical):



or in the following examples:



The three classes have also been termed amidogen bases (NH_2), imidogen bases (NH), and nitrile bases (N).

Formation of Some of the Artificial Organic Bases.—A few illustrations will suffice: Just as the addition of hydrogen iodide (HI) to ammonia (that is, the common trihydrogen ammonia, NH_3) gives common ammonium iodide ($NHHHHI$ or NH_4I), so the addition of ethyl iodide (C_2H_5I or EtI) (see p. 402) to ammonia (NH_3) gives ethyl-ammonium iodide ($NHHHETI$, or NH_3EtI , or $NH_3C_2H_5I$). A fixed alkali turns out common ammonia ($NHHH$) from the iodide (or any other salt) of common ammonium; it turns out ethyl-ammonia ($NHHEt$) from the iodide (or any other salt) of ethyl-ammonium. Ethyl-ammonia (or *ethylia*, or ethylamine), $NHHEt$, with ethyl iodide, EtI , gives diethyl-ammonium iodide [$NHHEtEtI$, or NH_2Et_2I , or $NH_2(C_2H_5)_2I$]. From the latter potash turns out diethyl-ammonia ($NHEt_2$). Diethyl-ammonia (*diethylia* or diethylamine) with ethyl iodide gives triethyl-ammonium iodide ($NHEt_3I$). The latter with alkali gives triethyl-ammonia, or *triethylia*, or triethylamine (NEt_3), and this with ethyl-iodide gives tetrethyl ammonium iodide, NEt_4I , or $N(C_2H_5)_4I$.

What has just been stated respecting ethyl iodide is true of other ethyl salts; and what is true of ethyl salts is true of salts of an immense number of other radicals—univalent, bivalent, etc.; so that a vast number of artificial organic bases and their salts can be produced. The reactions are not always so sharp as those just given. Mixtures of primary, secondary, and tertiary compounds, rather than either alone, often occur in an experiment, but the reactions are typically true. Some of these artificial bases not only resemble natural alkaloids, but are strong caustic liquids, like solution of ammonia.

Then the displacing radical in an artificial alkaloid or its salt may not only be of one kind, as indicated in the preceding paragraphs, but of different kinds; and while the radical displacing one atom of hydrogen is keeping its place, any of the many known radicals may occupy the position of one or all of the other atoms of hydrogen. Thus, for example, we have methyl-ethyl-amylamine ($C_8H_{19}N$, or $NCH_3C_2H_5C_5H_{11}$, or $NMeEtAy$), a colorless, oily body of agreeable aromatic odor. The empirical formulæ of the vegeto-alkaloids mor-

phine, quinine, etc. may some day be similarly resolvable into rational formulæ, either simply ammoniacal, benzenoid, or pyridinoid. Their artificial production will then quickly follow. In a few cases this has already been accomplished.

Analogues of Amines.—From the analogy of phosphorus, arsenum, and antimony to nitrogen there exist, as might be expected, *phosphines*, *arsines*, and *stibines*, bases resembling *amines*, but containing the respective elements (P, As, Sb) in place of the nitrogen (N) of the amines.

Methylamine (CH_3HHN) and *trimethylamine* ($\text{CH}_3)_3\text{N}$ are artificial ammoniacal alkaloids. The former was found by Schmidt in *Mercurialis annua* and *M. perennis*, and previously by Reichardt, who termed it *mercurialine*. Trimethylamine is also produced in large quantities in the dry distillation of the evaporated residue of the spent wash produced in beet-root spirit distilleries.

Propylamine, or *tritylia* ($\text{C}_3\text{H}_7\text{HHN}$), is a volatile oil, one product of the destructive distillation of bones and other animal matters.

The organic bases derived from one molecule of ammonia are termed *monamines*; from two molecules, *diamines*; from three, *triamines*; and from four, *tetramines*:



In these amines any bivalent, trivalent, or quadrivalent radical may occupy the place of two, three, or four univalent radicals (R). The diethylene-diamine is used medicinally under the name *piperazine*; its constitution is that of piperidine (*vide* Index), in which NH displaces CH_2 .

Wurtz first obtained methylamine and ethylamine in 1849; Hofmann, in 1850 and subsequently, added enormously to our knowledge of the secondary, tertiary, and other amines and of the directly ammoniacal type of bodies generally. Kekulé linked on the aromatic or benzene type in 1865. Dewar and Körner almost simultaneously, in 1870, demonstrated the benzenoid character of pyridine and quinoline (see p. 514), while no one has since been more active in alkaloidal research than Ladenburg.


Vegetal Alkaloids.—These are of great importance to the medical and pharmaceutical student. They are treated in considerable detail in the succeeding pages.

Animal Alkaloids.—Many well-known alkaloids occur in the juice of the flesh and in other parts of animals. Ordinary extract of meat contains abundance of crystals of *creatine*, $\text{C}_4\text{H}_9\text{N}_3\text{O}_2$, and some *creatinine*, $\text{C}_4\text{H}_7\text{N}_3\text{O}$. Creatine easily parts with the elements of water and yields creatinine; it takes up the elements of water and yields *sarkosine*, $\text{C}_3\text{H}_7\text{NO}_2$, and *urea*, $\text{CH}_4\text{N}_2\text{O}$. Sarkosine is methylglycocoll. *Taurine*, $\text{C}_2\text{H}_7\text{NSO}_3$, may be obtained from bile, and it can be constructed artificially from its elements. Some animal tissues, as of the spleen, brain, and pancreas, yield, as a product of


work, *leucine*, $C_6H_{13}NO_2$, which occurs in white, pearly crystals; also *tyrosine*, $C_9H_{11}NO_3$. Gautier recently obtained several new alkaloids from albumenoids, and hence termed them *leucomaines* (λείκωμα, *leucoma*, white of egg)—namely, *xanthocreatinine*, $C_5H_{10}N_4O$, *crusocreatinine*, $C_5H_8N_4O$, *amphicreatinine*, $C_9H_{19}N_7O_4$, and *pseudo-xanthine*, $C_4H_5N_5O$. The leucomaines and the animal alkaloids generally are of great physiological interest. Some of the leucomaines are toxic and indistinguishable from ptomaines; in fact, the three classes merge into one another.

Ptomaines.—A series of diamines, many of them toxic, have been isolated by Brieger from decaying nitrogenous animal principles, including the putrid albumenoids or proteids of the human body itself—hence the name *ptomaines* (πτῶμα, *ptoma*, a corpse.) These have some medico-legal importance, but, inasmuch as they may occur in life, poisoning the blood during the progress of disease, especially disease associated with the development of micro-organisms or microbes—that is, zymotic disease (ζύμη, *zumē*, ferment)—they have great pathological interest; indeed, physiological interest also, for one of a curaroid character seems to play a part in the process of digestion. The names of some of these bases are—*neurine* ($C_5H_{13}NO$) and *neuridene* ($C_5H_{14}N_2$), from putrid flesh; *muscarine* ($C_5H_{13}NO_2$) and *gadinine* ($C_7H_{16}NO_2$), from putrid fish; *cadaverine* ($C_5H_{16}N_2$), *saprine*, and *putrescine*, ($C_4H_{12}N_2$), from putrid human remains, *choline* being met with in the earlier stages of decay; and *tetanine* ($C_{13}H_{30}N_2O_4$), the administration of which to animals produced symptoms resembling those of tetanus in man, from beef putrified by the agency of the microbe which is associated with the cause of traumatic tetanus so distressing to the human subject. *Tyrototoxicicon* was the name given by Vaughan to a toxic ptomaine he isolated from poisonous cheese (τυρός, *tyros*, cheese; τοξικόν, *toxicon*, poison), afterward from poisonous milk and cream, which, taken as food, had caused more or less vomiting, headache, and diarrhœa. He afterward recognized it as *diazobenzene* hydrate, $C_6H_5 \cdot N : N \cdot OH$. Brieger states that when shell-fish is poisonous it is due to the presence of a ptomaine he has named *mytiloxine*, $C_6H_{15}NO_2$. Para- and meta-phenylene-diamine appear to have all the characters of leucomaines or ptomaines, the latter causing intense influenza.

Evidence of Constitution of the Natural Alkaloids.—Attempts to form artificially the more important natural organic bases commonly used in medicine have hitherto failed. Many artificial colorific alkaloids of the amidobenzene (aniline or phenylamine) type, and of a curious double nitrogen type (azo- or, rather, diazo-type; see the non-colorific diazobenzene, as above), have been obtained. But the type of the natural medicinal alkaloids seems rather to be found in *pyridine*, C_5H_5N . Pyridine is producible in various ways, but is contained in *bone oil* (from the distillation of bones—whence, also, *pyrrol*, C_4H_5N , and thence *iodopyrrol*, or *iodol*, C_4I_4HN , a rival of iodoform), together with the homologues *picoline*, C_6H_7N (or methylpyridine, ortho-, meta-, or para-); *lutidine*, C_7H_9N ; and *collidine*, $C_8H_{11}N$, forming an homologous series of *pyridine bases*, $C_nH_{2n-5}N$.



From quinine, cinchonine, and strychnine, by the disruptive action of caustic alkalis, not only pyridine and homologues, but *quinoline* or *chinoline*, $\text{C}_9\text{H}_7\text{N}$, have been obtained; hence pyridine and quinoline would seem to contribute to the construction of those and similar alkaloids. Quinoline can be made in various other ways, especially (Skraup) from nitrobenzene, aniline, and glycerin. Quinoline is closely related both to benzene and to pyridine (see the following formula). Its relation to naphthalene (two carbon-conjoined benzene residues) is the relation of pyridine to benzene.



Both pyridine and quinoline form additive compounds with hydrogen. (See Piperidine in Index.)

By adding six atoms of hydrogen to pyridine, piperidine is obtained, and conine, the alkaloid of hemlock, is piperidine with propyl (C_3H_7) replacing one of the hydrogen atoms. It has been formed artificially by Landenburg from picoline. (See also Ecgonine, Tropine, etc.)

Chemists, in the hope, doubtless, of discovering how to produce the valuable medicinal alkaloids artificially, have obtained several alkaloidal derivatives of quinoline. One, *kairine*, somewhat resembles quinine.


Again, alkaloids yield organic acids, and organic acids—notably those occurring in the nicotine-yielding and morphine-yielding plants—may be converted into pyridine compounds when the constituents of their molecules are interwoven with those of ammonia.

A careful consideration of the above and allied facts irresistibly leads to the inference that we are at last almost “within measurable distance” of the artificial production of most of the natural alkaloids. This is a subject of financial and general commercial weight; of considerable technological, including pharmaceutical, importance; of very great medical consequence, especially taken in connection

with its ramifications; and of transcendent interest as illustrating the working of the forces of nature within the molecules of matter.

Vegeto-animal Alkaloids.—*Choline*, $C_5H_{15}NO_2$, occurs in the bile and the brain, also in ergot and ipecacuanha, hops, areca-nut, cotton-seed cake, *Scopola Japonica*, etc. *Guanine*, $C_5H_5N_5O$, and *Sarkine*, $C_5H_4N_4O$, are found in flesh and in young plane-leaves. Fresh meat furnishes *Carnine*, $C_7H_8N_4O_3$; and *Betaine*, $C_5H_{11}NO_2$, is found in beet-root, cotton-seed cake, and in urine.

Hydroxylamine.—Beside the amide, imide, and nitrile bases already mentioned, ammonia may have one atom of its hydrogen displaced by hydroxyl, hydroxylamine (NH_2OH) resulting. It is often formed when nascent hydrogen acts on an oxide of nitrogen, as when zinc, diluted sulphuric acid, and a little nitric acid are brought together. It yields substitution-products, as ethylhydroxylamine (NHC_2H_5OH), and additive compounds, as hydroxylamine hydrochlorate (NH_2OH, HCl):



Hydroxylamine and aldehydes yield *aldoximes*. Hydroxylamine and acetones yield *acetoximes*. (*Vide* manuals not limited to the requirements of medical and pharmaceutical students.)

Hydrazine, H_2N-NH_2 .—Diethylamine, by action of nitrous acid, yields a nitroso-derivative which, on reduction, furnishes what apparently is a diamidic compound—diethylhydrazine ($C_2H_5)_2N-NH_2$. Hydrous hydrazine has the formula H_2N-NH_2, H_2O . Hydrazine itself cannot very easily be isolated. Its salts with ordinary acids are generally crystalline and isomorphous with corresponding ammonium salts. Acidified, they have very powerful reducing properties, and act as strong poisons toward the lower organisms.

Azoimide or *Imidazoic Acid*, HN_3 , is a body closely resembling the haloid acids, and was originally prepared by Curtius from hydrazine and ethyl hippurate; it may, however, be prepared more easily by a method proposed by Wislicenus, in which sodamide, $NaNH_2$, prepared by passing NH_3 over melted sodium, is heated with nitrous oxide.

Note on Nomenclature of Natural Alkaloids.—The first syllables of the names of the natural alkaloids recall the name of the plant whence they were obtained or some characteristic property. It is to be regretted that the last syllable is not either *ine* or *ia*, instead of sometimes one and sometimes the other. The termination in *ia* distinguishes the alkaloids from some other substances the names of which end in *ine*, as aniline, bromine, chlorine, fluorine, iodine, morphine, etc., but traders generally and the compilers of the American, British, French, and German Pharmacopœias adopt the termination in *ine*. The names of the salts of the alkaloids are given on the assumption that the acid unites with the alkaloid without decomposition. Thus hydrochlorate (sometimes termed “hydro-

chloride") of morphine is regarded as morphine with added hydrochloric acid; as we might assume sal-ammoniac to be ammonia (NH_3) with hydrochloric acid (HCl), and name it hydrochlorate of ammonia (NH_3HCl) instead of ammonium chloride (NH_4Cl). All acids, even sulphydric, unite with alkaloids and form additive salts having similar names.

Antidotes.—In cases of poisoning by alkaloids, emetics and the stomach-pump must be relied on rather than chemical agents. But astringent liquids may be administered, for tannic acid precipitates many of the alkaloids from their aqueous solution, absorption of the poison being thus possibly retarded.

MORPHINE AND OTHER OPIUM ALKALOIDS.

Formula of Morphine, $\text{C}_{17}\text{H}_{19}\text{NO}_3, \text{H}_2\text{O}$. Molecular weight, 303.

Occurrence.—Morphine, or morphia, occurs in opium (the inspissated juice of the fruit, termed the capsule, of the white poppy, *Papaver somniferum*) as morphine meconate [$(\text{C}_{17}\text{H}_{19}\text{NO}_3)_2, \text{C}_7\text{H}_4\text{O}_7, 5\text{H}_2\text{O}$, Dott] and sulphate. The dried poppy-capsule of pharmacy (*Papaveris Capsulae*, B. P.) contains opium principles, but varying much in nature and proportion: the presence of morphine, narcotine, and meconic acid has been demonstrated, and, by Groves, of codeine and narceine. Ordinary moist opium (*Opium*, U. S. P.) should contain not "less than 9 per cent. of crystalline morphine," and, when dried at 85°C . and powdered (*Opii Pulvis*, U. S. P.), not less than 13 nor more than 15 per cent. of crystallized morphine.

Deodorized Opium (*Opium Deodoratum*, U. S. P.), the old *denarcotized opium*, is dried and powdered opium from which narcotine has been washed out by fourteen times its weight of ether, the product being redried at 85°C . and made up to its original weight with powdered sugar of milk.

Morphina, U. S. P., may be made by adding to infusion of opium an equal bulk of alcohol, then slight excess of ammonia, and setting aside for crystalline morphine to separate. It is purified by recrystallization in colorless, shining, prismatic crystals.

Process for Hydrochlorate.—The hydrochlorate, $\text{C}_{17}\text{H}_{19}\text{NO}_3, \text{HCl}, 3\text{H}_2\text{O}$ (*Morphinæ Hydrochloras*, U. S. P.), occurs in slender white acicular crystals; it is prepared by simply decomposing an aqueous infusion of opium with calcium chloride, calcium meconate and morphine hydrochlorate being produced. (If the infusion, which is always acid, be first nearly neutralized by the cautious addition of small quantities of a very dilute solution of ammonia, the calcium chloride then at once causes a precipitate of calcium meconate, which can be filtered off, leaving a colored solution of morphine hydrochlorate. On the large scale—*vide* B. P.—the details are somewhat different.) The salt is partially purified by crystallization from the evap-

orated liquid, then by treatment of the solution of the impure hydrochlorate by animal charcoal, and lastly by precipitation of the morphine from the still colored liquid by ammonia and re-solution of the morphine in hot dilute hydrochloric acid; morphine hydrochlorate separates out on cooling.

Morphine hydrochlorate deposited from a hot solution in about twenty times its weight of alcohol is anhydrous.

Morphine may also, of course, be prepared by the methods given for the assay of opium. (See Index.)

Process for Acetate.—Morphine acetate ($C_{17}H_{19}NO_3, C_2H_4O_2, 3H_2O$) (*Morphinæ Acetas*, U. S. P.), a white pulverulent salt, is prepared by dissolving morphine in acetic acid, the morphine being obtained from a solution of the hydrochlorate by precipitation with ammonia or direct from opium. 15 parts of hydrochlorate yield, theoretically, 16 of acetate. 1 grain of acetate in 10 minims of water forms the *Injectio Morphinæ Hypodermica*, B. P.

Both morphine hydrochlorate and acetate are soluble in water, but the solution is not stable unless acidulated and containing alcohol; hence the official solutions, 1 per cent. (*Liquor Morphinæ Hydrochloratis*, B. P., and *Liquor Morphinæ Acetatis*, B. P.), consist of 3 parts of water and 1 part rectified spirit, a few minims per ounce of hydrochloric or acetic acid being added. Even solid morphine acetate is unstable, slowly dissociating into acetic acid and morphine; hence the acid odor of morphine acetate; hence, too, the necessity, when a solution of morphine acetate of perfectly definite strength is required, of preparing it from a weighed quantity of hydrochlorate or of pure crystalline morphine. Other preparations official in the British Pharmacopœia are—*Suppositoria Morphinæ*, *Trochisci Morphinæ*, *Trochisci Morphinæ et Ipecacuanhæ*, and *Liquor Morphinæ Bimeconatis*. The latter is made by dissolving morphine in a diluted spirituous solution of meconic acid. It contains about $1\frac{1}{2}$ per cent. of the so-called morphine bimeconate ($C_{17}H_{19}NO_3, C_7H_4O_7?$), and, so far, resembles tincture of opium.

Process for Sulphate.—Morphine sulphate ($C_{17}H_{19}NO_3)_2, H_2SO_4, 5H_2O$) (*Morphinæ Sulphas*, U. S. P.) is prepared by neutralizing precipitated morphine with diluted sulphuric acid. It occurs in white silky crystals, not very soluble in water. A 1 per cent. solution in weak spirit forms the *Liquor Morphinæ Sulphatis*, B. P. It is a constituent of *Pulvis Morphinæ Compositus*, U. S. P.

Solubility of Morphine Salts in Water at 60° F.—According to Dott, 1 part of the respective salts is soluble in the annexed numbers of parts of water: acetate, $2\frac{1}{2}$; tartrate, $9\frac{1}{4}$; sulphate, 23; hydrochlorate, 24; meconate, 34.

Morphine Tartrate has the formula ($C_{17}H_{19}NO_3)_2, C_4H_6O_6, 3H_2O$.

CODINE, or *Codeia* ($C_{18}H_{21}NO_3, H_2O$), is another officially recognized alkaloid of opium (*Codeina*, U. S. P.). It is soluble in the slight excess of ammonia employed in the foregoing process for the separation of morphine. It is obtained by evaporating the ammoniacal liquors, "treating the residue with water, precipitating with caustic potash, and purifying the precipitated alkaloid by recrystal-

lization from ether. It occurs in colorless or nearly colorless octahedral crystals; soluble in 80 parts of water and of solution of ammonia, readily soluble in spirit and in diluted acids. The aqueous solution has a bitter taste and an alkaline reaction. The alkaloid dissolves in sulphuric acid, forming a colorless solution, which, when gently warmed with ammonium molybdate or a trace of ferric chloride, assumes a deep-blue color. Ignited in air, it yields no ash."—*B. P.* The U. S. P. also requires codeine to be neutral to litmus-paper, and it should dissolve in nitric acid of sp. gr. 1.200 to a yellow liquid which should not become red (difference from and absence of morphine). It reduces a solution of 1 part of ammonium selenite in 20 of strong sulphuric acid, yielding a green color (Lafon). (See also p. 520.)

Other alkaloids exist in opium. In the above process for morphine a considerable quantity of an alkaloid of very weak basic properties, *narcotine*, ($C_{22}H_{23}NO_7$) or $C_{19}H_{14}(CH_3)_3NO_7$ (*Narcotina*, P. I.), remains in the exhausted opium, and may be extracted by digesting in acetic acid, filtering, and precipitating by ammonia. It crystallizes in brilliant needles from alcohol or ether. The formula of its hydrochlorate is $C_{22}H_{23}NO_7 \cdot HCl \cdot H_2O$. By oxidation it yields *cotarnine* and an acid termed *opianic*. From the mother-liquors there have also been obtained *thebaine* ($C_{19}H_{21}NO_3$), *papaverine* ($C_{21}H_{21}NO_4$, Hesse; $C_{20}H_{21}NO_4$, Merck), *opianine* ($C_{21}H_{21}NO_7$?), *narceine* ($C_{23}H_{29}NO_9$), *cryptopine* ($C_{21}H_{23}NO_5$), *meconin* ($C_{10}H_{10}O_4$), *meconoisin* ($C_8H_{10}O_2$), *laudanine* ($C_{20}H_{25}NO_4$), *codamine* ($C_{20}H_{25}NO_4$), *gnoscopine* ($C_{34}H_{36}N_2O_{11}$), *pseudomorphine* ($C_{17}H_{18}NO_3$), *protopine* ($C_{20}H_{19}NO_5$), *laudanoline* ($C_{21}H_{27}NO_4$), *hydrocotarnine* ($C_{20}H_{19}NO_5$), *rhæadine* ($C_{23}H_{21}NO_6$), *meconidine* ($C_{21}H_{23}NO_4$), *lanthopine* ($C_{23}H_{25}NO_4$).

A little acetic acid also exists in all opium (D. Brown).

ANALYTICAL REACTIONS.

First Analytical Reaction.—To a minute fragment of a morphine salt add 1 drop of water, and warm the mixture until the salt dissolves; then stir the liquid with a glass rod moistened by a strong *neutral* solution of ferric chloride; a dirty-blue color is produced.

Even in dilute solutions morphine reduces potassium ferri-cyanide to ferrocyanide, hence may be detected by the blue precipitate (prussian blue) produced on the addition of ferric chloride and ferri-cyanide. Other substances, but no other official alkaloids, give this reaction.

Second Analytical Reaction.—To a drop or two of a strong solution of a morphine salt in a test-tube add a minute fragment of iodic acid (HIO_3 , p. 296); iodine is set free. Into the upper part of the tube insert a glass rod covered with starch mucilage, and warm the solution; dark-blue "starch iodide" is produced. If the mixture of morphine and iodic acid be

shaken up with chloroform or carbon disulphide, a violet solution is obtained. This reaction is only confirmatory of others, as albuminous matters also reduce iodic acid.

Third Analytical Reaction.—To a few drops of an aqueous infusion of opium add a drop of *neutral* solution of ferric chloride; a red solution of the ferric meconate is produced. Add solution of corrosive sublimate; the color is not destroyed (as it is in the case of ferric sulphocyanate, a salt of similar tint). In cases of poisoning by a preparation of opium this test is almost as conclusive as a direct reaction of morphine (the poison itself), meconic acid being obtainable from opium only.

Other Reactions.—Add sodium carbonate to a solution of a salt of morphine; a white precipitate of morphine falls, slowly and of a crystalline character if the solution is dilute. Collect this precipitate and moisten it with neutral solution of ferric chloride; the bluish tint above referred to is produced.—Add an alkali to a solution of hydrochlorate or acetate of the alkaloid; morphine is precipitated, soluble in excess of the fixed alkali, far less readily so in ammonia.—Moisten a particle of a morphine salt with nitric acid; an orange-red coloration is produced. Warm a little morphine with strong sulphuric acid and sodium arsenate; blue-green tinges result.—To morphine add strong sulphuric acid, mix, and strew bismuth nitrate on the fluid; the mixture turns dark brown or black.—Heat morphine on platinum-foil; it burns entirely away.

APOMORPHINE ($C_{17}H_{17}NO_2$).

Apomorphine (*ἀπόδ*, *apo*, from, and *morphine*) is an alkaloid obtained from morphine by Matthiessen and Wright. It possesses remarkable physiological effects: one-tenth of a grain (in aqueous solution) injected under the skin, or a quarter of a grain taken into the stomach, produces vomiting in from four to ten minutes.

Process.—Morphine hydrochlorate is hermetically sealed in a thick tube with considerable excess of hydrochloric acid, and heated to nearly 300° F. for two or three hours. The product is purified by diluting the contents of the tube with water, precipitating with sodium bicarbonate, and treating the precipitate with ether or chloroform.—On shaking up the ethereal or chloroform solution with a very small quantity of strong hydrochloric acid, the sides of the vessel become covered with crystals of the hydrochlorate of the new base. These may be drained from the mother-liquor, washed with a little cold water, in which the salt is sparingly soluble, recrystallized from hot water, and dried on bibulous paper or over sulphuric acid. The formula ($C_{17}H_{17}NO_2HCl$) indicates that the new alkaloid is derived from morphine by abstraction of the elements of water.

Apomorphine Hydrochlorate (*Apomorphinæ Hydrochloras*, U. S.

P.) occurs in colorless or grayish-white shining crystals, turning greenish on exposure to light and air, odorless, having a bitter taste and a neutral or faintly acid reaction. Soluble in 45 parts of water and in 45 parts of alcohol at 15° C. (59° F.). "When heated to near 100° C. (212° F.) the salt is decomposed, rapidly if in solution, slowly when dry. At 270° C. (518° F.) it fuses to a black mass, and when ignited it is consumed without leaving a residue. Nitric acid colors the crystals blood-red to orange; sulphuric acid, violet to light-brown; dark-purple by a mixture of the two acids. A few crystals of manganese dioxide will color a saturated solution of apomorphine hydrochlorate green, which will be turned by a crystal of oxalic acid to a reddish-brown. "Addition of sodium-bicarbonate solution to the aqueous solution throws down the white amorphous alkaloid, which soon turns green on exposure to air, and imparts a violet or blue color to chloroform, in which it is very soluble (difference from morphine). If the salt impart at once an emerald-green color to 100 parts of water on being shaken with it a few times in a test-tube, it should be rejected."—*U. S. P.* 2 grains of the hydrochlorate dissolved in 100 minims of camphor-water constitute the *Injectio Apomorphinæ Hypodermica*, B. P.


Codeine also, according to the same chemists, yields apomorphine by similar treatment, a reaction that would seem to indicate that codeine is methyl-morphine; indeed, Grimaux (Hesse also) has since obtained codeine—or, possibly, an isomer of codeine, methyl-morphine—from morphine. Codeine may also be obtained by heating a sodium compound of morphine, $C_{17}H_{18}NaNO_3$, with methyl iodide, CH_3I ; sodium iodide and methyl-morphine or codeine result.

Codeine gives neither a blue color with ferric chloride nor a red with nitric acid. Both codeine and morphine when heated with a mixture of strong sulphuric acid and sodium arsenate give a blue color, the morphine yielding a greenish blue and the codeine a violet blue.

Constitution of the Opium Alkaloids.—The opium alkaloids, like the cinchona alkaloids, have been attacked by many highly-skilled chemists in the hope that analytical—or, in a sense, destructive—investigation would lead to synthetical or constructive knowledge, and many interesting and promising results have been obtained. But the subject is not yet sufficiently advanced for presentation before the younger students of medicine or pharmacy.

QUESTIONS AND EXERCISES.

Write some general formulæ of artificial alkaloids.—Name the substances represented by the following formulæ:



What is the assumed constitution of the salts of the alkaloids?—Describe the treatment in cases of poisoning by alkaloids.—Give the official process for the preparation of morphine hydrochlorate. In what form

does morphine occur in opium?—How is morphine acetate prepared?—What plan is adopted for preventing the decomposition of the official solutions of morphine?—Mention the analytical reactions of morphine.—In addition to the reactions of morphine, what test may be employed in searching for opium in a liquid or semi-fluid material?—How is apomorphine prepared? and what are its properties?—Describe the relation of morphine to codeine.

QUININE AND OTHER CINCHONA ALKALOIDS.

Formula of Quinine, $C_{20}H_{24}N_2O_2 \cdot 3H_2O$. Molecular weight, 378.

Source.—Quinine (*Quinina*, U. S. P.) and other alkaloids exist in the varieties of cinchona-bark as *kinates*, or, rather, *quinates*.* In the British Pharmacopœia cinchona-bark is defined as “the dried bark of *Cinchona Calisaya*, *Weddell*; *Cinchona officinalis*, *Linn.*; *Cinchona succirubra*, *Pavon*; *Cinchona lancifolia*, *Mutis*; and other species of *Cinchona* from which the peculiar alkaloids of the bark may be obtained.” The official galenical preparations are made with the succirubra or red cinchona-bark (*Cinchona Rubra*, U. S. P.). Quinine, quinidine, and cinchonine also occur, together with cupreine, in the bark of various species of *Remijia*, known as cuprea-bark.

Under *Cinchona* the United States Pharmacopœia recognizes “the bark of any species of *Cinchona* containing at least 5 per cent. of its peculiar alkaloids.”

Extraction of the Mixed Alkaloids.—Mix about 2 ounces of powdered bark with a quarter of its weight of lime and a little water, and extract with benzoated amylic alcohol. (For a description of this operation see “Quinine, Quantitative Estimation of,” in the Index.) Shake the product in a separating-funnel, with an ounce of water acidulated with sulphuric or hydrochloric acid. Draw off the aqueous liquid, which will contain the alkaloids as acid salts, and add to it a slight excess of ammonia. Collect the precipitated alkaloids on a filter, wash, and dry in the air or over a dish of sulphuric acid covered by a bell-glass. (For the separation of alkaloids see Index, “De Vrij’s Process”—an operation which should not be attempted at this stage of study.)

Process for Quinine Sulphate.—Quinine sulphate (*Quininæ Sulphas*, U. S. P.) may be prepared by treating the yellow bark with diluted hydrochloric acid, precipitating the resulting solution of quinine hydrochlorate by soda, and redissolving the precipitated quinine in the proper proportion of hot diluted sulphuric acid; or

* *Quinic acid*, $C_7H_{12}O_6$, occurs in cinchona, coffee, holly, ivy, oak, elm, etc. Heated, it yields *hydroquinone*, $C_6H_4(OH)_2$. Oxidized, it gives quinone, $C_6H_4O_2$, which is probably a di-ketone, $C_4H_4(CO)_2$, dicarbonyl benzene, or $C_2H_2 < \begin{smallmatrix} CO \\ CO \end{smallmatrix} > C_2H_2$. The homologues of benzene yield other “quinones.”

by extracting with spirit, etc. after the addition of lime. (See the sections on the quantitative analysis of cinchona-bark.) This, the common commercial sulphate, crystallizes out on cooling in silky acicular crystals, one molecule containing two atoms of quinine ($2C_{20}H_{24}N_2O_2$), one of sulphuric acid (H_2SO_4), and eight of water of crystallization ($8H_2O$).

In the process of an old United States Pharmacopœia (1870) lime was used instead of soda, the precipitated quinine dissolved in boiling alcohol, the latter recovered by distillation, the residual quinine neutralized by diluted sulphuric acid, the solution treated with animal charcoal, filtered while hot, set aside to crystallize, and re-crystallized if necessary.

Quinine sulphate, the *common* or so-called *disulphate*—($C_{20}H_{24}N_2O_2$)₂. H_2SO_4 . $8H_2O$)—is only slightly soluble in water; on the addition of diluted sulphuric acid the so-called *neutral sulphate* or *soluble sulphate* (*Quininæ Bisulphas*, U. S. P.) ($C_{20}H_{24}N_2O_2$. H_2SO_4 . $7H_2O$) is formed, which is freely soluble. The latter salt may be obtained in large rectangular prisms.* An *acid sulphate* ($C_{20}H_{24}N_2O_2$. $2H_2SO_4$. $7H_2O$) also exists. The *Infusum Cinchonæ Acidum*, B. P., contains the soluble sulphate.

The ordinary quinine disulphate is more soluble in alcohol or alcoholic liquids than in water. An ammoniated tincture (*Tinctura Quininæ Ammoniata*, B. P.) is made by dissolving the sulphate in proof spirit and adding a large excess of solution of ammonia. This tincture contains quinine itself liberated from combination by, and dissolved by aid of, the excess of the ammonia. Quinine wine (*Vinum Quininæ*, B. P.) is a solution of neutral quinine sulphate and citrate in orange wine, made by dissolving the disulphate (1 grain in the ounce) in orange wine by the help of citric acid. The remaining pharmacopœial preparations of quinine are the hydrochlorate and the mixed iron, ammonium, and quinine citrates (*Ferri et Quininæ Citras*, U. S. P.), the well-known scale compound. The latter is made by dissolving ferric hydrate, prepared from ferric sulphate, and quinine, prepared from the sulphate, in solution of citric acid; the liquid, evaporated to a syrupy consistence and dried in thin layers on glass plates, yields the usual greenish-yellow scales. (*Vide* p. 155.)

Quinine Hydrochlorate (*Quininæ Hydrochloras*, U. S. P.) is prepared by neutralizing quinine by hydrochloric acid. Its formula is $C_{20}H_{24}N_2O_2$. HCl . $2H_2O$. It is soluble in about 34 parts of water at common temperatures, the sulphate requiring 700 or 800. The two salts resemble each other in appearance, but the crystals of the

* We do not know whether or not these sulphates are ordinary sulphates, the hydrogen of the acid going over to the quinine molecule, nor whether or not the quinine molecule is univalent or bivalent; hence we cannot say whether the *common* sulphate or the *soluble* sulphate is, in constitution, the neutral sulphate. In the above paragraph the names *disulphate*, *neutral sulphate*, *acid sulphate* indicate nothing more than that the first sulphate contains in one molecule two atoms (chemical atoms) of quinine to one of sulphuric acid, the second one of each, and the third two of acid to one of quinine.

hydrochlorate are commonly somewhat larger than those of the sulphate. 8 grains of hydrochlorate of quinine dissolved in 1 ounce of tincture of orange-peel forms the *Tinctura Quininae*, B. P.

Quinine Hydrobromas, or *Quinine Hydrobromate*, U. S. P., has the formula $C_{20}H_{24}N_2O_2 \cdot HBr, 2H_2O$.

Basic Quinine Citrate has the formula $(C_{20}H_{24}N_2O_2)_2 \cdot H_3C_6H_5O_7, 5H_2O$. Other citrates contain three molecules of quinine to two of citric acid, and one of quinine to one of citric acid.

Quinine Valerianas, or *Quinine Valerianate*, U. S. P., may be made by dissolving precipitated quinine in warm aqueous solution of valerianic acid and setting aside to crystallize. Its formula is $C_{20}H_{24}N_2O_2 \cdot C_5H_{10}O_2, H_2O$.

REACTIONS.

First Analytical Reaction.—To a solution of quinine or its salts in acidulated water add fresh chlorine-water, shake, and then add solution of ammonia; a green coloration (*thalleioquin* or *dalleiochin*) is produced. Bromine-water or bromine-vapor may be used instead of chlorine.

Second Analytical Reaction.—Repeat the foregoing reaction, but precede the addition of solution of ammonia by that of solution of potassium ferrocyanide; an evanescent red coloration is produced (Livonius and Vogel).

Third Analytical Reaction.—To an aqueous solution of a soluble salt of quinine add solution of ammonium oxalate; a white crystalline precipitate of quinine oxalate falls. It is soluble in acids. If the solution to be tested be made from ordinary quinine sulphate, excess of the latter should be added to water very faintly acidulated with sulphuric acid, and the undissolved crystals removed by filtration.

Fourth Analytical Reaction.—A saturated aqueous solution of any neutral salt of quinine is made by dissolving so much of the salt in hot water as that some shall separate when the mixture has cooled to about 60° F. After standing for some time, filter. To the filtrate water-washed ether is added until a distinct layer of ether remains undissolved, and then ammonia in slight excess. After agitation and rest for fifteen minutes all precipitated quinine will have redissolved.

Note.—In the case of quinidine salts well-defined crystals appear at the junction of the aqueous and ethereal layers, especially after standing. In the case of cinchonidine salts a thick layer of small crystals appears at once. In the case of cinchonine salts the undissolved alkaloid makes the ethereal layer nearly solid. In testing quinine for other alkaloids evaporate the aqueous solution to one-fifth.

Fifth Analytical Reaction.—*Formation of Quinine Iodo-sul-*

phate.—Dissolve quinine sulphate in weak spirit of wine slightly acidulated with sulphuric acid, and add an alcoholic solution of iodine; a black precipitate forms. Allow the precipitate to settle, pour away the fluid, wash once or twice with cold alcohol, and then boil with alcohol; on cooling minute crystals separate, having the optical properties of the mineral tourmaline. This iodo-sulphate is sometimes termed herapathite, from the name of one of the chemists who discovered it (in 1852). Under the name of iodide of hydriodate of quinine Bouchardat described and used it in 1845. It is so slightly soluble in alcohol that by its means quinine can be fairly well separated from its admixture with the other cinchona alkaloids. According to Jörgensen, it has the formula $4C_{20}H_{24}N_2O_2, 3H_2SO_4, 2HI, I_4, xH_2O$.

Sixth Analytical Reaction.—Prepare a saturated solution of ordinary quinine sulphate in water at about 60° F., and add to five volumes of that solution seven volumes of solution of ammonia (sp. gr. 0.96). The alkaloid which at first precipitates redissolves upon slight agitation if the quinine sulphate is free from anything but traces of other cinchona alkaloids. If, however, more than traces of quinidine, cinchonidine, and cinchonine salts be present, a permanent precipitate remains. This is Kerner's method of testing quinine sulphate for other cinchona alkaloids. It turns upon the fact that the solubility of the sulphates of the cinchona alkaloids in water is in the opposite order to the solubility of the alkaloids themselves in solution of ammonia.

Other Characters.—Concentrated sulphuric acid dissolves quinine with production of only a faint yellow color, which is not increased by warmth.—Quinine and its salts, heated on platinum-foil, burn entirely away.—Most salts of quinine when in solution have a beautiful blue fluorescence. They twist the ray of polarized light to the left.—Quinine is soluble in alcohol, ether, benzol, and chloroform. Ordinary quinine sulphate is insoluble in chloroform, and but slightly soluble in water. Its solubility in chloroform is increased by the presence in solution of quinidine and cinchonine sulphates (Prescott), and its solubility in water is decreased by the presence in solution of ammonium sulphate (Carles). The slight solubility of its sulphate and iodo-sulphate in water distinguishes quinine from the other cinchona alkaloids, including the "amorphous alkaloid," or "quinoidine."

Quinidine ($C_{20}H_{24}N_2O_2$, the *conquinine* or *conchinine* of Hesse) is an isomer of quinine. Its salts are fluorescent, and give thalleioquin

with chlorine- or bromine-water and ammonia. They twist the ray of polarized light to the right. Quinidine is insoluble in water and sparingly soluble in ether (see Quinine, Fourth Analytical Reaction). It is soluble in alcohol, benzol, and chloroform. It is less soluble than quinine in ammonia, five volumes of a saturated aqueous solution of its ordinary sulphate requiring sixty to eighty volumes of ammonia solution (sp. gr. 0.96). *Quinidine Sulphate* (*Quinidinæ Sulphas*, U. S. P. $2(C_{20}H_{24}N_2O_2), H_2SO_4, 2H_2O$) is more soluble in water and chloroform than quinine sulphate. Quinidine tartrate is soluble in water. The hydriodate is insoluble in water and weak spirit, and occurs as sandy crystals. The hydriodates of the other cinchona alkaloids, though more soluble than quinidine hydriodate, are sometimes precipitated from neutral concentrated solutions as amorphous or semi-liquid precipitates. These, however, are soluble in weak spirit.

"If a small quantity of ammonia-water be added to 3 cc. of an aqueous solution of the salt saturated at 15° C. (59° F.) a white precipitate (quinidine) will be produced, which requires more than 30 cc. of ammonia or more than thirty times its weight of ether to dissolve it (absence of more than small proportions of other cinchona alkaloids)." — U. S. P.

Cinchonidine ($C_{20}H_{24}N_2O$).—The sulphate (*Cinchonidinæ Sulphas* ($C_{20}H_{24}N_2O$) $_2, H_2SO_4, 3H_2O$, U. S. P.) is official. It may be obtained from the mother-liquids of the crystallization of quinine sulphate. When perfectly pure, salts of cinchonidine do not give thalleioquin and are not fluorescent. Even good commercial salts, however, nearly always give both reactions. Salts of cinchonidine twist the polarized ray to the left. Cinchonidine is insoluble in water and nearly so in ether. (See Quinine, Fourth Analytical Reaction.) It is soluble in alcohol, benzol, and chloroform. It is less soluble in ammonia solution than quinine, five volumes of a saturated aqueous solution of cinchonidine sulphate requiring about eighty volumes of ammonia solution (sp. gr. 0.96). It is true cinchonidine is dissolved as readily as quinine if excess of strong ammonia is quickly mixed with the solution of the salt of cinchonidine, but from such a solution cinchonidine soon crystallizes out, while quinine remains dissolved for many hours. Cinchonidine sulphate and hydriodate are soluble in water, but the sulphate, like quinine sulphate, is insoluble in chloroform. Cinchonidine tartrate is insoluble in water, and in this form cinchonidine is usually separated from neutral solutions containing the other cinchona alkaloids except quinine, the filtrate from the precipitate of tartrate yielding cinchonine on the addition of ammonia.

Cinchonine ($C_{20}H_{24}N_2O$) (*Cinchonina*, U. S. P.) is an isomer of cinchonidine. When quite pure its salts are not fluorescent and do not give thalleioquin. As in the case of cinchonidine, even good commercial specimens of cinchonia salts nearly always give both reactions. Cinchonine salts twist the polarized ray to the right. Cinchonine is insoluble in water and nearly so in ether. (See Quinine, Fourth Analytical Reaction.) It is soluble in chloroform, benzol, and alcohol. Chloroform containing one-fourth of

its weight of 95 per cent. alcohol dissolves cinchonine much more readily than either alcohol or chloroform alone. Cinchonine is insoluble in ammonia solution.

The sulphate (*Cinchonine Sulphas*, B. P.) is official. "It may be obtained from the mother-liquors of the crystallization of the quinine, cinchonidine, and quinidine sulphates by precipitating the alkaloid with caustic soda, washing it with spirit until free from other alkaloids, dissolving in sulphuric acid, and, after purifying the solution with animal charcoal, allowing to crystallize."

Cinchonine Sulphate (*Cinchonine Sulphas*, U. S. P.) ($C_{20}H_{24}N_2O$) $\cdot$ $H_2SO_4 \cdot 2H_2O$, tartate and hydriodate of cinchonine, are soluble in water, and the sulphate, like quinidine sulphate, is soluble in chloroform. In mixtures of cinchona alkaloids this alkaloid is precipitated by alkali after the others have been successively removed by ether, sodium tartrate, and potassium iodide.

Constitution of the Cinchona Alkaloids.—This is not yet clear, though great advances have been made. In the course of the investigations derivatives of quinoline more or less resembling quinine have been obtained—namely, *kairine*, *kairoline*, and *thalline*; *antipyrene* also. (*Acetanilide*, or "antifebrin," has, too, been found to possess greater antipyretic powers than the derivatives just mentioned.) See also p. 513.

"*Quinoidine*," "*Chinoidine*," or the "*Amorphous Alkaloid*."—Cinchona-barks generally contain some alkaloid isomeric with quinine which, like quinine, is soluble in ether, but the ordinary sulphate and iodo-sulphate of which are not crystalline and are soluble. These salts are semi-solid resinous-looking substances. The iodo-sulphate is used in De Vrij's method for the separation of mixed alkaloids. Quinoidine is usually obtained along with quinine, etc. from the mixed alkaloids by ether, and remains in the mother-liquor, from which it is precipitated by an alkali.

Cinchovatine occurs in a particular variety of cinchona-bark. *Quinicine* and *cinchonine* are alkaloids produced by the action of heat on quinine or quinidine and on cinchonidine respectively. They, also, are isomers—Hesse says polymers—of the parent alkaloids. Both yield ordinary salts. *Quiniretin* is the name given to the brown or reddish-brown indifferent substance into which quinine in aqueous solution is converted when exposed to much light.

Quinamine ($C_{20}H_{26}N_2O_2$) is a fifth cinchona alkaloid obtained by Hesse in 1872 from the bark of *Cinchona Succirubra*. Its solution is not fluorescent, and does not give thalleioquin. The same chemist announces the presence in cinchona of a sixth alkaloid, *cinchamidine* ($C_{20}H_{26}N_2O$).

Cupreine ($C_{19}H_{22}N_2O_2$) is an alkaloid discovered simultaneously by Howard and Hodgkin, by Paul and Cownley, and by Whiffen, in the bark of a *Remijia* (allied to *Cinchona*) and termed *cuprea*-bark. It closely resembles quinine, but is sparingly soluble in ether. It may be converted into quinine by heating its sodium compound with methyl chloride; whence it appears that quinine is the methyl ether of cupreine. The alkaloid, at first termed *homoquinine* or *ultraquinine*, seems to have been a mixture of cupreine and quinine.

Hydroquinine, $C_{20}H_{26}N_2O_2$, containing two more atoms of hydrogen than are present in the quinine molecule, is an alkaloid associated with quinine, in minute amount, in cinchona-bark. It remains in the mother-liquor when quinine sulphate is crystallized from an acid solution. Its therapeutic action is similar to that of quinine. Its characters are closely allied to those of quinine. It was discovered by Hesse.

STRYCHNINE.

Formula, $C_{21}H_{22}N_2O_2$. Molecular weight, 334.

Source.—Strychnine or strychnia exists, to the extent of 0.2 to 0.5 per cent., in the seed of *Strychnos Nux-vomica* (*Nux Vomica*, U. S. P.), also (Shenstone) in minute quantity in the bark of the nux-vomica tree (false Angustura-bark), and to 1.0 or 1.5 per cent. in St. Ignatius's bean (*Strychnos Ignatia*), chiefly in combination with strychnic or igasuric acid, or, after slight fermentation when moistened, with lactic acid. Crow also found it in the bark of the *S. Ignatius*.

Process.—According to the British official process for its preparation (*Strychnina*, B. P.), the seeds, disintegrated by subjection to steam, and, after drying, grinding in a coffee-mill, are exhausted with spirit, the latter removed by distillation, the extract dissolved in water, coloring and acid matters precipitated by lead acetate, the filtered liquid evaporated to a small bulk, the strychnine precipitated by ammonia, the precipitate washed, dried, and exhausted with spirit, the spirit recovered by distillation, and the residual liquid set aside to crystallize. Crystals of strychnine having formed, the mother-liquor (which contains the brucine of the seeds) is poured away, and the crystals of strychnine washed with spirit (to remove any brucine) and recrystallized.

In the U. S. P. (1870) process the rasped nux vomica is exhausted by very dilute hydrochloric acid, milk of lime added to the evaporated decoction to decompose the strychnine hydrochlorate, the precipitated and dried mixture of strychnine and lime treated with diluted alcohol to remove brucine, and then with strong hot alcohol to dissolve out strychnine; the alcohol having been recovered by distillation, the residual impure strychnine is dissolved in very dilute sulphuric acid, the solution decolorized by animal charcoal, evaporated, and set aside to crystallize, the crystals of strychnine sulphate (*Strychnine Sulphas*, U. S. P.; $(C_{21}H_{22}N_2O_2)_2H_2SO_4 \cdot 5H_2O$; Coleman, $6H_2O$) redissolved in water, ammonia added to precipitate pure strychnine, and the latter dried.

Properties.—Strychnine occurs "in right square octahedrons or prisms, colorless and inodorous; sparingly soluble in water, but communicating to it an intensely bitter taste; soluble in boiling rectified spirit and in chloroform, but not in absolute alcohol or in ether." It forms salts with acids. The *citrate*, $(C_{21}H_{22}N_2O_2)_2C_6H_8O_7 \cdot 4H_2O$ (or $6H_2O$), dissolves, at 60° F., in about 40 parts of water and 115 parts of alcohol. The hydrochlorate seems to have the formula $C_{21}H_{22}N_2O_2HCl \cdot 3H_2O$, and (Dott) is soluble in 35½ parts of water.

REACTIONS.

First Analytical Reaction.—Place quite a small fragment of strychnine on a white plate, and near to it also a small piece of red potassium chromate; to each add one drop of concentrated sulphuric acid; after waiting a minute or so for the chromate to fairly tinge the acid, draw the latter, by a glass rod, over the strychnine spot; a beautiful purple color is produced, quickly fading into a yellowish red. The following oxidizing agents may be used in the place of the chromate: puce-colored lead oxide, fragments of black manganese oxide, potassium ferrieyanide, or potassium permanganate.

This reaction is highly characteristic: a minute fragment dissolved in much dilute alcohol—or, better, chloroform—and one drop of the liquor evaporated to dryness on a porcelain crucible-lid or other white surface, yields a residue which immediately gives the purple color on being oxidized in the manner directed.

Other Reactions.—Strong sulphuric acid does not act on strychnine, even at the temperature of boiling water—a fact of which advantage is taken in separating strychnine from other organic matter for the purposes of toxicological analysis.—Potassium sulphocyanate produces, even in dilute solutions of strychnine, a white precipitate, which under the microscope is seen to consist of tufts of acicular crystals.—Strong nitric acid does not color strychnine in the cold, and on heating only turns it yellow.

The Physiological Test.—A small frog placed in an ounce of water to which $\frac{1}{100}$ of a grain of strychnine salt (acetate) is added is in two or three hours seized with tetanic spasms on the slightest touch, and dies shortly afterward.

Strychnine has an intensely bitter taste. Cold water dissolves only $\frac{1}{2000}$ part, yet this solution, even when largely diluted, is distinctly bitter. Alcohol is a somewhat better solvent. The salts of the alkaloid are more soluble. The official solution (*Liquor Strychnine Hydrochloratis*, B. P.) contains 1 per cent. of strychnine, the solvent being 3 parts water, 1 part spirit, and a few minims (6 per ounce) of hydrochloric acid (rather more than sufficient to form strychnine hydrochlorate). If too much acid be present, the hydrochlorate will crystallize out in cold weather. A syrup of phosphates of iron, quinine, and strychnine is official (*Syrupus Ferri, Quinine et Strychnine Phosphatum*). It contains 1 part of strychnine in 5000.

BRUCINE, or BRUCIA ($C_{23}H_{26}N_2O_4 \cdot 4H_2O$), is an alkaloid accompanying strychnine in nux vomica and St. Ignatius's bean to the extent of about 2 per cent. It is readily distinguished by the intense red color produced when nitric acid is added to it. *Igasurine*, once supposed to be a third alkaloid of nux vomica, has been shown by Shenstone to be only a mixture of brucine and strychnine.

Curarine ($C_{10}H_{15}N$), the active principle of the arrow-poison termed *curari*, *urari*, *ourari*, *wourali*, or *woorara*, prepared from

a *Strychnos*, resembles strychnine in giving color by oxidation, but the color is more stable. Potassium iodide or platinocyanide does not with curarine afford precipitates which crystallize from alcohol like those of strychnine. Curarine also is soluble in water. Unlike strychnine, curarine is reddened by sulphuric acid; it also is not dissolved out by ether from an acid or alkaline liquid. *Curari* appears to vary much in strength and quality. It is probably a mixture of vegetable extracts.

Distinction of Brucine from Morphine.—The red coloration produced by the action of nitric acid on brucine is distinguished from that yielded with morphine by the action of reducing agents (such as stannous chloride, sodium hyposulphite, sodium sulphhydrate), which decolorize the morphine-red, but change that of the brucine to violet and green (Cotton). The solution of brucine in the nitric acid should be heated to the boiling-point, diluted with water, and the stannous chloride then be added.

Distinction of Free Alkaloids or their Salts from Each Other.—This is accomplished by remembering the appearance and other physical characters of the substances as met with in pharmacy, the effect of heat, the action of such solvents as water, alcohol, and ether, the influence of strong and diluted acids, strong and weak alkalies, oxidizing substances, and other reagents. (Tables to aid in the analysis of small quantities of official alkaloids, their salts, glucosides, and various "scale" compounds, will be found opposite p. 544.)

QUESTIONS AND EXERCISES.

What alkaloids are more or less characteristic of the different varieties of cinchona-bark? In what form do they occur?—By what method is sulphate of quinine obtained?—Give the characters of quinine sulphate.—Describe the tests for quinine.—How would you detect salicin in quinine sulphate?—Show how the quinidine or cinchonine sulphates may be proved to be present in commercial quinine sulphate.—How are cinchonine and quinine distinguished from morphine?—Whence is strychnine obtained?—Describe the official process for the isolation of strychnine.—Give the characters of strychnine.—Describe the tests for strychnine.—By what reagent is brucine distinguished from strychnine?—Distinguish between brucine and morphine.—By what general methods would you distinguish common alkaloids from each other?

ALKALOIDS OF LESS FREQUENT OCCURRENCE.

ACONITINE, ACONITINA, or ACONITIA is an alkaloid obtained from aconite (*Aconitum Napellus*) leaves (*Aconiti Folia*, B. P.) and root (*Aconitum*, U. S. P.). The alkaloid itself is only slightly soluble in water; it occurs in the plant in combination with a vegetable acid, forming a soluble salt.

Process.—The official process for its preparation (*Aconitina*, B. P.) consists in dissolving out the natural salt of the alkaloid from the root by rectified spirit, recovering the latter by distillation, mixing

the residue with water, filtering, precipitating the aconitine by ammonia, drying the precipitate and digesting it in ether (in which some of the accompanying impurities are insoluble), recovering the ether by distillation, dissolving the dry residue in the retort in water acidulated by sulphuric acid, again precipitating the alkaloid by ammonia, and finally washing and drying.

Properties.—Aconitine usually occurs as a white powder. It has been obtained and studied in the crystalline state by Groves, Wright, Williams, and others. It is very slightly soluble in cold water, more so in hot, and much more soluble in alcohol, in ether, and in chloroform. It is one of the most violent poisons known. "When rubbed on the skin it causes a tingling sensation, followed by prolonged numbness." The thousandth part of a grain on the tip of the tongue produces, after a minute or so, a characteristic tingling sensation and numbness; large quantities rubbed into the skin cause numbness and loss of feeling. Sulphuric acid turns it of a yellowish, and afterward dirty violet, color.

According to Wright, who, in conjunction with Groves and Williams, worked by the aid of grants from the British Pharmaceutical Conference, *Aconitum Napellus* yields, chiefly, crystalline aconitine, $C_{33}H_{43}NO_{12}$, with some crystalline pseud-aconitine, $C_{36}H_{49}NO_{12}$, and a little non-crystalline alkaloid.

In the research laboratories of the Pharmaceutical Society of Great Britain, Dunstan and Umney found, in addition to aconitine ($C_{33}H_{43}NO_{12}$, Dunstan and Ince), aconine and an amorphous alkaloid, *napelline* or *isaconitine*, whose salts are also uncrystallizable. It is isomeric with aconitine, and also yields the same products on hydrolysis. Aconitine is readily hydrolyzed into aconine, $C_{26}H_{41}NO_{11}$, and benzoic acid (Dunstan and Passmore).

According to Jürgens, the formula of aconitine is $C_{33}H_{47}NO_{12}$. On allowing an acetic solution containing potassium iodide to evaporate to dryness, and then adding water, crystals of aconitine hydriodate of characteristic appearance remain.

The tuberous roots of *Aconitum Ferox* and other species constitute the *bish* or *bikh* of India (*Aconiti Ferocis Radix*, P. I.). It chiefly contains the variety of aconitine termed *pseudaconitine*. Some of the aconitine of pharmacy is *pseudaconitine*.

According to Paul and Kingzett, the alkaloid of Japanese aconite has the formula $C_{29}H_{43}NO_9$, while Wright and Menke state that the formula is $C_{66}H_{88}N_2O_{21}$, and the name is *japaconitine*.

Unguentum Aconitinæ, B. P., contains 8 grains of the alkaloid to 1 ounce of benzoated lard.

Aconitum heterophyllum, *Atis*, or *Atees*, or *Wakhma* (*Aconiti Heterophylli Radix*, P. I.), contains no aconitine, but an alkaloid, *ateesine*, having the formula $C_{46}H_{74}N_2O_5$.

ASPIDOSPERMINE, $C_{22}H_{30}N_2O_2$, is an alkaloid of *Quebracho blanco* bark (Fraude). Another and different alkaloid is *quebrachine* ($C_{21}H_{26}N_2O_3$) (Hesse). The latter chemist has isolated four other closely related alkaloids, and two from *Quebracho colorado* bark.

ATROPINE, or ATROPIA ($C_{17}H_{23}NO_3$).—This alkaloid has hitherto been considered to exist ready formed in the belladonna or deadly

nightshade (*Atropa Belladonna*; *Belladonnæ Folia et Radix*, U. S. P.) as soluble acid atropine malate. But the observations of Messrs. Schering and the researches of Will indicate that not atropine, but an isomer of atropine—namely, hyoscyamine—is the alkaloid chiefly and often solely present, and that the alkaline treatment during the process of extraction converts the hyoscyamine into atropine. Hyoscyamine solutions rotate a plane-polarized ray to the left; atropine has no optical rotatory power. Each similarly affects the eye.

Process.—Atropine is obtained by exhausting the root with spirit, precipitating the acid and some coloring-matter by lime, filtering, adding sulphuric acid to form atropine sulphate (which is somewhat less liable to decomposition during subsequent operations than the alkaloid itself), recovering most of the spirit by distillation, adding water to the residue, and evaporating till the remaining spirit is removed; solution of potassium carbonate is then poured in till the liquid is nearly but not quite neutral, by which resinous matter is precipitated: the latter is filtered away, excess of potassium carbonate then added, and the liberated atropine dissolved out by shaking the liquid with chloroform. The latter solution, having subsided, is removed, the chloroform recovered by distillation, the residual atropine dissolved in warm spirit, coloring-matter separated by digesting the liquid with animal charcoal, the solution filtered, evaporated, and set aside to deposit crystals.

Solubility.—Atropine is sparingly soluble in water, the liquid giving an alkaline reaction—more soluble in alcohol and ether.

Tests.—Atropine solutions give with gold perchloride a yellow precipitate. 1 drop of a dilute aqueous solution (2 grains to the ounce) powerfully dilates the pupil of the eye. It is generally applied on a piece of thin tissue-paper or small disk (*Lamellæ Atropinæ*, B. P.) placed between the eyelid and the eye. *Homatropine hydrobromate* is official (*Homatropinæ Hydrobromas*, B. P.). It is a white crystalline powder or aggregation of minute prismatic crystals, soluble in 6 parts of cold water and in 133 of ethylic alcohol. The dilute aqueous solution powerfully dilates the pupil of the eye. A 2 per cent. aqueous solution is not precipitated by the cautious addition of solution of ammonia previously diluted with twice its volume of water [distinction from atropine]. About $\frac{1}{10}$ of a grain moistened with 2 minims of nitric acid and evaporated to dryness on the water-bath yields a residue which is colored yellow by an alcoholic solution of potash [distinction from atropine, hyoscine, and hyoscyamine]. If about $\frac{1}{10}$ of a grain be dissolved in a little water and the solution be made alkaline with ammonia and shaken with chloroform, the separated chloroform will leave on evaporation a residue which will turn yellow, and finally brick-red, when warmed with about 15 minims of a solution of 2 grains of mercury perchloride in 100 minims of proof spirit. For Gerrard, Schweissinger, and Flückiger have observed that *homatropine* (Ladenburg's oxytoluyltropine, a physiologically similar but less powerful, and therefore sometimes more useful, alkaloid than atropine), like hyoscyamine and atropine, has unusually powerful alkaline properties, precipitating mercuric

oxide from mercuric solutions, reddening phenolphthalein, and, with warmth, blackening calomel. No other ordinary alkaloids are so powerfully alkaline.

Baryta-water decomposes atropine into *tropine* ($C_8H_{15}NO$) and *tropic acid* ($C_9H_{10}O_3$), a molecule of water being absorbed; hence the atropine, so called, would seem to be *tropine tropate*, minus water, or *tropyl-tropine*. Indeed, Ladenburg by heating tropic acid and tropine in sealed tubes has produced a base indistinguishable from atropine. The same chemist by removing the elements of water from tropine gets *tropidine*, $C_8H_{13}N$, closely related to ecgonine (p. 534) and anhydro-ecgonine. This is, possibly, an intermediate member of a group of alkaloids, of which others are *conine*, $C_8H_{15}N$, and *collidine*, $C_8H_{11}N$, the latter a product of the destructive distillation of bone oil, coal, quinine, etc. Commercial atropine is said by Regnaud and Valmont to be a mixture of true atropine with hyoscyamine.

In the so-called Japanese belladonna (*Scopola Japonica*) occurs *scopoleine*, an alkaloid resembling but more powerful than atropine (Eyckmann), but Schmidt considers that only atropine, hyoscyamine, and hyoscyne are present.

Preparations.—The alkaloid itself (*Atropina*) and the Sulphate of Atropine (*Atropine Sulphas*, or *Atropine Sulphate*, $(C_{17}H_{23}NO_3)_2H_2SO_4$, a colorless powder soluble in water, made by neutralizing atropine with sulphuric acid), are official in the United States Pharmacopœia.

The fluorescence of alkaline solutions of extract of belladonna is caused by *chrysotropic acid*, $C_{12}H_{10}O_5$ (Kunz), probably allied to the fluorescent *scopoletin*, $C_{10}H_8O_4$, found in Japanese belladonna by Eyckmann.

BAPTITOXINE.—Schroeder gives this name to a poisonous alkaloid in *Baptisia tinctoria*, wild indigo, in which he also finds the glucosides *baptisin* and *baptin*.

BEBERINE, BEBIRINE, or BIBIRINE ($C_{36}H_{42}N_2O_6$) is an alkaloid in the bark of bebeeru or bibiru (*Nectandra Rodiaei*).

Process.—According to the British Pharmacopœia, it—or rather its sulphate, $C_{36}H_{42}N_2O_6 \cdot H_2SO_4$ (*Beberine Sulphas*, B. P.)—may be prepared by exhausting the bark (*Nectandra Cortex*, B. P.) with water acidulated by sulphuric acid, concentrating, removing most of the acid by lime, filtering, precipitating the alkaloid by ammonia, filtering, drying, dissolving in spirit (in which some accompanying matters are insoluble), recovering most of the spirit by distillation, neutralizing by diluted sulphuric acid, evaporating to dryness, dissolving the residual sulphate in water, evaporating to the consistence of a syrup, and spreading on glass plates, drying the product at $140^\circ F$. Thus obtained, it occurs in dark-brown translucent scales, yellow when powdered, strongly bitter, soluble in water and in alcohol. It is probably a mixture of sulphates of beberine, nectandrine, and other alkaloids.

Tests.—Alkalies give a pale-yellow precipitate of beberine when added to an aqueous solution of a salt of the alkaloid; the precipitate is soluble in ether. With red potassium chromate and sul-

phuric acid beberine gives a black resin, and with nitric acid a yellow resin.

Buxine, from the bark of *Buxus sempervirens*; *Pelosine*, or *Cissampeline*, from the root (*Pareira*, U. S. P.) of *Cissampelos Pareira*; and *Paricine*, from a false *Para cinchona*-bark,—are probably identical with beberine (Flückiger).

Nectandrine ($C_{40}H_{46}N_2O_8, 4H_2O$).—Drs. MacLagan and Gamgee a few years ago discovered this second alkaloid in bebeeru-bark. It differs from beberine in fusing when placed in boiling water, in being much less soluble in ether, in giving with strong sulphuric acid and black manganese oxide a beautiful green and then violet coloration, and in having a distinct molecular weight. They considered that two other alkaloids exist in bebeeru-bark.

BERBERINE ($C_{20}H_{17}NO_4$) is an alkaloid existing in several plants of the natural order *Berberideæ* (three species yield *Indian Barberry*, *Berberis Cortex*, P. I.), in calumba-root (*Calumbæ*, U. S. P.), in the root of *Coptis Teeta*, or *Mishmi Bitter* (*Coptidis Radix*, P. I.), an Indian tonic, and in many other yellow woods. *Hydrastis Canadensis*, or *Golden Seal* (*Hydrastis*, U. S. P.), contains berberine, though another alkaloid, *hydrastine* (*vide* page 537) ($C_{22}H_{23}NO_6$), related to narcotine (which seems to be methoxyl-hydrastine; Schmidt) and to papaverine, and even a third, are asserted to be present, all, in Perkin's opinion, benzene derivatives of iso-quinoline. The dried rhizome and rootlets are official, *Hydrastis Rhizoma*, B. P., and these are the source of *Extractum Hydrastis Liquidum*, B. P., *Tinctura Hydrastis*, B. P., and *Glyceritum Hydrastis*, U. S. P. The root of *Berberis vulgaris* contains berberine and (Wacker) *oxyacanthine* ($C_{18}H_{19}NO_3$; $C_{19}H_{21}NO_3$, Rüdel), as well as (Hesse) *berbamine* (also $C_{18}H_{19}NO_3$). *Xanthorrhiza apii folia*, an old American tonic, and, apparently, *Xanthoxylon Fraxineum*, or prickly-ash bark (*Xanthoxylum*, U. S. P.) also contain berberine. The rhizome of *Menispermum Canadense*, *Yellow Parilla*, or *Canadian Moonseed* (*Menispermum*, U. S. P.), contains, according to Maisch, a colorless alkaloid as well as berberine. The color of the tissues of these vegetables is apparently due to berberine, for the alkaloid itself is remarkable for its beautiful yellow color.

Tests.—When a dilute solution of iodine and potassium iodide is added to a solution of any salt of berberine in hot spirit, excess of iodine being carefully avoided, brilliant green spangles are deposited. The reaction is sufficiently delicate to form, according to Perrins, an excellent test of the presence of berberine. This iodo-compound polarizes light and has other analogies with herapathite.

Berberine itself is not official, but plants in which it occurs are used as medicinal agents in all parts of the world.

Process.—Berberine is readily extracted by boiling the raw material with water, evaporating the strained liquid to a soft extract, digesting the residue in alcohol, recovering the alcohol by distillation, boiling the residue with diluted sulphuric acid, filtering, and setting aside; berberine sulphate separates out, and may be purified by recrystallization from hot water. The neutral sulphate, ($C_{20}H_{17}NO_4$) $_2$ H $_2$ SO $_4$, is very soluble in water; the acid sulphate, $C_{20}H_{17}NO_4$,

H_2SO_4 , is less soluble. The alkaloid itself is obtained by shaking lead hydrate with a hot aqueous solution of the berberine sulphate (Procter).

CAFFEINE.—See THEINE.

CAPSICINE.—M. Felletár obtained from capsicum-fruits (*Capsicum*, U. S. P.), which when ground form *Cayenne pepper*, a volatile alkaloid having the smell of conine. Thresh has obtained crystalline hydrochlorate and sulphate. The latter chemist has also succeeded in isolating the active principle of capsicum, which he has termed *capsaicin*, ($\text{C}_9\text{H}_{14}\text{O}_2$), a crystalline non-alkaloidal, excessively acrid substance. Its exact chemical character is not yet made out. According to Dr. Thresh, a similar pungent principle occurs in ginger (gingerol) and in grains of paradise (paradol), bodies probably isomeric with capsaicin. (See also *Capsicin* in Index.)

CARPAINE, $\text{C}_{14}\text{H}_{27}\text{NO}_2$, occurs in *Carica papaya*.

CHELIDONINE ($\text{C}_{19}\text{H}_{17}\text{N}_3\text{O}_3$) and CHELERYTHRINE ($\text{C}_{19}\text{H}_{17}\text{NO}_4$), the latter identical, apparently, with *sanguinarine*, are two alkaloids occurring in *Celandine* (*Chelidonium*, U. S. P.), associated with citric, malic, and chelidonic ($\text{C}_7\text{H}_4\text{O}_6$) acids. (See p. 541.)

COCAINE ($\text{C}_{17}\text{H}_{21}\text{NO}_4$) is an alkaloid of *Erythroxylon Coca* (*Coca*, U. S. P.), the leaves of which act powerfully as a restorative to the human system. The hydrochlorate of the alkaloid is official (*Cocainæ Hydrochloras*, U. S. P.); also a 10 per cent. solution of the same (*Liquor Cocainæ Hydrochloratis*, B. P.), preserved by aid of salicylic acid. Cocaine and its salts may be prepared by agitating with petroleum spirit a strong, acidulated, aqueous extract of the leaves made alkaline with sodium carbonate, well shaking the separated spirit with acidulated water, treating the separated acid fluid with ether and excess of sodium carbonate, washing out the alkaloid from the ether by water acidulated with hydrochloric acid, and finally evaporating the resulting aqueous solution of the hydrochlorate to the crystallizing point. Cocaine may be precipitated with ammonia and recrystallized from alcohol, ether, or warm benzene. From this pure cocaine the pure and very soluble hydrochlorate may be prepared by neutralizing with hydrochloric acid and crystallizing.

Prolonged contact of cocaine with hot water, acids, alkalies, or even alcohol, is undesirable, as cocaine readily breaks up into benzoyl-ecgonine and methylic alcohol, $\text{C}_{17}\text{H}_{21}\text{NO}_4 + \text{H}_2\text{O} = \text{C}_{16}\text{H}_{19}\text{NO}_4 + \text{CH}_3\text{OH}$, benzoyl-ecgonine afterward yielding ecgonine and benzoyl-hydrate or benzoic acid, $\text{C}_{16}\text{H}_{19}\text{NO}_4 + \text{H}_2\text{O} = \text{C}_9\text{H}_{15}\text{NO}_3 + \text{C}_7\text{H}_6\text{O}_2$. In coca other bases occur with cocaine. Paul and Cownley, also Giesel, find *cinnamyl-cocaine*. Hesse finds *cocamine* and *cocaidine*, isomeric with cocaine. Liebermann finds several bases, one of which is poisonous—namely, *isatropylcocaine*, $\text{C}_{19}\text{H}_{23}\text{NO}_4$, containing isatropyl in place of the benzoyl group in ordinary cocaine. It has an amorphous and sticky appearance. All these bases are easily hydrolyzed, yielding ecgonine; the latter with benzoic anhydride yields benzoyl-ecgonine, and this, with methyl iodide or otherwise, yields benzoyl-methyl ecgonine or ordinary

cocaine. By thus building up with other acidulous bodies than the benzoic a whole chemical series of "cocaines" can be produced.

Another alkaloid (benzoyl-pseudotropeine), yielding instead of ecgonine a compound isomeric with tropine, also occurs in coca (Giesel, Liebermann). Cinnamyl-cocaine and other coca bases are officially detected by the following test: "If 1 drop of a mixture of one volume of decinormal potassium permanganate and two volumes of water be added to 5 cc. of a 2 per cent. solution of cocaine hydrochlorate mixed with 3 drops of diluted sulphuric acid, and contained in a small, clean, glass-stoppered vial, the pink tint produced by the permanganate should not entirely disappear within half an hour."—*U. S. P.*

Cocaine hydrochlorate occurs in colorless acicular crystals or a white crystalline powder, soluble in water, chloroform, alcohol, amyl alcohol, very slightly in ether; not readily decomposed even when boiled in water. On heating a small amount of the powdered cocaine hydrochlorate for twenty minutes on a water-bath, no material loss should occur (absence of water of crystallization). When ignited it should burn completely away. If a small quantity of the hydrochlorate be rubbed with a glass stirrer, on the bottom of a dry white porcelain dish, with an equal amount of mercuric chloride, and the mixture then breathed upon, it will turn to a grayish-white color. "On adding 5 drops of a 5 per cent. solution of chromic acid to 5 cc. of a 2 per cent. solution of cocaine hydrochlorate, a yellow precipitate is produced which redissolves on shaking; on now adding 1 cc. of hydrochloric acid a permanent orange-yellow precipitate will be formed."—*U. S. P.* The free alkaloid is readily decomposed by water, especially when the solution is warmed. The solution in water has a bitter taste, and produces on the tongue a tingling sensation followed by numbness. The aqueous solution dilates the pupil of the eye. It gives no color to cold strong acids, but chars with hot sulphuric acid, and gives a purple precipitate with permanganates and a white precipitate with ammonia. Evaporated to dryness on a water-bath with nitric acid, and treated with alcoholic potash, it develops an odor resembling peppermint. Besides its action as a restorative when taken internally, cocaine brought into contact with the mucous membrane of the eye, mouth, throat, etc. produces local anæsthesia. According to Squibb, good cocoa yields .5 per cent. of cocaine. Cocaine may be detected in presence of other alkaloids by giving a yellow precipitate of the chromate with potassium chromate or chromic acid in presence of free hydrochloric acid.

COLCHICINE, the active principle of *Colchicum autumnale* (*Colchici Cormus*, B. P.; *Colchici Semen*, U. S. P.), is said to be an alkaloid, though some investigators think it has more of the characters of a neutral substance, and give it the name *colchicin*. Hertel gives it the formula $C_7H_{23}NO_6$, and states that ebullition with acidulated water converts it into *colchicein*, $C_{16}H_{21}NO_5 \cdot 2H_2O$, and methyl alcohol. Zeisel says it may be crystallized from chloroform, and offers the following formulæ for it and its derivative: *colchicin*, $C_{21}H_{22}(OCH_3)NO_5$; *colchicein*, $C_{21}H_{22}(OH)NO_5$. The most active medicinal prep-

aration is an extract made from the *fresh* seeds by digestion in large volumes of alcohol of at least 90 per cent., and subsequent digestion of the marc in hot water. The extracts left on evaporating the two fluids separately are to be carefully mixed (Mols).

CONINE, CONIA, CONYLIA, CONICINE, or CICUTINE.—Formula, $C_8H_{17}N$ (Hofmann), or *α -normal-propyl-piperidine*, $C_5H_{10}N(C_3H_7)$. This alkaloid is a volatile liquid, occurring in hemlock (*Conium maculatum*) in combination with an acid (malic?). It is not official. According to Petit, its boiling-point is $170^\circ C$. and its density 0.846. It forms crystalline salts.

Process.—It may be obtained by distilling hemlock-fruit (*Conium*, U. S. P.) with water rendered slightly alkaline by caustic soda or potash, or by similarly treating the fresh juice of the leaves. The alkaloid is a yellow oily liquid floating on the water that distils over; by redistillation it is obtained colorless and transparent.

The salts of conine have no odor, but when moistened with solution of an alkali yield the alkaloid, the strong smell of which, at once recalling hemlock, is characteristic. Juice of hemlock-leaves (*Conii Folia*, B. P.), to which solution of potash and boiling water has been added, forms the official inhalation of conine (*Vapor Conine*, B. P.).

Tests.—Sulphuric acid turns conine purplish red, changing to olive-green; nitric acid a blood-red; gold perchloride produces a yellowish-white precipitate, platinum perchloride no precipitate, in aqueous solutions.

Hemlock also contains *methyl-conine*, $(C_8H_{16})''CH_3N?$ (Kekulé and Von Planta), and *conhydrine*, $C_8H_{17}NO$.

The latter by dehydration yields a base, $C_8H_{15}N$. Kekulé's base is probably a methylic derivative of this, $C_8H_{14}CH_3N$, and not true methyl-conine.

According to Schiff, conine, isomeric at least with the natural alkaloid, may be produced artificially by action of ammonia on butyric aldehyde and destructive distillation of the resulting compound. Ladenburg has produced conine identical with the natural alkaloid from *α -picoline*. Conine may now therefore be said to be a product of organic synthesis, producible from its elements.

CORYDALINE is an alkaloid obtained by Wenzell from "turkey corn," the tubes of *Dicentra* (*Corydalis*) *formosa*.

CUSPARINE ($C_{20}H_{19}NO_3$), with *cusparidine* ($C_{19}H_{17}NO_3$) and *galipine* ($C_{20}H_{21}NO_3$), are alkaloids occurring in the bark of *Galipea cusparia*, or true Angostura bark (*Cuspariæ Cortex*, B. P.). The bitter principle, *angosturin*, is not an alkaloid.

CYTISINE, or ULEXINE, is an alkaloid found in laburnum and furze, and is also probably identical with *sophorine*, from *Sophora tomentosa*.

DATURINE.—*Vide* HYOSCYAMINE.

DELPHINE or DELPHININE and DELPHINOIDINE are the poisonous alkaloids of stavesacre (*Delphinium Staphisagria*). The powdered seeds of the plant are employed to kill the *pediculi* of animals. The seeds (*Staphisagriæ Semina*, B. P.) contain about 25 per cent. of

oil. *Unguentum Staphisagriae*, B. P., contains about 10 per cent. of oil.

DITAMINE (Jobst and Hesse), present in the *ditain* of Gruppe, is an alkaloid of "dita," or bark of *Eschites scholaris* or *Alstonia scholaris* (*Alstonia Cortex*, P. I.), a reputed febrifuge. Others are *echitamine* and *echitenine*. Oberlin and Schlagdenhauffen state that the allied *Alstonia constricta* (the bark of which is said to have advantages over the hop as a dietetic bitter) contains a crystalline alkaloid, *alstonine*, and uncrystallizable *alstonicine*. *Alstonine* seems to be allied to strychnine.

DUBOISINE.—*Vide* HYOSCYAMINE.

EMETINE ($C_{30}H_{44}N_2O_4$, Glenard; $C_{30}H_{40}N_2O_5$, Kunz).—This alkaloid is the active emetic principle of the root of *Cephaelis Ipecacuanha* (*Ipecacuanha*, U. S. P.). It occurs to the extent of 1 to 2 per cent. in the root, less in the stems, in combination with ipecacuanhic acid. The nitrate is peculiarly slightly soluble in water (Lefort). In the *Pulvis Ipecacuanhae et Opii*, U. S. P., or "Dover's powder" (powdered ipecacuanha, 1 part; powdered opium, 1 part; and sugar of milk, 8 parts), minute division of the active ingredients is promoted by prolonged trituration with sugar of milk, which is a very hard salt. *Ipecacuanha Wine* (*Vinum Ipecacuanhae*, B. P.) is a solution of neutral emetine acetate and of other matters extracted (by aid of acetic acid) from the root in sherry wine. *Acetum Ipecacuanhae*, B. P., is obtained from 1 part of root in 20 of diluted acetic acid. Choline (Arndt, a "volatile base") has been found in ipecacuanha. Ipecacuanha is said to have pharmacological activity (in dysentery) even after all emetine has been extracted; hence it would seem to contain more than one active principle.

The Indian substitute of ipecacuanha is the dried leaf (*Tylophora Folia*, P. I.) of *Tylophora asthmatica*. Its active principle has not been satisfactorily isolated.

GELSEMINE ($C_{11}H_{19}NO_2$, Sonnenschein; $C_{12}H_{14}NO_2$, Gerrard; $C_{54}H_{69}N_4O_{12}$, Thompson) is one of the alkaloids of *Gelsemium nitidum*, or Carolina yellow jasmine (*Gelsemium*, U. S. P.), in the tissues of which plant the *gelseminic acid* of Wormley, and *æsculin* ($C_{15}H_{16}O_9$), the fluorescent glucoside of the horse-chestnut and of many other plants, are also present. Like strychnine, gelsemine is not apparently affected by strong sulphuric acid. Nitric acid does not color it. A mixture of sulphuric acid and manganese peroxide colors it a crimson red, changing to green. In *Gelsemium elegans* Crow finds an allied alkaloid which does not resist the action of strong sulphuric acid. *Gelseminine* is another alkaloid said to be more powerful than gelsemine.

GRINDELIN is the name given by Fischer to a bitter crystalline alkaloid he extracted from *Grindelia Robusta* (*Grindelia*; *Extractum Grindelia Fluidum*, U. S. P.). The plant also contains resin and volatile oil.

GUARINE.—*Vide* THEINE.

HYDRASTINE, $C_{21}H_{21}NO_6$, is a colorless alkaloid obtained from *Hydrastis* (*vide* p. 533); on oxidation with dilute nitric acid it yields opianic acid ($C_{10}H_{10}O_5$) and hydrastinine.

HYDRASTININE ($C_{11}H_{11}NO_2$) is an artificial alkaloid obtained from hydrastine, as above, in white needles. The hydrochlorate is official (*Hydrastininæ Hydrochloras*, U. S. P.); it is a light-yellow acid powder, without odor, having a strong bitter but saline taste, deliquescent when exposed to moist air, soluble in water (very dilute solution has a blue fluorescence) and alcohol, but only very slightly in chloroform and ether. Pure hydrastinine hydrochlorate should burn away without leaving a residue.

Test.—"On adding to 2 cc. of an aqueous solution of the salt (1 in 100) an excess of bromine-water a yellow precipitate is produced, which is dissolved by ammonia-water to a nearly colorless liquid (difference from hydrastine, with which the ammonia produces a brick-red precipitate)."—U. S. P.

HYOSCINE.—This is an alkaloid which is supposed to be identical with "scopolamine" ($C_{17}H_{21}NO_4$), from the *Scopola atropoides* and *S. carnialica*, a tropate of another alkaloid isomeric (but not identical, Hesse, Schmidt) with tropine. Hyoscine hydrobromate ($C_{17}H_{21}NO_4 \cdot HBr + 3H_2O$) is official (*Hyoscine Hydrobromas*, U. S. P.). It crystallizes in colorless rhombohedra, which are neutral and easily soluble in water and alcohol; very slightly so in chloroform and ether. Heated to $100^\circ C.$, the salt loses its water of crystallization, and on ignition burns completely.

Test.—"If 5 drops of fuming nitric acid be added to 0.01 grm. of the salt in a small porcelain capsule, and the mixture be evaporated to dryness on a water-bath, a scarcely tinted residue will be left, which, when treated after cooling with a few drops of an alcoholic solution of potassium hydrate, will assume a violet color."—U. S. P.

HYOSCYAMINE ($C_{17}H_{23}NO_3$) occurs in the leaves (*Hyoscyamus*, U. S. P.) and other parts of henbane, belladonna, stramonium, and various species of *Scopola*; also (Dymond) in lettuce. It forms brilliant colorless needles. Its salts also are crystalline. Its effect on the eye is similar to that of atropine. The researches of Ladenburg show that hyoscyamine is the tropate of an alkaloid isomeric with tropine. (See Atropine.) Ladenburg also finds in henbane some *hyoscine*. (See above.) The hyoscyamine hydrobromate is official (*Hyoscyaminæ Hydrobromas*, U. S. P.). It is a yellowish-white, neutral, deliquescent, resin-like mass, either amorphous or crystalline. It is soluble in water and alcohol, but less so in ether and chloroform, and is not precipitated from its solution by platinum chloride. Gold chloride gives a precipitate which, on being recrystallized from a small quantity of boiling weak hydrochloric acid, deposits in lustrous golden-yellow scales (difference from atropine).

Hyoscyamine sulphate, $(C_{17}H_{23}NO_3)_2 \cdot H_2SO_4$, is official (*Hyoscyaminæ Sulphas*, U. S. P.). It is yellowish-white.

The alkaloids which occur in *Datura Stramonium*, or thornapple (*Folia et Semina Stramonii*, U. S. P.), *Datura* (*Datura alba*; *Datura Folia et Semina*, P. I.), and in *Duboisia Myoporoides*, and were formerly supposed to be distinct alkaloids, called respectively *daturine* and *duboisine*, are identical with hyoscyamine, and the latter is isomeric with atropine (Ladenburg). Duboisine may, however, be identical with hyoscine. Indeed, according to Schmidt,

the alkaloid of *Duboisia myoporoides* is sometimes hyoscyamine and sometimes hyosine or scopolamine. Bees which sip from the flowers of stramonium are said to produce poisonous honey.

Hyoscyamine melts when heated to between 108° and 109° C., and then is soon converted into atropine. Its solutions in alcohol or ether are stable, but the presence of a very minute amount of fixed caustic alkali or a very little alkaline carbonate causes complete conversion of the hyoscyamine into atropine. With gold chloride its salts give a yellow crystalline precipitate.

JABORANDINE and JABORINE.—See PILOCARPINE.

JERVINE ($C_{30}H_{46}N_2O_3$) occurs in *Veratrum album*, white hellebore, and (*V. Viride*, U. S. P.)* American white hellebore, the root of which is officially recognized in Great Britain (*Veratri Viridis Rhizoma*, B. P.). Its salts are much less soluble in water than those of veratrine. According to Bullock, *Veratrum viride* contains still another alkaloid, *veratroidine*, and, according to Mitchell, *Veratrum album* also contains an alkaloid which he terms *veratralbine*. Tobien gives the formula of jervine as $C_{27}H_{47}N_2O_8$, and of veratroidine as $C_{51}H_{78}N_2O_{16}$ or $C_{24}H_{37}NO_7$. According to Wright, *Veratrum album* contains jervine, $C_{26}H_{37}NO_3$; pseudo-jervine, $C_{29}H_{43}NO_7$; rubijervine, $C_{26}H_{43}NO_3$; veratralbine, $C_{28}H_{43}NO_5$; and traces of veratrine, $C_{37}H_{63}NO_{11}$. The same author finds *Veratrum viride* to contain jervine, pseudojervine, cevadine, $C_{32}H_{42}NO_9$, rubijervine, and traces of veratrine and veratralbine. Pehkschen finds jervine, $C_{14}H_{22}NO_2$, pseudojervine, $C_{29}H_{49}NO_{12}$, and veratroidine, $C_{32}H_{53}NO_9$, while Salzberger, besides jervine, rubijervine, and pseudojervine, finds protoveratrine, $C_{32}H_{51}NO_{11}$ and protoveratradine, $C_{26}H_{45}NO_8$. Salzberger confirms Wright and Luff's formula for jervine.

JUGLANDINE is the name given by Tanret to an alkaloidal substance obtained from the leaves of the walnut, *Juglans regia*. In the root-bark of *Juglans cinerea*, or butternut (*Juglans*, U. S. P.), Thiebaud found a bitter substance and an acid resembling chrysophanic.

LOBELINE.—A volatile fluid alkaloid first isolated from the dried flowering herb *Lobelia inflata* (*Lobelia*, B. P.) by Procter. In the pure state it is inodorous; impure, it smells slightly of the plant, but mixed with ammonia it emits a strong and characteristic smell of the herb. With acids it forms salts. A solid alkaloid is said to be present also.

LUPULINE is stated by Greismayer to be a liquid volatile alkaloid contained in the hop, *Humulus lupulus* (*Lupulus*, B. P.).

NECTANDRINE.—*Vide* BEBERINE.

NICOTINE, $C_{10}H_{14}N_2$ or $(C_5H_7)'''_2N_2$, or hexahydrodipyridyl, $C_{10}H_8(N_2)_2$.—This also is a volatile liquid alkaloid, forming the powerful active principle of tobacco (*Nicotiana Tabacum*), malate and citrate of nicotine being the forms in which it occurs in the leaf (*Tabacum*, U. S. P.). Its odor is characteristic; like conine, it yields a precipitate with gold perchloride; but, unlike that alkaloid, its aqueous

* The name *Green Hellebore* is sometimes applied to this drug, but properly belongs to *Helleborus viridis* (*vide* "Helleborin" in Index), which is medicinal in some parts of Europe (Hanbury).

solutions are precipitated yellowish-white by platinum perchloride. It is not official. It is also contained in *pituri*, a drug "chewed by the natives of some parts of Australia as a stimulant narcotic," though, according to Liversedge, the latter alkaloid may have the formula $C_{12}H_{16}N_2$.

PHYSOSTIGMINE ($C_{15}H_{21}N_3O_2$) (*Physostigmina*, B. P.).—An alkali obtained from the Calabar bean (*Physostigma*, U. S. P.), the seed of *Physostigma venenosum* (Jobst and Hesse), "by dissolving the alcoholic extract in water, filtering, adding sodium bicarbonate, shaking the mixture with ether, and evaporating the ethereal liquid." It occurs "in colorless or pinkish crystals, slightly soluble in water, but readily soluble in alcohol and in diluted acids. The aqueous solution has an alkaline reaction, when warmed with or when shaken with dilute solution of potash becomes red, and when evaporated to dryness with ammonia over a water-bath leaves a bluish residue, the acidified solution of which is beautifully dichroic, being blue and red."—*B. P.* A trace of it powerfully contracts the pupil of the eye (*Lamellæ Physostigminæ*, B. P.); a small quantity is highly poisonous. *Physostigmine Salicylate* is official (*Physostigminæ Salicylas*, U. S. P.), $C_{15}H_{21}N_3O_2 \cdot C_7H_6O_3$. The sulphate is also official (*Physostigminæ Sulphas*, U. S. P.), $(C_{15}H_{21}N_3O_2)_2 \cdot H_2SO_4$. It is a fine, odorless white crystalline powder with a bitter taste, very soluble in water and alcohol. Fraser also isolated another (but, possibly, the same) principle from the bean, and termed it *eserine*, from *esere*, the name of this ordeal-poison at Calabar. Eber states that physostigmine by action of acids, etc. takes up the elements of water, and becomes *eseridine*, $C_{15}H_{23}N_3O_3$, an alkaloid one-sixth the strength of physostigmine, and occurring to some extent in the Calabar bean itself.

PILOCARPINE is, apparently, the active principle of the diaphoretic and sialagogue *jaborandi*, the leaflets of *Pilocarpus pennatifolius* (*Pilocarpus*, U. S. P.). The occurrence of an alkaloid in this plant was first announced by Hardy, followed almost immediately by Byasson. A crystalline nitrate and hydrochlorate were first obtained by Gerrard. The leaves also yield an essential oil, a terpene, $C_{10}H_{16}$ (Hardy). Harnack and Meyer state that the true formula for pilocarpine is $C_{11}H_{16}N_2O_2$, and that its effects resemble those of nicotine, but that *jaborandi* yields another alkaloid, *jaborine*, which probably closely approaches pilocarpine in composition, though allied to atropine in effects. One salt is official in each of the Pharmacopœias, the nitrate (*Pilocarpinæ Nitrates*, B. P.), $C_{11}H_{16}N_2O_2 \cdot HNO_3$, and pilocarpine hydrochlorate (*Pilocarpinæ Hydrochloras*, $C_{11}H_{16}N_2O_2 \cdot HCl$, U. S. P.). This alkaloid is obtained from extract of *jaborandi* by shaking it with chloroform and a little alkali, and evaporating the chloroformic solution. The product, neutralized by nitric acid and purified by recrystallization, yields the nitrate as a white granular powder or as prismatic crystals. It has a faintly bitter taste, and is soluble in water and in rectified spirit. Strong sulphuric acid forms with it a yellowish solution which, on the addition of red potassium chromate, gradually acquires an emerald-green color. It

leaves no ash when burned with free access of air. It causes contraction of the pupil of the eye. Merck states that a third alkaloid, *pilocarpidine*, $C_{10}H_{14}N_2O_2$, is present in jaborandi. Harnack thinks that pilocarpine is probably a methyl derivative of pilocarpidine. The suggestion also is offered that the formula for nicotine differing only by O_2 from pilocarpidine, the latter is perhaps only dihydroxyl-nicotine. According to Merck, confirmed by Hardy and Calmels, jaborine is derived from pilocarpine by natural oxidation, while pilocarpidine similarly yields *jaboridine*, $C_{10}H_{12}N_2O_3$. The latter chemists have obtained pilocarpine artificially, " β -pyridine α -lactic acid" being converted into pilocarpidine, and this into pilocarpine.

PIPERINE (*Piperinum*, U. S. P.) ($C_{17}H_{19}NO_3$) is a feeble alkaloid occurring in *White, Black* (*Piper*, U. S. P.), *Long Pepper* (*Chavica officinarum*, Mign.), and *Cubeb Pepper* (*Cubeba*, U. S. P.), associated with volatile oil and resin; to these three substances the odor, flavor, and acridity belong. Piperine is obtained on boiling white pepper with alcohol, and evaporating the liquid with solution of potash, which retains resin. Recrystallized from alcohol, piperine forms colorless prisms fusible at 212° F. With acids and certain metallic compounds it forms salts, and distilled with strong alkali yields *piperidine* or *piperidia* ($C_5H_{10}HN$), an alkaloid of strong chemical properties, and *piperic acid* ($C_{12}H_{10}O_4$). Piperidine is interesting as being one of the alkaloids that has been obtained artificially by Ladenburg. It is hexa-hydro-pyridine, and is obtained by the action of nascent hydrogen on pyridine. Johnstone finds it in long pepper and in ordinary pepper, more especially in the husk. According to Buchheim, the amorphous resin of the peppers is similar in constitution to piperine, alkalies breaking it up into piperidine and *chavicic acid*. *Pyrethrin* is also said to be a member of the series. The piperine of cubeb pepper is not to be confounded with *cubebin*, a neutral constituent and having the formula $C_{10}H_{10}O_3$ —a derivative, probably, of pyrocatechin.

SANGUINARINE is the alkaloid of *blood-root* (*Sanguinaria Canadensis*). Its salts are red. König and Tietz find five distinct alkaloids in the root of sanguinaria—viz. *chelerythrine*, $C_{21}H_{17}NO_4$; *sanguinarine*, $C_{20}H_{15}NO_4$; α -*homochelidonine*, $C_{21}H_{21}NO_5$; β -*homochelidonine*, $C_{21}H_{21}NO_5$; and *protopine*, $C_{20}H_{17}NO_5$. Protopine was found in opium by Hesse (who assigned it the formula $C_{20}H_{19}NO_5$). It also occurs in *Celandine Chelidonium*, U. S. P., p. 534, and probably is identical with *macleine*, obtained by Eyekmann from *Macleya cordata*.

SOLANINE, ($C_{43}H_{69}NO_{16}$).—An alkaloid existing in the woody nightshade or bitter-sweet (*Solanum dulcamara*). The dried young branches of the plant are official (*Dulcamara*, U. S. P.). It occurs also in the shoots and in minute amount in the skins of the tubers of the potato (*Solanum tuberosum*). This alkaloid is only slightly soluble in water, alcohol, or ether; nitric acid colors it yellow; sulphuric acid produces at first a yellow, then a violet, and finally a brown coloration. It is said to be a conjugated compound of sugar with *solanidine* ($C_{25}H_{39}NO$). Geissler finds *dulcamarin* ($C_{22}H_{34}O_{10}$), a glucoside, to be the bitter constituent of *Solanum dulcamara*. A mixture of sulphuric acid and alcohol, or either selenic acid or

sodium selenate and sulphuric acid, colors solanine or solanidine a dark red.

SPARTEINE ($C_{15}H_{26}N_2$) is a poisonous volatile alkaloid occurring in broom-tops (*Scoparius*, U. S. P.). Its discoverer, Stenhouse, considers that the diuretic principle of broom is *scoparin*, a non-poisonous body, sparingly soluble in cold water. Mills has obtained ethyl-sparteine ($C_{15}H_{25}C_2H_5N_2$) and diethyl-sparteine ($C_{15}H_{24}C_2H_5N_2$). Apparently sparteine contains two pyridine nuclei. The sulphate is official, *Sparteinae Sulphas*, U. S. P. ($C_{15}H_{26}N_2 \cdot H_2SO_4 \cdot 4H_2O$). It is a colorless crystalline or granular powder, devoid of smell, but having a bitter taste, soluble in water and alcohol. At a temperature of 83° – 100° C. the salt loses its water of crystallization, and on heating more strongly is finally burnt completely away. "25 cc. of ether added to about 0.1 gm. of sparteine sulphate in a test-tube, then a few drops of dilute ammonia, so that the latter shall not be in excess, and an ethereal solution of iodine (1 in 50) afterward added until the liquid, when shaken, turns from an orange to a dark reddish-brown color; the bottom and sides of the test-tube will after a short time be found coated with minute, dark greenish-brown crystals, distinctly seen with a lens after the liquid has been poured out."—U. S. P. A very small quantity of the salt added to a few drops of sodium hydrate solution and gently warmed should give no smell of ammonia (absence of ammonium compounds).

SPIGELINA.—According to Dudley, this is a volatile alkaloid and active principle of *Spigelia marilandica*, or pink-root (*Spigelia*, U. S. P.), a vermifuge and depressant.

STILLINGINE.—Bichy states that this alkaloid is present in *Stillingia sylvatica* or *queen's root*.

TAXINE, $C_{37}H_{52}O_{10}N$? is an alkaloid occurring in the yew.

THEINE, or CAFFEINE, or GUARANINE (methyl-theobromine) ($C_8H_{10}N_4O_2 \cdot H_2O$).—This alkaloid (*Caffeina*, U. S. P.) occurs in tea, 2 to $4\frac{1}{2}$ per cent.; coffee, 1.2 per cent.; maté or Paraguay tea, .2 to 2 per cent.; guarana (*Guarana*, U. S. P.), "a dried paste chiefly consisting of the crushed or pounded seeds of *Paullinia Cupana*, Kunth (*Paullinia sorbilis*, Martius)," 5 per cent.; and the kola-nut. Infusions and preparations of these vegetable products are used, chiefly as beverages, by three-fourths of the human race. It is remarkable that the instinct of man, even in his savage state, should have led him to select as the bases of common beverages just the four or five plants which out of many thousands are the only ones, so far as we know, containing theine.

Theine is volatile. Considerable quantities may be collected by condensing the vapors evolved during the roasting of coffee on the large scale. The infusion of tea from which astringent and coloring matters have been precipitated by solution of lead subacetate, and which has been evaporated to a small bulk, yields a precipitate of theine on the addition of a strong solution of potassium carbonate. It may be crystallized from alcohol or by sublimation. Theine forms salts with acids. With citric acid caffeine forms the so-called citrated caffeine (*Caffeina Citrata*, U. S. P.), $C_8H_{10}N_4O_2 \cdot H_3C_6H_5O_7$, which, in the dry state, is a citrate of the alkaloid, but on the addi-

tion of water is immediately decomposed. The citrate is a white, odorless powder with an acid taste and reaction, soluble in two volumes of chloroform and one volume of alcohol. It is used in the preparation of *Caffeina Citrata Effervescens*, U. S. P.

Tests.—Concentrated nitric acid, or, better, a mixture of potassium chlorate and hydrochloric acid, rapidly oxidizes theine, forming compounds which with ammonia yield a beautiful purple-red color, resembling the murexid obtained under similar circumstances from uric acid; the oxidation must not be carried too far. Theine boiled with caustic potash yields methylamine (CH_3HHN), the vapor of which has a peculiar characteristic odor.

The chemical action of theine on the system is not yet quite made out. It is probably a pure stimulant.

THEOBROMINE, $\text{C}_7\text{H}_8\text{N}_4\text{O}_2$, is an alkaloid occurring in *cocoa*, the seed of *Theobroma Cacao*, to the extent of 1 to 2 per cent. According to Schmidt, a little theine is present also. Theobromine is also present in kola-nut (Heckel and Schlagdenhauffen). The theine in cacao and kola, and probably in tea, is said to occur normally as a glucoside, which would explain why it is only partially extracted by chloroform from a mixture of tea, etc. and lime.

Relations between Theine and Theobromine.—Both theine and theobromine are methyl derivatives of xanthine, $\text{C}_5\text{H}_4\text{N}_4\text{O}_2$ (belonging to the uric-acid group, uric acid having the formula $\text{C}_5\text{H}_4\text{N}_4\text{O}_3$). Theobromine, or dimethylxanthine (? the "theophylline" found in tea by Kossel), may be obtained from a compound of xanthine and silver by the action of methyl iodide, and theine (methyltheobromine or trimethylxanthine) may be obtained by heating theobromine-silver with methyl iodide (Strecker).

TRIGONELLINE, $\text{C}_7\text{H}_7\text{NO}_2\cdot\text{H}_2\text{O}$.—Jahns states that this alkaloid, as well as one identical with choline, is present in the seeds of *fœnugreek* or *fenugreek* (*Trigonella Fœnum-græcum*), much used in veterinary medicine and in some varieties of cattle-food and curry-powder.

VERATRINE, OR VERATRIA ($\text{C}_{32}\text{H}_{50}\text{NO}_9$, Schmidt and Köppen; $\text{C}_{52}\text{H}_{86}\text{N}_2\text{O}_{15}$, Weigelin) (*Veratrina*, U. S. P.).—This alkaloid occurs in *Cevadilla* (*Sabadilla*, B. P., the seeds of *Schænocaulon officinale* of A. Gray, termed *Asagrea officinalis* by Lindley and *Veratrum officinale* by Schlecht). It is also said to occur in the leaves of *Sarracenia purpurea*. According to Weigelin, cevadilla contains two isomeric varieties of veratrine, the one soluble, the other insoluble, in water. He says there are also present *sabadilline* ($\text{C}_{41}\text{H}_{66}\text{N}_2\text{O}_{13}$) and *sabatrine* ($\text{C}_{51}\text{H}_{86}\text{N}_2\text{O}_{17}$). The veratrine of trade contains the two latter alkaloids (Weigelin). A mere trace of veratrine brought into contact with the mucous membrane of the nose causes violent fits of sneezing. These alkaloids, and those from the different species of *Veratrum*, are evidently very closely allied. Wright and Luff, by the use of tartaric acid, a solvent less likely than the stronger acids to decompose alkaloids, extract from cevadilla *veratrine*, $\text{C}_{37}\text{H}_{53}\text{NO}_{11}$; *cevadine*, $\text{C}_{32}\text{H}_{49}\text{NO}_9$; and *cevadilline*, $\text{C}_{34}\text{H}_{53}\text{NO}_8$. According to Merck, cevadilla contains two alkaloids—*sabadine*, $\text{C}_{29}\text{H}_{51}\text{NO}_8$, and *sabadinine*, $\text{C}_{27}\text{H}_{46}\text{NO}_8$.

The British official process for the preparation of the alkaloid consists in exhausting the disintegrated cevadilla-seeds by alcohol, recovering most of the spirit by distillation, pouring the residue into water, by which much resin is precipitated, filtering, and precipitating the veratrine from the aqueous solution by ammonia. It is purified by washing with water, solution in dilute hydrochloric acid, decolorization of the liquid by animal charcoal, reprecipitation by ammonia, washing, and drying. The U. S. P. (1870) process is similar, but includes treatment of the first crude veratrine by diluted sulphuric acid and precipitation of alkaloid by magnesia. Bosetti states that it is a mixture of crystalline cevadine, insoluble in water, with an amorphous isomeric soluble alkaloid, *veratridine*. According to Lissauer, their physiological action is identical.

Unguentum Veratrinæ, U. S. P., contains 4 per cent. of the alkaloid. The veratrine is rubbed in a mortar, with 6 per cent. of olive oil and 90 per cent. of benzoated lard gradually added and thoroughly mixed.

QUESTIONS AND EXERCISES.

How is aconitine prepared?—Give the strength of the official preparations of atropine.—Describe the properties of atropine.—What is the active principle of stramonium?—Mention official substances containing beberine and berberine.—Give the characters of beberine.—In what does nectandrine differ from beberine?—Mention the characteristics of conine.—What is the active principle of ipecacuanha?—Name the alkaloid of tobacco.—Give the properties of the alkaloid of Calabar bean.—What are the sources of piperine?—Whence is theine obtained? what is its relation to theobromine?—Describe the preparation of veratrine.—State the properties of veratrine.

HERE THE STUDENT IS RECOMMENDED TO QUALITATIVELY ANALYZE UNNAMED SPECIMENS (PREVIOUSLY SELECTED FOR HIM) OF THE FREE AND COMBINED ORGANIC SUBSTANCES INCLUDED IN THE APPENDED TABLES 1 AND 2.

PROXIMATE CONSTITUENTS OF ANIMAL ORGANISMS.

PROTEID PRINCIPLES, OR ALBUMENOIDS.

Albumen.—Agitate, thoroughly, white of egg (*Ovi Albumen*, B. P.) with water, and strain off the liquid from the flocculent membranous insoluble matter. One white of egg in 180 cc. of water forms the "Albumen Test-solution," U. S. P.

Test.—Heat a portion of this solution of albumen to the

L. ALKALOIDS, GLUCOSIDES, ETC.

7.)

a drop of Mayer's solution (mercuric iodide and potassium

r glucosides, etc.

y so, then—

chloride and by sodium bicarbonate and chloroform.

on of stannous chloride.

chloride and other tests.

uric acid and ferric chloride.

ng with strong hydrochloric acid, which gives a reddish-

on.

tion.

joquin, etc.

test, etc.

l tingling.

l = red precipitate.

y acidified with hydrochloric acid and cooled—crystalline

aporation to dryness; the residue dissolved in acid gives a

file or phenyl-carbamine, C_6H_5NC .

scarlet.

alcohol, partly in ether. Acid taste.

deep red.

uted sulphuric acid = deep green.

d.

bride with an equal volume of strong sulphuric acid gives a

chloric acid, and ferric chloride = deep red (Fe_26CNS).

ities of the substance.

eye is involved should be made with extremely dilute solu-
to 1 pint of water. If no effect is produced in an hour, it is
one of twice the latter strength, and so on. The chief
tracting agents (myotics) are physostigmine and pilocarpine.

following memoranda:

Dissolved in strong sulphuric acid, with addition of a few
or red color, which is changed to deep claret on addition of

ong sulphuric acid gives red-brown color.

ad precipitated by acids. Taste slightly bitter.

1. TABLE TO AID IN THE IDENTIFICATION OF OFFICIAL ALKALOIDS, GLUCOSIDES, ETC.

(Compiled by F. W. SHORT.)

To face page 544.]

Dissolve a grain or so in a few drops of water or dilute hydrochloric acid, and add a drop of Mayer's solution (mercuric iodide and potassium iodide) or of bis-niuth and potassium iodide.

A precipitate indicates the presence of an alkaloid.

No precipitate indicates the absence of an alkaloid. Search must then be made for glucosides, etc.

To a small quantity on a porcelain plate add strong nitric acid.	Purple-red color.	A. If the substance is colorless, or nearly so, then—	
	Blood-red "	Apomorphine.	Confirm by ferric chloride and by sodium bicarbonate and chloroform.
	Orange-red "	Brucine.	" " addition of stannous chloride.
	" "	Morphine.	" " ferric chloride and other tests.
To another portion add strong sulphuric acid.	" "	Cocaine.	" " sulphuric acid and ferric chloride.
	Dirty-red "	Veratrine.	See next section.
	Red or brown on plate, deep red if warmed in tube.	Veratrine.	Confirm by heating with strong hydrochloric acid, which gives a reddish-purple color.
	Bluish tinge.	Cocaine.	See previous section.
If not found by preceding sections, heat a little in a dry tube.	Blood-red color.	Salicin, glucosides.	Confirm by oxidation.
	" "	Quinine.	" " thalleioquin, etc.
	" "	Quinidine.	" " "
	Red vapors.	Cinchonine.	" " ether test, etc.
	" "	Cinchonidine.	" " "

If not found by aid of the preceding sections, test specially as follows:

1st. If an alkaloid:

Aconitine	Make a dilute solution and place a drop on the tongue—numbing and tingling.
Atropine	(Alcoholic sol. added to alcoholic sol. of mercuric chloride and warmed—red precipitate.
Homatropine	Add nitric acid, dry over water-bath, add alcoholic potash—yellow.
Caffeine	Murexid test.
Cocaine	Permanganate—purple precipitate. Boiled with potash, then <i>slightly</i> acidified with hydrochloric acid and cooled—crystalline precipitate of benzoic acid.
Physostigmine	Warmed with potash gives red color, which becomes bluish on evaporation to dryness; the residue dissolved in acid gives a dichroic solution.
Pilocarpine	See page 546.
Strychamine	Sulphuric acid and red potassium chromate.
2d. If not an alkaloid:	
Aconitine	Heat with potash and chloroform—unpleasant odor of phenyl-isocyanide or phenyl-carbamine, C_6H_5NC .
Elaterin	With phenol and strong sulphuric acid, a crimson color, changing to scarlet.
Gluside (saccharin).	Is extremely sweet.
Jalapin (purified jalap resin).	Insoluble in water or turpentine; soluble in alkalis or alcohol, partly in ether. Acrid taste.
Phenacetin	Boil with hydrochloric acid, dilute, cool, filter, add red chromate—deep red.
Phenazon (antipyrine, analgesin)	To an aqueous solution add sodium nitrite and diluted sulphuric acid—deep green.
Picrotoxin	Compare a microscopic slide with one of picrotoxin similarly prepared.
Santonin	Almost insoluble in water, but soluble in alkalis. Dilute ferric chloride with an equal volume of strong sulphuric acid gives a red or violet color.
Sulphonat	Heat with potassium cyanide (odor of mercaptan); add water, hydrochloric acid, and ferric chloride—deep red (Fe_2CNS).

Note. Acids are sought by the ordinary reactions carefully applied on small quantities of the substance.

Caution.—Any experiments in which contraction or dilatation of the pupil of the eye is involved should be made with extremely dilute solutions to be in with—say 1 drop of the solution of the substance under examination to 1 pint of water. If no effect is produced in an hour, it is easy then to make the experiment with a fluid of double this strength, afterward with one of twice the latter strength, and so on. The chief dilating agents (mydriatics) are atropine, its isomers and homologues, and the chief contracting agents (myotics) are physostigmine and pilocarpine.

B. If the substance is colored, seek the aid of the following, memoranda:

1st. If an alkaloid:

Boberine . . . Very bitter; soda gives a yellow precipitate, soluble in ether.

2d. If not an alkaloid:

Alcin . . . Very bitter; nitric acid gives a red color, with soclain brownish. Dissolved in strong sulphuric acid, with addition of a few drops of nitric acid, and diluted with water, it gives an orange or red color, which is changed to deep claret on addition of ammonia in excess.

Chrysarobin . . . Scarcely soluble in water, soluble in alkalis with fine red color. Strong sulphuric acid gives red-brown color.

Jalap resin . . . See above (may be almost white).

Podophyllin . . . Soluble in alcohol and precipitated by water. Soluble in ammonia and precipitated by acids. Taste slightly bitter.

Santonin . . . See above (white when fresh, but yellow after exposure to light).

DINARY SCALE COMPOUNDS.

(generally converted into pyrophosphoric). a contamination).		INORGANIC BASES.	{ AMMONIUM (often as a contamination). FERRIC SALT. POTASSIUM. SODIUM.	
y of the scale. Heat the ash with nitric ess of solution of ammonium molybdate in				
E.	NO YELLOW PRECIPITATE.		<i>Ammonium</i> .—Boil aqueous solution of scale with potash and test vapor for ammonia. Filter and dissolve precipi- tate in hydrochloric acid, and test the solution for <i>iron</i> by ferrocyanide, sul- phocyanate, etc. <i>Potassium and Sodium</i> .— Ignite a small quantity of the scale, and moisten the residue with water. Test moistened residue with lit- mus-paper. If alkaline, ex- amine for potassium and sodium by the color im- parted to flame, and for potassium by the platinum test.	
po- pi- ous er, id,	Precipitate some of the aqueous solution with potash, filter, and add to a portion of the filtrate a slight excess of nitric acid, divide into two parts. To one add barium chloride (ppt. = <i>sulphuric acid</i>). To the other add silver nitrate (ppt. = <i>hydrochloric acid</i>). Neutralize another portion of the potash filtrate with nitric acid and add silver nitrate.			
to ip- u- ric po- us	PRECIPITATE GRAY TO BLACK.	PRECIPITATE WHITE.		
	Add very little ammonia (not suf- ficient to dissolve the whole precipi- tate) and heat. A silver mirror = <i>tartaric acid</i> . Calcium chlo- ride and lime ppt. a neutral solution (if concentrated) in the cold, the precip. redissolv- ing on boiling.	<i>Citric acid</i> gives imperfect or no mirror. Calcium chloride and lime do not precipitate <i>citric acid</i> in the cold, but upon boiling (if solu- tion be sufficient- ly concentrated) precipitation oc- curs.		
Confirm Tartaric or Citric Acid.—To slightly acidified potash filtrate add ammonia in slight excess and considerable quantity of ammo- nium and calcium chlorides. <i>Tartrates</i> are precipitated completely in the cold with agitation and rest for about ten minutes. To the solution (or filtrate, if tartrates are present) add three volumes of spirit of wine, when <i>citrates</i> are precipitated. If sulphates have been found, disregard a slight precipitate with spirit of wine.				

Compiled by A. SENIER.

2. TABLE FOR THE QUALITATIVE ANALYSIS OF ORDINARY SCALE COMPOUNDS.

ALKALOIDS.	(QUININE, QUINIDINE, HEBERLINE.)	(CINCHONINE, CINCHONIDINE, STRYCHNINE.)	ACIDS.	PYROPHOSPHORIC. HYPOPHOSPHOROUS (generally converted into pyrophosphoric). SULPHURIC. HYDROCHLORIC (as a contamination). TARTARIC. CITRIC.	INORGANIC BASES.	AMMONIUM (often as a contamination). FERRIC SALT. POTASSIUM. SODIUM.
Dissolve a portion in water, and add ammonia cautiously. Precipitate alkaloïds (except strychnine) and sometimes ferric hydrate. Agitate the mixture with a little ether, and separate, by means of a pipette, the ethereal solution, aqueous solution and insoluble precipitate.						
ETHEREAL SOLUTION.	INSOLUBLE PRECIPITATE.	AQUEOUS SOLUTION.	A YELLOW PRECIPITATE.	NO YELLOW PRECIPITATE.		
May contain quinine, quinidine, or heberline. To solution in a test-tube add water very slightly acidulated with acetic acid, boil, burning off the ether. To a portion of the acetic solution add chlorine or bromine water, then ammonia.	Is cinchonine, cinchonidine, or ferric hydrate. Saturate a drop or two of acetic acid in a little water with the precipitate, and to part of the solution add sodium tartrate, a precipitate occurs in the case of <i>cinchonidine</i> , and no precipitate in that of <i>cinchonine</i> .	May contain strychnine. Agitate with chloroform and separate chloroformic solution. Evaporate chloroformic solution and moisten residue with strong sulphuric acid. Draw across the acid film a small crystal of red potassium chromate moistened with sulphuric acid, when a transient play of colors (violet, red, yellow) is observed. No colors, no strychnine. In case of doubt add ammonia to original solution and treat with chloroform, and proceed as before.	Pyrophosphoric or hypophosphorous acid. Precipitate some of the aqueous solution with potash, filter, neutralize with nitric acid, and add silver nitrate. White precipitate soluble in nitric acid— <i>pyrophosphoric acid</i> . White to black precipitate soluble in nitric acid— <i>hypophosphorous acid</i> .	Precipitate some of the aqueous solution with potash, filter, and add to a portion of the filtrate a slight excess of nitric acid, divide into two parts. To one add barium chloride (ppt. <i>sulphuric acid</i>). To the other add silver nitrate (ppt. <i>hydrochloric acid</i>). Neutralize another portion of the potash filtrate with nitric acid and add silver nitrate. PRECIPITATE GRAY TO BLACK. Add very little ammonia (not sufficient to dissolve the whole precipitate) and heat. A silver mirror <i>tartaric acid</i> . Calcium chloride and lime ppt. a neutral solution (if concentrated) in the cold, the precip. redissolves on boiling.	Ammonium.—Boil aqueous solution of scale with potash and test vapor for ammonia. Filter and dissolve precipitate in hydrochloric acid, and test the solution for iron by ferrocyanide, sulphocyanate, etc. Potassium and Sodium.—Ignite a small quantity of the scale, and moisten the residue with water. Test moistened residue with litmus-paper. If alkaline, examine for potassium and sodium by the color imparted to flame, and for potassium by the platinum test. PRECIPITATE WHITE. <i>Citric acid</i> gives imperfect or no mirror. Calcium chloride and lime do not precipitate citric acid in the cold, but upon boiling of solution be sufficiently concentrated precipitation occurs. Confirm Tartaric or Citric Acid.—To slightly acidified potash filtrate add ammonia in slight excess and considerable quantity of ammonium and calcium chlorides. <i>Tartrates</i> are precipitated completely in the cold with agitation and test for about ten minutes. To the solution or filtrate, if tartrates are present add three volumes of spirit of wine when <i>citrates</i> are precipitated. If sulphates have been found, disregard a slight precipitate with spirit of wine.	
GREEN COLOR (thalleioquin).	NO GREEN COLOR.					
Solution is fluorescent, and contains either quinine or quinidine. Concentrate the remainder of the solution and divide into two parts. To one add potassium iodide, and to the other add ammonium oxalate. The former precipitates <i>quinidine</i> not quinine. The latter precipitates <i>quinine</i> , not quinidine. For other methods see page 524.	To a portion of the acetic solution add potash, a yellowish-white precipitate is formed.					

Compiled by A. SENIER.

boiling-point; the albumen becomes insoluble, separating in clots or coagula of characteristic appearance.

Other Reactions.—Add to small quantities of aqueous solution of albumen solutions of corrosive sublimate, silver nitrate, copper sulphate, lead acetate, alum, stannic chloride; the various salts not only coagulate, but form insoluble compounds with albumen. Hence the value of an egg as a temporary antidote in cases of poisoning by many metallic salts, its administration retarding the absorption of the poison until the stomach-pump or other means can be applied. Sulphuric, nitric, and hydrochloric acids precipitate albumen; the coagulum is slowly redissolved by aid of heat, a brown, yellow, or purplish-red color being produced. Neither acetic, tartaric, nor organic acids generally, except picric and gallotannic, coagulate albumen. Alkalies prevent the precipitation of albumen.

Yolk or Yelk of Egg (Vitellus, U. S. P.) contains only 3 per cent. of albumen, the white $12\frac{1}{2}$. The yolk also contains 30 per cent. of yellow fat and 14 of casein, with what is stated to be another proteid, vitellin.

Albumen is met with in large quantity in the serum of blood, in smaller quantity in chyle and lymph, and in the brain, kidneys, liver, muscles, and pancreas. It is not a normal constituent of saliva, gastric juice, bile, or mucus, but may occur during inflammation. It is found in the urine and fæces only under certain diseased states of the system.

The cause of the coagulation of albumen by heat has not yet been discovered.

Albumen has never been obtained sufficiently pure to admit of its composition being expressed by a trustworthy formula; Gerhardts regarded it as a sodium compound ($\text{HNaC}_{72}\text{H}_{116}\text{N}_{18}\text{SO}_{22}\text{H}_2\text{O}$).

Egg-albumen—and, to some extent, blood-albumen—is largely used by calico-printers as a vehicle for colors, serving also, when dry, as a glaze. Curriers prize egg oil for softening leather.

Albumen coagulated by heat is said to be recoverable in a scarcely altered fluid condition by contact with a dilute aqueous solution of a very small proportion of pepsin.

Fibrin, Casein, Legumin.

Fibrin is the chief constituent of muscular tissue. It occurs in the blood in the form of a very unstable compound termed *fibrinogen*, and its liberation from this union and spontaneous solidification or coagulation is the cause of the *clotting* of blood shortly after being drawn from the body. The latter phenomenon cannot at present be explained satisfactorily. Fibrin may be obtained by whipping fresh blood with a bundle of twigs, separating the adherent fibres, and washing in water till colorless. It may be kept either dry or in spirit of wine.

Average Composition of Blood (in 1000 parts) (compiled by Kirkes).

	Water	784
	Albumen	70
	Fibrin	2.2
	Red corpuscles (globulin, 123; hæmatin, 7)	130
Fatty matters.	Cholesterin	0.08
	Cerebrin	0.40
	Serolin	0.02
	Oleic and margaric acids	—
	Volatile and odorous fatty acid	—
	Fat containing phosphorus	—
Inorganic salts.	Sodium chloride	3.6
	Potassium chloride	.35
	Sodium phosphate (Na_3PO_4)	.2
	Sodium carbonate	.82
	Sodium sulphate	.28
	Calcium and magnesium phosphates	.25
	Iron oxide and phosphate	.50
	Extractive matters, biliary coloring-matter, gases, and accidental substances	6.4
		1000.0

Percentage Proportion of the Chief Constituents of Blood.

Water	78.4
Red corpuscles (solid residue)	13.0
Albumen of serum	7.0
Inorganic salts	.6
Extractive, fatty, and other matters	.8
Fibrin	.2
	100.0

Casein occurs in cow's milk (*Lac*, B. P.) to the extent of about 4 per cent., dissolved by a trace of alkaline salt. Its solution does not spontaneously coagulate, like that of fibrin, nor by heat like albumen, but acids cause its precipitation from milk in the form of a curd (cheese) containing the fat- (butter-) globules previously suspended in the milk, a clear yellow liquid (or whey) remaining. *Curds and whey* are also produced on adding to milk a piece or an infusion of *rennet*, the salted and dried inner membrane of the fourth stomach of the calf. The exact action of *rennet* is not known, but it seems to be due to the presence of a milk-curdling ferment which is not an acid and not pepsin, and which appears also to occur in the pancreatic juice, the intestinal juice, and some vegetable juices. Respecting *rennet* Soxhlet says: "60 to 80 grammes of calf's stomach, steeped for five days in 1 litre of a 5 per cent. solution of common salt at ordinary temperatures, yield a solution of which one volume will coagulate ten thousand volumes of new milk at a temperature of 95° F. in forty minutes. If the filtered solution is

treated with 60 to 90 grms. more of stomach, a solution of double strength is obtained; another repetition gives a solution three times the strength of the original one. To prevent decomposition, about 0.3 per cent. of thymol may be added to the concentrated rennet-extract solution. Possibly a slight taste due to this may be detected in the finest cheese, but for the same reason oil of cloves is much more objectionable. Boric acid is on all accounts the best antiseptic to employ, and solutions to which it has been added may be kept in covered vessels for months. All extract solutions lose strength on keeping; during the first two months the solution may become 30 per cent. weaker, then the strength remains nearly constant for eight months in the case of a solution of 1 in 18,000. Alcohol is almost as good an antiseptic as boric acid if the solution be preserved in well-stoppered flasks."

Average Composition of 1000 Parts of Milk.

	Specific gravity.	Water.	Solid constituents.	Casein and extractive.	Sugar.	Butter.	Salts.
Woman . .	1.030 to 1.034	870	130	27	60	40	3
Cow	1.030 to 1.035	877	123	40	46	30	7

Leeds puts the average composition of human milk at 2 per cent. of albumenoids, 7 per cent. of milk-sugar, 4 per cent. of fat, and 0.2 per cent. of ash.

Specific gravity alone, as taken by the form of hydrometer termed a *lactometer* or even by more delicate means, is of little value as an indication of the richness of milk, the butter and the other solids exerting an influence in opposite directions. Good cow's milk affords from 10 to 12 per cent. by volume of cream and 3 to 3½ per cent. of butter. The water of milk seldom varies more than from 87 to 88 per cent., and the solid constituents from 13 to 12. Indeed, excluding its butter, milk is curiously regular in composition. The non-fatty solids in the mixed milk of a herd or dairy of healthy cows is almost a constant quantity—namely, 9.3 per cent. A lower proportion of non-fatty solids in a sample of milk points to the addition of water. Thus, supposing that 100 grains of a specimen of milk evaporated to dryness, and all butter extracted from the residue (previously disintegrated by help of 1 or 2 parts of dried gypsum or the dried infusorial earth termed *Kieselguhr*) by ether (or placed on blotting-paper and dried and exhausted by ether—Adams), yielded a non-fatty residue of 7.44 grains, the specimen would probably be four-fifths milk and one-fifth water.* For if 9.3 indicate 100, then 7.44 indicate 80. Occa-

* Soxhlet determines fat by noting the specific gravity of an ethereal solution, and then referring to tables showing percentage of fat in ethereal solutions of varying specific gravity.

sionally, under exceptional circumstances, a sample of genuine milk might be slightly poorer than that from a healthy herd, and therefore, for legal purposes, a standard of 9 per cent. by weight of non-fatty solids and 2.5 per cent. of butter-fat has been proposed. Only in the rare cases of milk containing an unusually large proportion of butter-fat would any milk yielding less than 9 per cent. of non-fatty solids be regarded as genuine. And, again, no milk would be considered genuine, under this standard, if it yielded less than 2.5 per cent. of fat, not even in the rare case of its containing an unusually large proportion of real non-fatty milk-solids. Half-starved cows might yield milk below these standards, but it could scarcely be considered to be normal or better fitted for food than milk watered *after* leaving the cow. If, however, such milk is to be regarded as genuine, the standard of 8.5 of non-fatty solids will not be too low.

Ewe's milk is much richer than either human or cow's milk.

Under the microscope milk is seen to consist of minute corpuscles floating in a transparent medium. These corpuscles consist of the fatty matter (butter) said to be contained in a filmy albumenoid envelope. The fat is fluid at the normal temperature of the animal, and remains so until the milk is well agitated by churning or otherwise or until the milk is frozen.

Legumin, or *vegetable casein*, is found in most leguminous seeds and in sweet and bitter almonds. Peas contain about 25 per cent. of legumin.

Vegetable albumen is contained in many plant-juices, and is deposited in flocculi on heating such liquids. *Vegetable fibrin* is the name given by Liebig and Dumas to that portion of the gluten of wheat which is insoluble in alcohol and ether. *Spongine*, the organic matter of sponge, appears to be a proteid.

Albumenoid substances are nearly identical in percentage composition. Albumen and fibrin contain 53.5 of carbon, 7 of hydrogen, 15.5 of nitrogen, 22 of oxygen, 1.6 of sulphur, and 4 of phosphorus. Casein contains no phosphorus. These three bodies Liebig termed the *plastic elements of nutrition*, under the assumption that animals directly assimilate them in forming muscles, nerves, and other tissues—starch, sugar, and similar matter forming the *respiratory* materials of food, because more immediately concerned in keeping up the temperature of the body by the combustion going on between them and their products and the oxygen of the air in the blood; but the classes are scarcely so well differentiated as the terms would imply.

The whole of the organic nitrogen in food must not, moreover, be regarded as representing true albumenoids, some existing as amidic and similar compounds—bodies having a simplicity of composition characteristic of the *products* of physiological action on food, rather than that complexity of composition characteristic of true nutrients. Albumenoids in decomposing yield much fatty matter as well as other substances. Possibly a portion, at least, of the *adipocere* (*adeps*, fat; *cera*, wax), or *corpse-fat*, characteristic of the remains of buried animals, is thus derived.

Albumenoids are divided, according to their solubility in water and certain saline solutions such as ammonium sulphate, into "albumens" or "albumins," "globulins," "albumoses," "peptones," etc. To the second of these classes the poisons of most venomous snakes probably belong, and a globulin and an albumose, each harmless when swallowed, but extremely poisonous when injected into the blood, occur in the seeds of *Abrus precatorius* (jequirity).

Musk (*Moschus*, U. S. P.), "the dried secretion from the preputial follicles of *Moschus moschiferus*" (the musk deer), is a mixture of albumenoid, fatty, and other animal matters with a volatile odorous substance of unknown composition. "Artificial musk," a synthetic compound having an odor resembling in quality and power that of natural musk, is trinitro-isobutyl-toluene, $C_6HCH_3, C(CH_3)_3(NO_2)_3$.

GELATIGENOUS SUBSTANCES.

These nitrogenous bodies differ, chemically, from the albumenoid in containing less carbon and sulphur and more nitrogen. They are contained in certain animal tissues, and on boiling with water yield a solution which has the remarkable property of solidifying to a jelly on cooling. The tendons, ligaments, bones, skin, and serous membranes afford *gelatin* proper; the cartilages give *chondrine*, which differs from gelatin in composition and in being precipitated by vegetable acids, alum, and lead acetates. The purest source of gelatin is *isinglass* (B. P.) (*Ichthyocolla*, U. S. P.), "the swimming-bladder or sound of various species of *Acipenser*, Linn., prepared and cut in fine shreds." Small quantities are more easily disintegrated by a file than a knife. 50 grains dissolved in 5 ounces of distilled water forms the official "Solution of Isinglass," B. P. Gelatin (*Gelatinum*, B. P.) is officially defined as "the air-dried product of the action of boiling water on gelatigenous animal tissues, such as skin, tendons, ligaments, and bones." *Glue* is an impure variety of gelatin made from the trimmings of hides; *size* is glue of inferior tenacity prepared from the parings of parchment and thin skins. "Among the varieties of gelatin derived from different tissues, and from the same sources at different ages, much diversity exists as to the firmness and other characters of the solid formed on the cooling of the solutions. The differences between isinglass, size, and glue in this respect are familiarly known, and afford good examples of the varieties called weak and strong or low and high gelatins. The differences are sometimes ascribed to the quantities of water combined in each case with the pure or anhydrous gelatin, part of which water seems to be intimately united with the gelatin; for no artificial addition of water to glue would give it the character of size, nor would any abstraction of water from isinglass or size convert it into the hard, dry substance of glue. But such a change is effected in the gradual process of nutrition of the tissues; for, as a general rule, the tissues of an old animal yield a much firmer or stronger jelly than the corresponding parts of a young animal of the same species" (Kirke's *Physiology*).

Gelatin is supposed by some to be a glucoside, yielding an ammonium salt when boiled with diluted acids. It appears to unite chemically with a portion of the water in which it is soaked when used for culinary or manufacturing purposes, for a solution of glue in hot anhydrous glycerin does not yield an ordinary jelly on cooling. From its solution in water gelatin is precipitated by alcohol, corrosive sublimate, platinum perchloride, and by tannic acid. Its aqueous solution is not, like that of albumen, coagulated by heat, nor is it precipitated by acids. By prolonged ebullition its gelatinizing power is destroyed.

PEPSIN.

Pepsin (from πέπω, *pepto*, I digest) is a nitrogenous substance existing in the gastric juice and as a viscid matter in the *peptic* glands and on the walls of the stomachs of animals. It appears to be a modification of a precursor termed *pro-pepsin*, stomachs not yielding so much pepsin when quite fresh as after twenty-four hours. To isolate pepsin, the cleansed mucous membrane of the stomach (of the hog, sheep, or calf, killed fasting) is scraped, and macerated, with the scrapings, in cold water for twelve hours; the pepsin in the strained liquid is then precipitated by lead acetate, the deposit washed once or twice by decantation, sulphuretted hydrogen passed through the mixture of the deposit with a little water to remove the whole of the lead, and the filtered liquid evaporated to dryness at a temperature not exceeding 105° F. Pepsin is a powerful promoter of digestion; its solution is hence frequently termed *artificial gastric juice*. As met with in pharmacy its strength varies greatly. It is often prepared by simply mixing with starch the thick liquid obtained on macerating the scraped stomach with water, and evaporating to dryness. (*Vide Pharmaceutical Journal*, 1865-66, p. 112, and 1871-72, pp. 785 and 843.) The official B. P. process simply consists in scraping the viscid pulp from the slightly washed inner surface of the stomach, and quickly evaporating it to dryness on glass or glazed earthenware at a temperature not exceeding 100° F. The product is powdered.

Official pepsin (*Pepsinum*, U. S. P.) is a yellowish-white to light-brown amorphous powder, sometimes in the form of scales, having a faint but not disagreeable odor, and a slightly saline taste, without any indications of putrescence, being somewhat hygroscopic and insoluble in alcohol, ether, and chloroform.

Valuation of Pepsin.—To 93 cc. of water add almost 2 cc. of dilute hydrochloric acid, and, having brought it to a temperature of 104° F. (40° C.), add 5 cc. of a solution of pepsin (obtained by dissolving 0.067 grm. of pepsin in 98 cc. of water and nearly 2 cc. of dilute hydrochloric acid); the resulting liquid will contain 0.2 cc. (0.21 grm.) of absolute hydrochloric acid, 0.00335 grm. of the pepsin to be tested, and 98 cc. of water.

“Immerse and keep a fresh hen’s egg during fifteen minutes in boiling water; then remove it and place it into cold water. When it is cold separate the white, coagulated albumen, and rub it through

a clean sieve having thirty meshes to the linear inch. Reject the first portion passing through the sieve. Weigh off 10 grms. of the second, cleaner portion, place it in a flask of the capacity of about 200 cc., then add one-half of solution, and shake well, so as to distribute the coherent albumen evenly throughout the liquid. Then add the second half of solution, and shake again, guarding against loss. Place the flask in a water-bath or thermostat kept at a temperature of 38° to 40° C. (100.4° to 104° F.) for six hours, and shake it gently every fifteen minutes. At the expiration of this time the albumen should have disappeared, leaving at most only a few thin, insoluble flakes. (Trustworthy results, particularly in comparative trials, will be obtained only if the temperature be strictly maintained between the prescribed limits, and if the contents of the flasks be agitated uniformly and in equal intervals of time.)—*U. S. P.*

The solvent or digestive action of pepsin on the albumenoids, etc. in the stomach results in a nutritive and digestive fluid termed *peptone*, forming a portion of the whole product of stomach-digestion, or *chyme*. It is thus that such food is prepared for conversion into blood. Artificial or alimentary peptone may be made by digesting blood-fibrin with pepsin in very weak hydrochloric acid. Clermont prepares alimentary peptone in solution by heating 40 parts of minced meat with 30 of water and 1 of sulphuric acid in a sealed tube, filtering, and evaporating the resulting fluid to dryness, and treating the residue with water. The solution is not precipitated by hydrochloric, acetic, or nitric acid; when diluted with strong 90 per cent. alcohol it gives an abundant precipitate; and it is precipitated by tannin, mercuric chloride, and platinic chloride. Peptone is not readily coagulated by heat, and it freely diffuses through membranes. It appears to be isomeric with albumen. *Pro-peptone*, *para-peptone*, or *hemialbumose*, is a mixture of substances intermediate between albumen and peptone. It readily diffuses through membranes. Some vegetables, notably the leaves of the papaw tree, *Carica papaya*, appear to contain a principle, "papain," analogous in properties to pepsin. According to Wurtz, papain is an albumenoid.

Pepsinum Saccharatum, *U. S. P.*, is prepared by triturating 1 part of pepsin and 9 of sugar of milk. "Saccharated pepsin, when tested by the process given above, with the modification that 0.67 gm. of it is to be taken, should digest three hundred times its own weight of freshly coagulated and disintegrated egg albumen."—*U. S. P.*

(For a *résumé* of the modes of preparing pepsin see an article by Petit in the *Pharmaceutical Journal* for July 17, 1880.)

PANCREATIN.

The pancreas (or "sweetbread") secretes a colorless fluid which contains 1½ to 2½ per cent. of an albumenoid substance, or mixture of enzymes, which has the power of converting starch into sugar, and, especially, of emulsifying fat. It may be precipitated by sodium chloride from an acidulated infusion of the pancreas. Stutzer obtains a powerful extract by digesting the pancreas in lime-water and

glycerin with free exposure to air. It is soluble in cold water. An extremely small proportion emulsifies a large volume of fat. Pancreatine is now official (*Pancreatinum*, U. S. P.); it is a yellowish to gray amorphous powder with a faint odor and taste, soluble in water, but not in alcohol. "If there be added to 100 cc. of tepid water contained in a flask 0.28 gm. of pancreatine and 1.5 grms. of sodium bicarbonate, and afterward 400 cc. of fresh cow's milk previously heated to 38° C. (100.4° F.), and if this mixture be maintained at the same temperature for thirty minutes, the milk should be so completely peptonized that, if a small portion of it be transferred to a test-tube and mixed with some nitric acid, no coagulation should occur. Peptonized milk, prepared in the manner just described, or even when the process is allowed to go on to the development of a very distinct bitter flavor, should not have an odor suggestive of rancidity."—U. S. P. The pancreatic juice would seem to contain four distinct ferments—namely, the *emulsifying principle*, the *milk-curdling ferment*, *pancreatic diastase*, and a pepsin-like substance termed *trypsin*, which, unlike pepsin, attacks albumenoids in neutral or even slightly alkaline fluids.

BILE.

Bile (*Fel Bovis*, U. S. P.) is officially the *gall* or *bile* of the ox (*Bos taurus*, Linn.), which, evaporated to one-fourth of its bulk and freed from mucus by agitating with twice its bulk of rectified spirit (in which mucus is insoluble), filtering and evaporating, yields the official Purified Ox-bile (*Fel Bovinum Purificatum*, U. S. P.): the latter has the appearance of a yellowish-green soft resin, but is chiefly composed of two crystalline substances having the constitution of a soap; the one is termed *sodium taurocholate* ($\text{NaC}_{26}\text{H}_{44}\text{NO}_7\text{S}$), the other is *sodium glycocholate*, or simply *cholate* ($\text{NaC}_{26}\text{H}_{42}\text{NO}_6$). Both taurocholates and glycocholates are conjugated bodies, readily yielding, the former *cholic* or *cholalic acid* ($\text{HC}_{24}\text{H}_{39}\text{O}_5$) and *taurine* ($\text{C}_2\text{H}_7\text{NO}_3\text{S}$), the latter *cholalic acid* and *glycocine* or *glycocoll* or *amidacetic acid*, $\text{CH}_2(\text{NH}_2)\text{COOH}$, a soluble crystalline body having interesting physiological relations, for it is obtainable from gelatin (hence the name *glycocoll* or *sugar of gelatin*, from $\gamma\lambda\upsilon\kappa\delta\varsigma$, *glucūs*, sweet, and $\kappa\acute{o}\lambda\lambda\alpha$, *kolla*, glue) and from hippuric acid. Choline ($\text{C}_5\text{H}_{15}\text{NO}_2$) is an alkaloid originally found in bile, hence its name ($\chi\acute{o}\lambda\eta$, *chole*, bile), but it occurs in the brain, etc., in cod-liver oil, and in plants—ergot, Indian hemp, ipecacuanha, etc. (*Vide Index*, "Choline.")

Tests for Bile.—The presence of bile in a liquid, such as urine, may be detected by the following tests: The fluid is gradually mixed with half its bulk of strong sulphuric acid in a test-tube, rise of temperature being prevented by partial immersion of the tube in water. A small quantity of powdered white sugar is then introduced and well mixed with the acid liquid, and more sulphuric acid then poured in; as the temperature rises a reddish or violet coloration is produced. The

cholalic acid liberated in the reaction furnishes the color. This is Pettenkofer's test. It is somewhat interfered with by albumen and volatile oils. Quinlan tests for bile by placing a three-millimetre stratum of the suspected fluid before the slit of the spectroscope, and observing the absorption, which extends, according to the amount present, from the violet of the spectrum to the Fraunhofer line D.

QUESTIONS AND EXERCISES.

In what form is albumen familiar?—Name the chief tests for albumen.—Why is the administration of albumen useful in cases of poisoning?—Mention the points of difference between yolk and white of egg.—From what sources other than egg may albumen be obtained?—In what respects does fibrin differ from albumen?—Enumerate the chief constituents of blood.—How may fibrin be obtained from blood?—State the difference between casein, fibrin, and albumen.—What are the relations of cream, butter, curds and whey, and cheese to milk?—Describe the microscopic appearances of blood and milk.—How much cream should be obtained from good milk?—What is the percentage of water in genuine milk?—Name the sources of vegetable albumen and vegetable casein.—Give the percentage of nitrogen in albumenoid substances.—Describe the chemical nature of musk.—In what lie the peculiarities of gelatigenous substances?—To what extent do isinglass, glue, and size differ?—Whence is pepsin obtained, and how prepared?—Give the proximate constituents of bile.—What are the tests for bile?

COLORING-MATTERS.

The animal, vegetable, and mineral kingdoms abound in substances or pigments which powerfully decompose light, absorbing certain of its constituent colors and reflecting the others. Thus, for example, most leaves contain a body termed chlorophyll, which has the property of absorbing red light and reflecting green; these reflected rays, entering the eye of an observer and striking on the retina (the expanded extremity of the optic nerve), always communicate the same impression to the brain; in popular language the leaf is said to be green. Art has richly supplemented the number of such natural coloring-matters.

YELLOW.—1. *Chrome yellow* occurs in more than a dozen shades. (See Lead Chromate.) 2. *Fustic or yellow wood* is the wood of the *Rhus cotinus*. 3. *Gamboge*. (See Gamboge.) 4. *Ochre* is met with of many tints, under the names of *yellow ochre*, *gold yellow*, *gold earth* or *ochre*, *yellow sienna*, *Chinese yellow*. It is chiefly a mixture of iron oxyhydrates with alumina and lime. It has been used from the earliest times. 5. *Orpiment* is an arsenum sulphide (As_2S_3). 6. *Persian berries*, or *Avignon grains*, contain a yellow principle termed *rhamnin* and other crystalline bodies; they are the product of two or three species of *Rhamnus*. 7. *Purree*, or *Indian yellow*, is said by Stenhouse to owe its color to *magnesium purrate* or *euxan-*

thate ($\text{MgC}_{42}\text{H}_{34}\text{O}_{22}$). 8. *Quercitron* is the bark of *Quercus tinctoria*: it contains the yellow glucoside *quercitrin* ($\text{C}_{36}\text{H}_{38}\text{O}_{20}$). 9. *Rhubarb*. (See Chrysophanic Acid, p. 338.) 10. *Saffron* (*Crocus*, B. P.), the dried stigma and part of the style of *Crocus sativus*, yields *saffranin* or *polychroite*, an orange-red glucoside, which, by the action of dilute acids and by other means, breaks up as shown in the following equation, yielding red *crocin* (Weiss):



Kayser, however, gives the formula of pure crocin as $\text{C}_{44}\text{H}_{70}\text{O}_{28}$, and states that by absorption of water, $7\text{H}_2\text{O}$, it yields pure *croctin*, $\text{C}_{34}\text{H}_{46}\text{O}_9$, and sugar, $9\text{C}_6\text{H}_{12}\text{O}_6$. Any admixture of calcium carbonate, barium or calcium sulphates, or similar powder with saffron is readily detected on placing a little in a glass of warm water and stirring, when insoluble powder is deposited. "Ignited with free access of air, it should yield not more than 6 per cent. of ash" (B. P.). 11. *Turmeric*, the rhizome of *Curcuma longa*, owes its yellow color to *curcumin*, a resin, the formula of which is said by Daube to be $\text{C}_{10}\text{H}_{10}\text{O}_3$, and by Iwanof, $\text{C}_{16}\text{H}_{16}\text{O}_4$. Jackson and Menke state that curcumin is an acid, and that its formula is $\text{H}_2\text{C}_{14}\text{H}_{12}\text{O}_4$. Apparently two yellow pigments are present. The coloring-matters of turmeric are readily dissolved by chloroform—not so those of saffron, mustard, or the best East-Indian rhubarb, on which fact methods of detecting turmeric in those substances have been founded (Howie, *Pharmaceutical Journal*, November 1, 1873). 12. *Weld* (*Reseda luteola*) contains a durable yellow matter termed *luteolin* ($\text{C}_{20}\text{H}_{14}\text{O}_3$). 13. *Picric* or *carbazotic acid* (p. 454) is a very powerful yellow dye. 14. Dried and powdered carrots yield to carbon disulphide a yellow coloring-matter, "*carotin*," which is obtained on evaporating the solvent. It is said to be used in coloring butter.

RED.—1. *Alkanet*, the root of *Alkanna tinctoria*, Tausch, *Anchusa tinctoria*, Desf., yields *anchusin* ($\text{C}_{35}\text{H}_{40}\text{O}_8$), a resinoid matter soluble in oils and fat. 2. *Annatto*, *Arnatto*, or *Arnotto*, a paste prepared by evaporating a strained aqueous extract of the seeds of *Bixa Orellana*, contains *bixin* ($\text{C}_{28}\text{H}_{34}\text{O}_5$), an orange-red, and *orellin*, a yellow principle. 3. *Brazil-wood* (*Cesalpinia Brasiliensis*) furnishes *brezil-in*, $\text{C}_{16}\text{H}_{14}\text{O}_5$, the basis of several lakes; *sapan-wood* and *cam-wood* probably contain the same substance. 4. *Cinnabar*, Chinese red, vermilion, or Paris red, is mercuric sulphide. It is a very ancient red pigment. 5. *Chrome-red*, is lead oxychromate. 6. *Cochineal* (p. 337). 7. *Madder*, the root of *Rubia tinctoria* powdered and treated with sulphuric acid and acidulated water to effect the removal of earthy and other inert matters, furnishes a residual powder termed *garancin*. *Garancin* yields to pure water *alizarin*, $\text{C}_{14}\text{H}_{10}\text{O}_4 \cdot 3\text{H}_2\text{O}$, the red, neutral, crystallizable coloring-matter of madder. *Alizarin* does not exist ready formed in the plant, but is derived by fermentation from a glucoside termed *rubianic acid*. *Alizarin* is now largely produced artificially from *anthracene*, one of the solid constituents of coal-tar (see p. 434). 8. *Mulberry juice* (*Mori Succus*, B. P.) contains a violet-red coloring-matter

which has not been chemically examined. 9. *Red lead* (p. 212). This and the following ochre are very ancient red coloring-matters. 10. *Red ferric oxide*, of shades varying from light to brown-red, is found native. The common names of it are Armenian bole, Berlin red, colcothar, English red, red ochre, burnt ochre, red earth, terra di Sienna, mineral purple, stone-red, and Indian red. 11. *Red sanders-wood* or *red sandal-wood* or *bar-wood* (*Pterocarpus lignum*, B. P.), the billets and chips of *Pterocarpus santalinus*, owes its color to *santalin* ($C_4H_{12}O_4$), a crystalline resinoid matter. Crystalline *pterocarpin*, $C_{10}H_8O_3$, and *homopterocarpin*, $C_{12}H_{12}O_3$, are also present (Cazeneuve). 12. *Red-poppy petals* (*Rhæados petala*, B. P.), from the *Papaver Rhæas*, contain a red coloring principle which has not yet been isolated in a state of purity. The author has sought for morphine in large quantities of the petals, but could not find a trace of that alkaloid. 13. *Red-rose petals* (*Rosæ Gallicæ Petala*, B. P.) and those of the cabbage rose (*Rosæ Centifoliæ Petala*, B. P.) also yield a red substance which has not been analyzed. 14. *Safflower*, *dyer's saffron*, or *bastard saffron*, the florets of *Carthamus tinctorius*, contains an unimportant yellow dye and 5 per cent. of *carthamin* ($C_{14}H_{16}O_7$), an uncrystallizable red dye, the pigment of the old *pink saucers*. Carthamin seems to possess acid characters, and (like silicic acid and other substances) to be soluble in water for a certain time after liberation from its alkaline solution; for fabrics are dyed with safflower by immersion in a bath made of an infusion in dilute alkali neutralized by citric acid immediately before use, the carthamin probably penetrating the cells and vessels of the fibres in a soluble form, there becoming insoluble and imprisoned, and thus giving permanent color to the wool, silk, or other material. Mixed with French chalk, carthamin is used as a cosmetic under the name of *vegetable rouge*—*carmine* being *animal rouge*, and *red oxide of iron* the *mineral rouge*. 15. *Lac-dye* is a cheap form of cochineal, and is also yielded by the species of *Coccus* whose resinous excretion constitutes *lac* (*stick-lac*, *seed-lac*, or *shell-lac*, according to its condition as gathered off the twigs on which it is deposited, or as roughly separated from impurities in seed-like powder or lumps, or as melted and squeezed through bags into shell-like pieces). 16. *Logwood* (*Hæmatoxylon*, U. S. P.) contains a yellow substance, *hæmatoxylin* ($C_{16}H_{14}O_6 \cdot H_2O$ or $3H_2O$), to which any medicinal usefulness of the wood is perhaps due, and which, under the influence of air and alkali or ferments, assumes a very intense red color—*hæmateïn*. Under the influence of ammonia and air also hæmatoxylin yields greenish-violet, iridescent scales of this *hæmateïn* ($C_{16}H_{12}O_3 \cdot 3H_2O$). 17. *Red enamel* colors, for glass-staining and ceramic operations, are produced either by cuprous silicate or purple of Cassius (p. 247).

BLUE.—1. *Cobalt oxide* precipitated in combination or admixture with alumina or calcium phosphate forms *Thénard's blue*, *cobalt-blue*, *Hoffner's blue*, and *cobaltic ultramarine*. 2. *Smalt*, *Saxony blue* or *king's blue*, is rough cobalt glass in fine powder (p. 235). 3. *Copper-blue*, *mountain-blue*, and *English* or *Hambro' blue* are chiefly copper oxycarbonates. 4. *Indigo*, $C_{16}H_{10}N_2O_2$, *sommaruga* (p. 291). 5. *Litmus*, *lichen-blue*, *turnsole*, *orchil* or *archil*, and *cudbear* are

products of the action of air and alkalis on certain colorless principles, as *orcin* ($C_6H_3(OH)_2CH_3$), derived from different species of lichen—*Roccella*, *Variolaria*, and *Lecanora*. 6. *Prussian blue* (p. 341) and *Turnbull's blue* (p. 342) are met with under the names of *Erlangen*, *Louisa*, *Saxon*, *Paris*, or *Berlin blue*. 7. *Ultramarine*, a very old blue pigment, formerly obtained from the rare mineral *lapis lazuli*, is now cheaply made on a large scale by roasting a mixture of fine white clay, sodium carbonate, sulphur, and charcoal or rosin. Its constitution is not well made out. Acids decompose it, sulphuretted hydrogen escaping.

PURPLE.—See Murexid, p. 362.

GREEN.—1. *Cupro-arsenical green* pigments (p. 178). Most of the ancient greens contain copper carbonate. The old *emerald green* is a hydrous chromium oxide, but copper aceto-arsenite is now sold under this name. 2. *Chlorophyll*, *leaf-green*, or *chromule*. A method of extracting chlorophyll is given under "Extracts" (*vide* Index). It is resinoid, soluble in alcohol and ether, insoluble in water, and, according to Frémy and to Schunck, consists of a blue substance, *phyllocyanin* ($C_{34}H_{68}N_4O_{17}$?), and a yellow, *phylloxanthin*: the yellow tints in fading autumnal leaves, he says, are due to the latter principle, the former being the first to fade. Chlorophyll would probably well repay extended investigation, as it has never been obtained pure. 3. *Sap-green*, *buckthorn-*, *vegetable-*, or *bladder-green*, known also as *Chinese green* or *lokas*, is obtained by evaporating to dryness a mixture of lime and the juice (*Rhamni Succus*, B. P., 1867) of the berries of buckthorn (*Rhamnus catharticus*). It is soluble in water, slightly in alcohol, and insoluble in ether and oils. 4. *Green ultramarine* is made by a process similar to that for blue ultramarine. 5. Mixtures of blue and yellow pigments and dyes are common sources of green colors. 6. Glass and earthenware are colored green by chromium oxide and black copper oxide.

BROWN.—1. *Umber*, *sienna*, or *chestnut brown* is found native. By heat it is darkened in tint, and is then known as *burnt umber*. It is a mixture of iron oxide, silica, and alumina. 2. *Sepia* is a dried fluid from the ink-bag of cuttle-fishes (*Sepiadæ*); by its ejection into adjacent water the animal is said to obtain opportunity of escape from enemies. 3. *Catechu* (p. 359) furnishes a brown coloring-matter.

BLACK.—1. *Black-lead* (p. 30), *bone-black* (p. 112), or *ivory-black* and *lamp-black*, the latter a deposited soot from the incomplete combustion of resin and tar, are varieties of carbon. 2. *Burnt sugar* or *caramel* (p. 471). 3. *Indian ink* is usually a dried mixture of fine lampblack and size or thin glue. 4. *Black ink* is essentially tannates and iron gallates suspended in water containing a little gum in solution. 5. *Printer's ink* is well-boiled linseed or other oil mixed with good lampblack, vermilion, or other pigment. 6. *Black dyes* are of the same nature as ink. 7. The old "*pigmentum nigrum*" of black feathers, such as those of the common rook, of dark hair, and probably also of the skin of the negro, is, doubtless, the black substance which remains undissolved when black feathers are

digested for some time in dilute sulphuric acid. It is said to have the formula $C_{18}H_{16}N_2O_8$ (Hodgkinson and Sorby).

WHITE PIGMENTS.—1. *Chalk* or *whiting* (p. 111). 2. *French chalk*, *steatite*, or *soapstone*, is largely magnesium silicate. 3. *Heavy white* (p. 103). 4. *Pearl-white* (p. 252). 5. *Plaster of Paris* (p. 106). 6. *Starch* (p. 472). 7. *White lead* or *Cremnitz white* (p. 210). 8. *Zinc white* or *Chinese white* (p. 134). 9. “*Constant*” *white* is barium tungstate. 10. *Flake white* is basic bismuth nitrate. 11. Tin and zinc oxides and calcium phosphate are employed for giving a white opacity to glass.

ANILINE COLORS.—*Coal-tar Colors*.—Within the last ten years nearly every shade of color seen in the animal and vegetable kingdoms has been successfully imitated by certain dyes and pigments primarily derived from a mineral, coal. Coal distilled for gas furnishes tar or gas-tar. Coal-tar contains some aniline; but especially it contains a liquid convertible into aniline—namely, benzene (C_6H_6), first discovered by Faraday in compressed oil-gas. From aniline, by oxidation, Runge obtained the violet-color reaction, the body producing which Perkin afterward studied and isolated, and manufactured under the name of *mauve*. *Aniline-red* (*fuchsine*, *magenta*, or *roseaniline*), *aniline-yellow*, *aniline-green*, *aniline-blue*, and, in short, aniline dyes, lakes, and pigments of every hue of the rainbow, are now common articles of trade. Their application has revolutionized the arts of the dyer and color-printer. Some of the coal-tar colors are not “aniline” colors, being derived from naphthalene, phthalic acid, phenol, etc.

QUESTIONS AND EXERCISES.

Explain the production of color by the various natural and artificial pigments.—Mention the chief yellow coloring-matters, and describe their chemical nature.—What is annatto? Name the colorific constituent of madder. Can it be made artificially?—State the source of litmus.—Distinguish between prussian blue and Turnbull’s blue, and state how they are manufactured.—How is blue ultramarine obtained? How is it affected by acids?—Describe the chemical nature of the coloring principles of leaves.—By what agents is glass colored green?—Whence is sepia obtained?—Describe the chemistry of black ink.—Write a few sentences on *aniline colors*.

QUALITATIVE ANALYSIS OF SUBSTANCES HAVING UNKNOWN PROPERTIES.

Substances are presented to the analyst in one of the three forms in which all matter exists—namely, solid, liquid, or gaseous—and they may contain animal or vegetable as well as mineral matter.

The method of analysis in the case of *solid mineral bodies* has been described on pp. 370 to 378.

Solid animal or vegetable substances (or mixtures of these with mineral bodies) may be indefinite and beyond the grasp of chem-

istry, or definite and quite within the range of proximate qualitative organic analysis. The presence of such substances is indicated in the preliminary examination of a solid (pp. 371 to 374) by charring and other characters. If no charring occurs and no volatile liquid is expelled by heat, the absence of such matter is indicated. But if organic matter is present, an endeavor is made to ascertain its precise character. The analyst's knowledge of the history of the substance or the circumstances under which it comes into his hands will probably afford a clue to its nature, and enable him to search directly for its proximate constituents. If no such information is at hand, the action of solvents may be employed, as likely to afford indication of the general, if not of the precise, nature of the substance. Water, alcohol, ether, chloroform, carbon bisulphide, each hot and cold, may in turn be agitated with the substance, the mixture be filtered, a portion of the filtrate evaporated, at first partially, setting the product aside, and afterward to dryness, and any deposit or residue be examined with and without the aid of a microscope. Other portions of the filtrate may be treated with acids, alkalies, and solutions of such metallic salts as are commonly used as group-tests for acidulous radicals (p. 369). The action of alkalies, as well as acids, weak and strong, hot and cold, may also be tried on the solid substance itself, and colors, odors, and, in short, any effect whatever, be duly noted. A portion of the substance should also be burnt in an open porcelain crucible until no carbon remains, and the ash, if any, be examined: its amount and nature may afford information leading to the identification of the substance.

The foregoing experiments having been carefully performed and all results entered in the note-book, a little reflection will possibly lead to recognition, or may suggest further direct experiments or confirmatory tests, or will, at least, have pointed to the absence of 90 or 95 per cent. of all possible substances, and thus have restricted the area of inquiry to narrow limits. The success attainable in qualitative proximate organic analysis by the medical or pharmaceutical student will of course largely depend on the thoroughness with which the operator has prosecuted his study of practical chemistry generally; but it also will be considerably affected by the extent to which he has cultivated the art of observation, and the opportunities he has had of acquiring a knowledge of the appearance, uses, and common properties of definite chemical substances and of articles of food, drink, and medicine. The most successful of several good analysts will be the one who has most common sense and most experience.

The pharmaceutical student, who has probably already had some years of experience in pharmacy, occupies an unusually favorable position for prosecuting the proximate analysis of organic and inorganic substances, or, at all events, of that large proportion of such bodies met with in the domain of hygiene and pharmacy. Many substances he will identify at sight or by aid of a lens, or after applying some simple physical or chemical test. Nor should he find much difficulty, after reaching the present point of practical

study, in deciding whether the solid substance under examination belongs to the class of organic acids, organic salts of metallic radicals, alkaloids, salts of alkaloids, amylaceous matter, gums, saccharine substances, glucosides, albumenoid matters, fats, soaps, resins, coloring-matters, etc. For instance, the pharmaceutical student will find less difficulty than the general student in successfully analyzing a substance occurring in "scales," because he has experience of the appearances of compounds commonly produced in that form, and because, even if the appearance is new to him, he knows what kind of substances most readily lend themselves to production in that form. While the general student is testing generally and proceeding cautiously, or searching for general information in books of reference, the pharmaceutical or medical student has incinerated some of the material, noticed whether or not the ash is red (iron) and strongly alkaline (potassium), treated more of the material with an alkali (for ammonium), added excess of ammonia, and examined the precipitate (for cinchonine or quinine), or shaken up the alkaline liquid successively with ether and chloroform, and tested the residue of these decanted and evaporated solvents (quinine, beberine, strychnine), and examined the aqueous solution of the material or one of the filtered alkaline liquids in the usual way for acidulous radicals (citric, tartaric, sulphuric, hypophosphorous). Or he has modified his methods to include search for some "scale preparation" which his special knowledge tells him has been newly introduced to, or is rare in, pharmacy.

In the case of liquids the solvents as well as the dissolved matters claim attention. A few drops are evaporated to dryness on platinum-foil to ascertain if solid matter of any kind is present; the liquid is tested by red and blue litmus-paper to ascertain if free alkalies, free acids, or neither are present; a few drops are heated in a test-tube and the odor of any vapor noticed, a piece of glass tubing bent to a right angle being, if necessary, adapted to the test-tube by a cork, and some of the distilled liquid collected and examined; finally, the usual group-reagents for the several basylous and acidulous radicals are consecutively applied.

Proceeding in this way, the student who has already had some experience in pharmacy will not be likely to overlook such solvents as water, acids, alkalies, alcohol, glycerin, ether, chloroform, benzene, fixed oils, and essential oils, or to miss the substances which these menstrua may hold in solution. He will probably also recognize such liquids as carbolic acid, formic acid, lactic acid, methylic alcohol, aldehyde, aniline, nitrobenzene. He must not, however, suppose that he will always be able to qualitatively analyze, say, a bottle of medicine, for the various infusions, decoctions, tinctures, wines, syrups, liniments, confections, extracts, pill-masses, and powders contain vegetable matters, most of which at present are quite beyond the reach of the analyst. Neither the highest skill in analysis nor the largest amount of experience concerning the odor, appearance, taste, and uses of drugs is sufficient for the detection of all these vegetable matters. Skill and experience combined, however, will do much; and in most cases even so difficult a task as the one just

mentioned may be accomplished with reasonable success. Obviously, qualitative analysis alone will not enable the operator to produce a mixture of substances similar to that analyzed; to this end recourse must be had to quantitative analysis, a subject treated subsequently.

Natural fluids, as "Milk" and "Urine" (*vide* Index), admit of special analytical treatment.

Gas-analysis, or *eudiometry* (from *εἰδία*, *eudia*, calm air, and *μέτρον*, *metron*, a measure, in allusion to the *eudiometer*, an instrument used in measuring the proportion and, as the early chemists thought, the salubrity of the gases of the air), is a branch of experimental investigation, chiefly of a quantitative character, concerning which information must be sought in other treatises. The analysis of atmospheric air from various localities, coal-gas, and gases obtained in chemical researches involves operations which are scarcely within the sphere of Chemistry applied to Medicine. Beyond the recognition, therefore, of oxygen, hydrogen, nitrogen, chlorine, carbonic, sulphurous, nitrous, and hydrosulphuric acid gases, etc., the experimental considerations of the chemistry of gaseous bodies may be omitted. Their study, however, should not be neglected, as existing conceptions of the constitution of chemical substances are largely dependent on the observed relations of the volumes of gaseous compounds to their elements. (See previous paragraphs, pp. 19-30, 42-46, 52-56, and 130.) A useful work on the latter part of the subject is a small book by Hofmann, "Introduction to Modern Chemistry."

Spectrum Analysis.—It may be well to state here that the preliminary and final examinations of minute quantities of solid matter may, in certain cases, profitably include their exposure to a temperature at which they emit light, the flame being physically analyzed by a spectroscope. A spectroscope consists essentially of a prism to decompose a ray of light into its constituent colors, with tubes and lenses to collect and transmit the ray or rays to the eye of an observer. The material to be examined is placed on the end of a platinum wire, which is then brought within the edge of a spirit-lamp or other smokeless flame; volatilization, attended usually in the case of a compound by decomposition, at once occurs, and the whole flame is tinged with a characteristic hue. A flat ribbon of rays is next cut off by bringing near to the flame a brass tube, the cap of which is pierced by a narrow slit. At the other end of the tube, at focal distance for parallel rays, is a lens through which the ribbon of light passes to a prism; the prism decomposes the ribbon, spreading out its constituent colors like a partially opened fan, and the colored beam or *spectrum* thus produced is then examined by help of a telescope attached by a movable joint to a stand which carries the prism and the object-tube. It is this combination of tubes, lenses, and prism or prisms which constitutes the spectroscope. Sodium compounds under the circumstances give yellow light only, indicated by a double band of light in a position corresponding to a portion of the yellow part of an ordinary solar spectrum. The potassium spectrum is mainly composed of a red and

violet band; lithium, a crimson and, at very high temperatures, a blue band. Most of the other elements give equally characteristic spectra.

By passing white light through a colored substance an "absorption spectrum" will be produced which is often characteristic, as in the case of blood or chlorophyll, while by aid of a combined microscope and spectroscope (microspectroscope) drops of colored fluids can be analyzed.

CHEMICAL TOXICOLOGY.

In cases of criminal and accidental poisoning the substances presented to the chemical analyst for examination are usually articles of food, medicines, or vomited matters, or the liver, kidneys, intestines, stomach, and contents, removed in course of post-mortem examination. In these cases some special operations are necessary before the poison can be isolated in a state of sufficient purity for the application of the usual tests; for in most instances the large quantity of animal and vegetable—or, in one word, organic—matter present prevents or masks the characteristic reactions on which the tests are founded. These operations will now be described;* they form the chemical part of the subject of Toxicology (*τοξικὸν*, *toxicon*, poison, and *λόγος*, *logos*, discourse).

Substances occurring apparently as definite salts or unmixed with organic matter need no special treatment. They are analyzed by the ordinary methods already given, attention being restricted to poisonous compounds.

EXAMINATION OF AN ORGANIC MIXTURE SUSPECTED TO CONTAIN MERCURY, ARSENUM, ANTIMONY, LEAD, COPPER, CHROMIUM, OR ZINC; SULPHURIC, NITRIC, HYDROCHLORIC, OXALIC, OR HYDROCYANIC ACIDS; CAUSTIC ALKALIES; PHOSPHORUS; STRYCHNINE, MORPHINE, OR OTHER POISONOUS ALKALOIDS.

Preliminary Examination.

Odor, Appearance, Taste.—Smell the mixture, with the view of ascertaining the presence or absence of any notable quantity of free hydrocyanic acid. Look carefully for any small solid particles, such as arsenic, corrosive sublimate, or verdigris, and for any appearance which may be regarded as abnormal, any character unusual to the coffee, tea, beer, medicine, vomit, coats

* Materials for these experiments are readily obtained for educational purposes by dissolving the poison in infusions of tea or coffee, in porter, or in water to which some mucilage of starch or linseed meal, pieces of bread, potato, and fat have been added.

of stomach, kidney, liver, or other organ, tissue, or solid matter under examination.

Poisonous Quantity of Acid.—Add to a small portion some solution of sodium carbonate, with the view of ascertaining by strong effervescence the presence of any large, poisonous quantity of sulphuric, nitric, or hydrochloric acid (p. 564).

Poisonous Quantity of Alkali.—If so excessively alkaline as to require the addition of a very large quantity of acid before neutralization is effected, a noxious quantity of a corrosive or caustic alkali is present. Whether soda or potash, etc. is ascertained by the usual tests.

Special Instructions may induce the operator to suspect the presence of one particular poison. Direct examination for the latter may then be made, either at once if the substance has an aqueous character, or when filtration or treatment with warm hydrochloric or acetic acid has afforded a more or less colorless liquid.

Fluids.—A vomit or the contents of a stomach, if set aside in a long narrow vessel (test-glass or ale-glass), or, better, exposed on a filter during a night, will often yield a more or less limpid portion at the bottom or top of the solid matter. This fluid (separated by a pipette or otherwise) will sometimes respond to tests without further preparation, and always requires less preparatory treatment than a semi-solid mixture. If none passes through a filter, a portion often collects in the upper part.

General Procedure.—If the preliminary examination does not indicate the method to be pursued, proceed as follows, treating a portion (not more than one-fourth) of the mixture for the poisonous metals, another for the acids, and a third for alkalis, reserving the remainder for any special experiments which may suggest themselves:

Examination for Mercury, Arsenum, Antimony, Lead, Copper, Chromium, Zinc.

If a liquid, acidulate with hydrochloric acid and boil for a short time. If solid or semi-solid, cut up the matter into small pieces, add enough water to form a fluid mixture, stir in 10 or 20 per cent. of ordinary liquid hydrochloric acid, and boil until, from partial aggregation and solution of the solid matter, filtration can easily be effected.

Heat a portion of the clear liquid with a thin piece of bright pure copper or copper gauze, about an inch long and a quarter of an inch broad, for about ten or twenty minutes; metallic

mercury, *arsenum*, or *antimony* will be deposited on the copper, darkening it considerably in color. Pour off the liquid from the copper, carefully rinse the latter with a little cold water, dry the piece of metal by holding it over or near a flame (using fingers, not tongs, or it may become sufficiently hot for a loss of mercury or arsenum to occur by volatilization), introduce it into a narrow test-tube or piece of glass tubing closed at one end, and heat the bottom of the tube in a flame, holding it horizontally, that the upper part of the tube may be kept cool, and partially closing the mouth with the finger to prevent escape of vapor. Under these circumstances any *mercury* will volatilize from the copper and condense on the cool part of the tube in a ring or patch of white sublimate, readily aggregating into visible globules on being pressed by the side of a thin glass rod inserted into the tube; *arsenum* will volatilize from the copper, and, absorbing oxygen from the air in the tube, condense on the cool part of the glass in a ring or patch of white sublimate of arsenic (gray or even darker if much arsenum as well as arsenic be present), not running into globules when rubbed, but occurring in small crystals, the characteristic octahedral form of which (*vide* p. 172) is readily seen by aid of a good hand-lens or the low power of a microscope; *antimony* volatilizes from the copper if strongly heated, and, absorbing oxygen, immediately condenses as a slight white deposit close to the metal.

Confirmatory Tests.—1. Nothing short of the production of globules should be accepted as evidence of the presence of mercury. It will usually have existed as corrosive sublimate. 2. To confirm indications of the presence of arsenum, a portion of the acid liquid may be subjected to the hydrogen tests (pp. 173, 175), or the tube containing the white crystalline arsenic may be broken, and the part on which the sublimate occurs boiled for some time in water, and the hydrosulphuric acid, silver ammonio-nitrate, and copper ammonio-sulphate tests (pp. 176, 177), applied to the aqueous solution. 3. For antimony a portion of the acid liquid must always be introduced into the hydrogen-apparatus with the usual precautions. (*Vide* p. 185.) 4. Any sulphur present may darken the copper, and such stained copper may subsequently yield a whitish sublimate of sulphur on the sides of the subliming tube; such appearances, therefore, are consistent with the entire absence of mercury, arsenum, and antimony.

Note.—Before finally concluding that arsenum is absent from a fluid the latter should be warmed with a little sulphurous acid, and ordinary tests then again applied, for arsenic acid and other arsenates are not readily affected by the usual reagents for arsenum.

For *lead* and *copper* pass hydrosulphuric acid through the

clear acidulated liquid for some time, warming the liquid if no precipitate is produced, or diluting and partially neutralizing the acid by ammonia if much acid has been added. Collect on a filter any black precipitate that may have formed; wash, dissolve in a few drops of aqua regia, dilute, and apply tests, such as ammonia for copper, sulphuric acid for lead, or any other of the ordinary reagents (pp. 193, 215).

Copper may be often at once detected in a small quantity of acidulated liquid by immersing the point of a penknife or a piece of bright iron wire—a deposit of copper, in its characteristic color, quickly or slowly appearing, according to the amount present (p. 193).

Chromium and Zinc.—To the acid liquid through which sulphuretted hydrogen has been passed add excess of ammonia (or to the original acid fluid add excess of ammonia, and then ammonium sulphhydrate); a precipitate falls which may contain alumina, phosphates, chromium, and zinc. (It is usually blackish, from the presence of ferrous sulphide.) Collect the precipitate on a filter, wash, dissolve in a little hydrochloric acid, add a few drops of nitric acid, boil, pour in excess of ammonia, filter, and test the filtrate with ammonium sulphhydrate; a white precipitate indicates zinc. A green precipitate would indicate chromium. Chromates also should be sought (p. 240).

Examination for Mineral Acids, and Oxalic and Hydrocyanic Acids.

To detect *hydrochloric, nitric, or sulphuric acid* in a liquid containing organic matter, dilute with water and apply to small portions the usual tests for each acid, disregarding indications of small quantities. (*Vide* pp. 268, 289, 310.)

Excessive sourness, copious evolution of carbonic acid gas on the addition of sodium carbonate, and abundant evidence of acid on applying the various tests to small portions of the fluid presented for analysis, collectively form sufficient evidence of the occurrence of a poisonous amount of either of the three common mineral acids. Small quantities of the hydrochloric, nitric, and sulphuric radicals, occurring as metallic salts or acids, are common normal constituents of food; hence the direction to disregard insignificant indications. If the fluid under examination be a vomit or the contents of a stomach, and an antidote has been administered, free acid will not be found, but, instead, a large amount of the corresponding salt.

For *oxalic acid*, filter or strain a portion of the liquid, if not already clear, and add solution of lead acetate so long as a

precipitate occurs; collect the precipitate, which in any case is only partly lead oxalate, on a filter, wash, transfer it to a test-tube or test-glass, add a little water, and pass hydrosulphuric acid through the mixture for a short time; the lead is thus converted into the insoluble form of sulphide, while any oxalic acid is set free in the solution. Filter, boil to get rid of hydrosulphuric acid, and apply the usual tests for oxalic acid (see p. 316) to the clear filtrate.

The contents of a stomach containing oxalic acid will often be of a dark-brown color with a tinge of green (altered blood and mucus), and the viscid mixture generally, though slowly, affords some clear, limpid, almost colorless liquid by filtration or on standing.

For *hydrocyanic acid* the three chief tests may be applied at once to the liquid or semi-liquid organic mixture, whether it has an odor of hydrocyanic acid or not. First: Half fill a small porcelain crucible with the material, add 8 or 10 drops of strong sulphuric acid, stir gently with a glass rod, and invert over the mouth of the crucible a watch-glass moistened with a small drop of solution of silver nitrate; a white film on the silver solution is probably silver cyanide, formed by the action of the gaseous hydrocyanic acid on the silver nitrate. Second: Prepare a small quantity of the organic mixture as before, slightly moistening the centre of the watch-glass with solution of potash; here, again, the heat generated by the action of the strong acid is sufficient to volatilize some of the hydrocyanic acid, which, reacting on the potash, forms potassium cyanide. On removing the watch-glass and stirring into it successively solution of a ferrous salt, a ferric salt, and hydrochloric acid, flocks of prussian blue are produced if hydrocyanic acid is present. Third: Proceed as before, moistening the watch-glass with ammonium sulphhydrate; after exposure to the hydrocyanic acid gas for five or ten minutes add a drop of solution of ammonia, evaporate to dryness at a low temperature, and add a drop of hydrochloric acid and of solution of ferric chloride; a blood-red color, due to ferric sulphocyanate, is produced if cyanogen is present.

If the above reactions are not well marked, the organic mixture may be carefully and slowly distilled in a small retort, the neck of which passes into a bottle and dips beneath the surface of a little water at the bottom of the bottle; the reagents may then be applied to separate portions of the distillate.

The examination of organic mixtures for hydrocyanic acid must be made without delay, as the poison soon begins to decompose, and in a day or two is usually destroyed.

Examination for Phosphorus.

A paste containing phosphorus is commonly employed for destroying vermin. In cases of poisoning the phosphorus is generally in sufficient quantity to be recognized by its characteristic unpleasant smell. A stomach in which it occurs not infrequently exhibits slight luminosity if opened in a dark room. When the phosphorus is too small in quantity or too much diffused to afford this appearance, a portion of the material is placed in a flask, water acidulated by sulphuric acid added, a long wide glass tube fitted to the neck of the flask by a cork, and the mixture gently boiled. If phosphorus is present (even 1 part in 2,000,000, according to De Vrij), the top of the column of steam as it condenses in the tube will appear distinctly phosphorescent when viewed in a dark room. From its liability to oxidation phosphorus cannot be detected after much exposure of an organic mixture to air.

Examination for Strychnine and Morphine.

Strychnine.—If solid or semi-solid, digest the matter with water and about 10 per cent. of hydrochloric acid till fluid, filter, evaporate to dryness over a water-bath. If the organic mixture is already liquid, it is simply acidulated with hydrochloric acid and evaporated to dryness. The acid residue is next treated with spirit of wine as long as anything is dissolved, the filtered tincture evaporated to dryness over the water-bath, and the residue digested in water and filtered. This slightly acid aqueous solution must now be rendered alkaline by ammonia, and well shaken in a closed bottle or long tube with about half an ounce of chloroform, and set by till the chloroform has subsided. The chloroform (which contains the strychnine) is then removed by a pipette, the presence of any aqueous liquid being carefully avoided, and evaporated to dryness in a small basin over a water-bath, the residue moistened with concentrated sulphuric acid, and the basin kept over the water-bath for several hours. (It is highly important that the sulphuric acid used in this operation should be free from nitrous compounds. Test the acid, therefore, by adding powdered ferrous sulphate, which becomes pink if nitrous bodies are present. If these are found, the acid should be purified by strongly heating with ammonium sulphate, 70 or 80 grains to a pint.) The charred material is exhausted with water, filtered, excess of ammonia added, the filtrate shaken with about a quarter of an ounce of chloroform, the mixture set aside for the chloroform to separate, and the chloroform again

removed. If on evaporating a small portion of this chloroform solution to dryness, adding a drop of sulphuric acid to the residue, and warming, any darkening of color or charring takes place, the strychnine is not sufficiently pure for chemical detection; in that case the rest of the chloroform must be removed by evaporation, and the residue redigested in warm sulphuric acid for two or three hours. Dilution, neutralization of acid by ammonia, and agitation with chloroform are again practised, and the residue of a small portion of the chloroform solution once more tested with sulphuric acid. If charring still occurs, the treatment must be repeated a third time. Finally, a part of the chloroform solution is taken up by a pipette, and drop after drop evaporated on one spot of a porcelain crucible-lid until a fairly distinct dry residue is obtained. A drop of sulphuric acid is placed on the spot, another drop placed near, a minute fragment of red potassium chromate placed in the second drop, and, when the acid has become tinged with the chromate, one drop drawn across the other; the characteristic evanescent purple color is then seen if strychnine is present. Other tests (*vide* p. 528) may be applied to similar spots.

This is Girdwood and Rogers's method for the detection of strychnine when mixed with organic matter. It is tedious, but trustworthy, and, though apparently complicated, very simple in principle, thus: Strychnine is soluble in acidulated water or alcohol or in chloroform, readily removed from an alkaline liquid by agitation with chloroform, and not charred or otherwise attacked when heated to 212° F. with sulphuric acid; much of the organic matter of the food is insoluble in water; of that soluble in water, much is insoluble in alcohol; and of that soluble in both menstrua, all is charred and destroyed by warm sulphuric acid in a shorter or longer time. (See also Stas's general process, p. 568.)

Morphine, and the Meconic Acid with which it is Associated in Opium.—To the liquid or the semi-fluid mixture, warmed for some time with a small quantity of acetic acid, filtered, and concentrated if necessary, add solution of lead acetate until no further precipitate is produced. Filter and examine the *precipitate* for meconic acid, reserving the *filtrate* for the detection of morphine.

The Precipitate.—Wash the *precipitate* (lead meconate, etc.) with water, place it in a test-tube or test-glass with a small quantity of water, pass hydrosulphuric acid gas through the mixture for a short time, filter, slightly warm in a small basin, well stirring to promote removal of excess of the gas, and add a drop of neutral solution of ferric chloride; a red color, due to the formation of ferric meconate, is produced if meconic acid is present. This color is not destroyed on boiling the

liquid after the addition of one drop of diluted hydrochloric acid, as is the case with ferric acetate, nor is it bleached by solution of corrosive sublimate, thus distinguishing it from ferric sulphocyanate. It is discharged by hydrochloric acid.

The Filtrate.—The solution from which meconic acid has been removed by lead acetate is evaporated to a small bulk over a water-bath, excess of potassium carbonate added, and evaporation continued to dryness. The residue is then treated with alcohol, which dissolves the morphine. The alcoholic solution, evaporated similarly, may leave the morphine sufficiently pure for the application of the usual tests (*vide* p. 518) to small portions of the residue. If no reaction is obtained, add a drop of sulphuric acid and a little water to the residue, and shake with ether, in which the salt of morphine is insoluble. The treatment with ether may be repeated until nothing more is removed, the acid aqueous liquid saturated with potassium carbonate, the mixture evaporated to dryness, the residue digested in alcohol, filtered, and portions of the alcoholic liquid evaporated to obtain spots of morphine for the application of the ordinary tests.

If much organic matter is believed to remain in the filtrate after the lead-acetate treatment, or if a considerable excess of lead acetate has been employed, the filtered liquid should be subjected to a current of sulphuretted hydrogen until no more lead sulphide is precipitated, the mixture filtered, and the filtrate, with the washings from the lead sulphide, evaporated to a small bulk, excess of potassium carbonate added, the whole well mixed and agitated with twice or thrice its bulk of a mixture of ether and acetic ether (ether alone might not dissolve the morphine). On standing, the ethereal liquid rises to the surface: it is carefully removed, evaporated to dryness, and the residue tested or further purified in the manner described in the preceding paragraph.

The examination for morphine must be conducted with great care, and with as large a quantity of material as can be spared; for its isolation from other organic matter is an operation of difficulty, especially when only a minute proportion of alkaloid is present. Fortunately, the detection of meconic acid does not include similar difficulties, and, as its reactions are quite characteristic, its presence is held to be strong evidence of the existence of opium in an organic mixture.

Examination for Other Poisonous Alkaloids.

Stas's Process.—Minutely subdivide any solid matter; to this and the liquid portion of the vomit, etc. add about twice their

weight of the strongest spirit of wine containing sufficient tartaric acid to fairly acidify the mixture. Digest the whole in a flask at a temperature of 150° or 160° F.; set aside to cool; filter. The solution, which will contain the whole of the alkaloid, should then be evaporated nearly to dryness *in vacuo*, or at all events at a temperature not exceeding 100° F., lest volatile alkaloids should be dissipated. The residue is next exhausted with cold anhydrous alcohol, filtered, and the filtrate evaporated to dryness with the precautions already stated. The extract is dissolved in a very small quantity of water, treated with excess of powdered sodium or potassium bicarbonate, and well shaken with five or six times its volume of pure ether (with perhaps a little acetic ether). This ethereal liquid contains the alkaloid. Small portions should be evaporated in watch-glasses and tasted, or tested physically and chemically, according as the knowledge of collateral circumstances by the operator or his experience or such reactions as are recorded on pp. 530, 543 may suggest.

If a volatile alkaloid (conine, nicotine, lobeline, sparteine) is indicated, the ethereal solution, which may contain animal matter, is removed, agitated with aqueous solution of potash, decanted, and shaken with pure diluted sulphuric acid. On standing, the aqueous portion, containing the alkaloid as acid sulphate, subsides; the upper ethereal portion, containing the animal matter, is rejected; the acid aqueous liquid is made alkaline with caustic potash or soda; ether added, well shaken; the ethereal liquid decanted, evaporated to dryness *in vacuo* or at a low temperature, and (to get rid of all traces of ammonia) again moistened with ether and dried. The residue is now tested for the suspected alkaloid by taste, smell, and the application of appropriate reagents (pp. 530-543).

If a non-volatile alkaloid (aconitine, atropine, brucine, colchicine, emetine, hyoscyamine, physostigmine, solanine, veratrine, as well as morphine, codeine, and strychnine, etc.) is indicated, further purify by decanting the ethereal liquid from the lower aqueous solution of sodium bicarbonate, removing the ether by evaporation, digesting the residue in alcohol, filtering, evaporating the alcohol, treating the residue with diluted sulphuric acid, setting aside for a few hours, filtering, concentrating, adding powdered potassium carbonate, and finally anhydrous alcohol. The alcoholic liquid, on evaporation, yields the alkaloid in a fit condition for testing in the manner already stated.

Sonnenschein's Process.—Digest with diluted hydrochloric acid, evaporate to the consistence of syrup, dilute, set aside for

some hours, filter. Add solution of phosphomolybdic acid so long as any precipitate falls or cloudiness occurs; collect the precipitate on a small filter; wash it with water containing phosphomolybdic and nitric acids, and, while still moist, place it in a flask. Decompose this compound of phosphomolybdic acid and alkaloid by adding caustic baryta until the stirred mixture is distinctly alkaline. Distil off volatile alkaloids, condensing and collecting by help of a long tube so bent that the apparatus shall act as a retort, the end of the tube being attached to a bulb or a series of bulbs containing diluted hydrochloric acid. The acid liquid evaporated gives a residue of hydrochlorates of alkaloids. The latter will afford characteristic reactions with the tests for the suspected alkaloid, and, on being moistened with baryta-water and warmed, will afford fumes of volatile alkaloids, the odor of which is usually characteristic. The residue in the flask will contain non-volatile alkaloids. It is treated with carbonic acid gas to neutralize and precipitate the excess of baryta as insoluble barium carbonate; the mixture is evaporated to dryness over a water-bath, and the residue digested in alcohol. The alcoholic solution evaporated generally yields the alkaloids in a fit state for testing.

Reagents for Alkaloids.

Phosphomolybdic acid forms with ammonia, in acid solutions, a remarkably insoluble compound, and it comports itself in a similar manner with those compounds which are analogous to ammonia—the nitrogenized organic bases—consequently forming an excellent reagent for their detection. It may be prepared in the following manner: Ammonium molybdate is precipitated by sodium phosphate; the yellow precipitate, having been washed, is diffused through water and heated with sufficient sodium carbonate to dissolve it. The solution is then evaporated to dryness, and calcined to drive off the ammonia. In case any of the molybdic compound be reduced by this operation, the residue must be moistened with nitric acid and again calcined. The dry mass is then dissolved in cold water, the solution strongly acidulated with nitric acid, and water added until 10 parts of the solution contain 1 of the dry salt. The liquid, which is of a golden-yellow color, must be preserved from ammoniacal fumes. It precipitates all the alkaloids (with the exception of urea) when a mere trace only is present. The precipitates are yellow, generally flocculent, insoluble in water, alcohol, ether, and the diluted mineral acids, with the exception of phosphoric acid. Nitric, acetic, and oxalic acids, concentrated and boiling, dissolve them. These compounds are decomposed by the alkalies, certain metallic oxides, and the alkaline salts, which separate the alkaloid. To give an idea of the sensitiveness of this reagent, it may be stated that 0.000071 gramme of strychnine gives an appre-

ciable precipitate with 1 cubic centimetre of the solution of phosphomolybdic acid.

Phospho-antimonic and phosphotungstic acids are also precipitants of alkaloids. Platinum, iridium, palladium, and gold chlorides are occasionally serviceable. Tannic and picric acids, too, may be used, and a solution of iodine and potassium iodide.

Other special reagents for alkaloids are "Mayer's;" "Nessler's" (see Index); the double potassium and cadmium iodide; and a solution of the double "iodide of bismuth and potassium." The latter is made (by Thresh) by adding together 1 ounce of *Liquor Bismuthi*, B. P., 90 grains of potassium iodide, and 90 grains of strong hydrochloric acid. This orange-colored solution gives a red precipitate with dilute cold solutions containing alkaloids.

Ptomaines (πτῶμα, a corpse) have already been alluded to as including poisonous alkaloids producible from putrefying animal matters, even the human body itself, during the ordinary processes of decay. They are distinguished, according to Brouardel and Boutmy, by a drop or two of a solution of their sulphate converting a drop of solution of potassium ferricyanide into ferrocyanide, the mixture then giving a dark-blue precipitate with a ferric salt. Some other substances also, as morphine, possess this converting power.

Tyrotaxon.—This ptomaine (p. 513) may be isolated and tested as follows: Prepare an aqueous extract of the cheese or filter the coagulated milk, etc. No heat should be applied, and undue exposure to air should be avoided by using stoppered bottles. Make the filtered fluid faintly alkaline with sodium carbonate, and well shake with half its bulk of ether. Allow the perfectly clear ethereal solution to evaporate spontaneously; and, if necessary, again extract this aqueous residue with water, shaking with ether and evaporating as before. The resulting residue may be tested in two or three ways. A little placed on the tongue and swallowed will cause more or less of nausea, vomiting, purging, and headache. Again, the residue is either characteristically crystalline or will become so after standing in a vacuum over sulphuric acid. Mix two or three drops of sulphuric acid and carbolic acid on a white plate, and add a few drops of the aqueous residue just mentioned; if an orange-red or purple color results, the presence of tyrotaxon may be suspected, but any nitrate or nitrite present may cause a similar color. To some of the aqueous residue add an equal volume of a saturated solution of caustic potash; the double potassium and diazobenzene hydrate is then formed, and appears in six-sided plates, whereas any potassium nitrate appears in prisms. This residue may be treated with absolute alcohol, filtered, and the filtrate evaporated, when the plates may again be observed or the color reaction again obtained with this purified product (Vaughan).

Obscure Poisons.—Many substances, the active principles of which are at present beyond the reach of the chemical analyst, are poisons of a more or less active character. (See the *Pharmaceutical Journal* for Sept. 6, 1879, p. 195, and for Dec. 20, 1879, p. 481.)

"Chloral Hydrate," p. 486, and "Chloroform," p. 399, are now

included in "Part 2 of Schedule A" as "poisons within the meaning of the Pharmacy Act, 1868." "Cantharides" is in "Part 1." Its active principle is isolated as described on p. 422, and is recognized by its blistering action on any thin spot on one's skin.

ANTIDOTES.—*Vide* "Antidotes" in the Index.

QUESTIONS AND EXERCISES.

In examining food and similar matter for poison, why must not the ordinary tests for the poison be at once applied?—What preliminary operations should be performed on a vomit in a case of suspected poisoning?—How would you search for corrosive sublimate in wine?—By what series of operations would you satisfy yourself of the presence or absence of arsenic in the contents of the stomach?—Describe the treatment to which decoction of coffee should be subjected in testing it for tartar emetic.—State how the occurrence of lead in water is demonstrated.—Give a process for the detection of copper in jam.—How would you detect zinc in a vomit?—How may the presence of much sulphuric acid in gin be proved?—In testing ale for nitric acid what reactions would you select?—Show how you would conclude that a dangerous quantity of hydrochloric acid had been added to cider.—Describe the manipulations necessary in testing for hydrocyanic acid in the contents of a stomach.—By what method is oxalic acid discovered in infusion of coffee?—How is phosphorus detected in organic mixtures?—Give the process by which strychnine is isolated from a vomit.—Mention the experiments by which the presence of laudanum in porter is demonstrated.—Name the antidotes in cases of poisoning by—*a*, alkaloids; *b*, antimonials; *c*, arsenic; *d*, barium salts; *e*, copper compounds; *f*, hydrochloric acid; *g*, hydrocyanic acid; *h*, lead salts; *i*, corrosive sublimate; *j*, nitric acid; *k*, oxalic acid; *l*, silver salts; *m*, oil of vitriol; *n*, tin liquors; *o*, zinc salts; *p*, carbolic acid.

EXAMINATION OF MORBID URINE AND CALCULI.

The various products of the natural and continuous decay of animal tissue and the refuse matter of food are eliminated from the system chiefly as fæces, urine, and expired air. Air exhaled from the lungs carries off from the blood much carbon (about eight ounces in twenty-four hours) in the form of carbonic acid gas, and some aqueous vapor—the latter, together with a small amount of oily matter, also escaping by the skin. Directing the breath to a cold surface renders moisture evident, and breathing through a tube into lime-water demonstrates the presence of a considerable quantity of carbonic acid gas. The fæces consist mainly of the insoluble *débris* of the system, the soluble matters and water forming the urine. These excretions vary considerably, according to the food and general habits of the individual and external temperature. But in disease the variations become excessive; hence their detection by the medical practitioner, or by the pharmacist for the medical man, is a matter of importance.

An analysis of fæces or air cannot be made with sufficient ease and rapidity to be practically available in medical diagnosis. But with regard to urine, certain abnormal substances and abnormal quantities of normal constituents may be chemically detected in the course of a few minutes by any one having already some knowledge of chemical manipulation.

Healthy human urine contains, in 1000 parts, 957 of water, 14 of urea, 1 of uric acid, 15 of other organic matter, and 13 of inorganic salts. The amount passed in twenty-four hours varies from two to three pints in an adult, and its specific gravity, if healthy, will range from 1.015 to 1.025. Any considerable deviation from these limits would suggest a possibly pathological condition. The average amount of solid matter passed by the urine in one day is $1\frac{1}{2}$ to 2 ounces.

PHYSICAL EXAMINATION OF URINE.

Normal urine is either of a pale-yellow color or faintly reddish-yellow, due to a pigment termed *urobilin*. Urine has a reddish-brown tint if blood is present or greenish-brown if bile is present. Both color and odor are much influenced by certain kinds of food and by some drugs.

Fresh urine is clear. Any turbidity may be due to urates, phosphates, or pus. Urates redissolve when the urine is warmed, phosphates by the addition of acetic acid; pus is detected by the microscope (*vide infra*). If the urine be turbid from the presence of phosphates when first voided, it may be due to conversion of urea into ammonium carbonate, which precipitates the phosphates within the bladder, in which case the fresh and warm urine will effervesce slightly on the addition of acetic acid. This condition is abnormal.

On standing, healthy urine commonly gives a slight cloud of mucus, and after severe exercise or after a hearty nitrogenous meal may give a sediment of urates.

The specific gravity of urine should be taken on a specimen removed from the whole bulk excreted in twenty or twenty-four hours. Many qualitative experiments and all quantitative operations should only be performed on the mixed urine of twenty-four hours.

Healthy urine when fresh is always slightly acid, the acidity being said to be due to the presence of acid sodium phosphate. Alkalinity is probably due to that conversion of urea into ammonium carbonate within the bladder already described.

EXAMINATION OF MORBID URINE FOR ALBUMEN, SUGAR, BILE, EXCESS OF UREA, OR DEFICIENCY OF CHLORIDES; AND URINARY SEDIMENT FOR URATES (OR LITHATES), PHOSPHATES, CALCIUM OXALATE, AND URIC ACID.

Albumen.—To detect albumen, acidulate a portion of the clear urine in a test-tube with a few drops of diluted nitric acid (to keep phosphates in solution), and boil; flocks or coagula

will separate if albumen be present. To detect small quantities, nearly fill a long test-tube with clear urine (filtered, if necessary) and faintly acidulated with acetic acid; then, holding the tube by its lower end, boil the upper portion of the urine.—A cloudiness in the boiled portion, which, on addition of a few drops of acetic acid, does not disappear, indicates the presence of albumen.—Or heat a little nitric acid in a test-tube, and carefully pour down the side a little of the urine, so as to overlies the acid. If albumen be present, a whitish ring or coagulum will, sooner or later, be formed at the junction of the fluids.

These experiments should first be made on normal urine containing a drop or two of solution of white of egg. The coagulum is white if it is only albumen, greenish if bile-pigment be present, and brownish-red if the urine contain blood. The influence of acids and alkalies on the precipitation of albumen is noticed on p. 545.

A saturated solution of picric acid at once precipitates any albumen from urine. Should the urine be alkaline, it must be acidified before applying this test. On warming the mixture the precipitate will become more pronounced if due to the albumen or globulin of blood or to any modifications of albumen caused by acidity or alkalinity of urine, but will disappear if due to peptone or pro-peptone. Potassium ferrocyanide also will precipitate the former varieties of albumen, but not the peptones.

The occurrence of albumen in the urine may be temporary and of but little importance, or it may indicate the existence of a serious affection known as Bright's disease. "Albumenuria is rarely a serious condition unless it is sufficiently pronounced to be made out by the cold nitric-acid test" (Steward).

For quantitative purposes Esbach employs the picric test, dissolving 10 parts of picric acid and 20 of citric acid in 900 of water by aid of heat, and, when the solution is cold, diluting with water to 1000 parts. This solution is added to a given volume of urine in a graduated Cetti's Esbach tube, and the height of the precipitate is noted after twenty-four hours. Johnson finds a simple solution of 5 grains of picric acid in 1 fluidounce of water better than Esbach's solution, because the excess of acid in the latter tends to precipitate much uric acid which would be reckoned as albumen. A standard value is given to the solution in the first instance by washing, drying, and weighing the albumen.

Sugar.—To a portion of the clear urine in a test-tube add five or ten drops of solution of copper sulphate; pour in solution of potash or soda until the precipitate first formed is redissolved; slowly heat the solution to near the boiling-point; a yellow, yellowish-red, or red precipitate (cuprous oxide) is formed if sugar be present. (The production of a rose-red or pink tint with the cold alkaline copper solution indicates the

presence of the altered non-coagulable albumenoids termed *peptones*.)

This experiment should be first made on urine containing a drop or two of solution of grape-sugar (p. 468). The copper hydrate precipitated by the alkali is insoluble in excess of pure potash or soda, but readily dissolves if organic matter, especially sugar, be present. The copper salt should not contain iron.

Other tests may be applied if necessary. (*Vide* p. 470. See also "Sugar, Quantitative Estimation of," in Index.)

A minute amount of sugar is said to occur in normal urine, and a distinct trace is occasionally present. In searching for small quantities, uric acid, which also reduces the copper solution, should first be removed. (See p. 361.) Normal urine rotates a ray of plane polarized light slightly to the left, but if even a small amount of sugar be present marked dextro-rotation results. In larger quantities (often 5 per cent.) sugar is a characteristic constituent of the urine of diabetic patients, greatly increasing the specific gravity of the secretion. Small hydrometers (termed *urinometers*) are commonly employed for quickly and readily ascertaining the specific gravity of urine; they range from 1.000 to 1.050, the interval of 1.015 to 1.025 being marked as "H.S." or "healthy state." (*Vide* "Specific Gravity" and "Hydrometers" in Index; also "Sugar, Quantitative Estimation of.")

Bile.—This is detected by the dark greenish-brown color of the urine and by the general test (Pettenkofer's, or, still better, Quinlan's) described on p. 552. Or a little of the urine may be placed on a white plate, and strong nitric acid, containing some nitrous acid, dropped on it; a peculiar play of colors—green, yellow, violet, etc.—occurs if (the coloring-matter of) bile be present (Gmelin). In doubtful cases the urine should be thoroughly shaken up with a little chloroform, which dissolves the bile-pigments, and the acid test applied to the separated chloroform. Oliver recommends that the urine be diluted to a sp. gr. of 1.008, and then one volume be added to three volumes of the following reagent, when more or less opalescence will be produced, according to the amount of bile acids present. For the reagent dissolve 30 grains of flesh peptone, 4 grains of salicylic acid, and 33 minims of official acetic acid in 8 ounces of water; filter.


Excess of Uric Acid.—A rough quantitative process consists in applying the qualitative method already described (p. 361) to a known volume of urine, and collecting on a filter, washing, and weighing the resulting uric acid. The result is always low. Hopkins saturates the urine with ammonium chloride, and after a couple of hours decomposes the separated ammonium urate by hydrochloric acid, and collects, washes, dries, and weighs the resulting uric acid. (*Vide Proceedings of the Royal Society*, vol. lii. p. 93.)

Excess of Urea.—About one-third of the solid matter in the

urine is urea. Its proportion varies considerably, but $1\frac{1}{2}$ per cent. may be regarded as an average amount. Concentrate urine slightly by evaporation in a small dish, pour the liquid into a test-tube, set the tube aside till cold or cool it by letting cold water run over the outside, add an equal bulk of strong nitric acid, and again set aside; scaly crystals of urea nitrate are deposited more or less quickly.

With regard to the amount of urea in urine it is impossible to sharply define excess or deficiency. If nitric acid gives crystals without concentration, excess is certainly present in the sample examined, though, if the amount of urine passed in the twenty-four hours is much below the average, the quantity of urea excreted may not be abnormal. A rough estimate may be formed by mixing a few drops of the urine and acid on a piece of glass and setting aside; the time which elapses before crystals form is an indication of the quantity in the specimen. The time will vary according to the temperature and state of moisture of the atmosphere, but with care some useful comparative results may in this way be obtained.

For trustworthy quantitative estimations the urine is shaken with an alkaline solution of recently prepared sodium hypobromite, and the nitrogen then liberated collected and measured. The reaction is of the following character:




The whole of the nitrogen of the urea, however, is not evolved, while, on the other hand, some of the produced nitrogen is yielded by the uric acid, hippuric acid, and creatinine of the urine. Neither fact is of any consequence in this examination of urine, for a given specimen of urine always yields the same quantity of gas; hence a given volume of gas always indicates the same percentage of urea; and once a measuring-tube is so divided as to indicate percentages of urea when a given volume of urine is used, it may be trusted to indicate the varying percentages of urea in the urine of one patient or the different proportions of urea in the urine of different patients. The method is Davy's, with improvements and modifications of apparatus by Knop, Huefner, Russell and West, Apjohn, Dupré, Gerrard, Gillet, and others. In the chemical laboratory appliances already at hand may be adapted for the operation. Thus, as shown in the woodcut on p. 577, any two-ounce or three-ounce bottle serves for the reaction between the hypobromite and urine; a 50 or 60 cc. burette, containing water, may be the measuring-tube; while a funnel, supported in the ring of a retort-stand, serves both as a reservoir for water displaced by the evolved nitrogen and as a means of getting that equal level of water within and without the measuring-tube which shall prevent misleading attenuation or compression of the nitrogen. Attach the funnel to the bottom of the burette by india-rubber tubing. Attach a short glass tube by a pierced cork to the top of the burette, and by more india-rubber

tubing connect this glass tube with a similar tube in the well-fitting india-rubber cork of the gas-generating bottle. Disconnect the latter. Into the funnel pour water until it rises to the zero-mark of the burette and a little water remains in the bottom of the funnel. Into the generating-bottle pour about 25 cc. of solution of soda (made by dissolving about 100 grms. of solid soda in 250 cc. of water) and about 2.5 cc. of bromine. Into the bottle containing the hypobromite solution thus prepared a short test-tube containing 5 cc. of urine is lowered, care being taken that no urine is spilt. The cork is re-inserted and the water-level again adjusted (best accomplished if into the cork of the generating bottle be fitted a short glass tube, the external orifice of which can be closed by a cork or a cap as soon as the apparatus is ready for use). The generating bottle is now inclined, when, the urine and hypobromite mixing, nitrogen is at once evolved (the carbonic acid produced at the same time being absorbed by the strongly alkaline fluid in the bottle). The funnel is lowered until the surfaces of the water inside and outside the measuring-tube are on a level; after ten or fifteen minutes the level is finally adjusted and the amount of produced gas noted. Every 55 cc. of gas indicates 0.15 of a gm. of urea, the temperature being about 66° F. and the height of the barometer being about thirty inches. In cases in which frothing interferes put a fragment of suet into the generating bottle.

Instead of sodium hypobromite, the hypochlorite, which is more stable, may be employed (Squibb).

Tests.—Urea in solution in water may be detected by the reaction with nitric acid, and by the readiness with which it yields ammonia on being boiled with alkalis. In putrid urine its conversion into an ammoniacal salt has already been effected by ammoniacal fermentation.



This transformation of the urea into ammonium carbonate is due to the action of a special ferment belonging to the genus *Torula*, formed of chaplets of globules similar in form to, but much smaller than, those of beer-yeast. It occurs as a white deposit in the urine. If some of this deposit be added to a saccharine solution containing urea, it rapidly multiplies, ammonium carbonate being formed.

Formula of Urea.—The empirical formula of urea is $\text{CH}_4\text{N}_2\text{O}$. Its rational formula may be thus written:





Fig. 50.



that is, it may be regarded as carbamide or as one of the organic bases already referred to—a primary diamine, in which the bivalent radical CO occupies the place of H_2 . The other atoms of hydrogen may be displaced by various radicals, and many *compound ureas* thus be obtained.

Artificial Urea.—Urea may be prepared artificially by Williams's modification of Wöhler's method. Potassium cyanide of the best commercial quality (containing about 90 per cent. of real cyanide) is fused at a very low red heat in a shallow iron vessel; red lead is added in small quantities at a time, the temperature being kept down by constant stirring. When the red lead ceases to cause further action, the mixture (lead and potassium cyanate) is allowed to cool, the product finely powdered, exhausted with cold water, barium nitrate added till no more precipitate (barium carbonate) falls, the mixture filtered, and the filtrate treated with lead nitrate so long as lead cyanate is thrown down. The latter is thoroughly washed, and dried at a low temperature. Equivalent quantities of lead cyanate and ammonium sulphate, digested in a small quantity of water with a little heat (*vide* p. 339) and filtered, yield a solution from which urea crystallizes on cooling.

Another Process.—Basaroff has found that urea is produced when ordinary ammonium carbonate is heated in *strong* hermetically sealed tubes to about $275^{\circ} F.$ for a few hours. The same chemist had previously obtained urea by similarly heating pure ammonium carbamate; so that the source of the urea in the former case is probably the ammonium carbamate believed to occur in the carbonate. (See p. 93.)



Deficiency of Chlorides.—Any given bulk of urine in a test-tube yields, of course, abundance of flocks of silver chloride on the addition of nitric acid and silver nitrate. Any markedly smaller bulk points to pathological conditions, such as those of acute fever.

Chromogens.—Urine may contain chromogens, which are substances which do not at the time color the urine, but which, on the addition of oxidizing reagents or after standing some time, develop a color. A blue color may be seen in urine on the addition of much nitric acid. This is due to the formation of indigo from its chromogen. The darkening often noticed on the addition of acid to urine is due to liberation of the pigment urobilin from its chromogen. Care must be taken not to confound the color-changes in urine due to the action of drugs with the effects produced by the action of oxidizing agents on chromogens. Thus rhubarb and santonin darken the natural yellow of urine, the addition of an alkali causing a red coloration. Carbolic acid taken internally makes the urine greenish-black, resembling urine with much bile in it. If potassium iodide or bromide is being taken internally, the addition of a strong acid will often cause separation of iodine or bromine, respectively, in the urine.

URINARY SEDIMENTS.

Warm the sediment with the supernatant urine, and filter.

<i>Insoluble.</i>		<i>Soluble.</i>	
Phosphates, calcium oxalate, and uric acid. Warm with acetic acid, and filter.		Ammonium, calcium, or sodium urates; chiefly the latter.	
<i>Insoluble.</i>		<i>Soluble.</i>	
Calcium oxalate and uric acid. Warm with hydrochloric acid, filter.		Phosphates. Add ammonia; white ppt. = calcium phosphate or ammonio-magnesium phosphate, or both.	
<i>Insoluble.</i>	<i>Soluble.</i>		
Uric acid. Apply the murexide test (p. 362).	Calcium oxalate. May be reprecipitated by ammonia.	They are redeposited as the liquid cools, and if sufficient in quantity may be further examined for ammonium, calcium, sodium, and the uric radical by the appropriate tests.	

Notes.—Urinary deposits are seldom of a complex character: the action of heat and acetic and hydrochloric acids generally at once indicates the character of the deposit, rendering filtration and precipitation unnecessary.

The Urates are often of a pink or red color, owing to the presence of a pigment termed *purpurin*; hence the common name of *red gravel* for such deposits. *Purpurin* is soluble in alcohol, and may be removed by digesting a red deposit in that solvent. It is seldom necessary to determine whether the urate be that of ammonium, calcium, or sodium. (See, also, Uric Acid, p. 368.) The deposited urate is a very acid urate, which slowly (more rapidly in urine diluted with water) breaks up into a less acid urate and uric acid (Bence Jones). The occurrence in the urine of the salines of our food, especially of dipotassic phosphate, is apparently (Roberts) what prevents this decomposition before the urine is exposed to the air.

Calcium phosphate and *ammonio-magnesium phosphate* are usually both present in a phosphatic deposit, the magnesium salt forming the larger proportion. They may, if necessary and if sufficient in quantity, be separated by collecting on a filter, washing, and boiling with solution of sodium carbonate. The calcium and magnesium carbonates thus formed are collected on a filter, washed, and dissolved in a drop or two of hydrochloric acid; ammonium chloride, ammonia, and ammonium carbonate are added, and the mixture boiled and filtered; any calcium originally present will then remain

insoluble as calcium carbonate; while any magnesium will be precipitated from the filtrate as ammonio-magnesium phosphate on the addition of sodium phosphate, the mixture being also well stirred. —The chief portion of excreted phosphates is carried off by the fæces, that remaining in the urine being kept in solution by the influence of acid sodium phosphate and, frequently, lactic acid. —Occasionally, an hour or two after a hearty meal, the urine becomes sufficiently alkaline for the phosphates to be deposited, and the urine when passed is turbid from their presence. —The ammoniacal constituent of the magnesium salt does not occur normally, but is produced from urea as soon as urine becomes alkaline.

Calcium oxalate is seldom met with in excessive amounts, but very often in small quantities mixed with phosphates. In one case of oxaluria the whole urine excreted by a patient in twenty-four hours furnished to the author only two-thirds of a grain of calcium oxalate.

Free uric acid is in most cases distinctly crystalline, and nearly always of a yellow, red, or brown color.

Artificial Sediments.—For educational practice these may be obtained as follows: 1. Rub up in a mortar a few grains of serpent's excrement (chiefly ammonium urate) with an ounce or two of urine; this represents a sediment of urates. 2. Add a few drops of solution of ammonia or solution of ammonium carbonate to urine; the deposit may be regarded as one of phosphates. 3. To an ounce or two of urine add very small quantities of calcium chloride and ammonium oxalate; the precipitate is calcium oxalate. 4. To urine acidulated by hydrochloric acid add a little serpent's excrement; the sediment is uric acid.

Other deposits than the foregoing are occasionally observed. Thus, *hippuric acid* ($\text{HC}_9\text{H}_8\text{NO}_3$), a normal constituent of human urine and largely contained in the urine of herbivorous animals, is sometimes found associated with uric acid in urinary sediment, especially in that of patients whose medicine contains benzoic acid (p. 340). Its appearance, as observed by aid of the microscope, is characteristic—namely, slender, four-sided prisms having pointed ends. *Cystin*, $\text{C}_3\text{H}_7\text{NSO}_2$ (from κύστις, *küstis*, a bladder, in allusion to its origin), rarely occurs as a deposit in urine. It is not soluble in warm urine or dilute acetic acid, and scarcely in diluted hydrochloric acid; hence would be met with in testing for free uric acid. It is very soluble in ammonia, recrystallizing from a drop of the solution placed on a piece of glass in characteristic microscopic six-sided plates. *Leucine* and *tyrosine* (see p. 513) occur in cases of phosphorus-poisoning and of acute yellow atrophy of the liver. *Organized sediments* may be due to the corpuscles of pus, mucus, or blood, fat-globules, spermatozoa, cylindrical casts of the tubes of the kidneys, epithelial cells from the walls of the bladder, or foreign matters, such as fibres of wool or of cotton or wood, small feathers, dust, starch, etc.; these are best recognized by the microscope. (See the accompanying figures and the following paragraphs on the microscopic appearances of both crystalline and organized urinary sediments.)


MICROSCOPIC EXAMINATION OF URINARY SEDIMENTS.

Urine containing insoluble matter is usually more or less opaque. For microscopical examination a few ounces should be set aside in a conical test-glass for an hour or two, the clear supernatant urine poured off from the sediment as far as possible, a small drop of the residue placed on a slip of glass and covered with a piece of thin glass, and examined under the microscope with different magnifying powers.

The respective appearances of the various crystalline and organized matters are given in Figs. 51-62, which were kindly drawn by the late H. B. Brady, F. R. S., from natural specimens in the collections of St. Bartholomew's Hospital, Dr. Sedgwick, the late Mr. W. W. Stoddart, Mr. Waddington, and the author.

Uric acid occurs in many forms, most of which are given in Figs. 51 and 52. Flat, more or less oval crystals, sometimes attached to each other, their outline then resembling an 8, a cross, or a star, are common. Single and grouped quadratic prisms, aigrettes, spicula, and crystals recalling dumb-bells are met with. From urine

Fig. 51.



Uric Acid.

Fig. 52.



Uric Acid.


acidulated by hydrochloric acid square crystals, two opposite sides smooth and two jagged, are generally deposited; acidulated by acetic acid, more typical forms are obtained. A drop of solution of potash or soda placed on the glass slip will dissolve a deposit of uric acid, a drop of any acid reprecipitating it in minute but characteristic crystals.

Cystin is very rarely met with as an urinary deposit; that from which Fig. 53 was taken was found in the urine of a patient in St. Bartholomew's Hospital. Lamellæ of cystin always assume the hexagonal character, but the angles are sometimes ill defined and the plates superposed: in the latter case a drop of solution of ammonia placed on the glass at once dissolves the deposit, well-marked six-sided crystals appearing as the drop dries up.

Triple phosphate (magnesium and ammonium phosphate) is de-


posited as soon as urine becomes alkaline, the ammoniacal constituent being furnished by the decomposition of urea. It occurs in large prismatic crystals, forming a beautiful object when viewed by

Fig. 53.



Cystin.

Fig. 54.



Triple Phosphate.

polarized light—sometimes also in ragged stellate or arborescent crystals, resembling those of snow. Both forms may be artificially prepared by adding a small lump of ammonium carbonate to a few ounces of urine and setting aside in a test-glass (Fig. 54).


Amorphous deposits are either *earthy phosphates* (a mixture of magnesium and calcium phosphates) or calcium, magnesium, ammonium, potassium, or sodium *urates*—chiefly the latter. They may be distinguished by the action of a drop of acetic acid placed near the sediment on the glass slip, the effect being watched under the microscope; phosphates dissolve, while urates gradually assume characteristic forms of uric acid. Urates redissolve when warmed with the supernatant urine.

Sodium and magnesium urates, though generally amorphous, occasionally take a crystalline form—bundles or tufts of small needles—as shown in Figs. 55 and 56. When pink or brick-red the color is due to *uroerythrin*.

Calcium oxalate commonly occurs in octahedra, requiring high magnifying power for their detection. The crystals are easily overlooked if other matters are present, but are more distinctly seen after phosphates have been removed by acetic acid. In certain aspects the smaller crystals look like square plates traversed by a cross. A dumb-bell form of this deposit is also sometimes seen, resembling certain forms of uric acid and the coalescing spherules of a much rarer sediment—calcium carbonate. Calcium oxalate is insoluble in acetic, but soluble in hydrochloric acid. The octahedra are frequently met with in the urine of persons who have partaken of garden rhubarb and certain other vegetables. The crystals may often be deposited artificially (according to Waddington) by dropping a fragment of oxalic acid into several ounces of urine and setting aside for a few hours.

Calcium carbonate is rarely found in the urine of man, but frequently in that of the horse and other herbivorous animals. Human urine containing calcium carbonate often reddens litmus-paper, and it is only after the removal, on standing, of the excess of carbonic


Fig. 55.



Urates.
a, of Sodium;
b, of Magnesium.

} Calcium Oxalate.

Fig. 56.



Calcium Carbonate. Hippuric Acid.

acid that the salt is deposited. It consists of minute spherules, varying in size, the smaller ones often in process of coalescence. The dumb-bell form thus produced is easily distinguished from similar groups of uric acid or calcium oxalate by showing a black cross in each spherule when viewed by polarized light. Acetic acid dissolves calcium carbonate, liberating carbonic acid gas, with visible effervescence (under the microscope) if the slide has been previously warmed and a group of crystals be attacked.

Hippuric Acid.—The pointed rhombic prisms and acicular crystals are characteristic and easily recognized. The broader crystals may possibly be mistaken for triple phosphate, and the narrower for certain forms of uric acid; but insolubility in acetic acid distinguishes them from the former, and solubility in alcohol from the latter. These tests may be applied while the deposit is under microscopic observation. An alcoholic solution of hippuric acid, evaporated to dryness and the residue treated with water, gives a solution from which characteristic crystalline forms of hippuric acid may be obtained on allowing a drop to dry up on a slip of glass.

The organized deposits in urine entail greater care in their determination, and usually require a higher magnifying power for their proper examination, than those of crystalline form. The figures are drawn to 230 diameters. The following note will assist the observer:

Casts of uriniferous tubuli are fibrinous masses of various forms, and often of considerable length—sometimes delicate and transparent, occasionally granular, and often beset with fat-globules. Epithelial debris is frequently present in urine in the form of


nucleated cells, regular and oval when full, but angular and unsymmetrical when partially emptied of their contents—sometimes perfect, but more frequently a good deal broken up. Casts are very readily discovered by the use of the microscope if, to a sample

Fig. 57.



Epithelial Cells and Tubuli.

Fig. 58.



Blood-corpuscles.

of the urine supposed to contain them, best in a conical glass, a few drops of an aniline dye be added. "Carbo-fuchsine" answers well. The casts rapidly stain, and are then quite easily seen in the field (Fig. 57).

Blood is easily recognized. Urine containing it is usually high-colored or "smoky," and the corpuscles appear under the microscope as reddish circular disks, either single or laid together in strings resembling piles of coin (Fig. 58). Their color and somewhat smaller size serve to distinguish them from pus-corpuscles. In doubtful cases a minute drop of blood taken from a finger by help of a needle should be diluted with water and used for comparison. After urine containing blood has stood for some time, the corpuscles lose their regular rounded outline and become crenated. (See *a* in Fig. 58.)


Dr. Day of Geelong tests for blood in urine or in stains on clothing by employing a recently-prepared alcoholic solution of the inner unoxidized portions of guaiacum resin and an aqueous or ethereal solution of hydrogen peroxide, when a dull-blue color results. In the case of urine add to a drachm or two in a test-tube nearly as much of the ethereal fluid, and then two or three drops of the guaiacum tincture; on gently agitating the tube a bluish-green layer appears at the junction of the fluids if blood is present. "If the stain is on a dark-colored fabric, the parts moistened by the fluids may be pressed with white blotting-paper, when blue impressions will be obtained. Contact with many substances causes the blue reaction or oxidation of guaiacum: the peculiarity of blood is that it does not produce this effect unless hydrogen peroxide or a similar 'antozonic' liquid is present. Bodies such as potassium permanganate, whose oxygen is apparently in the form

of ozone, also cause the production of a blue color with guaiacum; hydrogen peroxide and other compounds whose oxygen is in the opposite, positive, or, according to Schönbein, antozonic condition, produce no such effect. It would seem as if blood or some constituent of blood has the power of converting positive into negative oxygen, and thus bringing about an effect which negative oxygen alone is able to produce; for of all substances which, like blood, do not alone cause guaiacum to become blue, blood is the only one that so affects 'antozonides' (themselves inactive) as to enable them to act as ozonides; that is, to oxidize the guaiacum. Both the venous and the arterial fluid from any red-blooded animal will produce this blue reaction. Fruit-stains are darkened by ammonia, which does not alter the color of blood. Iron stains or iron moulds yield no color to water, whereas the red coloring-matter of blood is soluble in water. The hydrogen peroxide should be free from more than a trace of acid."

The blood-corpuscles of ordinary animals are much smaller than those of man, but a $\frac{1}{16}$ or $\frac{1}{25}$ inch lens is necessary for proper differentiation (J. G. Richardson).


Pus and Mucus.—Purulent urine deposits, on standing, a light-colored layer, easily diffused through the liquid by shaking. Acetic acid does not dissolve the sediment, and solution of potash of official strength converts it into a gelatinous mass. Under the microscope pus-corpuscles appear rounded and colorless, rather larger than blood-disks, and somewhat granular on the surface. They generally

Fig. 59.



Pus-corpuscles.

Fig. 60.



Fat-globules.

show minute nuclei, which are more distinctly seen after treatment with acetic acid. (See the portion of Fig. 59 marked a.) Mucus possesses no definite microscopic characters, but commonly has imbedded in it pus, epithelium, and air-bubbles. Mucus is coagulated in a characteristic manner by acetic acid; and this reaction, together with the ropy appearance it imparts to urine, prevents it being confounded with pus.

Day's test for pus consists in adding a drop or two of oxidized tincture of guaiacum to the urine or other liquid, when a clear blue color is produced. It is necessary to moisten dry pus with water before applying the test. The test-liquid is made by exposing a saturated alcoholic solution of guaiacum to the air until it has absorbed a sufficient quantity of oxygen to give it the property of turning green when placed in contact with potassium iodide. Day's test for mucus consists in the application, first, of oxidized tincture of guaiacum, which by itself undergoes no change in the presence of mucus, and then in the addition of carbolic acid or creasote, which quickly changes the color of the guaiacum to a bright blue. Neither carbolic acid nor creasote alone will render guaiacum blue. In testing for mucus on cloths or when it is mixed with blood, it is necessary to use the carbolic acid pure; but when the mucus is in a liquid state, it is better to use carbolic acid diluted with alcohol.


Saliva.—This secretion is an aqueous fluid containing less than 1 per cent. of solid matter, of which one-third is an albumenoid substance termed *ptyalin* (from *πτύαλον*, *spittle*), a ferment that has the power of converting starch into dextrin and grape-sugar: alkaline salts, including a trace of potassium sulphocyanate, and

Fig. 61.



Spermatozoa.

Fig. 62.



Sarcina ventriculi.

calcium compounds, are also present. Day's test for saliva in urine, etc. is similar to that for mucus, with the exception that the blue reaction produced by the oxidized tincture of guaiacum and alcoholic solution of carbolic acid is highly intensified by the addition of a little ethereal or aqueous solution of hydrogen peroxide.

Fatty matter occurs either as minute globules partially diffused through the urine (as shown at *a*) or in more intimate emulsion (as at *b* in Fig. 60). When present in larger quantity it collects as a sort of skim on the surface after standing.

Spermatozoa are liable to escape notice on account of their small size and extreme transparency. Suspected urine should be allowed to settle some hours in a conical test-glass, and the drop at the

bottom examined under a high power. The drawing (Fig. 61) shows their tadpole-like appearance.

Sarcinæ rarely occur in urine, but are not infrequent in vomited matters. The upper figures (*a*, Fig. 62) are copied from Dr. Thudichum's drawing (from urine); the larger groupings (*b*) are from vomited matter.

Extraneous bodies, such as starch, hair, wool, fibres of cotton or of deal, or fragments of feathers, are often found in urinary deposits, and ludicrous mistakes have been made by observers not on their guard in respect to such casual admixtures.

EXAMINATION OF URINARY CALCULI.

The term *calculus* is the diminutive of *calx*, a lime- or chalk-stone.

The following calculi have been met with: (1) Uric acid, (2) sodium urate, (3) calcium oxalate (mulberry), (4) fusible or mixed calcium and triple phosphates, (5) calcium phosphate, (6) calcium carbonate, (7) xanthine, (8) cystin, (9) urostealith (fatty matter), (10) indigo (one case).

Knowledge of the composition of a calculus or urinary deposit affords valuable diagnostic aid to the physician; hence the importance of a trustworthy analysis of these substances.

Nature of Calculi.—Urinary calculi have the same composition as unorganized urinary sediments. They consist, in short, of sediments that have been deposited slowly within the bladder, particle on particle, layer on layer, the several substances becoming so compact as to be less easily acted on by reagents than when deposited after the urine has been passed—the urates less readily soluble in warm water, the calcic phosphate insoluble in acetic acid until it has been dissolved in hydrochloric acid and reprecipitated by an alkali.

Preliminary Treatment.—If the calculus is whole, saw it in two through the centre, and notice whether it is built up of distinct layers or apparently consists of one substance. If the latter, use about a grain of the sawdust for analysis; if the former, carefully scrape off portions of each layer and examine them separately. If the calculus is in fragments, select fair specimens of about half a grain or a grain each, and reduce to a fine powder by placing on a hard surface and crushing under the blade of a knife.

Analysis.—Commence the analysis by heating a portion, about the size of a pin's head, on platinum-foil, in order to ascertain whether organic matter, inorganic matter, or both, are present. If both, the ash is examined for inorganic substances, and a fresh portion of the calculus for uric acid by the murexid test. (In the absence of uric acid any slight charring may be considered to be due to indefinite animal matter.) If composed of organic matter only, the calculus will in nearly all cases be uric acid, the indication being confirmed by applying the murexid test in a watch-glass to another fragment half the

size of a small pin's head. If inorganic only, the ash on the platinum-foil may be examined for phosphates, and a separate portion of the calculus for oxalates. Even a single drop of liquid obtained in any of these experiments may be filtered by placing it on a filter not larger than a sixpence and previously moistened with water, and adding three or four drops of water, one after the other, as each passes through the paper; or a drop of mixture may be placed on a fragment of damped filter-paper on a glass slide, the latter then tilted, and a clear drop be drained off from the paper on to the slide ready for the addition of a reagent. If the calculus is suspected to contain more than one substance, boil about half a grain of the powder in half a test-tubeful of distilled water for a few minutes and pour it on a small filter; then proceed according to the following table:

<i>Insoluble.</i>		<i>Soluble.</i>	
Phosphates, calcium oxalate, and free uric acid.		Urates.	
Boil with two or three drops of hydrochloric acid, and filter.		These will probably be redeposited as the solution cools. Small quantities may be detected by evaporating the solution to dryness. They are tested for ammonium, sodium, calcium, and the uric radical by the appropriate reagents.	
Uric acid. Apply the murexid test (p. 362).			
	<i>Insoluble.</i>	<i>Soluble.</i>	
	Phosphates and calcium oxalate. Add excess of ammonia, and then excess of acetic acid; filter.		
	<i>Insoluble.</i>	<i>Soluble.</i>	
	Calcium oxalate.	Phosphates. They may be reprecipitated by ammonia.	

Varieties of Calculi.—Calculi composed entirely of *uric acid* are common; a minute portion heated on platinum-foil chars, burns, and leaves scarcely a trace of ash. The phosphates frequently occur together, forming what is known as the *fusible calculus*, from the readiness with which a fragment aggregates, and even fuses to a bead, when heated on a loop of platinum-wire in the blowpipe flame. The phosphates may, if necessary, be further examined by the method described in connection with urinary deposits. Calcium oxalate often occurs alone, forming a dark-colored calculus having a very rough surface, hence termed the *mulberry calculus*. Smaller calculi of the same substance are called, from their appearance,

hempseed calculi. Calculi of *cystin* are rarely met with. *Xanthine* (from ξάνθος, *xanthos*, yellow, in allusion to the color it yields with nitric acid) still less often occurs as a calculus. The earthy concretions, or "*chalk-stones*," which frequently form in the joints of gouty persons are composed chiefly of urates, the sodium salt being that most commonly met with. *Gall-stones*, or *biliary calculi*, occasionally form in the gall-bladder; they contain *cholesterin* (from χολή, *cholē*, bile, and στερεός, *stereos*, solid), a fatty substance of alcoholoid constitution, soluble in rectified spirit or ether, and crystallizing from such solutions in well-defined, square, scaly crystals. Phosphatic and other calculi of many pounds' weight are occasionally found in the stomach and larger intestines of animals.

QUESTIONS AND EXERCISES.

In breathing how much carbon (in the form of carbonic acid gas) is exhaled from the lungs every twenty-four hours?—How may the presence of carbonic acid gas in expired air be demonstrated?—Mention an experiment showing the escape of moisture from the lungs during breathing.—State the method of testing for albumen in urine.—Give the tests for sugar in urine.—What is the average composition of healthy urine?—Give the tests for urea.—Write the rational formulæ of some compound ureas in which methyl or ethyl displaces hydrogen.—Describe an artificial process for the production of urea, giving equations.—Sketch out a plan for the chemical examination of urinary sediments.—A deposit is insoluble in the supernatant urine or in acetic acid; of what substances may it consist?—Which compounds are indicated when a deposit redissolves on warming it with the supernatant urine?—Name the salts insoluble in warmed urine, but dissolved on the addition of acetic acid.—Mention the chemical characters of *cystin*. At what stage of analysis would it be recognized?—Describe the microscopical appearances of the following urinary deposits: Uric acid, *cystin*, triple phosphate, earthy phosphates, urates, calcium oxalate, calcium carbonate, hippuric acid, tube-casts, epithelial *débris*, blood, pus, mucus, fat, spermatozoa, sarcinæ, extraneous bodies.—How are Day's tests for blood, pus, and saliva applied?—State the physical and chemical characters of urinary calculi.—How are urinary calculi prepared for chemical examination?—Draw out a chart for the chemical examination of urinary calculi.—Why is the "*fusible calculus*" so called? and what is its composition?—State the characters of "*mulberry*" and "*hempseed*" calculi.—What are "*chalk-stones*" of gout, and "*gall-stones*," or "*biliary calculi*"?

THE GALENICAL PREPARATIONS OF THE PHARMACOPŒIAS.

The preparation of Cerates, Confections, Decoctions, Enemas, Extracts, Glycerins, Infusions, Inhalations, Juices, Liniments, Lozenges, Mixtures, Ointments, Pills, Plasters, Poultices, Powders, Spirits, Suppositories, Syrups, Tinctures, and Wines includes a number of mechanical rather than chemical operations, and belongs to the domain of pure Pharmacy. The medical or pharmaceutical pupil will probably have had some

opportunity of practically studying these compounds before working at experimental chemistry, and may have prepared many of them according to the directions of the pharmacopœias; if not, he is referred to the pages of the last edition of those works for details.

Among the extracts of the British Pharmacopœia, however, there are five (namely, those of aconite, belladonna, hemlock, henbane, and lettuce) which are not simply evaporated infusions, decoctions, or tinctures, like most others, but are evaporated juices from which vegetable albumen, the supposed source of fermentation and decay, has been removed, and chlorophyll (the green coloring-matter of plant-juice) retained, practically unimpaired in tint. For educational practice either of the above-named five raw materials may be employed; but in order that attention may be concentrated on the process by which the extracts are prepared, rather than on any one of the extracts themselves, it suffices to make an extract of some ordinary green vegetable, such as cabbage or turnip-tops. Bruise the green leaves of a good-sized cabbage in a mortar, and press out the juice; heat it gradually to 130° F. and remove the green flocks of chlorophyll which separate by filtration through calico. When the liquor has all passed through the filter, set the chlorophyll aside for a time, heat the strained liquor to 200° F. to coagulate albumen; remove the latter by filtration and throw it away; evaporate the filtrate on a water-bath to the consistence of thin syrup; then add to it the chlorophyll, first passing the latter through a hair sieve to break up clots, and, stirring the whole together assiduously, continue the evaporation at a temperature not exceeding 140° F. until the extract is of a suitable consistence for forming pills. A higher temperature than that indicated would cause the alteration of the chlorophyll to a dark-brown substance, any such extract used in pharmacy no longer having the green tint which custom and the British Pharmacopœia demand.

QUESTIONS AND EXERCISES.

Enumerate the different classes into which official galenical preparations may be divided.—Describe the general process for the preparation of green extracts—aconite, belladonna, hemlock, henbane, lettuce.—Why is vegetable albumen excluded in the preparation of green extracts?—How may chlorophyll be removed from vegetable juices, and again be introduced into their evaporated residues without destroying its color?—For what reason is exposure of chlorophyll to a boiling temperature avoided in the manufacture of green extracts?

THE CHEMICAL PREPARATIONS OF THE PHARMACOPŒIAS.

The process by which every official chemical substance is prepared has already been described, and the strictly chemical character of the processes illustrated by experiments and explained by aid of equations. Should the reader, in addition, desire an intimate acquaintance with the details of manipulation on which the successful and economic manufacture of chemical substances depends, he is advised to prepare, if he has not done so already, a few ounces of each of the salts mentioned in the pharmacopœias or commonly used in pharmacy. An additional guide in these operations will be the pharmacopœias themselves.

The production of many chemical and galenical substances on a commercial scale can only be successfully carried on in manufacturing laboratories and with some knowledge of the circumstances of supply and demand, value of raw material and by-products, etc.; for the technical preparation of such substances requires much knowledge beyond even a thorough acquaintance with chemistry. Still, in the present day, commercial chemistry and pharmacy can best hope for success when founded on the working out of abstract scientific principles. The problem of manufacturing success is now only solved with certainty by sound and wisely-applied science.

Memorandum.—The next subjects of experimental study will be determined by the nature of the student's future pursuits. In most cases the operations of quantitative analysis will engage attention. These should be of a volumetric and gravimetric character; for details concerning them see the following pages.

QUANTITATIVE ANALYSIS.

INTRODUCTORY REMARKS.

General Principles.—The proportions in which chemical substances unite with each other in forming compounds are definite and invariable (p. 48). Quantitative analysis is based on this law. When, for example, aqueous solutions of a salt of silver and a chloride are mixed, a white curdy precipitate is produced containing chlorine and silver in atomic proportions; that is, 35.5 parts of

chlorine to 108 of silver. No matter what the chloride or what the salt of silver, the resulting silver chloride is invariable in composition. The formula AgCl is a convenient picture of this compound in these proportions. The weight of a definite compound being given, therefore, the proportional amounts of its constituents can be ascertained by simple calculation. Suppose, for instance, 8.53 parts of silver chloride have been obtained in some analytical operation: this amount will contain 2.11 parts of chlorine and 6.42 of silver. For if 143.5 (the molecular weight) of silver chloride contain 35.5 (the atomic weight) of chlorine, 8.53 of silver chloride will be found to contain 2.11 of chlorine:

$$\begin{array}{rcl}
 143.5 & : & 35.5 \\
 & & \hline
 & & 8.53 \\
 & & 1.065 \\
 & & 17.75 \\
 & & \hline
 & & 284.0 \\
 143.5 & \times & 2.11 \\
 & & \hline
 & & 287.0 \\
 & & 15.81 \\
 & & \hline
 & & 14.35 \\
 & & \hline
 & & 1.465 \\
 & & \hline
 & & 1.435
 \end{array}
 \qquad x = 2.11$$

And if 143.5 of silver chloride contain 108 of silver, 8.53 of silver chloride will contain 6.42 of silver. To ascertain, for example, the amount of silver in a substance containing, say, silver nitrate, all that is necessary is to take a weighed quantity of the substance, dissolve it, precipitate the whole of the silver by adding hydrochloric acid or other chloride till no more silver chloride falls, collect the precipitate on a filter, wash, dry, and weigh. The amount of silver in the dried chloride, ascertained by calculation, is the amount of silver in the quantity of substance on which the operation was conducted; a rule-of-three sum gives the quantity per cent., the form in which the results of quantitative analyses are usually stated. Occasionally a constituent of a substance admits of being isolated and weighed in the uncombined state. Thus the amount of mercury in a substance may be determined by separating and weighing the mercury in the metallic condition; if occurring as calomel (HgCl) or corrosive sublimate (HgCl_2), the proportion of chlorine may then be ascertained by calculation ($\text{Hg} = 200$; $\text{Cl} = 35.5$).

Nature of Gravimetric Quantitative Analysis.—As above stated, a body may be isolated and weighed and its quantity thus ascertained, or it may be separated and weighed in combination with another body whose combining proportion is well known; this is quantitative analysis by the *gravimetric* method.

Nature of Volumetric Quantitative Analysis.—Volumetric operations depend for success on some accurate initial gravimetric operation. A weighed amount of a pure salt is dissolved in a given volume of water or other fluid, and thus forms a *standard solution*. Accurately measured quantities of such a solution will obviously contain

just as definite amounts of the dissolved salt as if those amounts were actually *weighed* in a balance; and as measuring occupies less time than weighing, the volumetric operations can be conducted with great economy of time as compared with the corresponding gravimetric operations. Quantitative analysis by the *volumetric* method consists in noting the volume of the standard liquor required to be added to the substance under examination before a given effect is produced. Thus, for instance, a solution of silver nitrate of known strength may be used in experimentally ascertaining an unknown amount of a chloride in any substance. The silver solution is added to a solution of a definite quantity of the substance until flocks of silver chloride cease to be precipitated: every 108 parts of silver added (or 170 of silver nitrate: $\text{Ag} = 108$, $\text{N} = 14$, $\text{O}_3 = 48$; total 170) indicates the presence of 35.5 of chlorine or an equivalent quantity of any chloride. The preparation of a standard solution, such as that of the silver nitrate to which allusion is here made, requires much care; but once it is prepared, certain analyses can, as already indicated, be executed with far more rapidity and ease than by gravimetric processes.

Quantitative Determination of (a) Atmospheric Pressure, (b) Temperature, and (c) Weight.—The analysis of solids and liquids often involves quantitative determinations not only of weight, as just indicated, but of temperature and atmospheric pressure. The two latter processes will now be explained, after which an outline of volumetric and gravimetric quantitative analysis will be given. The scope of this work precludes any attempt to describe all the little mechanical details observed by quantitative analysts; essential operations, however, are so fully treated that expert manipulators will meet with little difficulty.

QUANTITATIVE DETERMINATION OF ATMOSPHERIC PRESSURE.

The Barometer.—The analysis of gases and vapors involves determinations of the varying pressure of the atmosphere as indicated by the *barometer* (from $\beta\acute{\alpha}\rho\omicron\varsigma$, *baros*, weight, and $\mu\acute{\epsilon}\tau\rho\omicron\nu$, *metron*, measure).

The ordinary mercurial barometer is a glass tube thirty-three or thirty-four inches long, closed at one end, filled with mercury, and inverted in a small cistern or cup of mercury (Fig. 63). The mercury remains in the tube, owing to the weight or pressure of the atmosphere on the exposed surface of the liquid, the average height of the column being nearly thirty inches. In the popular form of the instrument, the wheel barometer, the cistern is formed by a recurvature of the tube (Fig. 64); on the exposed surface of the mercury a float is placed, from which a thread passes over a pulley and moves an index whenever the column of mercury rises or falls. As supplied to the public, these barometers are usually enclosed in ornamental frames with thermometers attached. In the wheel barometer the glass tube and contained column of mercury are alto-

gether enclosed, the index alone being visible. In the other variety the upper end of the glass tube and mercurial column are exposed, and the height of the mercury

Fig. 63. is ascertained by direct observation.




Barometer.

The aneroid barometer (from *à*, *a*, without, and *νῆρος*, *nēros*, fluid) consists of a small shallow vacuous metal drum, the sides of which approach each other when an increase of atmospheric pressure occurs, their elasticity enabling them to recede toward their former position on a decrease of pressure. This motion is so multiplied and altered in direction by levers, etc. as to act on a hand traversing a plate on which are marked numbers corresponding with those showing the height of the mercurial column of the ordinary barometer by which the aneroid was adjusted. The Bourdon barometer (from the name of the inventor) is a modified aneroid, containing in the place of the round metal box a flattened vacuous tube of metal bent nearly to a circle. These barometers are also useful for measuring the pressure in steam-boilers, etc. Under the name of *pressure-gauges* they are sold to indicate pressure of five hundred pounds per square inch and upward. From their portability (they can

be made of one to two inches in diameter and less than an inch thick) they are handy companions for travellers wishing to know the heights of hills, mountains, and other elevations.

(For further information concerning the influence of pressure on the volume of a gas or vapor see p. 619; and for descriptions of the methods of analyzing gases refer to Ganot's "Physics" (translated by Atkinson), Miller's "Chemical Physics," and "Analysis of Gases" in Watts's "Dictionary of Chemistry.")

Fig. 64.



Barometer.

QUANTITATIVE DETERMINATION OF TEMPERATURE.

General Principles.—As a rule, all bodies expand on the addition, and contract on the abstraction, of heat, the alteration in volume being constant and regular for equal increments or decrements of temperature. The extent of this alteration in a given substance, expressed in parts or degrees, constitutes the usual method of intelligibly stating with accuracy, precision, and minuteness a particular condition of warmth or temperature; that is, of sensible heat. The substance commonly employed for this purpose is mercury, the chief advantages of which are that it will bear a high temperature without boiling, a low temperature without freezing, does not ad-

here to glass to a sufficient extent to "wet" the sides of any tube in which it may be enclosed, and, from its good conducting power for heat, responds rapidly to changes of temperature. Platinum, earthenware, alcohol, and air are also occasionally used for thermometric purposes.

The Thermometer.—The construction of an accurate thermometer is a matter of great difficulty, but the following are the leading steps in the operation: Select a piece of glass tubing having a fine capillary (*capillus*, a hair) bore and about a foot long; heat one extremity in the blowpipe flame until the orifice closes and the glass is sufficiently soft to admit of a bulb being blown; heat the bulb to expel air, immediately plunging the open extremity of the tube into mercury; the bulb having cooled, and some mercury having entered and taken the place of expelled air, again heat the bulb and tube until the mercury boils and its vapor escapes through the bore of the tube; again plunge the extremity under mercury, which will probably now completely fill the bulb and tube. When cold the bulb is placed in melting ice. The top of the column of mercury in the capillary tube should then be within an inch or two of the bulb; if higher, some of the mercury must be expelled by heat; if lower, more metal must be introduced as before. The tube is now heated near the open end and a portion drawn out until the diameter is reduced to about one-tenth. The bulb is next warmed until the mercurial column rises above the constricted part of the tube, which is then rapidly fused in the blowpipe flame and the extremity of the tube removed.


The instrument is now ready for *graduation*. The bulb is placed in the steam just above some rapidly-boiling water (a medium having, *cæteris paribus*, an invariable temperature), and when the position of the top of the mercurial column is constant (the flask containing the water and steam being jacketed to prevent loss of heat by radiation), a mark is made on the stem by a scratching diamond or a file. This operation is repeated with melting ice (also a medium having an invariable temperature). The space between these two marks is divided into a certain number of intervals termed *degrees*. Unfortunately, this number is not uniform in all countries: in England it is 180, as proposed by Fahrenheit; in France 100, as proposed by Celsius (the Centigrade scale), a number generally adopted by scientific men; in some parts of the Continent the divisions are 80 for the same interval, as suggested by Réaumur. Whichever be the number selected, similar markings should be continued beyond the boiling- and freezing-points as far as the length of the stem admits. They may be made on

the stem itself or on any wood, metal, or earthenware frame on which the stem is mounted.

Thermometric Scales. (Fig. 65).—On the Centigrade (C.) and Réaumur (R.) scales the freezing-point of water is made zero, and the boiling-point 100 and 80 respectively; on the Fahrenheit (F.) scale the zero is placed 32 degrees below the congealing-point of water, the boiling-point of which becomes, consequently, 212. Even on the Fahrenheit system temperatures below the freezing-point of water are often spoken of as “degrees of frost;” thus 19° as marked on the thermometer would be regarded as “13 degrees of frost.” It is to be regretted that the freezing-point of water is not universally regarded as the zero-point, and that the number of intervals between that and the boiling-point is not everywhere the same.

Fig 65.

THERMOMETRIC SCALES.



The degrees of one scale are easily converted into those of another if their relations be remembered—namely, 180 (F.), 100 (C.), 80 (R.); or 18, 10, and 8; or, best, 9, 5, and 4.

The degrees of one scale are easily converted into those of another if their relations be remembered—namely, 180 (F.), 100 (C.), 80 (R.); or 18, 10, and 8; or, best, 9, 5, and 4.

Formule for the Conversion of Degrees of one Thermometric Scale into those of Another.

F. = Fahrenheit.
R. = Réaumur.

C. = Centigrade.
D. = The observed degree.

If above the freezing-point of water (32° F.; 0° C.; 0° R.):

F. into C.	$(D - 32) \div 9 \times 5.$
F. " R.	$(D - 32) \div 9 \times 4.$
C. " F.	$D \div 5 \times 9 + 32.$
R. " F.	$D \div 4 \times 9 + 32.$

If below freezing, but above 0° F. (− 17.77° C.; − 14.22° R.):

F. into C.	$-(32 - D) \div 9 \times 5.$
F. " R.	$-(32 - D) \div 9 \times 4.$
C. " F.	$32 - (D \div 5 \times 9).$
R. " F.	$32 - (D \div 4 \times 9).$

If below 0° F. (− 17.77° C.; − 14.22° R.):

F. into C.	$-(D + 32) \div 9 \times 5.$
F. " R.	$-(D + 32) \div 9 \times 4.$
C. " F.	$-(D \div 5 \times 9) - 32.$
R. " F.	$-(D \div 4 \times 9) - 32.$

For all degrees:

C. into R.	$D \div 5 \times 4.$
R. " C.	$D \div 4 \times 5.$

In ascertaining the temperature of a liquid the bulb of a thermometer is simply inserted and the degree noted. In determining the boiling-point also the bulb is inserted in the liquid if a pure substance. In taking the boiling-point of a substance which is being distilled from a mixture, the bulb of the thermometer should be in the vapor, but not beneath nor very near to the surface of the boiling liquid.

The "boiling-point" of a liquid is the temperature at which the elasticity of the vapor of the substance overcomes the atmospheric or other pressure to which the liquid is exposed. If the pressure is equal to 760 mm. (29.92 inches) of mercury, water will boil at 100° C. (212° F.). The boiling-point of a drop of a fluid is taken by introducing it into the closed extremity of a small U tube, the remaining portion of the closed limb being filled with mercury. The tube is lowered into a bath, the open limb being above the surface of the fluid of the bath. The bath is slowly and equally heated, and the boiling-point of the liquid, indicated by the mercury falling until it is level in the two limbs, taken by a thermometer whose bulb is close to the U tube.

The following are the boiling-points of a few substances met with in pharmacy :

	Centigrade.	Fahrenheit.
Alcohol, absolute	78.3	173
" 84 per cent.	79.5	175
" 49 per cent. (proof spirit)	81.4	178.5
" amylic	132.2	270
Benzol	80.6	177
Bromine (below)	63	145.4
Benzoic acid	239.0	462
Carbolic acid	187.8	370
Chloroform	61	142
Ether (B. P.) (below)	40.5	105
" pure	35	95
Mercury, <i>in vacuo</i> (as in a thermometer) .	304	580
" in air (barometer at 30 inches) .	350	662
Water (barometer at 29.92 inches)	100	212
" (" at 29.33 ")	99.5	211
" (" at 28.74 ")	99	210
Saturated solutions of —		
Cream of tartar	101	214
Common salt	106.6	224
Sal-ammoniac	113.3	236
Sodium nitrate	119	246
Sodium acetate	124.4	256
Calcium chloride	179.4	355

To Determine Melting-points of Fats.—To melt at a given temperature is a *constant* property of a substance; therefore, a melting-point, once it is accurately determined, becomes a valuable indicator of purity in a substance. Heat a fragment of the substance (spermaceti or wax, for example) till it liquefies, and then draw up a small portion into a thin glass tube about the size of a knitting-needle. Immerse the tube in cold water contained in a beaker, and slowly heat the vessel till the thin opaque cylinder of solid fat melts and becomes transparent: a delicate thermometer placed in the water indicates the point of change to the fifth of a degree. Remove the source of heat and note the congealing-point of the substance; it will, in nearly all cases, be identical with or close to the melting-point.

The following are melting-points of substances official in the British Pharmacopœia:

	In degrees Centigrade.	In degrees Fahrenheit.
Acetic acid, glacial	8.9	48
“ “ “ congeals at	1.1	34
Benzoic acid	120	248
Carbolic acid	35	95
Oil of theobroma (about)	32	90
Phosphorus	43.3	110
Prepared lard (about)	38	100
“ suet	39.5	103
Spermaceti	44 to 50	111 to 122
White wax	63.3	146
Yellow wax	63.3	146

Pyrometers.—Temperatures above the boiling-point of mercury are determined by ascertaining to what extent a bar of platinum or porcelain has elongated. The bar is enclosed in a cavity of a suitable case, a plug of platinum or porcelain placed at one end of the bar, and the whole exposed in the region the temperature of which is to be found. After cooling, the distance to which the bar has forced the plug along the cavity is accurately measured and the corresponding degree of temperature noted. The value of the distance is fixed for low temperatures by comparison with a mercurial thermometer, and the scale carried upward through intervals of equivalent length. Such thermometers are conventionally distinguished from or-

dinary instruments by the name *pyrometer* (from $\pi\upsilon\rho$, *pur*, fire, and $\mu\acute{\epsilon}\tau\rho\omicron\nu$, *metron*, measure).

The order of fusibility of a few of the metals is as follows :

	In degrees Centigrade.	In degrees Fahrenheit.
Mercury	— 39.4	— 39
Potassium	+ 62.5	+ 144.5
Sodium	97.6	207.7
Tin	227.8	442
Bismuth	264	507
Lead	325	617
Zinc	411.6	773
Antimony	621	1150
Silver	1023	1873
Copper	1091	1996
Gold	1102	2016
Cast iron	1530	2786

QUESTIONS AND EXERCISES.

On what fundamental laws are the operations of quantitative analysis based?—What is the general nature of *gravimetric* quantitative analysis?—Describe the general principle of *volumetric* quantitative analysis.—How are variations in atmospheric pressure determined?—Explain the construction and mode of action of barometers.—In what respect does a wheel-barometer differ from an instrument in which the readings are taken from the top of the column of mercury?—Describe the principles of action of an aneroid barometer.—On what general principles are thermometers constructed?—What material is employed in making thermometers?—Why is mercury selected as a thermometric indicator?—Describe the manufacture of a mercurial thermometer.—How are thermometers graduated?—Give formulæ for the conversion of the degrees of one thermometric scale into those of another, (*a*) when the temperature is above the freezing-point of water; (*b*) below 32° F., but above 0° F., and (*c*) below 0° F.—Name the degree C. equivalent to 60° F.—What degree C. is represented by —4° F.?—Mention the degree F. indicated by 20° C.—Convert 100° R. into degrees C. and F.—State the boiling-points of alcohol, chloroform, ether, mercury, and water on either thermometric scale.—Describe the details of manipulation in estimating the melting-points of fats.—In what respect do pyrometers differ from thermometers?—Mention the melting-points of glacial acetic acid, oil of theobroma, lard, suet, and wax.—Give the fusing-points of tin, lead, zinc, copper, and cast iron.

QUANTITATIVE DETERMINATION OF WEIGHT.

All bodies, celestial and terrestrial, attract each other, the amount of attraction being in direct proportion to the quantity of matter of which they consist, and in inverse proportion to the squares of their distances. This is *gravitation*. When gravitation in certain directions is exactly counterbalanced by gravitation in opposite directions, a body (*e. g.* the earth) remains suspended in space. Such a body, in relation to other bodies, has gravity, but not weight. *Weight* is the effect of gravity, being the excess of gravitation in one direction over and above that exerted in the opposite direction. Weight, truly, in any terrestrial substance, is the excess of attraction which it and the earth have for each other over and above the attraction of each in opposite directions by the various heavenly bodies. But, practically, the weight of any terrestrial substance is the effect of the attraction of the earth only. *Specific weight* is the definite or precise weight of a body in relation to its bulk; it is more usually, but not quite correctly, termed *specific gravity*—gravity belonging to the earth, and not, in any sensible degree, to the substance.*

QUESTIONS AND EXERCISES.

What is understood by gravitation?—State the difference between weight and gravity.—Mention a case in which a body has gravity, but no apparent weight.—Practically, what causes the weight of terrestrial substances?

Weights and Measures.

The Balance.—The balance used in the quantitative operations of analytical chemistry must be accurate and sensitive. The points of suspension of the beam and pan should be polished steel or agate knife-edges working on agate planes. It should turn easily and quickly, without too much oscillation, to $\frac{1}{500}$ or $\frac{1}{600}$ of a grain, or $\frac{1}{10}$ of a milligramme, when 1000 grains, or 50 or 60 grms., are placed in each pan. (Grammes are weights of the metric system, a description of which is given on the next two or three pages.) The beam should be light but strong, capable of supporting a load of 1500 grains or 100 grammes: its oscillations are observed by the help of a long index attached to its centre, and continued downward for some distance in front of the supporting pillar of the balance. The instrument should be provided with screws for purposes of adjustment, a mechanical contrivance for supporting the

* It must be remembered, also, that centrifugal influence and gravitation are antagonistic.

beam above its bearings when not in use or during the removal or addition of weights, spirit-levels to enable the operator to give it a horizontal position, and be enclosed in a glass case to protect it from dust. It should be placed in a room the atmosphere of which is not liable to be contaminated by acid fumes, in a situation as free as possible from vibration; and a vessel containing lumps of quicklime should be placed in the case to keep the enclosed air dry and prevent the formation of rust on any steel knife-edges or other parts. During weighing the doors of the balance-case should be shut, in order that currents of air may not unequally influence the pans.

The Weights.—These should be preserved in a box having a separate compartment for each. They must not be lifted directly with the fingers, but by a small pair of forceps. If grain weights, they should range from 1000 gr. to $\frac{1}{10}$ gr., a $\frac{1}{10}$ weight being fashioned of gold wire to act as a “rider” on the divided beam, and thus indicate by its position 100ths and 1000ths of a grain. From $\frac{1}{10}$ to 10 grs. the weights may be of platinum or aluminium; thence upward to 1000 grains of brass. The relation of the weights to each other should be decimal. Metric decimal weights may range from 100 grammes to 1 gramme, of brass, and thence downward to 1 centigramme, of platinum or aluminium, a gold centigramme rider being employed to indicate milligrammes and tenths of a milligramme.

“The weights and measures referred to by physicians in prescribing, and used by pharmacists in dispensing, medicines are, in the United States, either those of the ‘apothecaries’ or Troy system of weights and the wine measure, or those of the metric system.”

Troy Weights.—These are derived from the *Troy pound*, and are exhibited in the following table, with their signs annexed :

One pound,	℔ = 12 ounces	= 5760 grains.
One ounce,	℥ = 8 drachms	= 480 grains.
One drachm,	ʒ = 3 scruples	= 60 grains.
One scruple,	ʒ	= 20 grains.
One grain,	gr.	= 1 grain.

It is highly important that persons engaged in preparing medicines should be provided with Troy weights. But those who are not so provided can make their avoirdupois weights available as substitutes for Troy weights by bearing in mind that 42.5 grains, added to the avoirdupois ounce, will make it equal to the Troy ounce, and that 1240 grains, deducted from the avoirdupois pound, will reduce it to the Troy pound.

Measures.—These are derived from the *wine gallon*, and are given in the following table, with their signs annexed :

One gallon,	C = 8 pints	= 61,440 minims.
One pint,	O = 16 fluidounces	= 7,680 minims.
One fluidounce,	f℥ = 8 fluidrachms	= 480 minims.
One fluidrachm,	fʒ	= 60 minims.
One minim,	℥	= 1 minim.

Relation of Troy Weight and Wine Measure.

1 minim =	0.95 grains.	1 grain =	1.05 minims.
1 f̄3 =	56.96 "	1 3̄ =	63.2 "
1 f̄3 =	455.69 "	1 3̄ =	505.6 "

Weights and Measures of the United States Pharmacopœia.

"Weight and volume are expressed in the units of the international system based on the mètre. In cases where only relative quantities are stated, the proportions are expressed in *parts by weight or by volume*. All weights and measures used now (1893) are derived from the United States national prototype standards of the mètre and the kilogramme. The actual litre is the volume of one kilogramme of pure water at the temperature of its maximum density *in vacuo*. Theoretically, the litre is equal to one cubic décimètre or 1000 cubic centimètres. The United States *yard* is defined to be equal to $\frac{3600000}{393700}$ mètre; the commercial *pound* (avoir-dupois) is defined as being equal to $\frac{7000000000}{1543235639}$ kilogramme; and the liquid *gallon* is the volume of 3785.434 grammes (58418.1444 grains) of water at the temperature of its maximum density, weighed *in vacuo*."

The Metric System of Weights (the word *metric* is from the Greek μέτρον, *metron*, measure) is greatly to be preferred to the British, the relation of the metric weights of all denominations to measures of length, capacity, and surface being so simple as to be within the perfect comprehension of a child; while under the British plan the weights have no such relation either with each other or with the various measures. Moreover, the metric system is in perfect harmony with the universal method of counting; it is a decimal system.

[It is perhaps impossible to realize, much more express, the advantages we enjoy from the fact that in every country of the world the system of numeration is identical. That system is the decimal. Whatever language a man speaks, his method of numbering is decimal; his talk concerning numbers is decimal; his written or printed signs signifying number are decimal. With the figures 1, 2, 3, 4, 5, 6, 7, 8, 9, 0 he represents all possible variations in number, the position of a figure in reference to its companions alone determining its value, a figure on the left hand of any other figure in an allocation of numeral symbols (for example, 1894) having ten times the value of that figure, while the figure on the right hand of any other has a tenth of the value of that other. When the youngest pupil is asked how many units there are in 1894, he smiles at the simplicity of the question, and says, 1894. How many tens? 189 and 4 over. How many hundreds? 18 and 94 over. How many thousands? 1 and 894 over. But if he is asked how many scruples there are in 1894 grains, how many drachms, how many ounces, he brings out his slate and pencil. And so with the pints or gallons in 1894 fluidounces, or the feet and yards in 1894 inches, or the pence, shil-

lings, and pounds in 1894 farthings; to say nothing of cross-questions, such as the value of 1894 articles at 9s. 6d. per dozen, or of the perplexity caused by the varying values of several individual weights or of measures of length, capacity, and surface in different parts of the country. What is desired is, that there should be an equally simple decimal relation among weights and measures and coins as already universally exists among numbers. This condition of things having already been introduced into most other countries, there is no good reason why it should not, in due time, be accomplished in the United States and Great Britain. The difficulty in the way consists in requiring, more or less suddenly, every one of the many millions of persons in the country to relinquish the life-long use of words expressive of the hourly wants of life—such words as pounds, pints, pence, etc., for words such as kilos, litres, cents, etc., which will not express those wants until after weeks or months, or perhaps years. The difficulty is one that explanation cannot meet, for it is one of *association* which time alone can resolve, at all events for the great mass of people. Still, “what man has done man can do.”]

The system of weights and measures legalized in Great Britain by “the Metric Weights and Measures Act, 1864,” is founded on the *mètre*. Fig. 66 represents a pocket folding-measure the tenth part of a *mètre* in length, divided into 10 centimètres, and each centimètre into 10 millimètres.

Fig. 66.



The Décimètre.

The units of the system, with their multiples and submultiples, are as follows:

Length.—The *Unit of Length* is the *MÈTRE*, derived from the measurement of the quadrant of a meridian of the earth. (Practically, it is the length of certain carefully-preserved bars of metal, from which copies have been taken.)

Surface.—The *Unit of Surface* is the *ARE*, which is the square of ten mètres.

Capacity.—The *Unit of Capacity* is the *LITRE*, which is the cube of a tenth part of a *mètre*.

Weight.—The *Unit of Weight* is the *GRAMME*, which is the weight of that quantity of distilled water, at its maximum density, which fills a cube of the one-hundredth part of the *mètre*.

TABLE.

Note.—Multiples are denoted by the Greek words “Deka,” ten,
“Hecto,” hundred, “Kilo,” thousand.

Subdivisions by the Latin words, “Deci,” one-tenth,
“Centi,” one-hundredth, “Milli,” one-thousandth.

Quantities.	Length.	Surface.	Capacity.	Weight.
1000	Kilo-mètre	. . .	Kilo-litre	Kilo-gramme.
100	Hecto-mètre	Hectare	Hecto-litre	Hecto-gramme.
10	Déca-mètre	. . .	Déca-litre	Déca-gramme.
1 (Units)	MÈTRE	ARE	LITRE	GRAMME.
.1	Déca-mètre	. . .	Déci-litre	Déci-gramme.
.01	Centi-mètre	Centiare	Centi-litre	Centi-gramme.
.001	Milli-mètre	. . .	Milli-litre	Milligramme.

When the metric method is exclusively adopted, these units and table, comprising the entire system of weights and measures, represent all that will be essential to be learnt in lieu of the numerous and complicated tables hitherto in use. Adopting the style of elementary books on arithmetic, the table may be expanded thus :

10 Milligrammes	make	1 Centigramme.
10 Centigrammes	“	1 Decigramme.
10 Decigrammes	“	1 Gramme.
10 Grammes	“	1 Decagramme.
10 Decagrammes	“	1 Hectogramme.
10 Hectogrammes	“	1 Kilogramme.
10 Millilitres	make	1 Centilitre,
	etc.	
10 Millimetres	make	1 Centimetre,
	etc.	

Abbreviations.—Mètre = *m* ; décimètre = *dm* ; centimètre = *cm* ; millimètre = *mm* ; kilomètre = *km*. Square mètre = *m*² ; cubic mètre = *m*³ ; and so on. Litre = *l* ; décilitre = *dl* ; etc. Kilo-gramme = *kg* ; décagramme = *dkg* ; gramme = *g* ; décigramme = *dg* ; centigramme = *cg* ; and milligramme = *mg*.

The following approximate British equivalents of metrical units should be committed to memory :

1 Mètre	= 3 feet 3 inches and 3 eighths.
1 Are	= a square whose side is 11 yards.
1 Litre	= 1 $\frac{3}{4}$ pints.
1 Gramme	= 15 $\frac{1}{2}$ grains.
The Kilomètre	is equal to 1100 yards.
The Hectare	= 2 $\frac{1}{2}$ acres nearly.

The Metric Ton of 1000 Kilogrammes = 19 cwt. 2 qrs. 20 lbs. 10 oz.

The Kilogramme = 2 lbs. 3 $\frac{1}{4}$ oz. nearly.

(For *exact* equivalents, in many forms, see pp. 606 and 607.) A litre of water at 39° F. weighs 15,432 grains ; at 50° F., 15,429 grains ; at 60° F. it weighs 15,418 grains ; at 70° F., 15,403 grains ; and at 80° F., 15,383 grains (Pile). (The word *gramme* is sometimes,

unfortunately, written *gram*, which too closely resembles the word *grain*.)

Decimal Coinage.—In most countries where the metric system of weights and measures is employed a decimal division of coins is also adopted. This course, conjoined with the ordinary decimal method of enumerating, which, fortunately, is in universal use, renders calculations of all kinds most simple—easy to an extent which cannot be conceived in countries like England where the operations of weighing, measuring, paying, and counting have only the most absurdly intricate relations to each other.

The General Council under whose authority the British Pharmacopœia is issued encourages medical practitioners and pharmacists in the adoption of the metric system, and gives the annexed statement of metric weights and measures:

WEIGHTS AND MEASURES OF THE METRICAL SYSTEM.

WEIGHTS.

1 Milligramme	= the thousandth part of one grm., or 0.001 grm.	
1 Centigramme	= the hundredth	0.01 “
1 Décigramme	= the tenth	0.1 “
1 Gramme	= weight of a cubic centimètre of water at 4° C.	1.0 “
1 Décagramme	= ten grammes	10.0 “
1 Hectogramme	= one hundred grammes	100.0 “
1 Kilogramme	= one thousand grammes	1000.0 (1 kilo).

MEASURES OF CAPACITY.

1 Millilitre	=	1 cubic centim., or the mea. of 1 gramme of water.	
1 Centilitre	=	10	“ “
1 Décilitre	=	100	“ “
1 Litre	=	1000	“ (1 kilo.)

MEASURES OF LENGTH.

1 Millimètre	= the thousandth part of one mètre, or 0.001 mètre.	
1 Centimètre	= the hundredth	0.01 “
1 Décimètre	= the tenth	0.1 “
1 Mètre	= the ten-millionth part of a quarter of the meridian of the earth.	

METRICAL MEASURES OF LENGTH.

	In English inches.	In English feet = 12 inches.	In English yards = 3 feet.	In English fathoms = 6 feet.	In English miles = 1760 yards.
Millimetre.....	0.03937	0.003251	0.0010936	0.0005468	0.0000006
Centimetre.....	0.39371	0.032809	0.0109363	0.0054682	0.0000062
Decimetre	3.93708	0.328090	0.1093633	0.0546816	0.0000621
Metre	39.37079	3.280899	1.0936331	0.5468165	0.0006214
Decimetre.....	393.70790	32.808992	10.9363316	5.4681653	0.0062138
Hectometre	3937.07900	328.089917	109.3633156	54.6816528	0.0621382
Kilometre	39370.79000	3280.899167	1093.6330556	546.8165278	0.6213824
Myriametre.....	393707.90000	32808.991667	10936.3305556	5468.1652778	6.2138242

1 inch = 2.539954 centimetres.
1 foot = 3.0479449 decimetres.

1 yard = 0.9143835 metre.
1 mile = 1.6093149 kilometres.

METRICAL MEASURES OF SURFACE.

	In English square feet.	In Eng. sq. yards = 9 square feet.	In English poles = 272.25 sq. feet.	In English roods = 10890 sq. feet.	In English acres = 43560 sq. feet.
Centiare, or square metre.....	10.764299	1.196033	0.0395383	0.0009885	0.0002471
Are, or 100 square metres.....	1076.429934	119.603326	3.9538290	0.0988457	0.0247114
Hectare, or 10,000 square metres....	107642.993418	11960.332602	395.3828959	9.8845724	2.4711431

1 square inch = 6.4513669 square centimetres.
1 square foot = 9.2899683 square decimetres.

1 square yard = 0.83609715 square metre or centiare.
1 acre = 0.40467102 hectare.
1 square mile = 2.58989451 square kilometres.

METRICAL MEASURES OF CAPACITY.

	In cubic inches.	In cubic feet = 1728 cubic inches.	In pints = 34.65923 cubic inches.	In gallons = 8 pints = 277.27384 cubic inches.	In bushels = 8 gal- lons = 2218.19072 cubic inches.
Millilitre, or cubic centimetre.....	0.06103	0.000035	0.00176	0.0002201	0.0000275
Centilitre, or 10 cubic centimetres..	0.61027	0.000353	0.01761	0.0022010	0.0002751
Decilitre, or 100 cubic centimetres..	6.10271	0.003532	0.17608	0.0220097	0.0027512
Litre, or cubic decimetre.....	61.02705	0.035317	1.76077	0.2200967	0.0275121
Decalitre, or centistere.....	610.27052	0.353166	17.60773	2.2009668	0.2751208
Hectolitre, or decistere.....	6102.70515	3.531658	176.07734	22.0096677	2.7512085
Kilolitre, or stere, or cubic metre...	61027.05152	35.316581	1760.77341	220.0966767	27.5120846
Myrialitre, or decastere.....	610270.51519	353.165807	17607.73414	2200.9667675	275.1208459
1 cubic inch = 16.386176 cubic centimetres.	1 cubic foot = 28.315312 cubic decimetres.	1 gallon = 4.535926 litres.			

METRICAL MEASURES OF WEIGHT.

	In English grains.	In troy ounces = 480 grains.	In avoirdupois lbs. = 7000 grains.	In cwt. = 112 lbs. = 784000 grains.	Tons = 20 cwt. = 15680000 grains.
Milligramme	0.01543	0.000032	0.0000022	0.0000000	0.0000000
Centigramme	0.15432	0.000322	0.0000220	0.0000002	0.0000000
Decigramme.....	1.54323	0.003215	0.0002205	0.0000020	0.0000001
Gramme.....	15.43235	0.032151	0.0022046	0.0000197	0.0000010
Decigramme	154.32349	0.321507	0.0220462	0.0001968	0.0000098
Hectogramme.....	1543.23488	3.215073	0.2204621	0.0019684	0.0000984
Kilogramme	15432.34880	32.150727	2.2046213	0.0196841	0.0009842
Myriagramme.....	154323.48800	321.507267	22.0462126	0.1968412	0.0098421
1 grain = 0.064799 gramme. 1 troy oz. = 31.103496 grammes. 1 lb. avd. = 453.593 kilogr. 1 cwt. = 50.802377 kilogra.					

WEIGHTS AND MEASURES OF THE BRITISH PHARMACOPŒIA OF 1885.

WEIGHTS.

1 Grain	gr.			
1 Ounce	oz.		=	437.5 grains.
1 Pound	lb.	=	16 ounces	= 7000.0 "

MEASURES OF CAPACITY.

1 Minim	min.			
1 Fluidrachm	fl. dr.		=	60 minims.
1 Fluidounce	fl. oz.		=	8 fluidrachms.
1 Pint	O.		=	20 fluidounces.
1 Gallon	C.		=	8 pints.

MEASURES OF LENGTH.

1 inch	=	
12 inches	=	1 foot.
36 "	=	3 feet = 1 yard.

(1 cubic inch of distilled water at 62° F. and 30 inch barom.
= 252.458 grains.)

RELATION OF MEASURES TO WEIGHTS.

1 Minim is the measure of		0.9114583 grain	of water.
1 Fluidrachm	"	54.6875	grains "
1 Fluidounce	"	1 ounce or 437.5	" "
1 Pint	"	1.25 pound, or 8750.0	" "
1 Gallon	"	10 pounds, or 70,000.0	" "

(Gtt. = *guttæ*, drops. The term "drop" indicates a quantity which is indefinite, and should only be used when approximateness is alone desired.)

The preceding (De La Rue's) tables, pp. 606, 607, will be found useful.

WEIGHTS AND MEASURES OF THE U. S. PHARMACOPŒIA.

The following tables, from the United States Pharmacopœia of 1880, especially taken together with those of the Pharmacopœia of 1890, will be found extremely useful :

A.—MEASURES OF LENGTH.

I. RELATION OF METRIC TO THE UNITED STATES MEASURES OF LENGTH.

1 Mètre	=	39.370432 inches.
1 Décimètre	=	3.937043 "
1 Centimètre	=	0.393704 "
1 Millimètre	=	0.039370 "

II. RELATION OF UNITED STATES TO METRIC MEASURES OF LENGTH.

1 Yard (or 36 Inches) = 0.91439 Metre.

1 Foot (or 12 Inches) = 30.40 Centimetres.

Feet.		Centimetres.	Inches.		Centimetres.	Inch.		Centimetre.
11	=	27.9	5	=	12.7	$\frac{1}{2}$	=	12.5
10	=	25.4	4	=	10.2	$\frac{1}{4}$	=	6.25
9	=	22.9	3	=	7.6	$\frac{1}{8}$	=	3.12
8	=	20.3	2	=	5.1	$\frac{1}{16}$	=	1.54
7	=	17.8	1	=	2.5	$\frac{1}{32}$	=	1.00
6	=	15.2						

B.—MEASURES OF CAPACITY.

III. RELATION OF METRIC TO UNITED STATES FLUID MEASURES.

Cubic Centim.	Fluidounces.	Cubic Centim.	Fluidrachms.	Cubic Centim.	Minims.			
1,000	=	33.81	15	=	4.06	0.40	=	6.49
950	=	32.12	10	=	2.71	0.35	=	5.68
900	=	30.43	9	=	2.43	0.30	=	4.87
850	=	28.74	8	=	2.16	0.25	=	4.06
800	=	27.05	7	=	1.89	0.20	=	3.25
750	=	25.36	6	=	1.62	0.19	=	3.08
700	=	23.67	5	=	1.35	0.18	=	2.92
650	=	21.98	4	=	1.08	0.17	=	2.76
600	=	20.29				0.16	=	2.60
550	=	18.59	Cubic Centim.	Minims.	0.15	=	2.43	
500	=	16.90	3	=	48.69	0.14	=	2.27
450	=	15.22	2	=	32.46	0.13	=	2.11
400	=	13.53	1	=	16.23	0.12	=	1.95
350	=	11.84	0.95	=	15.42	0.11	=	1.79
300	=	10.14	0.90	=	14.61	0.10	=	1.62
250	=	8.45	0.85	=	13.80	0.09	=	1.46
200	=	6.76	0.80	=	12.98	0.08	=	1.30
150	=	5.07	0.75	=	12.17	0.07	=	1.14
100	=	3.38	0.70	=	11.36	0.06	=	0.97
30	=	1.01	0.65	=	10.55	0.05	=	0.81
			0.60	=	9.74	0.04	=	0.65
Cubic Centim.	Fluidrachms.	0.55	=	8.93	0.03	=	0.49	
25	=	6.76	0.50	=	8.12	0.02	=	0.32
20	=	5.41	0.45	=	7.30	0.01	=	0.16

IV. RELATION OF UNITED STATES TO METRIC FLUID MEASURES.

Minims.	Cubic Centim.	Minims.	Cubic Centim.	Minims.	Cubic Centim.
1	= 0.06	8	= 0.49	15	= 0.92
2	= 0.12	9	= 0.55	16	= 0.99
3	= 0.18	10	= 0.62	17	= 1.05
4	= 0.25	11	= 0.68	18	= 1.11
5	= 0.31	12	= 0.74	19	= 1.17
6	= 0.37	13	= 0.80	20	= 1.23
7	= 0.43	14	= 0.86	21	= 1.29

RELATION OF UNITED STATES TO METRIC FLUID MEASURES.—*Cont.*

Minims.	Cubic Centim.	Fluidrachms.	Cubic Centim.	Fluidounces.	Cub. Centim.
22	=	1.36	3	=	11.09
23	=	1.42	4	=	14.79
24	=	1.48	5	=	18.48
25	=	1.54	6	=	22.18
26	=	1.60	7	=	25.88
27	=	1.66	8	=	29.57
28	=	1.73	9	=	33.27
29	=	1.79	10	=	36.97
30	=	1.85	11	=	40.66
35	=	2.16	12	=	44.36
40	=	2.46	13	=	48.06
45	=	2.77	14	=	51.75
50	=	3.08	15	=	55.45
55	=	3.39	16	=	59.10
60	=	3.70	Fluidounces.		
70	=	4.31			
80	=	4.93	3	=	88.67
90	=	5.54	4	=	118.24
100	=	6.16	5	=	147.81
110	=	6.78	6	=	177.39
120	=	7.39	7	=	206.96
			8	=	236.53
			9	=	266.10
			10	=	295.68
				=	325.25
				=	354.82
				=	384.40
				=	413.97
				=	443.54
				=	473.11
				=	502.69
				=	532.26
				=	561.93
				=	591.50
				=	621.08
				=	650.65
				=	680.22
				=	709.80
				=	739.37
				=	768.94
				=	798.51
				=	828.09
				=	857.66
				=	887.23
				=	916.80
				=	946.38
				=	1892.75
				=	3785.51

C.—WEIGHTS.

V. RELATION OF METRIC TO APOTHECARIES' OR TROY WEIGHT.

Grammes.		Grains.		Grammes.		Grains.
0.0010	==	0.015	==	0.193	==	1.852
0.0013	==	0.019	==	0.231	==	2.006
0.0015	==	0.023	==	0.309	==	2.161
0.0020	==	0.031	==	0.386	==	2.315
0.0025	==	0.039	==	0.463	==	2.469
0.0030	==	0.046	==	0.540	==	2.623
0.0035	==	0.054	==	0.617	==	2.778
0.0040	==	0.062	==	0.694	==	2.932
0.0045	==	0.069	==	0.772	==	3.086
0.0050	==	0.077	==	0.849	==	3.241
0.0055	==	0.085	==	0.926	==	3.395
0.0060	==	0.093	==	1.003	==	3.549
0.0065	==	0.100	==	1.080	==	3.704
0.0070	==	0.108	==	1.157	==	3.858
0.0075	==	0.116	==	1.235	==	4.012
0.0080	==	0.123	==	1.312	==	4.167
0.0085	==	0.131	==	1.389	==	4.321
0.0090	==	0.139	==	1.466	==	4.475
0.0095	==	0.147	==	1.543	==	4.630
0.0100	==	0.154	==	1.698	==	4.784

RELATION OF METRIC TO APOTHECARIES' OR TROY WEIGHT.—*Cont.*

Grammes.		Grains.	Grammes.		Grains.	Grammes.		Grains.
0.320	=	4.938	13	=	200.621	39	=	601.862
0.330	=	5.093	14	=	216.053	40	=	617.294
0.340	=	5.247	15	=	231.485	50	=	771.617
0.350	=	5.401	16	=	246.918	60	=	925.941
0.360	=	5.556	17	=	262.350	70	=	1080.264
0.370	=	5.710	18	=	277.782	80	=	1234.588
0.380	=	5.864	19	=	293.215	90	=	1388.911
0.390	=	6.019	20	=	308.647	100	=	1543.235
0.400	=	6.173	21	=	324.079	125	=	1929.044
0.500	=	7.716	22	=	339.512	150	=	2314.852
0.600	=	9.259	23	=	354.944	200	=	3086.470
0.700	=	10.803	24	=	370.376	250	=	3858.087
0.800	=	12.346	25	=	385.809	300	=	4629.705
0.900	=	13.889	26	=	401.241	333	=	5144.118
1	=	15.432	27	=	416.673	350	=	5401.322
2	=	30.865	28	=	432.106	400	=	6172.940
3	=	46.297	29	=	447.538	450	=	6944.557
4	=	61.729	30	=	462.970	500	=	7716.174
5	=	77.162	31	=	478.403	600	=	9259.409
6	=	92.594	32	=	493.835	700	=	10802.644
7	=	108.026	33	=	509.268	750	=	11574.262
8	=	123.459	34	=	524.700	800	=	12345.879
9	=	138.891	35	=	540.132	900	=	13889.114
10	=	154.323	36	=	555.565	1000	=	15432.350
11	=	169.756	37	=	570.997			
12	=	185.188	38	=	586.429			

VI. THE RELATION OF APOTHECARIES' (OR TROY) TO METRIC WEIGHT.

Grains.		Grammes.	Grains.		Grammes.	Grains.		Grammes.
$\frac{1}{64}$	=	0.00101	$\frac{1}{4}$	=	0.01620	14	=	0.90718
$\frac{1}{60}$	=	0.00108	$\frac{1}{3}$	=	0.02160	15	=	0.97198
$\frac{1}{50}$	=	0.00130	$\frac{1}{2}$	=	0.03240	16	=	1.037
$\frac{1}{48}$	=	0.00135	$\frac{3}{4}$	=	0.04860	17	=	1.102
$\frac{1}{40}$	=	0.00162	1	=	0.06480	18	=	1.166
$\frac{1}{36}$	=	0.00180	$1\frac{1}{2}$	=	0.09720	19	=	1.231
$\frac{1}{32}$	=	0.00202	2	=	0.12960	20	=	1.296
$\frac{1}{30}$	=	0.00216	$2\frac{1}{2}$	=	0.16200	21	=	1.361
$\frac{1}{25}$	=	0.00259	3	=	0.19440	22	=	1.426
$\frac{1}{24}$	=	0.00270	4	=	0.25920	23	=	1.458
$\frac{1}{20}$	=	0.00324	5	=	0.32399	24	=	1.555
$\frac{1}{18}$	=	0.00360	6	=	0.38879	25	=	1.620
$\frac{1}{16}$	=	0.00405	7	=	0.45359	26	=	1.685
$\frac{1}{15}$	=	0.00432	8	=	0.51839	27	=	1.749
$\frac{1}{12}$	=	0.00540	9	=	0.58319	28	=	1.814
$\frac{1}{10}$	=	0.00648	10	=	0.64799	29	=	1.869
$\frac{1}{8}$	=	0.00810	11	=	0.71297	30	=	1.944
$\frac{1}{6}$	=	0.01080	12	=	0.77759	40	=	2.592
$\frac{1}{5}$	=	0.01296	13	=	0.84239	50	=	3.240

RELATION OF APOTHECARIES' (OR TROY) TO METRIC WEIGHT.—*Cont.*

Drachms.		Grammes.	Ounces.		Grammes.	Ounces.		Grammes.
1	=	3.888	1½	=	46.655	11	=	342.138
2	=	7.776	2	=	62.207	12	=	373.250
3	=	11.664	3	=	93.310	13	=	404.345
4	=	15.552	4	=	124.414	14	=	435.449
5	=	19.440	5	=	155.517	15	=	466.552
6	=	23.328	6	=	186.621	16	=	497.656
7	=	27.216	7	=	217.724	17	=	528.759
			8	=	248.823	18	=	559.863
Ounces.			9	=	279.931	19	=	590.966
1	=	31.103	10	=	311.035	20	=	622.070

VII. RELATION OF METRIC TO AVOIRDUPOIS WEIGHT.

Avoirdupois Ounces and Grains.			Avoirdupois Ounces and Grains.			Avoirdupois Ounces and Grains.		
Grammes.	Oz.	Grs.	Grammes.	Oz.	Grs.	Grammes.	Oz.	Grs.
28.35	=	1	50	=	1 334	500	=	17 279
29	=	1 10	60	=	2 50½	550	=	19 175
30	=	1 25½	70	=	2 205	600	=	21 72
31	=	1 41	80	=	2 359	650	=	22 405½
32	=	1 56½	90	=	3 76½	700	=	24 303
33	=	1 72	100	=	3 230½	750	=	26 198½
34	=	1 87½	150	=	5 127	800	=	28 96
35	=	1 103	200	=	7 24	850	=	29 429
36	=	1 118	250	=	8 358	900	=	31 326½
37	=	1 133½	300	=	10 255	950	=	33 222
38	=	1 149	350	=	12 151½	1000	=	35 120
39	=	1 164½	400	=	14 48			
40	=	1 180	450	=	15 382			

VIII. RELATION OF AVOIRDUPOIS TO METRIC WEIGHT.

Avoirdupois Ounces.		Grammes.	Avoirdupois Ounces.		Grammes.	Avoirdupois Pounds.		Grammes.
1/16	=	1.772	7	=	198.447	1	=	453.592
1/8	=	3.544	8	=	226.796	2	=	907.18
1/4	=	7.088	9	=	255.146	3	=	1360.78
1/2	=	14.175	10	=	283.496	4	=	1814.37
1	=	28.350	11	=	311.846	5	=	2267.96
2	=	56.699	12	=	340.195	6	=	2721.55
3	=	85.049	13	=	368.544	7	=	3175.14
4	=	113.398	14	=	396.894	8	=	3628.74
5	=	141.748	15	=	425.243	9	=	4082.33
6	=	170.098				10	=	4535.92

QUESTIONS AND EXERCISES.

Mention some advantages of decimal weights and measures.—What is the name of the chief unit of the metric decimal system of weights and measures?—Mention the names of the metric units of surface, capacity, and weight, and state how they are derived from the unit of length.—How are multiples of metric units indicated?—State the designations of submultiples of metric units.—How many mètres are there in a kilomètre?—How many millimètres in a mètre?—How many grammes in 5 kilogrammes?—How many milligrammes in $13\frac{1}{2}$ grammes?—In 1894 centigrammes how many grammes?—In a mètre measure 5 centimètres wide and 1 centimètre thick how many cubic centimètres?—How many litres are contained in a cubic mètre of any liquid?—State the British equivalent of the mètre.—How many square yards in an are?—How many fluidounces in a litre?—How many ounces in a kilogramme?—Give the relation of a metric ton (1000 kilos) to a British ton.—How many grains are there in 1 British ton?—How many ounces in 1 ton?—How many grains of water in 1 fluidrachm?—How many minims in 1 pint?—How many grains in 1 pint of water?—Whence is the British unit of length derived?


Specific Weight, or Specific Gravity.

The *specific* weight of a substance is its weight in comparison with the weights of similar bulks of other substances. This comparative heaviness of *solids and liquids* is conventionally expressed in relation to water: they are considered as being lighter or heavier than water. Thus, water being regarded as unity = 1, the relative weight, or specific weight, of ether is represented by the figures .720 (it is nearly three-fourths, .750, the weight of water), oil of vitriol by 1.843 (it is nearly twice, 2.000, as heavy as water). The specific weight of substances is, moreover, the weight of similar volumes *at sixty degrees* (60° F.); for the weight of a definite volume of any substance will vary according to temperature, becoming heavier when cooled and lighter when heated, different bodies (gases excepted) differing in their rate of contraction and expansion. While, then, specific weight (or, conventionally, *specific gravity*) is truly the comparative weight of equal bulks, the numbers which in Great Britain commonly represent specific gravities are the comparative weights of equal bulks at 60° F., water being taken as unity.* The standard of comparison for gases was formerly air, but is now usually hydrogen.

* The true weight of a body is its weight in air plus the weight of an equal bulk of air, and minus the weight of a bulk of air equal to the bulk of the brass or other weights employed; or, in other words, its weight *in vacuo* uninfluenced by the buoyancy of the air; but such a correction of the weight of a body is seldom necessary, or, indeed, desirable. *Density* is sometimes improperly regarded as synonymous with *specific gravity*. It is true that the density of a body is in exact proportion to its specific gravity, but the former is more correctly the comparative bulk of equal weights, while specific gravity is the comparative weight of equal bulks.

SPECIFIC GRAVITY OF LIQUIDS.

Procure any small bottle holding from 100 to 1000 grains (Fig. 67), and having a narrow neck; counterpoise it in a delicate balance; fill it to about halfway up the neck with pure distilled water having a temperature of 60° F.; ascertain the weight of the water, and, for convenience, add or subtract a drop or two, so that the weight shall be a round number of grains; mark the neck by a diamond or file-point at the part cut by the lower edge of the curved surface of the water.



Specific-gravity Bottles.

Consecutively fill up the bottle to the neck-mark with several other liquids, cooled or warmed to 60° F., first rinsing out the bottle once or twice with a small quantity of each liquid, and note the weights; the respective figures will represent the relative weights of equal bulks of the liquids. If the capacity of the bottle is 10, 100, or 1000 grains, the resulting weights will, without calculation, show the specific gravities of the liquids; if any other number, a proportional sum must be worked out to ascertain the weight of the liquids as compared with 1 (or 1000) of water. Bottles conveniently adjusted to contain 250, 500, or 1000 grains, or 100 or 50 grammes, of water, when filled to the top of their perforated stopper (Fig. 69), and other forms of the instrument (Figs. 68 and 70), are sold by all chemical-apparatus makers. Fig. 70 is that of a bottle extremely useful in ascertaining the specific gravities of very volatile liquids.

Verify some of the following stated specific gravities of substances official in the U. S. Pharmacopœia 1890:

Acid, Acetic	1.048	Acid, Lactic	1.213
" " dil.	1.008	" Nitric	1.414
" " Glacial	1.056-1.058	" " dil.	1.057
" Hydrobromic dil.	1.077	" Oleic	.900
" Hydrochloric	1.163	" Phosphoric	1.710
" " dil.	1.050	" " dil.	1.057


Acid, Sulphuric	1.835	Oleum Caryoph.	1.060-1.067
“ “ Aromat.	.939	“ Chenapodii	.970
“ “ dil.	1.070	“ Cinnamomi.	1.055-1.065
“ Sulphurous	1.035	“ Copaiba	.890
Æther	.725-.728	“ Coriandri	.870
Alcohol	.797	“ Cubebæ	.920
“ dil.	.936	“ Erigerontis	.850
Amyl Nitris	.870-.880	“ Eucalypti	.915-.925
Aq. Ammon.	.960	“ Fœniculi	.960
“ “ Fort.	.901	“ Gaultheriæ	1.175-1.185
Bals. Peru	1.135-1.150	“ Gossypii Sem.	.920-.930
Benzinum	.670-.675	“ Hedeomæ	.930-.940
Bromum	2.990	“ Juniperi	.850-.890
Camphora	.995	“ Lavandulæ Flor.	.885-.897
Carbonei Disulphidum {	1.268	“ Limonis	.858-.859
	1.269	“ Lini	.830-.840
Cera Alba	0.965-0.975	“ Menth. Pip.	.900-.920
Cera Flava	.955-.967	“ “ Virid.	.930-.940
Cetaceum	0.945	“ Morrhuæ	.920-.925
Chloroform	1.490	“ Myrciæ	.975-.990
Copaiba	.940-.990	“ Myristicæ	.870-.900
Creasote	1.070	“ Olivæ	.915-.918
Fel Bovis	1.018-1.028	“ Picis Liquida	.970
Glycerinum	1.250	“ Pimentæ	1.045-1.055
Hydrargyrum	13.558	“ Ricini	.950-.970
Iodoformum	2.000	“ Rosæ	.865-.880
Liq. Calcis	1.0015	“ Rosmarini	.895-.915
“ Ferri Chloridi	1.387	“ Sabinæ	.910-.940
“ “ Citratis	1.250	“ Santali	.970-.978
“ “ Nitratis	1.050	“ Sassafras	1.070-1.090
“ “ Subsulph.	1.550	“ Sesami	.919-.923
“ “ Tersulph.	1.320	“ Sinapis vol.	1.018-1.029
“ Hydrarg. Nit.	2.100	“ Terebinthinæ	.855-.870
“ Plumbi Subacetatis	1.195	“ Thymi	.900-.930
“ Potassæ	1.036	“ Tiglii	.940-.960
“ Potassii Citratis	1.059	Petrolatum liquid.	.875-.945
“ Sodæ	1.059	Phosphorus (at 50° F.)	1.830
“ “ Chloratæ	1.052	Resina	1.070-1.080
“ Sodii Silicatis	1.300-1.400	Sp. Ætheris Nitrosi	.842-.836
“ Zinci Chlor.	1.535	“ Ammoniaæ	.810
Mel	1.101-1.105	“ “ Aromat.	.905
Oleum Adipis	0.910-0.920	“ Frumenti	.930-.917
“ Æthereum	0.910	“ Vini Gallici	.941-.925
“ Amygd. Amar.	1.060-1.070	Syrupus	1.317
“ “ Express.	.915-.920	Syr. Acidi Hydriodici	1.313
“ Anisi	.980-.990	Thymol	1.069
“ Aurantii Cort.	.850	Tinct. Ferri Chloridi	0.960
“ “ Flor.	.875-.890	Vinum Album	.990-1.010
“ Bergamottæ	.880-.885	“ Rubrum	.989-1.010
“ Cajuputi	.922-.929	Zincum	6.9-7.2
“ Cari	.910-.920		

Hydrometers, formerly termed Areometers.—The specific gravity of liquids may be ascertained, without scales and weights, by means of an *hydrometer*, an instrument, usually of glass, having a graduated stem and a bulb or bulbs at the lower part. The specific gravity of a liquid is indicated by the depth to which the hydrometer sinks in the liquid, the zero of the scale marking the depth to which it sinks in pure water. Hydrometers constructed for special purposes are known under the names of saccharometer, lactometer, elæometer, urinometer, alcoholometer. Hydrometers require a considerable quantity of liquid fairly to float them, and specific gravities observed with them are less delicate and trustworthy than those obtained by the balance; nevertheless, they are exceedingly useful for many practical purposes where the employment of a delicate balance would be inadmissible.

SPECIFIC GRAVITY OF SOLIDS IN MASS.

Weigh a piece (50 to 250 grains) of any solid substance heavier than water in the usual manner. Then weigh it in water, by suspending it from a shortened balance-pan by a fine thread or hair and immersing in a vessel of water (Fig. 71). The buoyant properties of the water will cause the solid apparently to lose weight; *this loss in weight is the exact weight of*

Fig. 71.



Weighing a Solid in Water.

an equal bulk of water. The weight of the substance and the weight of an equal bulk of water being thus ascertained, a proportional sum shows the relative weight of the substance to 1.000 of water. To express the same thing by rule, divide the weight in air by the loss of weight in water; the resulting number is the specific gravity in relation to 1 part of water, the conventional standard of comparison.

Verify some of the following specific gravities :

Aluminium	2.56
Antimony	6.71
Bismuth	9.83
Coins, English, gold	17.69
“ “ silver	10.30
“ “ bronze	8.70
Copper	8.95
Gold	19.34
Iron	7.84
Lead	11.36
Magnesium	1.74
Marble	2.70
Phosphorus	1.77
Platinum	21.53
Silver	10.53
Sulphur	2.05
Tin	7.29
Zinc	7.14

Specific gravities of solid substances should be taken in water having a temperature of about 60° F. The body should be immersed about half an inch below the surface of the water; adhering air-bubbles must be carefully removed; the body must be quite insoluble in water.

(For a table of the specific gravities of a large number of fatty and resinoid substances see the *Pharmaceutical Journal* for Oct. 11, 1879.)

SPECIFIC GRAVITY OF SOLIDS IN POWDER OR SMALL FRAGMENTS.

Weigh the particles; place them in a counterpoised specific-gravity bottle of known capacity, and fill up with water, taking care that the substance is thoroughly wetted; again weigh. From the combined weights of water and substance subtract the amount due to the substance: the residue is the weight of the water. Subtract this weight of water from the quantity which the bottle normally contains: the residue is the amount of water displaced by the substance. Having thus obtained the weights of equal bulks of water and substance, a proportional sum shows the relation of the weight of the substance to 1 part of water—*i. e.* the specific gravity.

Or, suspend a cup, short glass tube, or bucket from a shortened balance-pan; immerse in water; counterpoise; place the weighed powder in the cup, and proceed as directed for taking the specific gravity of a solid in mass.

This operation may be conducted on fragments of any of the substances the specific gravities of which are given in the foregoing

table, or on the powdered piece of marble the specific gravity of which has been taken in mass. The specific gravity of one piece of glass, first in mass then in powder, may be ascertained; the result should be identical. The specific gravity of shot is about 11.350; sand, 2.600; mercury, 13.56.

SPECIFIC GRAVITY OF SOLIDS SOLUBLE IN WATER.

Weigh a piece of sugar or other substance soluble in water; then suspend it from a balance in the usual manner, and weigh it in turpentine, benzol, or petroleum the specific gravity of which is known or has been previously determined; the loss in weight is the weight of an equal bulk of the turpentine. Ascertain the weight of an equal bulk of water by calculation :

Sp. gr. of	:	sp. gr. of	::	observed	:	equal bulk of
turpentine	:	water	::	bulk of turp.	:	water.

The exact weights of equal bulks of sugar and water being obtained, the weight of a bulk of sugar corresponding to 1.000 of water is shown by a proportional sum; in other words, divide the weight of sugar by that of the equal bulk of water; the quotient is the specific gravity of sugar. The stated specific gravity of sugar ranges from 1.590 to 1.607.

SPECIFIC GRAVITY OF SOLIDS LIGHTER THAN WATER.

This is obtained in a manner similar to that for solids heavier than water; but the light body is sunk by help of a piece of heavy metal, the bulk of water which the latter displaces being deducted from the bulk displaced by both; the remainder is the weight of a bulk of water equal to the bulk of the light body. For instance, a piece of wood weighing 12 grammes (or grains, for it is assumed that the student works equally well with metric as with imperial weights) is tied to a piece of metal weighing 22 grammes, the loss of weight of the metal in water previously having been found to be 3 grammes. The two, weighing 34 grammes, are now immersed, and the loss in weight found to be 26 grammes. But of this loss 3 grammes have been proved to be due to the buoyant action of the water on the metal; the remaining 23 therefore represent the same effect on the wood; 23 and 12 therefore represent the weights of equal bulks of water and wood. As 23 are to 12, so is 1 to .5217. Or, shortly, as before, divide the weight in air by the weight of an equal bulk of water; .5217 is the specific gravity of the wood. Another specimen of wood may be found to be three-fourths (.750) the weight of water, and others heavier. Cork varies from .100 to .300.

The specific gravity of a very minute quantity of a heavy or light substance may be ascertained by noting the specific gravity of a fluid in which it, being insoluble, neither sinks nor swims; or by immersing it in a weighed piece of paraffin whose specific gravity is known, noting the specific gravity of the whole and deducting the influence of the paraffin.

SPECIFIC GRAVITY OF GASES.

This operation is similar to that for liquids. A globe exhausted of air and holding from 1 to 4 litres (or quarts) is suspended from the arm of a balance and counterpoised by a similar flask. Gases are introduced in succession and their weights noted. A proportional sum shows their specific gravity in relation to air or hydrogen, whichever be taken as a standard,

Correction of the Volume of Gases for Pressure.—The height of the barometer at the time of manipulation is noted. Remembering that “the bulk of a gas is inversely as the pressure to which it is subjected” (Boyle and Mariotte), a simple calculation shows the volume which the gas would occupy at 760 millimètres (or 29.992 inches), the standard pressure (30 inches is sometimes adopted as the standard in England).* Thus, 40 volumes of a gas at 740 millims. pressure are reduced to 39 when the pressure becomes 760 millimètres (or 90 vols. at 29 ins. barom. become 87 vols. at 30 inches).

Correction of the Volume of Gases for Temperature.—This is done in order to ascertain what volume the gas would occupy at 0° C. (32° F.) or 15.5° C. (60° F.), according to the standard taken. Gases are equally affected by equal variations in temperature (Charles). They expand about 0.3665† per cent. ($\frac{1}{273}$) of their volume at the freezing-point of water for every C. degree (0.2036, or $\frac{1}{491}$, for every F. degree) (Regnault). Thus 8 volumes of gas at 0° C. will become 8.293 at 10° C.; for if 100 become 103.665 on being increased in temperature 10° C., 8 will become 8.293 (or if 100 become 102.036 on being increased 10° F., 8 will become 8.1629).

Vapor-density.—Vapors are those gases which condense to liquids at common temperatures. By the density of a vapor is meant its specific gravity. The density of a vapor is the ratio of any given volume to a similar volume of air or hydrogen at the same temperature and pressure. But, for convenience of comparison, this experimental specific gravity is referred, by calculation as just described

* In France the conventional standard height of the barometer is 760 millimètres at 0° C. (32° F.): in England it is 30 inches, the temperature being 60° F. 760 millims. is equivalent to 28.922 inches; but the expansion of the metal between 32° F. and 60° F. increases the length of the column to 30.005 inches. The standards are therefore almost identical, difference in true length being counterbalanced by the temperature at which the length is observed.

† Corrected for the difference between the mercurial and air thermometers, the coefficient of expansion of air is 0.003656 (Miller). The coefficient of expansion of different gases varies very slightly, being somewhat higher for the more liquefiable gases.

for permanent gases, to a temperature of 0°C. , and 760 millimètres barom. A teaspoonful or so of liquid is placed in a weighed flask of about the capacity of a common tumbler and having a capillary neck: the flask is heated in an oil-bath to a temperature considerably above the boiling-point of the liquid; at the moment vapor ceases to escape the neck is sealed by a blowpipe flame and the temperature of the bath noted; the flask is then removed, cooled, cleaned, and weighed; the height of the barometer is also taken. The neck of the flask is next broken off beneath the surface of water (or of mercury), which rushes in and fills it, and again weighed, by which its capacity in cub. centims. is found. From these data the volume of vapor yielded by a given weight of liquid is ascertained by a few obvious calculations. The capacity of the globe having been ascertained, the weight of an equal bulk of air* is obtained by a rule-of-three sum. This weight of air is deducted from the original weight of the flask, which gives the true weight of the glass. The weight of the glass is next subtracted from the weight of the flask and contained vapor (now condensed), which gives the weight of material used in the experiment. The volume which this weight of material occupied at the time of experiment is next corrected for temperature (to 0°C.) and pressure (760 millimètres) in the manner just described. The weight of a similar volume of hydrogen is next found.† The weights of equal volumes of hydrogen and vapor being thus determined, the amount of vapor corresponding to 1 of hydrogen (the specific gravity or vapor-density) is shown by a short calculation. This process of finding the weight of a given volume of vapor is by Dumas. Gay-Lussac's consists in determining the volume of a given weight: it has been improved by Hofmann. An easy and excellent method by V. and C. Meyer consists, like that of Gay-Lussac, in determining the volume of the vapor of a given weight of a fluid or solid, but differs in the volume of the vapor being ascertained from an equal volume of air which the vapor is made to displace. (For a detailed description of this method, and a drawing of the apparatus, see *Pharmaceutical Journal*, May 17, 1879.)

Experiment shows that the specific gravities of many gases and vapors on the hydrogen scale and the proportions in which they combine by weight are identical. Thus, chlorine is 35.5 times as heavy as hydrogen, and 35.5 parts unite with 1 of hydrogen to form hydrochloric acid gas. Hence, if the specific gravity of a gas or vapor is known, its combining proportion may be predicated with

* 1 cubic centimètre of air at 0°C. and 760 millim. weighs 0.001293 gramme.

† 1 litre (1000 cub. centims.) of hydrogen at 0°C. and 760 millimètres (the barometer being at 0°C.) weighs 0.0896 gramme—a quantity sometimes termed a *crith* (from κριθή, *krithē*, a barley-corn—figuratively, a small weight); thus a litre of oxygen weighs 16 criths, chlorine 35.5 criths, etc. 100 cubic inches of hydrogen at 32°F. weigh 2.265 grains; at 60°F. 2.143 grains (the barometer being 30 ins. at 60°F. in both cases). 100 cubic inches of air at 32°F. weigh 32.698 grains; at 60°F. , 30.935 (barom. 30 ins. at 60°F.). 1 cubic inch of water weighs 252.458 grains (Chaney, 252.279) at 62°F. and 30 in. bar. 1 gallon of water contains 277½ (277.274 at 62°F.) cubic ins. 1 cubic foot contains about 6½ gallons.

reasonable certainty, and *vice versâ*. In applying this rule to gaseous or vaporous compounds attention must be paid to the extent to which their constituent gases contract at the moment of combination or expand at the moment of decomposition. Thus steam is found to be composed of two volumes of hydrogen and one of oxygen, the three volumes of constituents condensing to two at the moment of combination. Hence steam may be expected to be nine times as heavy as hydrogen, which experiment confirms.

These relations may be so expressed as to include both elementary and compound gases and vapors, thus: *molecular weights and specific weights are identical*. Molecular weights represent two volumes of a gas: specific gravity conventionally represents the relative weight of a gas compared with one volume of hydrogen or air; hence the specific gravity of a gas or vapor on the H scale is found by calculation on simply dividing the molecular weight by 2; on the air-scale, by dividing the hydrogen numbers by 14.44. For example:

Name.	Molecular formula.	Molecular weight.	Specific gravity.		
			H=2.	H=1.	Air=1.
Hydrogen	H ₂	2	2	1	.069
Chlorine	Cl ₂	71	71	35.5	2.460
Oxygen	O ₂	32	32	16	1.108
Nitrogen	N ₂	28	28	14	.970
Steam	H ₂ O	18	18	9	.624
Ammonia gas	NH ₃	17	17	8.5	.589
Carbonic acid gas . .	CO ₂	44	44	22	1.524
Alcohol (vapor) . . .	C ₂ H ₆ O	46	46	23	1.593
Air	—	—	28.88	14.44	1.000

In other words, it follows, as an arithmetical necessity, that if the specific gravities of gases or vapors have been rightly determined, and the molecular weights of those gases and vapors have been quite accurately ascertained, the product of the division of the figures showing the molecular weights by the figures showing the specific gravities will be quotients that will always be the same number. If the specific gravity be in relation to two volumes of hydrogen, the quotient will in all cases be the figure 1, or, obviously, the figures for specific gravity and molecular weight will be identical. If the specific-gravity figure be in relation to one volume of hydrogen, the quotient will be the figure 2; if in relation to air, the quotient will be 28.88 (air being 14.44 times heavier than hydrogen). By multiplying 28.88 by the specific gravity on the air-scale the experimental figure obtained for molecular weight can be checked. So by dividing the molecular weight by 28.88 the specific gravity on the air-scale can be checked. Once more, divide the molecular weight by the specific gravity on the air-scale, and the quotient will not be far from 28.88 if the two figures have been ascertained with all attainable experimental exactitude.

The foregoing columns of specific gravities closely correspond with those obtained by actual experiment. The specific gravity of any gas or vapor may therefore be calculated if the following data are at hand: (a) formula, (b) atomic weight of constituent elements:

these give the molecular weight, and *the molecular weight divided by 2 is the specific gravity* on the hydrogen scale. Specific gravity on the air-scale is then deducible, if (c) the specific gravity of air (14.44) in relation to hydrogen be remembered. The absolute weight of any volume of a gas or vapor on the metric system is then obtainable if (d) the weight of a litre of hydrogen (0.0896 gramme) be known, or on the English plan by remembering (e) that 100 cubic inches of hydrogen at 60° F. weigh 2.143 grains (100 cubic inches of air at 60° F. weigh 30.935 grains).

In confirmation of these statements regarding the mutual relation of specific gravity and atomic weight a remarkable fact may be mentioned. Regnault several years ago found the weight of 1 litre of hydrogen and oxygen to be respectively .089578 gramme and 1.429802 grammes. The latter number divided by the former gives 15.96 as the specific gravity of oxygen. Stas, in recent experimental researches on combining proportion, finds the atomic weight of oxygen to be not 16, but 15.96.

Exceptions to the law occur in a few compounds and in arsenum and phosphorus, whose vapor densities are twice that indicated by the rule. Possibly in these cases the temperature employed is insufficient to dissociate an unusually complex molecule into molecules of usual complexity. As regards compounds, and possibly as regards those elements in which the observed density is only half that indicated by this rule, heat may, and in some cases probably does, produce molecular *dissociation* (*thermolysis*) into free atoms (uniatomic molecules) or into less complex molecules. (See p. 195.)

Relation of the Specific Heat of Elements to their Atomic Weights.—Reference may here appropriately be made to a physical fact of great importance as regards molecular and atomic weights. In the earlier pages of this Manual it was stated that elements do not combine chemically in haphazard proportions, but in fixed weights; and abundant evidence of the truth of the statement has already been afforded, and will also be found in this section on quantitative analysis. Secondly, it has been shown that elements do not combine in haphazard proportions by volume, but in certain constant bulks; and the weights of these bulks have been found to be identical with the combining weights themselves. Thirdly (this is the point to which attention is now drawn), if equal amounts of heat be given to the elements in the solid state (that is, to solid elements or to solid compounds of volatile elements), and the quantity of the element be increased or diminished until each is thus heated through an equal number of degrees, it will be found that the different weights of elements required are (in relation to a common standard) identical with the combining weights of the elements and with the weights of the combining volumes of the elements. Thus, where 108 parts of silver would be employed 207 of lead would be necessary.* Hence, in the determination of (a) combining

* Obviously, if equal weights of silver and lead were heated through an equal number of degrees, the silver would absorb nearly twice as much heat as the lead. In fact, as regards all the solid elements "specific heats and atomic weights are inversely proportional." This law

proportion, (*b*) specific gravity in gaseous state, and (*c*) specific heat, three distinct methods of ascertaining atomic weight are available. In cases where one method is inapplicable recourse is had to either, or, if practicable, both of the others, and thus the trustworthiness of observations and generalizations placed more or less beyond question. The specific heat of a solid element is the same in the free as in the combined condition; therefore the specific heat of a molecule is the sum of the specific heats of its constituent atoms. From the specific heat of a solid compound of a volatile element (chlorine, for example) can thus be calculated the specific heat of an element in the solid state, even though the free element cannot itself be solidified. For the processes by which experimentally to determine specific heat the reader is referred to books on Physics.

There is equivalency also between *electrical* and chemical action. The amount of electricity which, passed through an iodide, would, by the resulting *electrolysis* (electric *loosing*, from *λύω, luo*, to decompose), set free 127 parts of iodine, would set free 80 parts of bromine.

Again, Raoult, reflecting that the degree to which the solidifying point of a fluid is affected by a dissolved substance is dependent on the molecular weight of that substance, has shown that determinations of such degrees point to molecular weights. (Further researches in this direction will probably throw much light on the nature of the phenomena of *solution*.) Optical activity and molecular weight would also seem to be interdependent.

QUESTIONS AND EXERCISES.

Define specific weight, or, as it is commonly termed, specific gravity.—In speaking of light and heavy bodies specifically, what standard of comparison is conventionally employed?—How are specific gravities expressed in figures?—Why should specific gravities be taken at one constant temperature?—How does the buoyancy of air affect the real weight of any material?—Describe the difference between density and specific gravity.—Give a direct method for taking the specific gravity of liquids.—A certain bottle holds 150 parts, by weight, of water, or 135.7 of spirit of wine; show that the specific gravity of the latter is 0.9046.—An imperial fluidounce of a liquid weighs $366\frac{1}{2}$ grains; prove that its specific gravity is 0.838.—Equal volumes of benzol and glycerin weigh 34 and 49 parts respectively, and the specific gravity of the benzol is 0.850; show that the specific gravity of the glycerin is 1.225.—Explain the process employed in taking the specific gravity of solid substances in mass and in powder.—State the method by which the specific gravity of a light body, such as cork, is obtained.—What modifications of the usual method are necessary in ascertaining the specific gravity of substances soluble in water?—How is the specific gravity of gases deter-

was discovered by Dulong and Petit. It follows that the product of the multiplication of the figures representing the specific heat of an element with the figures representing its atomic weight is in the case of nearly every such element the same number—namely 6, more or less according to the care with which the experiments have been conducted. Some exceptions to the law have disappeared when the experiments have been conducted at a sufficiently high temperature.

mined?—By what law can the volume of a gas, at any required pressure, be deduced from its observed volume at another pressure?—To what extent will 78 volumes of a gas at 29.3 inches barom. alter in bulk when the pressure is 30.2 inches?—Write a short account of the means by which the volumes of gases are corrected for temperature.—At the temperature of 15° C. 40 volumes (litres, pints, ounces, cubic feet, or other quantity) of a gas are measured. To what extent will this amount of gas contract on being cooled to the freezing-point of water (0° C.)? *Answer:* As 1 vol. of any gas at zero expands or contracts .003665 of a vol. for each rise or fall of 1° C., 1 vol. at 0° C. if heated to 15° C. will become increased by .054975 (that is, .003665 multiplied by 15); 1 vol. will expand to 1.054975. Conversely, 1.054975 vol. will contract to 1 vol. if cooled from 15° C. to 0° C. And if 1.054975 becomes 1 in cooling through 15° C., 40 vols. will (as found by rule of three) contract to 37.916.—10 litres of oxygen are measured off at 14° F. Required the volume of the gas at 15° C. *Answer:* The first operation must be to reduce the temperature quoted in Fahrenheit's degrees to an equivalent value on the Centigrade scale. 14° F. is 18° below 32° F., the freezing-point of water; and a range of 9° on the Fahrenheit scale is equal to a range of 5° on the Centigrade scale, so that the temperature at which the oxygen is measured off is -10° C. The rise of temperature up to 0° expands the gas in such proportion that its volume at 0° is to its volume at -10° as 1 is to 1 - 0.03665, *i. e.* as 1 to 0.96335. The further rise of temperature from 0° C. to 15° expands the gas in the proportion of 1 to 1 + 15 × 0.003665, *i. e.* 1 to 1.054975. The total rise of temperature therefore expands the gas in the proportion of 0.96335 to 1.054975.

$$0.96335 : 1.054975 :: 10 : x;$$

$$\therefore x = \frac{10 \times 1.054975}{0.96335} = 10.95.$$

—230 cubic centimètres of oxygen are measured off at 14° C. and 740 millimètres mercurial pressure. Required the volume of the gas at the normal temperature and pressure (0° C. and 760 millimètres). *Answer:* Let the reduction for change of temperature be made first. The proportion—

$$1 + (14 \times 0.003665) : 1 :: 230 : x$$

gives

$$x = \frac{230}{1.05131} = 218.774.$$

To reduce this volume at 740 millimètres pressure to the volume corresponding to the pressure of 760 millimètres, we have the proportion—

$$760 : 740 :: 218.77 : x;$$

whence

$$x = \frac{740 \times 218.77}{760} = 213.02.$$

—A litre of oxygen is confined in a glass flask at 10° C. by the atmospheric pressure, added to that of a column of mercury 60 millimètres high. The flask must be heated to 300° C. without any increase of volume taking place in the oxygen. How high must the column of mercury then be which presses on the gas, supposing the atmospheric pressure to remain constant at 760 millimètres? *Answer:* The oxygen is given at 10° C. and 820 millimètres pressure. If the pressure remain constant, the rise of temperature from 10° C. to 300° C. would expand the gas in such proportion that 1.03665 volumes would expand to 2.0995

volumes. In order to prevent any expansion the pressure must be increased in the same proportion, whence

$$1.03665 : 2.0995 :: 820 : x;$$

$$\therefore x = \frac{820 \times 2.0995}{1.03665} = 1660.6.$$

From this total pressure the atmospheric pressure of 760 millimètres has to be deducted, leaving 900.6 millimètres as the height of the required mercurial column.—A litre of oxygen is required of the density of 100 at 0° C. What weight of potassic chlorate must be used for its preparation, and what total pressure must be applied to it? *Answer:* The pressure required to compress oxygen from the density of 16 to that of 100 is found by the proportion—

$$16 : 100 :: 760 : x;$$

$$\therefore x = \frac{76000}{16} = 4750.$$

At the pressure of 4750 millimètres of mercury the weight of a litre of oxygen (16 grammes measure 11.2 litres at 0° C. and 760 millims. pressure) is found by the proportion—

$$760 : 4750 :: \frac{16}{11.2} : x;$$

whence

$$x = \frac{16 \times 4750}{11.2 \times 760} = 8.93 \text{ grammes.}$$

The weight of chlorate required for the evolution of 8.93 grammes of oxygen is found from the proportion—

$$48 : 122.5 :: 8.93 : x;$$

$$\therefore x = 22.8 \text{ grammes.}$$

—What is the volume of 12 grammes of hydrogen at 15° C.? *Answer:* 1 gramme of hydrogen measures 11.2 litres at 0° C.; therefore 12 grammes measure $12 \times 11.2 = 134.4$ litres at 0°. To find their volume at 15° C. we have the proportion—

$$1 : 1 + 15 \times 0.003665 :: 134.4 : x;$$

whence

$$x = 134.4 \times 1.054975 = 141.788 \text{ litres.}$$

(The foregoing five problems are from Williamson's "Chemistry.")

—What interest for chemists have the specific heats of substances?

VOLUMETRIC QUANTITATIVE ANALYSIS.

Preliminary Note.—Great care should be observed in selecting a *fair* sample of any bulk of material that is to be examined either by volumetric or gravimetric quantitative analysis. If the whole quantity is in separate parcels, and there is any ground for believing that the parcels differ in quality, they should, if practicable, be carefully mixed, or, technically,

"bulked." Small portions should be taken from different parts of the resulting heap and well mixed in a mortar or other vessel, or, in certain cases, dissolved and the solution well stirred or shaken. A specimen of the powder or a portion of the solution may then be selected for analysis.

Introduction.—The operations of volumetric analysis consist (a) in carrying out some definite *chemical reaction*, already well known to the operator, with (b) *definite quantities* of substances or salts; (c) the exact termination of the reaction between the two salts or substances being ascertained, usually by some *chemical indicator* (litmus, starch, etc.). A portion of the substance or salt, etc. to be tested is carefully weighed. To this is gradually added the second substance or salt contained in the testing fluid, commonly termed the Standard Volumetric Solution. The usefulness, and, indeed, the preparation, of this standard solution is founded (as already indicated on p. 592) on some accurate initial gravimetric operation. A *weighed* amount of a pure salt is dissolved in a given *volume* of water. "Accurately *measured* quantities of such a standard volumetric solution will obviously contain just as definite amounts of the dissolved salt as if those amounts were actually *weighed* in a balance, and as measuring occupies less time than weighing, the volumetric operations can be conducted with great economy of time as compared with the corresponding gravimetric operations."

Normal solutions contain (a) one molecular weight in grammes, per litre, of substances, the molecular weights of which are chemically comparable with one atomic weight of hydrogen (e. g. $\text{HCl} = 36.37$, $\text{KOH} = 55.99$, etc.), or (b) such a proportion of the molecular weight as in the volumetric reactions is comparable with one atomic weight of hydrogen (e. g. $\text{H}_2\text{SO}_4 = 97.82 \div 2 = 48.91$, or $\text{K}_2\text{Mn}_2\text{O}_8 = 315.34 \div 10 = 31.534$). *Decinormal solutions* are one-tenth ($\frac{N}{10}$) the strength of normal solutions; *centinormal* ($\frac{N}{100}$), one-hundredth; *seminormal* ($\frac{N}{2}$), one-half; *double-normal* ($\frac{2}{N}$), twice; and so on.


APPARATUS.

The only special vessels necessary in volumetric quantitative operations are—1. A *litre flask* (Fig. 73), which, when filled to a mark on the neck, contains, at about 15°C . or 60°F ., 1 litre (1000 cubic centimètres—i. e. 1000 grammes) of water;* it serves for preparing solutions in quantities of 1 litre. 2. A tall, cylindrical *graduated litre jar* (Fig. 72) divided into one hundred equal parts; it serves for the measurement and admixture of decimal or centesimal parts of 1 litre. 3. A graduated tube or *burette* (Fig. 74), the marked portion of which, when filled to "0," holds 100 cubic centi-

* A cubic centimètre is, strictly speaking, the volume occupied by 1 gramme of distilled water at its point of greatest density—namely, 4°C . Metric measurements, however, are uniformly taken at 15° or 15.55°C . (59° or 60°F .).

mètres, and is divided into one hundred equal parts, with subdivisions; it is used for accurately measuring small volumes of liquids.

The best form of burette is Mohr's. It consists of a glass tube, commonly about the width of a little finger and the length of an arm



A Litre Jar.

A Litre Flask.

A Burette, etc.

from the elbow, contracted at the lower extremity and graduated. The width and length of burettes, however, as well as the extent and fineness of their graduation, vary considerably. To the contracted portion is fitted a small piece of vulcanized caoutchouc tubing, into the other end of which a small spout made of narrow glass tube is tightly inserted. A strong wire clamp effectually prevents any liquid from passing out of the burette unless the knobs of the clamp are pressed by the finger and thumb of the operator, when a stream or drops flow at will. Avoid accumulation of air behind the india-rubber. In place of the india-rubber tubing and clamp, a stopcock is often employed, and is to be preferred, though other modes of arresting the flow of liquid may be adopted. The accurate reading of the height of a solution in the burette is a matter of great importance: it should be taken from the bottom of the curved surface of the liquid. It may be still more exactly measured by the employment of a hollow glass float or bulb (Erdmann's float, see Fig. 74), of such a width that it can move freely in the tube without undue friction, and so adjusted in weight that it shall sink to more than half its length in any ordinary liquid. A fine line is scratched round the centre of the float; this line must always be regarded as marking the height of the fluid in the burette. In charging the burette a solution is poured in, not until its surface is coincident with 0, but until the mark on the float is coincident with 0.

ESTIMATION OF ALKALIES, ETC.

(Volumetric Solutions of Sulphuric Acid, $\text{H}_2\text{SO}_4 = 97.82$.)

The sulphuric radical being bivalent, and most of the metals contained in the salts which are estimated by sulphuric acid being univalent, it is convenient that each litre of this solution should contain half a molecular weight in grammes of the acid ($\text{H}_2\text{SO}_4 = 97.82 \div 2 = 48.91$). This is the "*Normal Sulphuric Acid*."

The standard sulphuric-acid solution is prepared by diluting oil of vitriol with three or four times its bulk of distilled water, and then determining the strength of this solution by titration with pure sodium carbonate, making use of the following memoranda:

$$\begin{array}{r} \text{Na}_2\text{CO}_3 + \text{H}_2\text{SO}_4 = \text{Na}_2\text{SO}_4 + \text{CO}_2 + \text{H}_2\text{O} \\ \hline \underbrace{2)106}_{53} \qquad \underbrace{2)97.82}_{48.91} \end{array}$$

Pure anhydrous sodium carbonate is easy to obtain, for commercial bicarbonate is usually of such purity that when a few grammes are heated to redness for a quarter of an hour the resulting carbonate is practically free from impurity. The bicarbonate should, however, be tested, and if more than traces of chlorides and sulphates are present, these may be removed by washing a few hundred grammes, first with a saturated solution of sodium bicarbonate, and afterward with pure distilled water. After drying, the salt is ready for ignition.

About half a gramme of the sodium carbonate is accurately weighed and placed in a half-litre flask, around the neck of which is tied calico or leather to protect the fingers when the heated vessel is shaken by the operator. The salt is dissolved in water to about one-third the capacity of the flask, and a few drops of the indicator, blue tincture of litmus, is added. The acid solution to be "set" or "standardized" is then poured into a burette, and run therefrom into the flask until the reddened litmus indicates the presence of free acid. This will be due in the first place to carbonic acid liberated and remaining dissolved in the solution. The contents of the flask are therefore boiled for several minutes, when the blue color will have returned. More acid is then run in until the mixture, after boiling, remains of a neutral color, indicating that just enough acid has been added to complete the reaction expressed in the foregoing equation.

Let it be supposed that .6 of a gramme of sodium carbonate was taken, and that this required 11 cc. of the sulphuric-acid solution; how many cc. of this solution would contain 48.91 grammes of pure sulphuric acid? or, what is equivalent in the reaction, how many cc. would be required to neutralize 53 grammes of sodium carbonate? As .6 of a gramme of the carbonate is to 11 cc. of solution, so are 53 grammes of the carbonate to x cc. of solution; therefore $x = 972$ cc. 972 cc. (nearly) are equivalent to 53 grammes of sodium carbonate, and contain 48.91 grammes of pure sulphuric acid.

This sulphuric-acid solution may be used as it is, or may be

diluted with water, every 972 cc. to be diluted to 1000 cc., so that 1000 cc. shall contain 48.91 grammes of sulphuric acid.

The following official substances are tested by this solution according to the United States Pharmacopœia:


Solutions of Ammonia.—2 or 3 grammes of dilute, or about 1 gramme of strong, solution of ammonia is a convenient quantity to operate upon. The weighing is most conveniently accomplished by taking a small stoppered bottle containing half an ounce or so of the substance, and having ascertained its total weight, transfer about the quantity desired to the flask in which the estimation is to be conducted, and again weigh the bottle with what remains in it. The difference is the exact quantity taken. The weighing of the ammonia solution having been accomplished, water is added, to about one-third the capacity of the flask (or, better, the ammonia is added to water already in the flask), and a few drops of tincture of litmus are introduced. The titration is then conducted as described before, except that no heat is employed.



$$\begin{array}{r} 2)70 \\ \hline 35 \end{array}$$

$$\begin{array}{r} 2)97.82 \\ \hline 48.91 \end{array}$$

48.91 = grammes in 1000 cc. of standard solution.



$$\begin{array}{r} 2)34 \\ \hline 17 \end{array}$$

$$\begin{array}{r} 2)97.82 \\ \hline 48.91 \end{array}$$

48.91 = grammes in 1000 cc. of standard solution.

1000 cc. of standard solution, or its equivalent of a solution of any other strength, would, according to this reaction, neutralize 17 grammes of ammonia gas (NH_3) or 35 grammes of ammonium hydrate (NH_4HO). If 3 grammes of ammonia solution had been taken, and it had required 15 cc. of standard oxalic-acid solution, then the amount of ammonia gas or ammonium hydrate it contained would be seen by the following calculations:

$$1000 \text{ cc.} : 17\text{NH}_3 :: 15 \text{ cc.} : x = .255 \text{ grammes NH}_3$$

$$1000 \text{ cc.} : 35\text{NH}_4\text{HO} : : 15 \text{ cc.} : x = .525 \text{ grammes NH}_4\text{HO}$$

3 grammes, then, would contain .255 grammes of the gas, or .525 grammes of ammonium hydrate. Or in percentage:

$$3 \text{ gr. sol.} : .255 \text{ gr. NH}_3 :: 100 \text{ gr. sol.} : x \text{ gr. NH}_3 = 8.5\% \text{ NH}_3$$

$$3 \text{ gr. sol.} : .525 \text{ gr. NH}_4\text{HO} : : 100 \text{ gr. sol.} : x \text{ gr. NH}_4\text{HO} = 17.5\% \text{ NH}_4\text{HO}$$

The solution would therefore contain 8.5 per cent. of ammonia gas (NH_3) or 17.5 per cent. of ammonium hydrate (NH_4HO). If the sulphuric-acid solution was not of the full standard, the number of cc. which contained 48.91 grammes of pure sulphuric acid—which was, in fact, equivalent to 1000 cc. of standard solution—would be substituted for 1000 cc. in the preceding proportions.


A comparison should now be made with the requirements of the Pharmacopœia. It is useful to express results as percentage of substance of pharmacopœial strength in the material examined. Thus

the U. S. Pharmacopœia requires dilute ammonia solution (both *Aqua Ammoniacæ* and *Spiritus Ammoniacæ*) to contain 10 per cent. of the gas (NH_3). The solution supposed to have been operated on contained 8.5 per cent. NH_3 ($10:8.5::100:x=85$). Therefore it contains 85 per cent. of the dilute ammonia of the U. S. Pharmacopœia.*

Strong Solution of Ammonia, U. S. P., contains 28 per cent. of ammonia gas (NH_3).

Note.—The calculations just described for ammonia are similar to those employed throughout Volumetric Analysis; they will not be repeated, therefore, in the case of every substance.


Ammonium Carbonate.—The reactions indicated by the following equations occur between commercial ammonium carbonate and sulphuric acid:



About 1 gramme is a convenient quantity to operate upon. If tincture of litmus be the indicator, the titration should be conducted at a temperature just short of boiling. The estimation is not very satisfactory, because the heat employed, while scarcely sufficient to expel the carbonic acid gas, is enough to occasion loss of ammoniacal salt. Practised analysts usually add excess of the standard acid, and thus fix every trace of ammonia; then gently boil to get rid of carbonic acid gas; bring back the liquid to neutrality by an observed volume of standard alkaline solution, and deduct an equivalent volume of acid from the quantity first added. The United States Pharmacopœia, however, orders the use of methyl-orange in place of litmus, which obviates the difficulty just mentioned. According

* Extremely minute quantities of ammonia—1 part in many millions of water—may be estimated volumetrically by adding excess of a colorless, strongly alkaline solution of red mercury iodide (Nessler's test), then in a similar vessel, containing an equal amount of pure water with excess of the Nessler reagent, imitating the depth of yellow or reddish-yellow color thus produced by adding an ammoniacal solution of *known strength*. The amount of ammonia thus added represents the amount in the original liquid.



The Nessler Reagent.—A litre may be made by dissolving 30 or 40 grammes of potassium iodide in a small quantity of hot water, adding a strong hot solution of mercury perchloride until the precipitate of mercuric iodide ceases to redissolve even by the aid of rapid stirring and heat, slightly diluting, filtering, adding a strong solution of (120 to 140 grammes) caustic soda or (160 to 180 grammes) caustic potash, and diluting to 1 litre. A few cc. (5 or 6 or more) of a strong solution of mercury perchloride are finally stirred in, the whole set aside till all precipitated red iodide has deposited, and the clear liquid decanted for use. The reaction of this Nessler test with ammonia is as follows:



Potassio-mercuric iodide, without alkali, is commonly known as *Mayer's Reagent*, HgI_2KI .




to official requirements, 5.23 grammes neutralize 100 cc. of the standard solution of sulphuric acid. This corresponds to 100 per cent. of carbonate having the formula $\text{N}_3\text{H}_{11}\text{C}_2\text{O}_5$.

Lead Acetate and Solution of Subacetate.—Operate upon about 3 grammes of lead acetate and from 5 to 10 grammes of solution of subacetate.






The flask in which the estimation is being conducted should previously contain one-third of water. In the case of both lead acetate and solution of subacetate a little acetic acid should be added to prevent precipitation of basic salt on dilution. The only indicator of complete reaction is cessation of production of the precipitate—lead sulphate. The British Pharmacopœia requires lead acetate to be pure (100 per cent.), and the U. S. P. requires the subacetate solution to contain 25 per cent.

Sodium, Caustic Potash and Soda, Potassium and Sodium Carbonates and Bicarbonates.—Methyl-orange is the indicator throughout, for the caustic alkalies always contain some carbonate.



Or,






Convenient quantities to operate with are—of caustic potash, 1 gramme; caustic soda, .5 to 1 gramme; potassium carbonate or bicarbonate, 1 to 2 grammes; sodium carbonate or bicarbonate, 2 to 3 grammes; dried sodium carbonate, .5 to 1 gramme; and of solutions a corresponding quantity. The United States Pharmacopœial requirements are—caustic potash or soda, 90 per cent. of KHO or NaHO; potassium carbonate, 95 per cent. of the anhydrous salt; anhydrous sodium carbonate, 98.9 per cent.; dried sodium carbonate, 73 per cent.; sodium bicarbonate, 98.6 per cent. *Liquor Potassæ* and *Liquor Sodæ*, U. S. P., must contain 5 per cent. of pure hydrate.

The strength of soda-ash is often reported in terms of "soda"—that is, oxide of sodium ($\text{Na}_2\text{O} = 62$). The old molecular weight of sodium carbonate, 54 (it should have been 53), derived from that of "soda," 32 (it should have been 31), is still employed in Great Britain in reporting the strength of soda-ash. The true amount of soda equivalent to 54 parts of carbonate is 31.41 parts. A modern analyst, having found the true amount of soda in a sample of soda-ash, is expected by some manufacturers to report 31 as 31.41 parts, or 53 of carbonate as 54, and other quantities in proportion to these figures.

Potassium and Sodium Tartrates and Citrates.—When tartrates or citrates of alkali-metals are burned in the open air, the whole of the metal remains in the form of carbonate. Each molecular weight of a neutral tartrate gives one molecular weight of carbonate, and every two molecular weights of an acid tartrate give one molecular weight of carbonate. Advantage is taken of these reactions to estimate indirectly the quantity of citrate or tartrate in presence of substances with which they are generally associated. 1 or 2 grammes of any of these salts is a convenient quantity to operate upon. The ignition may be conducted in a platinum or porcelain crucible. A low red heat only should be used, and the vessel removed when complete carbonization has been effected; that is to say, when nothing remains but the carbonate and free carbon. The mixture is in this case treated with hot water and the carbon separated by filtration. If too little heat has been used and carbonization is not complete, the filtrate will be more or less colored. If


this should be the case, the operation must be repeated with a fresh quantity of material. The carbonate is titrated in the usual way. The following equations, etc. explain the reactions :



It will be readily understood that in the first (for example) of the reactions just expressed 122 weights of potassium tartrate are equivalent to 69 weights of potassium carbonate; and as in a previous reaction it has been shown that 69 weights of potassium carbonate are equivalent to 48.91 grammes of pure H_2SO_4 , it follows that 122 weights of potassium tartrate are equivalent to 48.91 grammes of sulphuric acid. Let these weights be grammes, and then 122 grammes of potassium tartrate are equivalent to 48.91 grammes of sulphuric acid or to 1000 cc. of the standard solution of sulphuric acid. If the substance estimated be a crude sample of potassium tartrate, and the number of cc. of sulphuric acid used has been 15 cc., then as 1000 cc. of the acid solution are to 122 grammes of potassium tartrate, so are 15 cc. of the solution to 1.830 grammes of potassium tartrate. Now, if the weight of the sample taken was 2 grammes, then as 2 grammes of the sample contain 1.830 of real potassium tartrate, 100 will contain $x = 91.5$ per cent. of real tartrate. These salts are required to be 100 per cent. pure by the United States Pharmacopœia, except potassium acetate, which is to have 98 per cent. of real acetate. Trade samples are practically pure, as a rule. If calcium sulphate be present in tartrates or citrates, loss of potassium carbonate will ensue, potassium sulphate being formed. In estimating potassium acid tartrate, which is the salt most likely to contain calcium sulphate, direct titration without ignition may be followed.

Strontium Lactate.—To estimate this substance the strontium lactate is dried at 100°F. , and a small quantity taken—say about $1\frac{1}{2}$ grammes. This is then ignited, as in the case of Rochelle salt, and the residue dissolved in water and titrated with normal sulphuric acid,

methyl-orange as the indicator. The equation will explain the reaction :




$$\begin{array}{r} 2)265.3 \\ \hline 132.65 \end{array}$$

$$\begin{array}{r} 2)147.3 \\ \hline 73.65 \end{array}$$

73.65 equiv. to 1000 cc. of normal sulphuric acid.

The official substance should contain 98.6 per cent. of pure strontium lactate.


Lithium Carbonate, Benzoate, Citrate, and Salicylate.—All these substances may be estimated on the principles (by ignition, etc.) which have been explained in the case of sodium and potassium salts. The following equations will aid the student :



$$\begin{array}{r} 2)73.87 \\ \hline 36.94 \end{array}$$

$$\begin{array}{r} 2)97.82 \\ \hline 48.91 \end{array}$$

36.94 48.91 = grammes in 1000 cc. of standard solution.



$$\begin{array}{r} 2)127.72 \\ \hline 63.86 \end{array}$$

$$\begin{array}{r} 2)73.87 \\ \hline 36.94 \end{array}$$

36.94 equiv. to 1000 cc. of normal sulphuric acid.



$$\begin{array}{r} 6)419.14 \\ \hline 69.86 \end{array}$$

$$\begin{array}{r} 6)321.61 \\ \hline 53.60 \end{array}$$

36.94 equiv. to 1000 cc. of normal sulphuric acid.



$$\begin{array}{r} 2)287.36 \\ \hline 143.68 \end{array}$$

$$\begin{array}{r} 2)73.87 \\ \hline 36.94 \end{array}$$

36.94 equiv. to 1000 cc. of normal sulphuric acid.

In the case of the last three the residues, after ignition, are dissolved in water and titrated. It will be easily seen, on looking through the above equations, that in all four cases the lithium carbonate is being directly estimated, and indirectly the respective salts. The Pharmacopœia (U. S. P.) requires the carbonate to contain 98.98 per cent. of the pure salt, the benzoate 99.6, the citrate 99.2, and the salicylate 99.13 per cent. of the pure salt.

Decinormal Sulphuric Acid is sometimes more convenient to use than normal 1000 cc. = 4.891 grammes of sulphuric acid.


VOLUMETRIC SOLUTION OF OXALIC ACID.

(Crystallized Oxalic Acid, $\text{H}_2\text{C}_2\text{O}_4 \cdot 2\text{H}_2\text{O} = 125.7$.)

Standard Oxalic Acid may be used in the place of sulphuric acid. If pure crystallized oxalic acid be at hand, the *normal* solution is made by dissolving 62.85 grammes in water, and making the volume up with more water to exactly 1 litre.



Pure oxalic acid, however, not being easy to obtain, the solution may be made from the commercial acid by dissolving 65 to 70

grammes in enough water to make a litre of solution, and then determining the strength of this solution by a titration with pure sodium carbonate, making use of the following memoranda:




The *decinormal solution* will be one-tenth the strength, 6.285.

Liquor Calcis.—Measure out about 100 cc. of the lime-water, and weigh. The following equations, etc. are quantitative expressions of the reactions:



Phenolphthalein is used as an indicator. The official lime-water is to contain .14 per cent. $\text{Ca}(\text{OH})_2$.

Potassium Permanganate.—The reaction is shown in the following equation: $\text{K}_2\text{Mn}_2\text{O}_8 + 3\text{H}_2\text{SO}_4 + 5(\text{H}_2\text{C}_2\text{O}_4 \cdot 2\text{H}_2\text{O}) = \text{K}_2\text{SO}_4 + 2\text{MnSO}_4 + 18\text{H}_2\text{O} + 10\text{CO}_2$.



The salt satisfies official requirements if it contains 98.7 per cent of real potassium permanganate.

Normal Hydrochloric Acid Solution may be used in some cases, instead of sulphuric or oxalic acid. It is so made that 1000 cc. contain 36.37 grms. of pure HCl.

Weighing.—In the case of substances which are liable to alter by exposure to air it is important that a selected quantity should be quickly weighed, rather than selected weights be accurately balanced by material, the former operation occupying much the shorter time.

Salts other than the official may be quantitatively analyzed by the volumetric solutions of the Pharmacopœia, slight modifications of manipulation even enabling the processes to be adapted to fresh classes of salts.

QUESTIONS AND EXERCISES.

Describe the apparatus used in volumetric determination.—100 cubic centimetres of solution of oxalic acid contain 6.3 grammes of the crystallized salt; work sums showing what weights of potassium bicarbonate

and anhydrous sodium carbonate that volume will saturate. *Ans.* 10 grammes and 5.3 grammes.—Show what weight of potassium hydrate is contained in solution of potash 48.02 grammes of which are saturated by 50 cc. of the standard solution of sulphuric acid. *Ans.* 5.83 per cent.—Calculate the percentage of calcium hydrate in lime-water, 438 grammes of which are neutralized by 20 cc. of the volumetric solution of sulphuric acid. *Ans.* 0.1689.—8 grammes of a sample of Rochelle salt, after ignition, etc., require 54.3 cc. of the official sulphuric acid solution for complete saturation; work sums showing what is the centesimal proportion of real salt present. *Ans.* 95.7.


ESTIMATION OF ACIDS.

In the previous experiments a known amount of an acid has been used in determining unknown amounts of alkalies. In those about to be described a known amount of an alkali is employed in estimating unknown amounts of acids. The alkaline salt selected may be either a hydrate or a carbonate, but the former is to be preferred; for the carbonic acid set free when a strong acid is added to a carbonate interferes to some extent with the indications of alkalinity, acidity, or neutrality afforded by litmus. The alkali most convenient for use is either potash or soda, a solution of which has probably already been made the subject of experiment in operations with the standard solution of sulphuric acid. It should be kept in a stoppered bottle and exposed to air as little as possible.

VOLUMETRIC SOLUTION OF POTASH.

(Potassium Hydrate, KHO = 55.99, say 56.)


This aqueous solution of potash is most conveniently made of such a strength that each 1000 cc. contains one molecular weight in grammes of the alkali (KHO = 56). It will be seen from the following equation that 56 grammes of potash convert 48.91 grammes of sulphuric acid into the neutral potassium sulphate:



If pure caustic potash were at hand, it would only be necessary to weigh 56 grammes, dissolve this in water, and dilute to 1 litre. But pure potash cannot be readily produced. Therefore weigh about 60 grammes of commercial caustic potash, and add water to 1 litre. When dissolved take, say, 14 cc., dilute with more water in a flask, add a few drops of tincture of litmus, and titrate with sulphuric-acid solution of known strength. Suppose that the volume of standard acid solution required to neutralize the 14 cc. of potash solution, the strength of which is to be estimated, has been 15 cc., or an equivalent amount of acid solution of another strength; then, how many cc. of potash solution are equivalent to 1000 cc. of stand-


ard acid solution? or, what comes to the same thing, how many cc. of potash solution contain 66 grammes of real potash (KHO)? As 15 cc. standard acid are to 14 cc. potash solution, so are 1000 cc. standard acid to x cc. $x = 933$ cc. of the potash solution contains, therefore, 56 grammes of potash. This may either be diluted, every 933 cc. to 1000 cc., so that it may be standard (1000 cc. = 56 grammes KHO), or the solution may be used without dilution (933 cc. = 56 grammes KHO). It has already been mentioned that soda nearly always contains carbonate. To remove resulting carbonic acid, therefore, gentle heat should be employed toward the close of each titration in all the estimations with this solution if litmus is used as an indicator of completion of the reaction. There is, however, no need to warm if phenolphthalein be used instead of litmus, as is ordered in the United States Pharmacopœia. The following substances are officially estimated with this solution. The list admits of considerable extension:

Acetic Acid.—Operate upon about 1 gramme of glacial acid, about 20 grammes of diluted acid, or about 3 grammes of ordinary acetic acid.




Acetic Acid, U. S. P., should contain 36 per cent. of real acid ($\text{HC}_2\text{H}_3\text{O}_2$); Diluted Acetic Acid, U. S. P., 6 per cent.; Glacial Acetic Acid, U. S. P., 99 per cent.

Citric Acid.—Operate on about 1 gramme. The reaction is expressed by the following equation, etc.:




Citric Acid, U. S. P., should be pure (= 100 per cent. $\text{H}_3\text{C}_6\text{H}_5\text{O}_7\text{H}_2\text{O}$).

Hydrochloric Acid.—Operate on from 1 to 2 grammes of the concentrated acid or on about 4 grammes of the diluted acid.




Hydrochloric Acid, U. S. P., should contain 31.9 per cent. of real acid (HCl), and Diluted Hydrochloric Acid, U. S. P., 10 per cent.; Diluted Hydrobromic Acid, 10 per cent. (HBr).

Lactic Acid.—Operate on 1.5 to 2 grammes. The reaction is expressed by the following equation:




Lactic Acid, U. S. P., should represent 75 per cent. of absolute lactic acid ($\text{HC}_3\text{H}_5\text{O}_3$).

Nitric Acid.—Operate on from 1 to 2 grammes of concentrated or on from 4 to 5 grammes of dilute acid.




Nitric Acid, U. S. P., should contain 68 per cent., and Diluted Nitric Acid, U. S. P., 10 per cent. of absolute acid (HNO_3).

Sulphuric Acid.—Operate upon from .5 to 1 gramme of concentrated acid, or from 4 to 5 grammes of either diluted or aromatic sulphuric acid.




Sulphuric Acid, U. S. P., should contain 92.5 per cent.; Diluted, U. S. P., 10 per cent.; and Aromatic, U. S. P., 18.5 per cent. of absolute acid (H_2SO_4).

Phosphoric Acid.—Operate upon about 1 to 1.5 grammes of the concentrated acid or from 5 to 6 grammes of the diluted phosphoric acid.




Phosphoric Acid, U. S. P., should contain 85 per cent., and Diluted Phosphoric Acid, U. S. P., 10 per cent., of absolute acid (H_3PO_4).

Hypophosphorous Acid.—Operate upon 6 to 7 grammes of the official acid.



Diluted Hypophosphorus Acid, U. S. P., should contain 10 per cent. of absolute acid (H_3PO_2).

Tartaric Acid.—Operate upon about 1 gramme of the acid. The following equation, etc. represents the reaction:



Tartaric Acid, U. S. P., should contain 100 per cent. of $\text{H}_2\text{C}_4\text{H}_4\text{O}_6$.

Normal Soda Solution may be used instead of the potash. It is prepared similarly, making 1000 cc. contain 40 grammes (NaOH).

Notes.—1. Pure acetates, citrates more especially, tartrates, and some other organic salts, have an alkaline action on litmus, but not to an important extent. If the potash solution be added to acetic, citric, or tartaric acid containing litmus until the liquid is fairly blue, the operator will obtain trustworthy results. It is best to use

phenolphthalein, as recommended in the United States Pharmacopœia. It is produced by reaction of phenol and phthalic anhydride. Its tincture yields an intense red color with potash or soda; hence U. S. P. ordering it to be used as an indicator of the termination of volumetric reactions; it is especially useful with organic acids. The "Tincture of Phenolphthalein," B. P., is made by dissolving 1 gramme of phenolphthalein in 500 of proof spirit.

2. The operations for the quantitative analysis or measurement of acids are often collectively spoken of under the name of *acidimetry*.

QUESTIONS AND EXERCISES.

Calculate the percentage of real acid present in diluted sulphuric acid 30 grammes of which are neutralized by 84 cc. of the official volumetric solution of potash. *Ans.* 13.72.—Show how much real nitric acid is contained in a solution 36 grammes of which are saturated by 94 cc. of the standard solution of potash. *Ans.* 16.45 per cent.

ESTIMATION OF ACIDULOUS RADICALS PRECIPITATED BY SILVER NITRATE.


The purity of many salts and the strength of their solutions may be determined by this process; but only eighteen official substances (including diluted hydrocyanic acid, other cyanides, and some bromides and iodides) are quantitatively analyzed by standard solution of silver nitrate.

STANDARD SOLUTION OF SILVER NITRATE.

(Silver Nitrate, $\text{AgNO}_3 = 169.55$, say 170.)

Dissolve 17 grammes of crystals of pure silver nitrate in 1 litre of water. 1000 cc. of this solution contain $\frac{1}{10}$ of the molecular weight in grammes of silver nitrate. It is therefore a *decinormal solution*.


If pure dry crystals of silver nitrate are not at disposal, and pure dry crystals of sodium chloride are at hand, a solution may be made of approximate strength, and then be standardized by means of that salt. The method may be thus indicated:



Take rather less than .1 gramme of sodium chloride (NaCl), and dissolve in water. The salt (AgCl) precipitated in the reaction is an insoluble salt, and the end of its precipitation will serve as a good indication of the completion of the reaction. A better indicator, however, is a few drops of neutral potassium chromate (which previously was well purified by recrystallization). The silver nitrate


does not act upon the chromate until all the chloride is converted into silver chloride, after which a deep red precipitate of silver chromate is produced. This indication is extremely delicate, and in practice is noticed when the white color due to silver chloride changes to yellowish from formation of the first traces of silver chromate. Solutions should be cool and not very dilute. The titration being accomplished, suppose that .1 gramme of the sodium chloride has taken 17 cc. of the silver-nitrate solution of unknown strength; how many cc. of the solution are equivalent to 5.85 of the sodium chloride; that is, how many cc. of solution contain 17 grammes of silver nitrate? As .1 gramme of NaCl is to 17 cc., so are 5.85 NaCl to x cc. = 994 cc. 994 cc. of the solution of silver nitrate are equivalent, therefore, to 1000 cc. of official standard solution, and contain 17 grammes of the silver nitrate. They may be diluted to 1000 cc. if desired.

Hydrocyanic Acid.—3 to 4 grammes of *diluted* acid form a convenient quantity to operate upon. The HCN is first converted into KCN or NaCN with potash or soda. The following equations, etc. explain the reactions:



It is seen that 5.4 grammes of real hydrocyanic acid (HCy) are equivalent to 9.8 grammes of sodium cyanide, and represent 17 grammes of silver nitrate or 1000 cc. of standard solution of silver nitrate.


The sodium cyanide having been obtained, the titration is carried on until the salt is converted into the double salt (NaCy, AgCy), immediately after which a permanent turbidity occurs, due to precipitation of cyanide of silver, thus:



This turbidity forms a delicate and satisfactory proof of the completion of the volumetric reaction.

There is, however, a difficulty in the conversion of the acid into the cyanide (Siebold), to which it is necessary to pay particular attention. Tincture of litmus is added to the acid diluted largely with water, and the soda poured in. Owing to the strong alkaline reaction of the sodium cyanide formed, the mixture becomes blue when only a small proportion of the acid has been converted. If, then, the titration be conducted until the turbidity appears, only the sodium cyanide will be estimated, leaving free hydrocyanic acid still unacted upon. Indeed, sodium cyanide may be estimated in presence

of hydrocyanic acid in this way. Thus the following reaction (expressed approximately) might occur:




In this case only one-fifth of the cyanogen originally present would be estimated. The mixture would, however, become acid. If this acidity be prevented, all difficulty is overcome. The following details (Senier) will be found to answer well: To the diluted hydrocyanic acid add soda solution until a strong alkaline reaction is shown by the tincture of litmus. Then add the silver solution, drop by drop, from the burette, when in most cases the mixture will become acid. When it does so, add more soda solution, and repeat this process until the final reading, when the solution must be alkaline. In this way the addition of too much soda at the commencement, which would use up silver solution and make the reading a trifle too high, is avoided.

Diluted Hydrocyanic Acid, B. P. and U. S. P., should contain 2 per cent. of real acid (HCN)—*Aqua Laurocerasi*, B. P., 0.1 per cent.


The following is the quantitative test of purity ordered by the United States Pharmacopœia: "Mix in a flask (of the capacity of about 100 cc.) 0.27 gm. of hydrocyanic acid (obtained by distillation as above directed) with sufficient water and magnesia to make an opaque mixture of about 10 cc. Add to this 2 or 3 drops of potassium chromate, and then, from a burette, decinormal silver nitrate V. S., until a red tint is produced which does not again disappear by shaking. Each cc. of silver nitrate used indicates 1 per cent. of absolute hydrocyanic acid. After ascertaining the strength of the distillate, dilute it with distilled water so as to bring it to the strength of 2 per cent. of absolute acid. Lastly, test the finished product again, when 1.35 grms. of it should require, for complete precipitation, 10 cc. of decinormal silver nitrate."

Potassium Cyanide.—65 gm., in dilute solution, requires 45 cc. of standard decinormal silver nitrate = 90 per cent. of pure salt.

Ammonium Bromide.—Operate upon .075 to .1 gramme of the salt, using potassium chromate (or Bichromate, U. S. P.) as an indicator of the close of the reaction:



Ammonium Bromide, U. S. P., should be of 99 per cent. purity, but as the impurity is ammonium chloride, this too will be precipitated by the silver nitrate, and must be calculated in finding the percentage of bromide.




The amount of the salt equivalent to 1000 cc. of standard solution is first calculated by simple proportion: Let x represent this; then $9.78 - x = y$, the excess of standard solution used up by the ammonium chloride, reckoned in terms of bromide (NH_4Br); and since 5.35 grammes of $\text{NH}_4\text{Cl} = 9.78$ grammes of NH_4Br , the excess which ammonium chloride can consume is represented by $9.78 - 5.35 = 4.43$; therefore, as $4.43 : 5.35 :: y : z =$ the amount of ammonium chloride present in x grammes of the sample taken; lastly, the percentage is calculated by simple proportion:

As $x : 100 :: z : p =$ percentage. For example: .075 gramme of the salt required, 7.8 cc. of standard solution,

1. $7.8 : 1000 :: .075 : x$;
 $x = 9.615$.
2. $9.78 - 9.615 = y$;
 $y = .165$.
3. $4.43 : 5.35 :: .165 : z$;
 $z = .19926$.
4. $9.615 : 100 :: .19926 : p$;
 $p = 2.072$ per cent. of NH_4Cl .


Potassium Bromide.—Operate upon rather less than .1 gramme, and conduct the titration in the same manner as with sodium chloride, using potassium chromate as an indicator of the close of the reaction.



The United States Pharmacopœia requires potassium bromide to contain 97 per cent. of the pure salt, and *sodium bromide* 97.29 per cent.


Remembering that 170 parts of silver nitrate ($\text{AgNO}_3 = 170$) decompose 119 of potassium bromide ($\text{KBr} = 119$), while, on the one hand, they decompose as little as 74.5 of potassium chloride ($\text{KCl} = 74.5$), and, on the other hand, as much as 166 parts of potassium iodide ($\text{KI} = 166$), it will be seen that the quantitative operation of the chloride as an impurity may neutralize the quantitative operation of the iodide. Hence the necessity to test the bromide qualitatively as well as quantitatively, and, as regards either impurity singly, of fixing maximum as well as minimum limits of the action of the volumetric solution of silver nitrate on potassium bromide.

Lithium Bromide.—Operate on about half a gramme.




Lithium Bromide, U. S. P., should contain 98 per cent. of the pure salt.

Calcium Bromide.—Operate on about $\frac{1}{2}$ a gramme.




The United States Pharmacopœia requires calcium bromide to contain 99.7 per cent. of the pure salt.

Strontium Bromide would be estimated similarly, and the official substance should contain 98 per cent. of the pure salt.



Potassium Iodide.—Operate on about $\frac{1}{2}$ a gramme of the substance.




The United States Pharmacopœia requires potassium iodide to contain 99.5 per cent. of the pure salt.

The sodium iodide is estimated in the same way by making the alterations shown previously, and should contain 98 per cent. of the pure salt (NaI).


The sodium chloride is estimated, as shown on p. 639, and the official salt should contain 99.9 per cent. of pure NaCl.

Potassium Iodide may be volumetrically estimated by a decinormal solution of mercuric chloride, the termination of the operation being indicated by the commencement of the formation of a red precipitate:



The author of this process, M. Personne, stated (in 1875) that neither chlorides, bromides, nor carbonates interfere. Carles dissolves the iodide in spirit of wine of $17\frac{1}{2}$ per cent.; as much excess of water may decompose the double iodides.




Ferrous Iodide—Messrs. Naylor and Hooper (in 1881) demonstrated that Personne's solution is applicable to ferrous iodide, even in the state of syrup:




The use of mercuric chloride for estimating the strength of syrup of iodide of iron was first suggested by E. Smith in 1859. The process was improved by T. and H. Smith in 1860.

Zinc Chloride, Bromide, and Iodide.—Operate on about $\frac{1}{2}$ a gramme of the substance, as described in the case of Calcium, etc. The official salts should contain, respectively, 99.84 per cent., 99.95

per cent., and 98.62 per cent. of the pure salts. The equations will explain any difficulty that may occur:




Syrup of Hydriodic Acid.—Operate upon 10 to 15 grammes. The reaction which occurs is as follows:



The close of the reaction is shown by the cessation of the formation of silver iodide, the nitric acid liberated rendering potassium chromate inadmissible as an indicator.

Syrupus Acidi Hydriodici, U. S. P., should contain 1 per cent. of anhydrous hydriodic acid (HI).

Syrup of Ferrous Iodide.—Operate upon 1 to 2 grammes of the syrup until no further precipitate is formed:



Syrupus Ferri Iodidi, U. S. P., should contain 10 per cent. of iodide of iron (FeI_2).

Spirit of Wine (*Spiritus Rectificatus*, B. P.) may contain traces of amyl alcohol and aldehyde; these may be detected by silver nitrate, which is reduced by them to the metallic state. Any quantity beyond a mere trace of such bodies renders spirit of wine too impure for use in medicine. "4 fluidounces with 30 grain-measures (about 2 cc.) of the volumetric solution of silver nitrate, exposed for twenty-four hours to bright light, and then decanted from the black powder which has formed, undergoes no further change when again exposed to light with more of the test."—*B. P.* "If 20 cc. of alcohol be shaken in a clean glass-stoppered vial with 1 cc. of silver nitrate, the mixture should not become more than faintly opalescent, or acquire more than a faint brownish tint when standing during six hours in diffused daylight (limit of organic impurities, amyl alcohol, etc.)."—*U. S. P.*

QUESTIONS AND EXERCISES.

Explain the volumetric method of estimating the strength of aqueous solutions of hydrocyanic acid.—Work a sum showing how much silver nitrate will indicate, by the official volumetric process, the presence of 1 part of real hydrocyanic acid. *Ans.* 3.148 parts.

ESTIMATION OF SUBSTANCES READILY OXIDIZED.

Any deoxidizer—that is, any substance which quickly absorbs a definite amount of oxygen or is susceptible of any equivalent action—may be quantitatively tested by ascertaining how much of an oxidizing agent of known power must be added to a given quantity before complete oxidation is effected. The oxidizing agents employed for this purpose in the United States Pharmacopœia are iodine, the red potassium chromate, and potassium permanganate. Iodine acts indirectly by taking hydrogen from water and liberating oxygen; the red potassium chromate, directly by the facility with which it yields three-sevenths of its oxygen—as indicated by the equations and statements given on p. 651; potassium permanganate, by affording five-eighths of its oxygen in presence of acid,



STANDARD SOLUTION OF IODINE.

(Iodine, I = 126.53.)

The Pharmacopœia allows 127 to be taken as the atomic weight of iodine for rough purposes, but recommends 126.53 as the datum for accurate work. This fact should be borne in mind in the following simplified operations and calculations:

If pure iodine be not at hand, it may be prepared by mixing the commercial article with about a fourth of its weight of potassium iodide and subliming. Sublimation may be effected by gently warming the mixture in a beaker the mouth of which is closed by a funnel; the iodine vapor condenses on the funnel, while fixed impurities are left behind, and any chlorine which the iodine may contain is absorbed by the potassium iodide, an equivalent quantity of iodine being liberated. Small quantities may be similarly treated between two watch-glasses, placed edge to edge. Any trace of moisture in the resublimed iodine is removed by exposure for a few hours under a glass shade near a vessel containing oil of vitriol.


Place 12.7 grammes of pure iodine and about 18 grammes of pure potassium iodide (an aqueous solution of which is the best solvent of iodine; the salt plays no other part in these operations) in a litre flask; add a *little* water and agitate until the iodine is dissolved; dilute to 1 litre.

The following substances are officially estimated by this standard decinormal volumetric iodine solution:

Sulphurous Acid.—Operate on about .5 of a gramme of the acid, and dilute with water as usual. If the sulphurous acid be diluted to a less degree than .04 or .05 per cent., there will be some risk of


the sulphuric acid formed being again reduced to sulphurous acid, with liberation of iodine. In delicate experiments the distilled water used for dilution should previously be freed from air by boiling, to prevent the small amount of oxidizing action which dissolved air would exert. The solution of iodine is then added until a slight permanent brown tint is produced, showing the presence of free iodine. A better indicator of the termination of the reaction is starch mucilage, which gives a blue color with the slightest trace of free iodine.

The following equations, etc. show the reaction that takes place:



The official (U. S. P.) sulphurous acid should contain 64 per cent. of sulphurous anhydride (SO_2).

Arsenic.—About .1 of a gramme of *solid arsenic*, accurately weighed, should be dissolved in the usual quantity of water, heated to boiling, by aid of about .5 of a gramme of sodium bicarbonate. The arsenous acid is only partly, if at all, converted into sodium arsenite or arsenate; but the iodine reaction occurs more readily in an alkaline solution. When the liquid is quite cold, mucilage of starch is added, and the iodine solution allowed to flow in until, after well stirring, a permanent blue color is produced. The official *Liquor Potassi Arsenitis*, U. S. P., already containing some potassium carbonate, requires somewhat less sodium bicarbonate. 10 grammes is a convenient quantity to operate upon. To this should be added the usual quantity of water and about .3 of a gramme of sodium bicarbonate. After boiling and cooling the titration is carried on as before. About 10 grammes of the official *Liquor Acidi Arsenosi*, U. S. P., is also a convenient quantity to operate upon. This quantity requires about .6 of a gramme of sodium bicarbonate. The usual quantity of water is added and the titration performed as before. The following equation exhibits the reaction:




The United States Pharmacopœia requires both the above solutions of arsenic to contain 1 per cent. of As_2O_3 , and the arsenous acid 98.8 per cent. of As_2O_3 .

In the foregoing operation, if ebullition be continued longer than is necessary for the solution of the arsenic, more mon carbonate of sodium may be formed than will be reconverted into bicarbonate by

the liberated carbonic acid; loss of iodine will then ensue. E. J. Woolley has shown that borax may be usefully employed in the place of the sodium bicarbonate.


Antimony also passes from lower to higher active quantivalence under the influence of nascent oxygen, iodine, or an equivalent acidulous radical. The following equation illustrates the reaction with tartar emetic and iodine. The student should make several determinations on, say, 20 cc. of a solution of 2 grammes of pure crystals of tartar emetic in 200 cc. of water. To the 20 cc. add about an equal amount of strong solution of sodium bicarbonate, a couple of cc. of starch mucilage, and then the iodine solution, until, after stirring, the blue color is fairly persistent. The whole operation should be quickly conducted, or a precipitate of antimonious hydrate will be formed, and it is only when in solution that the antimony is properly attacked. This process is by Mohr. It has been tested by Fresenius and in the Research Laboratory of the Pharmaceutical Society, and is trustworthy.



The official Tartar Emetic, U. S. P. should be absolutely pure (100 per cent.).


Hyposulphite of Sodium.—About .4 of a gramme is a convenient quantity to employ. It is dissolved in water, starch mucilage added, and the iodine solution slowly run in, the whole being frequently stirred, until a permanent blue color is produced.

In the previous reactions iodine has acted as an indirect oxidizing agent by uniting with the hydrogen and thus liberating the oxygen of water. In the present case it unites with an analogue of hydrogen—namely, sodium—a new salt (tetrathionate of sodium) being also produced, thus:




The United States Pharmacopœia requires "Sodium Hyposulphite," now generally termed sodium thiosulphate, to contain 98.1 per cent of the crystalline salt.

Sodium Sulphite.—About .7 grammes is a convenient quantity to take, using starch mucilage as an indicator, as before. The reaction as below occurs:



Sodium sulphite should contain 96 per cent. of the pure crystalline salt ($\text{Na}_2\text{SO}_3, 7\text{H}_2\text{O}$).

Sodium Bisulphite.—Operate upon .2 to .3 gramme, as before :



The United States Pharmacopœia requires bisulphite of sodium to contain 90 per cent. of the pure salt (NaHSO_3).

QUESTIONS AND EXERCISES.


Give equations illustrative of the reactions on which the use of a standard volumetric solution of iodine is based.—From what point of view is iodine an oxidizing agent?—What reagent indicates the termination of the reaction between deoxidizing substances and moist iodine?—How much sulphurous acid gas will cause the absorption of 2.54 parts of iodine in the volumetric reaction? *Ans.* 0.64.—What quantity of iodine will be required, under appropriate conditions, to oxidize 5 parts of arsenic? *Ans.* 12.828.—Find by calculation the amount of sodium hypsulphite which will react with 13 parts of iodine in volumetric analysis. *Ans.* 25.386.

VOLUMETRIC SOLUTION OF RED POTASSIUM CHROMATE.

(Red Potassium Chromate, $\text{K}_2\text{Cr}_2\text{O}_7 = 293.78$.)

One molecule of red potassium chromate in presence of an acid, under favorable circumstances, yields four atoms of oxygen to the hydrogen of the acid, leaving three available either for direct oxidation or for combination with the six atoms of hydrogen of more acid, an equivalent proportion of acidulous radical being liberated for any required purpose. One-sixth, therefore, of the molecular weight of red chromate, taken in grammes ($293.78 \div 6 = 48.963$), and dissolved in 1000 cc., would form a *normal* solution (for that is the amount which is equivalent to one atomic weight, in grammes, 1.0, of hydrogen; see p. 626), while one-sixtieth (4.8963) in 1 litre would form the official *decinormal* solution. For all practical purposes 294 may be taken as the molecular weight = 4.9 grammes per litre.


When used as a volumetric agent the red chromate always yields the whole of its oxygen to the hydrogen of the accompanying acid, a corresponding quantity of acidulous radical being set free—four-sevenths of this radical immediately combining with the potassium and chromium of the red chromate, three-sevenths becoming available. Ferrous salts may thus be converted into ferric with sufficient rapidity and exactitude to admit of the estimation of an unknown quantity of iron by a known quantity of the red chromate. As one atom of any liberated bivalent acidulous radical will convert two molecules of ferrous salt into one of ferric, one molecule of red chromate causes six of ferrous to become three of ferric, as shown in the following equation :



The volumetric solution is made by dissolving 14.75 grammes ($\frac{1}{20}$ of a molecular weight in grammes) of red potassium chromate in water, and diluting to 1 litre. It is used in determining the strength of the ferrous preparations. It is known that the whole of the ferrous has been converted to ferric salt when a small drop of the liquid placed in contact with a drop of a fresh and very dilute solution of potassium ferricyanide on a white plate ceases to strike a blue color.

If the red chromate employed in making this standard solution is not known to be pure and dry, the strength of the solution may be checked by dissolving a small accurately weighed piece of pianoforte wire (0.4 or 0.5) in diluted sulphuric acid in a small flask, warming, and then running in the solution of red chromate until conversion is effected.

The reactions which take place may be thus expressed :



It is evident that 16.8 grammes of iron are equivalent in the reactions to 14.75 of red chromate or 1000 cc. of standard solution of the chromate. Now suppose that 0.5 of a gramme of pianoforte wire has been employed, and the quantity of solution of red chromate of unknown strength used has been 28 cc. How many cc. of this solution contains 1.475 of red chromate?—that is, how many cc. must be required to oxidize ferrous salt containing 16.8 of iron? As .5 of iron is to 28 cc. sol., so are 16.8 of iron to x cc. sol. = 941 cc. Of the supposed solution, then, 941 cc. would contain 14.75 grammes of red chromate, and would be equivalent to 1000 cc. of standard solution. It might be employed without being diluted, or, better, be diluted to official standard strength.

For standardizing the solution of red chromate, instead of iron wire the light-green crystals of the double *ferrous ammonium sulphate* ($\text{FeSO}_4 \cdot (\text{NH}_4)_2\text{SO}_4 \cdot 6\text{H}_2\text{O} = 392$) may be employed, for it is a very stable salt.

Special care should be taken in all these estimations of substances readily oxidized to avoid atmospheric oxidation. Flasks may usually be loosely corked, or corked closely with a gas exit-tube passing just beneath a little mercury, and in all cases the estimation should be performed quickly. When standardizing with iron wire any slight oxidation may be remedied by a fragment of zinc, the last portions of which must be removed or dissolved before the titration is commenced.



The "Decinormal Potassium Dichromate Volumetric Solution," U. S. P., is made from 4.89 (say 4.9) grammes of $\text{K}_2\text{Cr}_2\text{O}_7$ dissolved in 1000 cc. of water.

When used with phenolphthalein as indicator to neutralize alkalies, the volumetric solution of potassium dichromate is decinormal when it contains 14.689 grms. in 1 litre. It is then the exact equivalent of any decinormal acid, corresponding to the amounts of alkalies quoted, for instance, under Decinormal Oxalic Acid.

When used as an oxidizing agent to convert ferrous into ferric salts or to liberate iodine from potassium iodide, the solution just mentioned (containing 14.689 grms. in 1 litre) has the effect of a ($\frac{3N}{10}$) volumetric solution, and a solution of one-third of this strength, containing 4.896 grms. in 1 litre, has the value of a decinormal solution, and is the equivalent of equal volumes of decinormal potassium permanganate, or, in the case of iodide liberated from potassium iodide, it is the equivalent of equal volumes of decinormal sodium hyposulphite.—U. S. P.

The ferrous salt in the following substances is estimated officially by this solution:


Saccharated Ferrous Carbonate.—Operate upon 1 to 2 grammes. Dissolve in excess of diluted sulphuric or hydrochloric acid. Sulphuric acid is preferable in most cases, because ferrous sulphate absorbs oxygen much less readily than ferrous chloride. The reaction that occurs is shown in the following equation, the ferrous carbonate being converted into ferric sulphate:



The official (U. S. P.) strength in iron is 15 per cent. Trade samples yield from 20 to 30, and sometimes 35 per cent., according to the care with which oxidation has been prevented. The theoretical percentage obtainable from the ingredients is 45.5, the quantity that would be present if the compounds were anhydrous and unoxidized—conditions never obtained in practice. Howie has suggested that as hydrochloric acid is known to so rapidly convert ordinary sugar into inverted sugar as to render it easily attacked by chromic acid, while phosphoric acid very slowly affects sugar, the latter acid instead of the former should be employed in dissolving the saccharated ferrous carbonate for volumetric analysis. Another mode of eliminating the action of sugar is to char with sulphuric acid before analyzing.


Ferrous Sulphate.—Take about 1 to 2 grammes of the substance and titrate in the usual way. The equations are given on p. 614. Both the official ferrous sulphates are required to be absolutely pure (100 per cent.).

Ferrous Arsenate.—Operate upon 1 to 2 grammes; proceed as with the carbonate. The reaction that occurs is shown in the following equation, the ferrous arsenate being converted into ferric arsenate:



Arsenate of Iron, B. P., is supposed to contain about 10 per cent. of ferrous arsenate. The compound is more nearly a ferric than a ferrous arsenate.


Ferrous Phosphate.—Operate upon 1 to 2 grammes. Proceed as with the carbonate. The following equation indicates the reaction, the ferrous phosphate being converted into ferric phosphate:




The official (B. P.) requirement is about one-third of its weight of anhydrous ferrous phosphate, or 47 per cent of $\text{Fe}_3(\text{PO}_4)_2 \cdot 8\text{H}_2\text{O}$.

Phosphate of Iron, U. S. P., is *Ferric Phosphate*, and therefore cannot be estimated by this solution.

Magnetic Iron Oxide.—Operate upon 1 or 2 grammes, and proceed as with arsenate or phosphate. The reaction may thus be shown:



Or,



Absolutely pure magnetic iron oxide contains 31 per cent. of ferrous oxide. Oxidation occurs, however, during manufacture, as in the case of the ferrous salts just described.

Note.—The use of this volumetric solution in quantitative analysis admits of great extension. The student should at least employ it in the case of a few iron ores.

QUESTIONS AND EXERCISES.


Write equations explanatory of the oxidizing power of red potassium chromate.—100 cub. cent. of an aqueous solution of red potassium chro-

mate contain $\frac{1}{200}$ of the molecular weight of the salt in grammes; with what weight of metallic iron, dissolved in hydrochloric acid, will this volume react? *Ans.* 1.68 gramme.—If 8.34 grammes of impure crystallized ferrous sulphate, dissolved in acidulated water, require 93 cc. of the standard solution of chromate for complete conversion into ferric salt, what percentage of ferrous sulphate is present? *Ans.* 93.—Work a sum showing how much red potassium chromate is required for the conversion of 10 parts of ferrous sulphate into ferric salt. *Ans.* 1.768.—Show what quantity of pure ferrous carbonate is indicated by 1.475 part of red chromate as applied in volumetric analysis. *Ans.* 3.48.—Prove what amount of official saccharated ferrous carbonate is equivalent to .7375 part of red chromate in the volumetric reaction. *Ans.* 4.7.

STANDARD SOLUTION OF POTASSIUM PERMANGANATE.*

(Potassium Permanganate, $K_2Mn_2O_8 = 315.34$.)

Dissolve about 3.5 grammes of potassium permanganate crystals in distilled water. Boil the solution in a large basin for a quarter of an hour. Allow to stand for a couple of days, and then pour off the clear solution and make up to 1000 cc. This solution must be now standardized by the following reaction:



Weigh out carefully about 1 gramme of polished pianoforte wire; place this in a 250 cc. flask with about 70 cc. of dilute sulphuric acid; then add about a gramme of sodium carbonate. Close the flask with a Bunsen valve (a Bunsen valve is easily made by making a longitudinal slit about $\frac{1}{3}$ of an inch long in a short piece of india-rubber tubing, closing one end with a piece of glass rod, and placing the other on the tube passing through the cork; gases can escape by the slit thus formed, but cannot re-enter), and heat on the sand-bath until all the wire is dissolved. Allow the flask to cool, and only when perfectly cold remove the cork and fill up to the mark with water. Titrate 25 cc. of this solution with the permanganate solution. The strength can be easily calculated from the equation

* Potassium permanganate affords five-eighths of its oxygen in presence of acid. Thus:




the five liberated atomic weights of oxygen (O_5) at once performing chemical work equivalent to the oxidation of ten atomic weights of hydrogen (H_{10}). Hence, one-tenth (31.534) of the atomic weight of potassium permanganate ($K_2Mn_2O_8 = 315.34$), taken in grammes (31.534), dissolved in water, and made up to 1 litre (1000 cc.), would form a *normal* solution, while a solution of one-tenth of this strength (3.1534 grammes in 1000 cc.), the "Decinormal Potassium Permanganate Volumetric Solution," U. S. P., K_2MnO_4 and $K_2Mn_2O_8$, are the probable formulæ of potassium manganate and permanganate, but the latter is sometimes written $KMnO_4$.

given above. Care must be taken to deduct 0.4 per cent. from the weight of iron wire taken, for impurities. When the strength has been found—that is, how many cc. of the permanganate solution contains 3.15 grms. of potassium permanganate—make this number of cc. up to 1000 by adding water.

The United States calls this “Decinormal Potassium Permanganate.”


The following substances are officially estimated with this solution:

Hydrogen Dioxide Solution.—Take about 2 cc. and place in a beaker, dilute and add permanganate until a faint pink color is produced. The equation will explain the reaction:




The United States Pharmacopœia requires *Aqua Hydrogenii Dioxidii* to contain 3 per cent. of H_2O_2 .

Barium Dioxide.—Operate on about .5 of a gramme; the reaction is similar to hydrogen dioxide:





Barium dioxide should contain 80 per cent. of pure BaO_2 .

Ferrous Carbonate and Sulphate.—Operate on about 1 to 2 grammes. The two equations will show the action with the carbonate, and the second one for the sulphate. The action of the sulphuric acid on the ferrous carbonate is to produce ferrous sulphate, which then gets oxidized to ferric sulphate. 11.59 grammes of the carbonate are equivalent to 3.15 grms. of potassium permanganate.






Then



The United States Pharmacopœia requires the saccharated ferrous carbonate to contain 15 per cent. of FeCO_3 ; the crystalline and the granulated ferrous sulphate should both be pure—100 per cent.


Reduced Iron.—Use about 1 to 1.5 grammes, and place the “reduced iron into a glass-stoppered bottle; add 50 cc. of mercuric chloride, and heat the bottle, well stoppered, during one hour on a water-bath, frequently agitating. Then allow it to cool, dilute the contents with water to the volume of 100 cc., and filter. To 10 cc. of the filtrate, contained in a glass-stoppered bottle (having a capacity of about 100 cc.), add 10 cc. of diluted sulphuric acid, and subsequently decinormal potassium permanganate, until a permanent red color is produced. The number of cc. of the volumetric solution required, when multiplied by 10, will indicate the percentage of metallic iron” (*U. S. P.*), and should contain 80 per cent. of iron as metal.



“To confirm the assay, decolorize the liquid by a few drops of alcohol, then add 1 grm. of potassium iodide, and digest for half an hour at a temperature of 40° C. (104° F.). The cooled solution, mixed with a few drops of starch, should require not less than 8 cc. of decinormal sodium hyposulphite to discharge the blue or greenish color (each cc. of the volumetric solution indicating 10 per cent. of metallic iron).”—*U. S. P.*


Hypophosphorous Acid is officially estimated with standard potassium-permanganate solution. Take about .7 gramme of the diluted acid and add water; proceed to titrate in the usual way until a pink color is just produced after shaking.


The following equation will explain the reaction :



The United States Pharmacopœia required *Acidum Hypophosphorum Dilutum* to contain 10 per cent. of pure acid.

Calcium, Ferric, Potassium, and Sodium Hypophosphites.—Operate on from .1 to .2 grammes and proceed as above. The following equations correspond with the official data :





The official requirements are—calcium salt to contain 99.68 per cent. of the pure salt; ferric salt, 98.1 per cent.; potassium salt, 98.7 per cent.; sodium salt, 97.9 per cent.

Centinormal Potassium Permanganate is a solution, 1000 cc. of which contain .315 gramme of permanganate. It is useful for estimating small quantities of iron, etc.

ESTIMATION OF SUBSTANCES READILY DEOXIDIZED.

Any substance which quickly yields a definite amount of oxygen may be quantitatively tested by ascertaining how much of a deoxidizing agent of known power must be added to a given quantity before complete deoxidation is effected. The chief compounds which may be used for this absorption of oxygen (deoxidizers or reducing agents, as they are commonly termed) are sodium hyposulphite, sulphurous acid, ferrous sulphate,* oxalic acid, arsenous acid. The first-named is officially employed; it is only used in the estimation of free iodine, and, indirectly, of chlorine and chlorinated compounds. Iodine and chlorine are regarded as oxidizing agents, because their great affinity for hydrogen enables them to become powerful indirect oxidizers in presence of water.


STANDARD SOLUTION OF HYPOSULPHITE OF SODIUM.

(Crystallized Hyposulphite of Sodium, $\text{Na}_2\text{S}_2\text{O}_3 \cdot 5\text{H}_2\text{O} = 247.64$, or, roughly, 248.)

Dissolve about 27 grammes of sodium hyposulphide in a litre or less of water. Fill a burette with this solution, and allow it to flow into a beaker containing, say, 15 cc. of the volumetric solution of


* Five grains of permanganate of potassium dissolved in water require for decoloration a solution of 44 grains of granulated sulphate of iron acidulated with 2 fluidrachms of diluted sulphuric acid.

iodine until the brown color of the iodine is just discharged—or, starch being added, until the blue iodide of starch is decolorized. (The latter affords the more delicate indication.) When iodine and sodium hyposulphite react, two atoms of iodine remove two of sodium from two molecules of the hyposulphite, sodium tetrathionate being formed, thus:



Now suppose the number of cc. required to fully attack the 15 cc. of standard iodine were 14 cc., how many cc. of this hyposulphite solution would be equivalent to 1000 cc. of standard iodine solution? In other words, how many cc. would contain 24.8 grammes of hyposulphite? As 15 cc. iodine solution are to 14 cc. hyposulph. solution, so are 1000 iodine solution to x hyposulph. solution = 933 cc. Therefore 933 cc. of this solution of hyposulphite would contain 24.8 grammes of the salt, and be equivalent to 1000 cc. of the official standard solution. The 933 cc. would be diluted to 1000 cc. or be used without dilution. In either case its strength would, as usual, be recorded on the label. The following substances are estimated officially by means of this solution:

Chlorine Solution.—About 10 grammes are operated upon. Excess of potassium iodide is added—that is, to 10 grammes of solution of chlorine about $\frac{1}{2}$ a gramme of iodide. An amount of iodine is set free by the chlorine exactly in proportion to their atomic weights. The titration is then conducted as already described. The following equations show the reactions:




It is evident, then, that 1000 cc. of standard solution of sodium hyposulphite, or a corresponding quantity of a solution of different strength, is equivalent to 3.37 grammes of chlorine gas. Chlorine solution of the United States Pharmacopœia contains 0.4 per cent. of chlorine.

Iodine.—Solid iodine is dissolved in solution of potassium iodide and titrated as already described. About .2 of a gramme is a convenient quantity to employ. 1000 cc. of standard hyposulphite solution is equivalent, as seen in the equation, to 12.65 of iodine. The United States Pharmacopœia requires "iodine" to contain 98.85 per cent. of real iodine. It is assumed in this operation that the iodine has been shown by qualitative analysis to be free from chlorine and

bromine. These elements resemble iodine in reacting upon sodium hyposulphite, hence would reckon as iodine in a volumetric assay.

Chlorinated Lime.—Operate on from .1 to .2 of a gramme. Dissolve in the usual quantity of water, and add excess of potassium iodide and diluted hydrochloric acid. .1 to .2 of a gramme of chlorinated lime would require .4 to .8 of a gramme of potassium iodide. The following equations show the reactions:



The chlorine thus set free liberates an equivalent amount of iodine, and this is titrated as before. (See the equations for solution of chlorine, p. 656.) This chlorine, liberated from chlorinated lime by acids, is its available chlorine for indirect oxidizing action. It should correspond (U. S. P.) to 35 per cent. of chlorine.

Solution of Chlorinated Soda.—About 2 grammes are mixed with the usual quantity of water and about 1 gramme of potassium iodide, and excess of acid added. The available chlorine is estimated as in the case of chlorinated lime. The reaction by which the chlorine is evolved is familiar:



The action of the liberated chlorine on the potassium iodide and the iodine on the hyposulphite solution has been described under "Solution of Chlorine." The official (U. S. P.) requirement is about 2.6 per cent. of available chlorine.

Compound Solution of Iodine.—Process as before, using 1 or 2 grammes; the reaction has already been given. The requirements of the United States Pharmacopœia are 5 per cent. of free iodine.

Tincture of Iodine.—Use about 1 gramme. It will contain 7 per cent. of free iodine in 100 cc. when of official strength.

QUESTIONS AND EXERCISES.

For what purpose is the official volumetric solution of sodium hyposulphite used?—On what reaction is based the quantitative employment of sodium hyposulphite?—How much sodium hyposulphite is required to show the presence of 10 parts of iodine?—*Ans.* 19,527.—Calculate the amount of chlorine 4.96 parts of sodium hyposulphite are equivalent to in volumetric analysis. *Ans.* 71.—Describe the operations included in the estimation of the strength of bleaching-powders.—By what reagent is the complete absorption of free iodine by sodium hyposulphite indicated?

Other volumetric solutions are official.

DECINORMAL SODIUM CHLORIDE VOLUMETRIC SOLUTION.

(Sodium Chloride, $\text{NaCl} = 58.37$.)

Dissolve 5.84 grammes of pure sodium chloride* in 1000 cc. of

* Pure sodium chloride may be prepared by passing dry hydrochloric

water. This solution may be used as it is or standardized with decinormal silver nitrate.

$$\underbrace{\text{NaCl}}_{10)58.37} + \underbrace{\text{AgNO}_3}_{10)169.55} = \text{AgCl} + \text{NaNO}_3$$

5.837 16.955 = grammes in 1000 cc. of stand. solution.

The salts to be estimated with this standard solution of sodium chloride are the three official nitrates. The above equation will explain the reactions. Operate on about .5 to 1 gramme. Argentie nitrate should contain 99.97 per cent. of the pure salt, the fused nitrate 95 per cent., and the mitigated caustic 95 per cent., to fulfil the requirements of the United States Pharmacopœia.

“Decinormal Bromine Volumetric Solution” (Kappeschaar’s Solution) contains 7.976 grammes of bromine in 1000 cc. It is only used to test *Acidum Carbolicum*, U. S. P., which should contain 96 per cent. of pure phenol.

“Decinormal Mercuric Potassium Iodide Volumetric Solution” (Mayer’s Solution) contains 39.2 grms. of a mixture of mercuric iodide and potassium iodide, in the proportion of one molecular weight to two.

“Decinormal Potassium Sulphocyanate Volumetric Solution” (Volhard’s Solution) contains 9.699 grms. of KSCN in 1000 cc.

QUESTIONS, WITH ANSWERS FOR VERIFICATION.

Work sums showing how much potassium bicarbonate is contained in an eight-ounce bottle of medicine, 7 fluidrachms of which are saturated by $2\frac{1}{2}$ grains of crystallized oxalic acid. *Ans.* 36.3 grains.—A sample of soda-ash is said to contain 78 per cent. of pure anhydrous sodium carbonate: if the statement be true, how much of the official volumetric solution of oxalic acid will saturate 5 grammes of the specimen? *Ans.* 73.6.—2.69 grammes of common brown sulphuric acid are saturated by 43.5 cubic centimètres of the official volumetric solution of soda; how much acid of 96.8 per cent. is present? *Ans.* The 2.69 contain 2.2.—4 grammes of $1\frac{1}{2}$ litres of concentrated hydrocyanic acid are equivalent to 89 cubic centimètres of volumetric solution of silver nitrate of official strength; to what volume must the bulk of the acid be diluted for the production of acid of pharmacopœial strength? *Ans.* 9 litres.—3.18 grammes of a powder containing arsenic require for complete reaction 84 cubic centimètres of a volumetric solution of iodine, which is 1.43 per cent. weaker than the standard solution of the United States Pharmacopœia; what percentage of pure arsenic is contained in the powder? *Ans.* 12.86.—How much pure metal is present in a sample of iron 1.68 grammes of which, dissolved in diluted sulphuric acid, are exactly attacked by 95.7 cubic centimètres of an official volumetric solution of red potassium chromate which is .6 per cent. too strong?

acid gas into a saturated solution of common salt; separate the crystals which fall, and heat them in a porcelain basin until they are perfectly neutral; that is, until all the HCl gas has been driven off.

GRAVIMETRIC QUANTITATIVE ANALYSIS.

(For preliminary remarks on the general principles of gravimetric analysis and the relation of gravimetric and volumetric analyses to each other see pp. 591 and 592.)

ESTIMATION OF METALS.

POTASSIUM.

Outline of the Process.—This element is usually estimated in the form of double chloride of potassium and platinum. Qualitative analysis having proved the presence of potassium and other radicals in a substance, a small quantity of the material is accurately weighed, dissolved, and the other elements removed by appropriate reagents: the precipitates are well washed, in order that no trace of the potassium salt shall be lost, the resulting liquid concentrated over a water-bath (to avoid loss that would occur mechanically during ebullition), hydrochloric acid added if necessary, solution of platinum perchloride poured in, and evaporation continued to dryness: excess of the perchloride is then dissolved out by adding to the dried residue spirit of wine containing half its bulk of ether (a liquid in which the double chloride is insoluble); the mixture is carefully poured on to a tared and dried filter, washed with the spirit till every trace of free platinum perchloride is removed, and the whole dried and weighed; from the resulting amount the proportion of potassium or equivalent quantity of a salt of potassium is ascertained by calculation.

Note.—From this short description it will be seen, first, that the chemistry of quantitative analysis is the same as that of qualitative; and secondly, that the principle of gravimetric is the same as that of volumetric quantitative analysis: the combining proportions of substances being known, unknown quantities of elements may be ascertained by calculation from known quantities of their compounds.

Apparatus.—In addition to a delicate balance and weights and the common utensils, a few special instruments are used in quantitative manipulation; some of these may be prepared before proceeding with the estimation of potassium.

Filtering-paper may be of the kind known as "Swedish," the texture of which is of the requisite degree of closeness and its ash small in amount. A large number of circular pieces of one size, six to eight centimètres in diameter, should be cut ready for use. In delicate experiments, where a precipitate on a filter has to be ignited and the paper subsequently burnt, the weight of the ash of the filter must be deducted from the weight of the residue. The ash is estimated after burning ten or twenty of the cut filters. These are folded into a small

compass, a portion of a piece of platinum wire twisted a few times round the packet, so as to form a cage, the whole held by the free end of the wire over a weighed porcelain crucible placed in the centre of a sheet of glazed paper, the bundle ignited by a spirit-lamp or smokeless gas flame, the flame allowed to impinge against the charred mass till it falls into the crucible below, any stray fragments on the sheet carefully shaken into the crucible, the latter placed over a flame till carbon has all burnt off, and nothing but ash remains; the whole cooled, weighed, and the weight of the crucible deducted; the weight of the residue, divided by the number of pieces used, gives the average amount of ash in each filter.

Fig. 75.



A Pair of Weighing-tubes.

Fig. 76.



Clamped Watch-glasses for Weighing.

A pair of weighing-tubes (Fig. 75), for holding dried filters during operations at the balance, may be made from two test-tubes, one fitting closely within the other. About five centimètres of the closed end of the outer and seven of the inner are cut off by leading a crack round the tube with a pencil of incandescent charcoal, and the sharp edges fused in the blow-pipe flame. A filter, after drying, is quickly folded and placed in the narrower tube, the mouth of which is then closed by the wider tube. This prevents reabsorption of moisture from the air. A pair of watch-glasses, having accurately ground edges and clamped as shown in Fig. 76, also forms a convenient arrangement for weighing filters, etc.

The washing-bottle (Fig. 77), holding the spirit of wine and ether, is a common flask through the cork of which a short straight tube passes. The outer end of the tube should be sufficiently narrowed to enable it to deliver a very fine stream of the liquid. The flask being inverted, the warmth of the hand expands the air and vapor to a sufficient extent to force out the liquid.

Fig 77.



The ordinary washing-bottle for quantitative operations should be formed of a flask in which water may be boiled, fitted up as usual.

(Vide p. 109.)

A water-oven is the best form of drying apparatus. It is a small square copper vessel, jacketed on five sides and having a door on

the sixth; water is poured into the space between the inner and outer casing, and the whole placed over a gas-lamp or other source of heat, moist air and steam escaping by appropriate apertures. Desiccation at higher temperatures than the boiling-point of water may be practised by using oil or paraffin instead of water, inserting a thermometer in the fat. The apparatus may be purchased of any maker of chemical instruments.

Pure distilled water must be used in all quantitative determinations.

Note—In practising the operations of quantitative analysis, experiments should at first be conducted on definite salts of known composition, for the accuracy of results may then be tested by calculation.

Estimation of Potassium in the Form of Double Potassium and Platinum Chloride.—Select two or three crystals of pure potassium nitrate, powder them in a clean mortar, dry the powder by gently heating in a porcelain crucible over a flame for a few seconds, place about a couple of décigrammes (0.2 gm.) of the powder in a counterpoised watch-glass, accurately weigh the selected quantity, transfer to a small dish, letting water from a wash-bottle flow over the watch-glass and run into the dish; warm the dish till the nitrate is dissolved, acidulate with hydrochloric acid, add excess of aqueous solution of platinum perchloride (a quantity containing about 0.4 of solid salt), evaporate to dryness over a vapor-bath. While evaporation is going on place a filter and the weighing-tubes in the water-oven, exposing them to a temperature of 100° C. for about half an hour; fold the filter and insert it in the tubes, place them on a plate under a glass shade, and when cold accurately note their weight. Arrange the weighed filter in a funnel over a beaker. Transfer the dried and cooled platinum salt from the dish to the filter by moistening the residue with the mixture of alcohol and ether, and, when the salt is loosened, pouring the contents of the dish into the paper cone. Any salt still adhering may be freed by the finger, which, together with the dish, should be washed in the stream of spirit, the rinsings at once flowing into the filter. The filtrate should have a yellowish-brown color, due to the excess of platinum perchloride. If it is colorless, an insufficient amount of perchloride has been added, and the whole operation must be repeated. The washed precipitate and filter are finally dried in the water-oven, folded and placed in the weighing-tubes, the drying and weighing when cold being repeated until the whole ceases to alter, the final weight being noted.

Desiccators.—Highly-dried substances are very hygroscopic; hence before being weighed should be cooled under a bell-jar which also encloses a vessel containing sulphuric acid or calcium chloride—a

desiccator. (If filters are not freed from all traces of acid by thorough washing, the paper will be brittle when dry, falling to pieces on being folded.)

Analytical memoranda may have the following form :

Watch-glass and substance	
Watch-glass	
	Substance
Weighing-tubes, filter, and Pt. salt . .	
Weighing-tubes and filter	
	PtCl ₄ , 2KCl

The calculations are simple :

As $\left\{ \begin{array}{l} \text{PtCl}_4, 2\text{KCl} \\ = 484.8 \end{array} \right\}$ is equivalent to $\left\{ \begin{array}{l} 2\text{KNO}_3 \\ = 202 \end{array} \right\}$,
 so $\left\{ \begin{array}{l} \text{the weight of} \\ \text{double chloride} \\ \text{obtained} \end{array} \right\}$ is equivalent to x .

x will be the amount of pure potassium nitrate in the quantity of substance operated on. x should in the present instance be identical with the weight of substance taken, because, for educational purposes, pure nitre is under examination. Only after analyses of pure substances have yielded the operator results practically identical with those by calculation can analyses of substances of unknown degree of purity be undertaken with confidence. A table of atomic weights, from which to find molecular weights, is given in the Appendix. Cl = 35.5, or, more exactly, 35.4.

Platinum residues should be preserved, and the metal recovered from them from time to time. (*Vide* p. 249.)

Hot alcohol sometimes reduces platinum perchloride, the metal being thrown out of solution in a finely-divided form, known as *platinum black*; only aqueous solutions, therefore, of the salt should be used where heat is employed. Hence, also, in washing out excess of platinum perchloride from the double platinum and potassium chloride by spirit the application of heat should be avoided.


Effervescing Potash-water (*Liquor Potassæ Effervescens*, B. P.) is most easily estimated volumetrically (p. 634). Any adulteration by an equivalent amount of sodium bicarbonate would, however, by that process be undetected; hence the Pharmacopœia directs that "5 fluidounces, evaporated to one-fifth, and 12 grains of tartaric acid added, yield a crystalline precipitate which when dried weighs not less than 12 grains." 5 fluidounces of this preparation should contain 7.5 grains of bicarbonate, convertible into 14.1 grains of acid potassium tartrate by 11.25 grains of tartaric acid. The method is somewhat rough, but quite efficient for "potash-water" containing nothing but bicarbonate of alkali-metals.

Proportional Weights of Equivalent Quantities of Potassium and its Salts.

Metal	K ₂	78
Oxide ("Potash")	K ₂ O	94
Hydrate ("Caustic Potash")	2KHO	112
Carbonate (anhydrous)	K ₂ CO ₃	138
Carbonate (crystalline)	K ₂ CO ₃ + 16% Aq	164.285
Bicarbonate	2KHCO ₃	200
Nitrate	2KNO ₃	202
Platinum salt	PtCl ₄ .2KCl	484.8

SODIUM.

Sodium is usually estimated as sulphate. Accurately weigh a porcelain crucible and lid; place within about .3 of pure rock-salt and again weigh, making a memorandum of the weights in a note-book. Add rather more strong sulphuric acid than may be considered sufficient to convert the chloride into acid sodium sulphate. Heat the crucible gradually, the flame being first directed against the side of the crucible to avoid violent ebullition, until fumes of acid cease to be evolved, toward the end of the operation dropping in one or two fragments of ammonium carbonate to facilitate complete expulsion of all excess of acid. When cold weigh the crucible and contents. The weight of the crucible having been deducted, the amount of sulphate obtained should be the exact equivalent of the quantity of sodium chloride employed.

*Proportional Weights of Equivalent Quantities of Sodium and its Salts.*

Metal	Na ₂	46
Oxide ("Soda")	Na ₂ O	62
Hydrate ("Caustic Soda")	2NaHO	80
Carbonate (anhydrous)	Na ₂ CO ₃	106
Carbonate (crystals)	Na ₂ CO ₃ .10H ₂ O	286
Bicarbonate	2NaHCO ₃	168
Chloride	2NaCl	116.8
Sulphate (anhydrous)	Na ₂ SO ₄	142
Sulphate (crystals)	Na ₂ SO ₄ .10H ₂ O	322

AMMONIUM.

Salts of ammonium are, for purposes of quantitative analysis, generally converted into the double ammonium and platinum chloride (PtCl₄.2NH₄Cl), the details of manipulation being

the same as those observed in the case of potassium (p. 661). About 0.15 grm. of pure, white, dry ammonium may be taken for experiment.

Composition of the Platinum Salt.

		In 1 molec. wt.	In 100 parts.
Pt	194.4	194.4	43.903
Cl ₆	35.4 × 6	212.4	47.967
N ₂	14.0 × 2	28	6.324
H ₈	1.0 × 8	8	1.806
		442.8	100.00
		In 1 molec. wt.	In 100 parts.
or, PtCl ₄	336	336	75.88
2NH ₄ Cl	53.4 × 2	106.8	24.12
		442.8	100.00

The proportion of nitrogen, ammonium, or ammonium chloride in the double chloride may also be ascertained from the weight of platinum left on igniting the double chloride; indeed, this operation *must* be performed if any variety of ammonium other than the ordinary hydrogen ammonium may be present. The heat must be applied slowly or platinum will be mechanically carried off with the gaseous products of decomposition.

Proportional Weights of Equivalent Quantities of Ammoniacal Compounds.

Ammonia (gas)	2NH ₃	34
Ammonium	(NH ₄) ₂ ?	36
Ammonium chloride	2NH ₄ Cl	106.8
Platinum salt	PtCl ₄ ·2NH ₄ Cl	442.8
"Carbonate of ammonium"	(N ₃ H ₁₁ C ₂ O ₅) ÷ 3 × 2	104.7
Ammonium sulphate	(NH ₄) ₂ SO ₄	132

BARIUM.

Barium is estimated in the form of anhydrous barium sulphate (BaSO₄), Ba = 136.8.

Process.—Dissolve 0.3 or 0.4 of pure crystallized and dried barium chloride or nitrate in about $\frac{1}{2}$ a litre of water in a beaker, heating to incipient ebullition, and slightly acidulating with hydrochloric or nitric acid. Add diluted sulphuric acid (prepared some days previously, so that lead sulphate may have deposited) so long as a precipitate forms, keep the mixture hot for some time, set aside for half an hour, pass the supernatant liquid through a filter, gently boil the residue twice or thrice with acidulated water; finally, collect the precipitate on the filter, removing adherent particles from the beaker by the finger and cleansing by a stream of hot water from the wash-bottle. The precipitate must be washed with hot water

until the filtrate ceases to turn litmus-paper red or give any cloudiness when tested with barium chloride. The filter and barium sulphate, having thoroughly drained, are dried in a warm place, commonly by supporting the funnel in an inverted bottomless beaker over a sand-bath or hot plate. The barium sulphate is now removed from the filter, heated to drive off every trace of moisture, and weighed. This is accomplished by placing a weighed porcelain crucible (and cover) on a sheet of glazed paper, holding the filter over it, and carefully transferring the precipitate; the sides of the filter are then gently rubbed together and detached powder dropped into the crucible, the paper folded, encased in two or three coils of one end of a platinum wire, and burnt over the crucible, ash and any particles on the sheet of paper dropped into the barium sulphate, the open crucible exposed over a flame till its contents are quite white, covered, cooled, and weighed.

	Formulae.	Molecular weights.
Barium chloride	$\text{BaCl}_2 \cdot 2\text{H}_2\text{O}$	243.6
Barium nitrate	Ba_2NO_3	260.8
Barium sulphate	BaSO_4	232.8

Composition of Barium Sulphate.

		In 1 molec. wt.	In 100 parts.
Ba	136.8	136.8	58.77
S	32	32	13.73
O ₄	16×4	64	27.50
		232.8	100.00

In the first four or five educational experiments it is not essential to take filter-ash into account. Mistakes of manipulation due to inexperience may cause far greater errors.

CALCIUM.

Calcium is usually thrown out of solution in the form of oxalate, the precipitate ignited, and the resulting carbonate weighed.

Process.—Dissolve 0.3 or 0.4 of dried colorless crystals of calc-spar in about a third of a litre of water acidulated with hydrochloric acid, heat the solution to near the boiling-point, add excess of solution of ammonium oxalate, then ammonia until, after stirring, the liquid smells strongly ammoniacal; set aside in a warm place for twelve hours. Carefully pour off the supernatant liquid, passing it through a filter; add hot water to the precipitate, set aside for half an hour, again decant, and, after once more washing, transfer the precipitate to the filter,

allowing all contained fluid to pass through before a fresh portion is added. Wash the precipitate with hot water, avoiding a rapid stream or the precipitate may be driven through the pores of the paper. Dry, transfer to a weighed crucible, and incinerate, as described for barium sulphate, and slowly heat the precipitate till the bottom of the crucible is just visibly red when seen in the dark. As soon as the residue is white or only faintly gray remove the lamp, cool, and weigh.

The resulting calcium carbonate should have the same weight as the calc-spar from which it was obtained. If loss has occurred, carbonic acid gas has probably escaped. In that case moisten the residue with water, and after a few minutes test the liquid with red litmus- or turmeric-paper; if an alkaline reaction is noticed, it is due to the presence of caustic lime. Add a small lump of ammonium carbonate, evaporate to dryness over a water-bath, and again ignite, this time being careful not to go beyond the prescribed temperature. The treatment may, if necessary, be repeated.

Proportional Weights of Equivalent Quantities of Calcium and its Salts.

Metal	Ca	40
Oxide (quicklime)	CaO	56
Hydrate (slaked lime)	Ca ₂ HO	74
Carbonate	CaCO ₃	100
Sulphate (anhydrous)	CaSO ₄	136
Sulphate (crystalline or precipitated)	CaSO ₄ , 2H ₂ O	172
Chloride	CaCl ₂	110.8
Phosphate (of bones)	(Ca ₃ PO ₄) ÷ 3	103.3

MAGNESIUM.

Process 1.—The light or heavy magnesium carbonate of pharmacy may be estimated by heating a weighed quantity to redness in a porcelain crucible. If it has the composition indicated by the formula given in the British Pharmacopœia (3MgCO₃, Mg₂HO, 4H₂O), it will yield 42 per cent. of magnesia (MgO). According to that work, the purity of even magnesium sulphate (MgSO₄, 7H₂O) may be determined by boiling a weighed quantity with excess of sodium carbonate, collecting the precipitate, washing, drying, igniting, and weighing the resulting magnesia (MgO). The crystallized sulphate should afford 16.26 per cent. of oxide. The official solution of magnesium carbonate in carbonic-acid water (*Liquor Magnesii Carbonatis*, B. P.) should yield about 4 grains of pure magnesium oxide per fluidounce.

Process 2.—The general form in which magnesium is precipitated is as ammonium and magnesium phosphate ($\text{MgNH}_4\text{PO}_4 \cdot 6\text{H}_2\text{O}$); this, by heat, is converted into magnesium pyrophosphate ($\text{Mg}_2\text{P}_2\text{O}_7$). Accurately weigh a small quantity (0.4 to 0.5) of pure dry crystals of magnesium sulphate, dissolve in 200 to 300 cubic centimetres of cold water in a beaker, add ammonium chloride, ammonia, and sodium or ammonium phosphate, agitate with a glass rod (without touching the sides of the vessel, or crystals will firmly adhere to the rubbed portions), and set aside for twelve hours. Collect on a filter, wash the precipitate with water containing a tenth of its volume of the strongest solution of ammonia until the filtrate ceases to give a precipitate with an acidulated solution of silver nitrate. Dry, transfer to a crucible, burn the filter in the usual way, heat slowly to redness, cool, and weigh.

Proportional Weights of Equivalent Quantities of Magnesium Salts.

Pyrophosphate . . . $\text{Mg}_2\text{P}_2\text{O}_7$	222
Sulphate $2(\text{MgSO}_4 \cdot 7\text{H}_2\text{O})$	492
Oxide $2(\text{MgO})$	80
Official carbonate . $(3\text{MgCO}_3, \text{Mg}2\text{HIO}, 4\text{H}_2\text{O}) \div 2$.	191

ZINC.

Zinc is usually estimated as oxide (ZnO), occasionally as sulphide (ZnS), $\text{Zn} = 64.9$.


Process.—Dissolve a weighed quantity (0.5 to 0.6) of zinc sulphate in about $\frac{1}{2}$ a litre of water in a beaker, heat to near the boiling-point, add sodium carbonate in slight excess, boil, set aside for a short time; pass the supernatant liquid through a filter, gently boil the precipitate with more water, again decant; repeat these operations two or three times; collect the precipitate on the filter, wash, dry, transfer to a crucible, incinerate, ignite, cool, and weigh. 286.9 (= molec. weight) of sulphate should yield 80.9 (= molec. weight) of oxide.

MANGANESE.

To ascertain its value for evolving chlorine from hydrochloric acid, a weighed quantity of finely-powdered black manganese oxide is heated in a small flask with pure hydrochloric acid (contained in an inner tube, as for "oxalates" and "carbonates," p. 683), and the resulting chlorine conveyed into a U-tube containing solution of potassium iodide. (See Fig. 78.) The amount of iodine thus freed is estimated by the volumet-

ric solution of sodium hyposulphite. 126.6 of iodine indicate 35.4 of chlorine.

Fig. 78.



(Manganese may also be estimated by the reaction and apparatus described under "Oxalates, p. 684.)

ALUMINIUM.

Aluminium is always precipitated as hydrate ($\text{Al}_2\text{6HO}$) and weighed as oxide (Al_2O_3).

Process.—Dissolve about 2 grammes of pure dry ammonium-alum in $\frac{1}{2}$ a litre of water, heat the solution, add ammonium chloride and a slight excess of ammonia, boil gently till the odor of ammonia has nearly disappeared, set aside for the hydrate to deposit, pass the supernatant liquid through a filter, wash the precipitate three or four times by decantation, transfer to the filter, finish the washing, dry, burn the filter, ignite in a covered crucible, cool, and weigh.

$\text{Al}_2\text{3SO}_4\text{K}_2\text{SO}_4, 24\text{H}_2\text{O}$	948
$\text{Al}_2\text{3SO}_4, (\text{NH}_4)_2\text{SO}_4, 24\text{H}_2\text{O}$	906
Al_2O_3	102
Per cent. of Al_2O_3 yielded by ammonium-alum	11.26

QUESTIONS AND EXERCISES.

Give details of the manipulations observed in gravimetrically estimating salts of potassium or of ammonium.—What quantity of sodium chloride is contained in a sample of rock-salt, 0.351 of a gramme of which yields 0.426 of sodium sulphate? *Ans.* 99.83 per cent.—To what amount of the ammonium-alum is 0.888 of a gramme of the double platinum and ammonium chloride equivalent?—*Ans.* 1.817 grammes.—Find the weight of barium sulphate obtainable from 0.522 of nitrate? *Ans.* 0.466.—Describe the usual method by which salts of calcium are estimated.—By what quantitative process may the official salts of magnesium be analyzed?—Calculate the proportion of pure zinc sulphate in a sample of

crystals 0.574 of which yield 0.161 of oxide. *Ans.* 99.4 per cent.—Ascertain the weight of alumina (Al_2O_3) which should be obtained from 1.812 grammes of ammonium-alum.

IRON.

Iron and its salts are gravimetrically estimated in the form of ferric oxide (Fe_2O_3).

Compounds containing organic acidulous radicals are simply incinerated and the resulting oxide weighed. Thus 1 gramme of the official citrate of iron and ammonium (*Ferri et Ammonii Citras*, B. P.), incinerated, with exposure to air, leaves not less than .27 of ferric oxide. A small quantity of the salt is weighed in a tared covered porcelain crucible, flame cautiously applied until vapors cease to be evolved, the lid then removed, the crucible slightly inclined and exposed to a red heat until all carbonaceous matter has disappeared. The residual ferric oxide is then weighed. Potassium and iron tartrate (*Ferrum Tartaratum*, B. P.) is treated in the same manner, except that the ash must be washed and again heated before weighing, in order to remove potassium carbonate produced during incineration; 5 grammes should yield about 1.5 grammes of ferric oxide.

From other compounds of iron, soluble in water or acid, the metal is precipitated in the form of hydrate (Fe_26HO) by solution of ammonia, and converted into oxide (Fe_2O_3) by ignition. Dissolve a piece (about 0.2) of the purest iron obtainable (piano wire), accurately weighed, in water acidulated with hydrochloric acid; add a few drops of nitric acid, and gently boil; pour in excess of ammonia, stir, set aside till the ferric hydrate has deposited, pass the supernatant liquid through a filter, treat the precipitate three or four times with boiling water; transfer to the filter, wash till the filtrate yields no trace of chlorine (for ammonium chloride will decompose ignited ferric oxide, with volatilization of ferric chloride); dry, and ignite as usual, and weigh. Iron in the official solutions (*Liquor Ferri Acetatis*, *Liquor Ferri Nitratis*, and *Liquor Ferri Tersulphatis*) is estimated by this general process.

The proportion of metallic iron in a mixture of iron and oxides of iron may be determined by digestion in a strong solution of iodine and potassium iodide, which attacks the metal only. The reduced iron of pharmacy (*Ferrum Reductum*, U. S. P.) is in good condition so long as it contains, as shown by this method, half its weight of free metal.

Another Method.—Reduced iron is converted into ferrous chloride by a hot, strong solution of corrosive sublimate, while

the oxides are not affected. The filtrate may be treated gravimetrically or volumetrically (Wilner).

Proportional Weights of Equivalent Quantities of Iron and its Salts.

Metal	Fe ₂	112
Ferric oxide	Fe ₂ O ₃	160
Ferric hydrate	Fe ₂ 6HO	214
Ferric chloride	Fe ₂ Cl ₆	324.4
Ferric sulphate	Fe ₂ 3SO ₄	400
Ferrous sulphate	2(FeSO ₄ ,7H ₂ O)	556

ARSENUM.

Arsenic (As₂O₃) is usually estimated volumetrically (*vide* p. 651), and sometimes arsenates also (*vide* p. 646), but the latter are best precipitated as a lead salt; an "aqueous solution of 12.4 grains of anhydrous sodium arsenate, acidulated with acetic acid, requires not less than 34 grains of lead acetate for complete precipitation." With certain precautions arsenum may be precipitated and weighed as sulphide (As₂S₃).

Process 1.—The pure white arsenic in lump (about 0.2) is dissolved in a flask in a small quantity of water containing sodium or potassium bicarbonate, the liquid being heated. A slight excess of hydrochloric acid is then added, and sulphuretted hydrogen gas passed through the solution so long as a precipitate falls, the mouth of the flask being stopped by a plug of cotton-wool (to prevent undue access of air and consequent decomposition of the gas, resulting in precipitation of sulphur). The mixture is warmed in the flask, and carbonic acid gas passed through it until the odor of sulphuretted hydrogen has nearly disappeared; the precipitate is collected on a tared filter, washed as quickly as possible with hot water containing a little sulphuretted hydrogen, dried in a water-oven, and weighed. 198 parts of arsenic should yield 246 of arsenum sulphide.

Process 2.—The arsenum must be present in the arsenic condition. If the operator is not certain that this is the case, the solution must be warmed with a little hydrochloric acid and a few grains of potassium chlorate added until a distinct odor of chlorous vapor is evolved, which is then allowed to escape by continued application of heat. To the solution thus obtained ammonia, which must produce no turbidity, is added in excess, and then magnesia mixture. (See under "Phosphates," p. 684.) The solution is set aside for twenty-four or forty-eight hours. The precipitate is collected on a filter and washed with as little ammonia-water (1 to 3) as possible until the filtrate ceases to

give a reaction for chlorides. The precipitate is then dried on the filter, the precipitate and filter-paper burned, and the whole gently ignited in a crucible and weighed. This residue is magnesium pyroarsenate and has the formula $\text{Mg}_2\text{As}_2\text{O}_7$.

ANTIMONY.

The metal is precipitated in the form of sulphide (Sb_2S_3), with the precautions observed in estimating arsenum—a small quantity of tartaric acid, as well as hydrochloric, being added to prevent the precipitation of an oxysalt. If the sulphuretted hydrogen be passed through a *hot* solution, the particles of precipitate aggregate better, and the latter may be more quickly filtered and washed. The experiment may be performed on about $\frac{1}{2}$ a gramme of pure tartar emetic; the salt should yield nearly half its weight of sulphide. According to Fresenius, the sulphide dried at 100°C . still contains 2 per cent. of water, and must be heated in a current of carbonic acid gas until it turns from an orange to a black color before all moisture is expelled. In the British Pharmacopœia the purity of tartar emetic (*Antimonium Tartaratum*), and the strength of solution of antimony chloride (*Liquor Antimonii Chloridi*), are determined by the above process. Of the *Antimonium Sulphuratum*, B. P., it is stated “that 60 grains moistened and warmed with successive portions of nitric acid until red fumes cease to be evolved, and then dried and heated to redness, give a white residue [Sb_2O_4] weighing about 40 grains.”

(For the volumetric estimation of antimony in antimonious salts see p. 647.)

COPPER.

Copper is precipitated from its solutions and weighed either (1) as metal (Cu_2) or (2) as oxide (CuO).

Process 1.—Dissolve about $\frac{1}{2}$ a gramme of dry crystallized copper sulphate in a small quantity of water in a tared porcelain crucible or beaker, acidulate with hydrochloric acid, introduce a fragment or two of pure zinc, cover the vessel with a watch-glass, and set aside till evolution of hydrogen has ceased and the still acid liquid is colorless. The copper is then washed with hot water by decantation until no trace of acid remains, the precipitate drained, rinsed with strong spirit of wine, dried in the water-oven, cooled, and weighed.

Process 2.—From a solution acidulated by sulphuric acid and placed in a platinum crucible copper may be entirely deposited in a coherent form by a weak current of electricity, the crucible being connected with the zinc pole of the battery, a platinum

spatula suspended in the solution forming the positive pole. The crucible may afterward be freed from the deposited copper by nitric acid.

Process 3.—About three-fourths of a gramme of copper sulphate is accurately weighed, dissolved in $\frac{1}{2}$ a litre of water, the liquid boiled; dilute solution of potash or soda is then added till no more precipitate falls, ebullition continued for a short time, and the beaker set aside; the supernatant liquid is decanted, the precipitate boiled with water twice or thrice, collected on a filter, washed, dried, transferred to a crucible, the filter incinerated, and its ash moistened with a drop of nitric acid; the whole is finally heated strongly, cooled, and weighed.

249.3 parts of copper sulphate yield 79.3 of oxide or 63.3 of metal.

Other Processes.—*Vide Pharmaceutical Journal* for April 3, 1880, p. 801.

BISMUTH.


Dissolve 0.3 or 0.4 of the pure bismuth oxycarbonate ($\text{Bi}_2\text{O}_3 \cdot \text{CO}_2$), H_2O (*Bismuthi Subcarbonas*, U. S. P.) in a very small quantity of hydrochloric acid, dilute with water slightly acidulated by hydrochloric acid, pass excess of sulphuretted hydrogen through the liquid, collect the precipitate on a tared filter, wash, dry at 100°C ., and weigh. The sulphide must not be exposed too long in the water-oven or it will increase in weight, owing to absorption of oxygen; hence it should be tested in the balance every half-hour during desiccation. 519 parts of oxycarbonate should yield 514 of sulphide (Bi_2S_3). The strength of the official solution (B. P.) of bismuth and ammonium citrate (*Liquor Bismuthi et Ammonii Citratis*, B. P.) is determined by this process. "Two fluidrachms of the solution, mixed with an ounce of distilled water, and treated with sulphuretted hydrogen in excess, yield a black precipitate, which, collected, washed, and dried, weighs about 7 grains. 1 fluidrachm yields 3 grains of bismuth oxide" (B. P.). Of the official Citrate of Bismuth, B. P., it is stated that "10 grains dissolved in solution of ammonia and treated with sulphuretted hydrogen in excess, yield a precipitate which, when washed and dried, weighs about 7 grains;" and of the Bismuth and Ammonium Citrate, B. P., "10 grains dissolved in water and treated with sulphuretted hydrogen in excess, yield a precipitate which, when washed and dried, weighs about $6\frac{1}{2}$ grains" (B. P.). The atomic weight of bismuth is 208 if oxygen = 16, or 207.5 if oxygen = 15.96. The U. S. P. bismuth substances may be estimated similarly.

MERCURY.

This element may be (1) isolated and estimated in the form of metal, or precipitated and weighed as (2) mercurous chloride or (3) mercuric sulphide.

Process 1.—The process by which the metal itself is separated is one of distillation into a bulb surrounded by water. About $\frac{1}{2}$ a mètre of the difficultly fusible German glass known as *combustion-tubing* is sealed at one end after the manner of a test-tube (Fig. 79); a mixture of sodium bicarbonate and dry chalk is then dropped into the tube to the height of two or three centimètres, and, next, several small fragments of quicklime, so as to occupy another centimètre; a mixture of about


Figs. 79, 80, 81.



a gramme of pure calomel or corrosive sublimate with enough quicklime to occupy ten or twelve centimètres of the tube is added, then the lime-rinsings of the mixing-mortar, a layer of a few centimètres of powdered quicklime, and, finally, a plug of *asbestos*. The whole should occupy two-thirds of the tube. The part of the tube just above the asbestos is now softened in the blowpipe flame and drawn out about a décimètre to the diameter of a narrow quill (Fig. 80); again drawn out to the same extent at a point two or three centimètres nearer the mouth (Fig. 80), and any excess of tubing cut off. The bulb thus formed may be enlarged by softening and blowing. The tube is next softened at a point close to but anterior to the asbestos, and bent to form an obtuse angle; the tube is then softened close to the bulb and slightly bent so that the bulb may be parallel with the large tube; then softened on the other side of the bulb, and the terminal tube bent to an obtuse angle, so that, the tube being held in a horizontal position, the bulb may be sunk in water and the terminal tube point upward (Fig. 81). The long tube is now laid in the gas-furnace found in most laboratories (Fig. 82), a basin so placed that the bulb of the apparatus may be cooled by being surrounded by water, the part of the tube occupied by asbestos heated to redness, and the flame slowly lengthened until the whole tube is red hot. Under the circumstances just described the mercurial com-

pound volatilizes, is decomposed by the lime, and its acidulous radical fixed, the mercury carried in vapor to and condensed in the bulb, the carbonic acid gas evolved from the sodium bicarbonate and chalk washing out the last portions of mercury-

Fig. 82.



Distillation of Mercury for Quantitative Purposes.

vapor from the tube. When the distillation is considered to be complete, the dish of water is removed, the bulb dried, and then detached by help of a file at a point beyond any sublimate of mercury. The dried bulb is weighed, the mercury shaken or dissolved out, and the tube again dried and weighed. The difference between the weights gives the weight of the mercury. "Ammoniated Mercury," B. P., should yield, theoretically, 79.52 per cent., practically about 77.5 per cent., of metallic mercury.

Process 2.—The process by which mercury is separated in the form of calomel consists in adding solution of hydrochloric and of phosphorous acids to an aqueous or even acid solution of a weighed quantity of the mercurial compound, setting the mixture aside for twelve hours, collecting the precipitate on a tared filter, washing, drying at 100° C., and weighing (Rose). The experiment may be tried on $\frac{1}{2}$ a gramme to 1 gramme of corrosive sublimate.

Process 3.—2 or 3 décigrammes of corrosive sublimate are dissolved in water, the solution acidulated with hydrochloric acid, excess of sulphuretted hydrogen passed through it, the precipitate collected on a tared filter, washed with cold water, dried at 100° C., and weighed.

Proportional Weights of Equivalent Quantities of Mercury and its Salts.

Metal	Hg	200
Mercurous chloride	HgCl	235.4
Mercuric chloride	HgCl ₂	270.8
Mercuric sulphide	HgS	232

LEAD.

Lead is generally estimated either as (1) oxide, (2) sulphate, (3) chromate, or (4) metal.

Process 1.—Weigh out 1 or 2 grammes of pure lead acetate in a covered crucible, previously tared, and heat slowly until no more vapors are evolved. Remove the lid, stir down the carbonaceous mass with a clean iron wire, and keep the crucible in the flame so long as any carbon remains unconsumed. Introduce some fragments of fused ammonium nitrate, and again ignite until no metallic lead remains and all excess of the nitrate has been decomposed. Cool and weigh the resulting oxide (PbO).

Process 2.—Dissolve 0.4 or 0.5 of a gramme of lead acetate in a small quantity of water, drop in diluted sulphuric acid, add to the mixture twice its bulk of methylated spirit of wine, and set aside. Decant the supernatant liquid, collect the sulphate on a filter, wash with spirit, dry, transfer to a porcelain crucible, removing as much of the sulphate as possible from the paper, incinerate on the crucible-lid (not in the platinum coil, for the fused particles of reduced lead would alloy with the platinum), ignite, cool, and weigh.

Process 3.—About $\frac{1}{2}$ a gramme of lead acetate is dissolved in 200 or 300 cc. of water, acetic acid added, and then solution of red potassium chromate. Collect the precipitate on a tared filter, wash, dry at 100°C ., and weigh.

Process 4.—In certain cases, notably in that of commercial "white lead," the lead may be estimated in the metallic state by means of potassium cyanide. The lead paint (about 20 grammes) is weighed and carefully incinerated. The residue, a mixture of metallic lead and lead oxide, is then mixed with several times its bulk of potassium cyanide and the whole heated to fusion. With careful manipulation the lead collects in one globule, which, after cooling, may readily be separated from the mixed cyanide and cyanate and weighed. Commercially pure white lead should yield 74 per cent. of lead.

Proportional Weights of Equivalent Quantities of Lead and its Salts.

Metal	Pb	206.4
Acetate	$\text{Pb2C}_2\text{H}_3\text{O}_2, 2\text{H}_2\text{O}$	378.4
Oxide	PbO	222.4
Sulphate	PbSO_4	302.4
Chromate	PbCrO_4	322.9

SILVER.

Compounds of silver which are readily decomposed by heat

are estimated in the form of (1) metal, others usually as (2) chloride (AgCl), but sometimes as (3) cyanide (AgCN).

Process 1.—Heat about a gramme of silver oxide (Ag_2O) in a tared crucible, cool, and weigh. 232 parts of oxide yield 216 of metal. “29 grains heated to redness yield 27 grains of metallic silver.”—*B. P.*

Process 2.—Dissolve 0.4 or 0.5 grm. of pure dry crystals of silver nitrate in water, acidulate with 2 or 3 drops of nitric acid, slowly add hydrochloric acid, stirring rapidly, until no more precipitate falls. Pour off the supernatant liquid through a filter, wash the silver chloride once or twice with hot water, transfer to the filter, complete the washing, and dry. After removing as much as possible of the precipitate from the paper to the crucible, burn the filter, not in a wire helix, but on the inverted lid of the crucible, moisten with a drop of nitric acid, warm, add a drop of hydrochloric acid, evaporate to dryness, replace the lid of the crucible, ignite the whole until the edges of the mass of chloride begin to fuse; cool, and weigh. 170 parts of nitrate yield 143.4 of chloride. According to the United States Pharmacopœia, all silver salts should be estimated volumetrically with standard sodium chloride solution. (*Vide p. 657.*) 30 parts of “Mitigated Caustic” (*Argenti et Potassii Nitrates*, *B. P.*) should give 8.44 parts of silver chloride, and the nitrate yields potassium nitrate and chloride on evaporation. “Toughened Caustic,” *B. P.*, contains 5 per cent. of potassium nitrate; hence 10 parts will give only 8 parts of silver chloride, and the filtrate will yield a little potassium salt. 10 parts of silver, if pure (*Argentum Purificatum*, *B. P.*), will give 13.285 of silver chloride.

Process 3.—Silver cyanide may be collected on a tared filter and dried at 100°C . 170 parts of nitrate yield 134 of cyanide. Silver and its salts may be volumetrically estimated by a standard solution of sodium chloride.

Cupellation.—The amount of silver in an alloy may also be determined by a dry method. The metal is folded in a piece of thin sheet lead, placed on a *cupel* (*cupella*, little cup, made of compressed bone-earth) and heated in a furnace, the cupel being protected from the direct action of the flame by a muff-shaped—or, rather, oven-shaped—case, termed a muffle. The metals melt, the baser become oxidized, the lead oxide fusing and dissolving the other oxides; the fluid oxides are absorbed by the porous cupel, a button of pure silver remaining. An alloy suspected to contain 95 per cent. of silver requires about three times its weight of lead for successful cupellation; if $92\frac{1}{2}$ per cent. (English silver coin), between five and six times its weight of lead is necessary.

QUESTIONS AND EXERCISES.

Explain the gravimetric process by which the strength of the official solutions of ferric chloride, nitrate, and sulphate are determined.—Mention the various amounts of ferrous and ferric salts equivalent to 100 parts of metal.—State the precautions necessary to be observed in estimating arsenum or antimony in the form of sulphide.—In what form are the official compounds of bismuth weighed for quantitative purposes?—Give an outline of the process by which mercury may be isolated from its official preparations and weighed in the metallic condition.—Describe three methods for the quantitative analysis of salts of lead, and the weights of the respective precipitates, supposing 0.56 of crystallized acetate to have been operated on in each case.—Describe the processes by which silver is estimated in the forms of metal, chloride, and cyanide.—What proportions of silver nitrate are indicated, respectively, by 15 of metal, 9.8 of chloride, and 8.1 of cyanide?—Describe cupellation.

GRAVIMETRIC ESTIMATION OF THE ACIDULOUS RADICALS OF SALTS.

CHLORIDES.

Free chlorine (chlorine-water) and compounds which by action of acids yield free chlorine (chlorinated lime, chlorinated soda, and their official solutions) are estimated volumetrically by a standard solution of sodium hyposulphite. (*Vide* p. 655.) The amount of combined chlorine in pure chlorides (HCl , NaCl) may also be determined by volumetric analysis with a standard solution of silver nitrate (p. 639).

Combined chlorine is gravimetrically estimated in the form of silver chloride, the operation being identical with that just described for silver salts (p. 676). 58.4 parts of pure, colorless, crystallized sodium chloride (rock-salt) yield 143.4 of silver chloride.

IODIDES.

Free iodine is estimated volumetrically by solution of sodium hyposulphite. (*Vide* p. 655.)

Combined iodine is determined gravimetrically in the form of silver iodide, the operations being conducted as with silver chloride. Potassium iodide may be used for an experimental determination: $\text{KI} = 166$ should yield $\text{AgI} = 235$. Of cadmium iodide (*Cadmii Iodidum*, B. P. 1867) it is stated that "10 grains dissolved in water, and silver nitrate added in excess, give a precipitate which, when washed with water and afterward with $\frac{1}{2}$ an ounce of solution of ammonia, and dried, weighs 12.5 grains."

In presence of chlorides and bromides the iodine in iodides may be precipitated and weighed as palladium iodide.

Moisture in iodine is estimated by loss on exposing a weighed quantity of iodine in a capsule over a dish of sulphuric acid under a small bell jar, or by adding to a weighed sample five or six times as much mercury or twice as much zinc, and a little water, drying, and weighing. The product is the amount of metal employed plus that of the *dry* iodine in the sample.

BROMIDES.

Free bromine may be estimated by shaking with excess of solution of potassium iodide, and then determining the equivalent quantity of liberated iodine by a standard solution of sodium hyposulphite (p. 655).

The bromine in bromides may be precipitated and weighed as silver bromide, the manipulations being the same as those for silver chloride: 0.2 to 0.3 of pure potassium bromide may be used for an experimental analysis.

Ammonium bromide and calcium bromide are estimated volumetrically. (*Vide* p. 642.)

CYANIDES.

The hydrogen cyanide (hydrocyanic acid) is usually estimated volumetrically. (*Vide* p. 640.)

From all soluble cyanides cyanogen may be precipitated by silver nitrate after acidulating with nitric acid, the silver cyanide being collected on a tared filter, dried at 100°, and weighed.

Silver Cyanide.

		In 1 molec. wt.	In 100 parts.
Silver Ag		107.66	80.55
Cyanogen CN		26.00	19.45
		133.66	100.00


NITRATES.

Nitrates cannot be estimated by direct gravimetric analysis, none of the basylous radicals yielding a definite nitrate insoluble in water. With some difficulty they may be determined by indirect volumetric methods.

Process.—The following (Thorpe's) method depends upon the fact (Gladstone and Tribe) that when zinc upon which copper is deposited in a spongy form is boiled with water, hydrogen is evolved. Thorpe found that in a solution containing nitrates the nascent hydrogen converts the whole of the nitrogen of

the nitrates into ammonia, which may be collected and estimated. (The oxygen of the nitrate is simultaneously converted into water, the nitrate-metal into hydrate, and the zinc into zinc hydrate. The power of the copper-zinc couple is considered to depend largely on the hydrogen absorbed by the finely-divided metal.)

Fig. 83.



Estimation of Nitrates.

An apparatus such as shown in Fig. 83 should be constructed. A flask (about 100 cc.) is fitted with a clean sound cork perforated for a delivery-tube, which should be of strong glass tubing of about a quarter-inch bore, and for a stoppered funnel, which should have about half the capacity of the flask. The whole is supported by a clamp or on wire gauze. The outer jar shown in the figure should have a capacity of 2 or 3 litres, and the inner receiving-jar should be capable of holding 200 cc. The latter is fitted with a cork perforated for the delivery-tube, and perhaps for another tube containing fragments of glass moistened with acidulated water to prevent possible loss of ammonia, though the latter tube is practically found to be almost unnecessary. The addition of washing-bottle tubes is also recommended, as convenient for obtaining the distillate from the jar without dismounting the apparatus from time to time.

A few strips of clean zinc (*granulated* zinc recently cleansed with diluted acid is best) are boiled in a beaker with a 3 per cent. solution of copper sulphate, the operation being repeated with a fresh portion of solution until an adherent and fairly

thick coating of finely-divided copper is deposited. The pieces of metal are well washed and introduced into the flask, which is then half filled with pure water free from ammonia. To avoid transference, the flask itself may be used instead of the beaker. The funnel also of the apparatus is filled with pure water. Water is now placed around the inner receiver in the outer jar, and, the connections being sound, heat is applied with the view of freeing the apparatus itself from any trace of ammonia. When the contents of the flask are evaporated nearly to dryness, pure water is admitted from the funnel until the flask is again about half full (the funnel should be filled again at once), and the distillation carried on as before. This must be repeated until no further trace of ammonia is evolved, when the apparatus is ready for use. On each occasion that the apparatus is used it must be freed from ammonia in this way. A suitable quantity of the substance to be estimated is now introduced (in the case of potable waters the prepared solid residue from 100 cc.), and water added, if necessary, until the flask is half full. Heat is now applied and the operation conducted in the manner already described until ammonia ceases to come over—a point which usually occurs in the case of water-residues when the flask has been twice or thrice charged with water and the distillate is about 100 cc. The warm water from the upper part of the cooling-jar may be removed by a siphon or otherwise, cold water being introduced from time to time.

The ammonia being all evolved, disconnect the flask and receiver simultaneously (unless washing-bottle tubes are fitted), and treat the contents of the latter by the Nessler method described on p. 630.—Urea yields but traces of ammonia by this process, and neither the sulphates nor chlorides of the alkali-metals affect the result.—The method is only applicable to highly dilute solutions of nitrates, for with stronger solutions oxides of nitrogen are formed and escape.

Another process (Pelouze's, improved by Fresenius) consists in adding the nitrate to an acid solution of a ferrous salt of known strength, and, when reaction is complete, estimating the amount of ferrous salt unattacked by volumetric solution of red chromate or of permanganate. Three molecular weights of converted ferrous salt indicate one molecular weight of nitric acid. Regeneration of nitric or nitrous acids by aërial oxidation of the nitric oxide evolved is prevented either by a current of carbonic acid gas or by using a closed flask in which is a *Bunsen valve*. (*Vide* p. 652.)

SULPHIDES.

Process 1.—Soluble sulphides (H_2S , NaHS , *e. g.*) may be estimated volumetrically by adding to the aqueous liquid a measured excess of an alkaline solution of arsenic of known strength, neutralizing by hydrochloric acid, diluting to any given volume, filtering off the arsenum sulphide precipitated, taking a portion of the filtrate equal to a half or a third of the original volume, and, after neutralizing by acid sodium carbonate, estimating the residual arsenic by the standard iodine solution (*vide* p. 646). The process may be tried on a measured volume of sulphuretted hydrogen (the weight of which is easily calculated; 1 litre of hydrogen = 0.0896 gramme) absorbed by a strong solution of soda or potash.

Process 2.—Sulphur and sulphides may also be quantitatively analyzed by oxidizing to sulphuric acid and precipitating in the form of barium sulphate. A couple of décigrammes of a pure metallic sulphide may be decomposed by careful deflagration with a mixture of potassium chlorate and sodium carbonate, the product dissolved in water, acidulated with hydrochloric acid, solution of barium chloride added, and the precipitated barium sulphate washed and collected as described in connection with the estimation of barium (p. 665). Many sulphides may be oxidized in a flask by potassium chlorate and hydrochloric acid, and then precipitated by barium chloride. Experimental determinations may also be made on a weighed fragment of sulphur, about 0.1, cautiously fused with a little solid caustic alkali, and the product oxidized while hot by the slow addition of powdered potassium nitrate or chlorate, or, when cold, by treatment with potassium chlorate and hydrochloric acid, and subsequent precipitation by barium chloride.

Note.—Fusions performed by help of a gas-lamp must be carefully conducted, for any alkali that may creep over the side of a crucible will certainly absorb sulphurous acid from the products of combustion of the gas, and error result.

Process 3.—Soluble sulphides may also be treated with excess of an alkaline arsenite, arsenous sulphide then be precipitated by the addition of hydrochloric acid, and the precipitate collected and weighed with the usual precautions. (*Vide* p. 670.)

Weights of Equivalent Quantities of Sulphur and its Compounds.

Sulphur	S	32
Sulphuretted hydrogen . .	H_2S	34
Barium sulphate	BaSO_4	232.8
Arsenous sulphide	$(\text{As}_2\text{S}_3) \div 3$	82
Iron bisulphide	$(\text{FeS}_2) \div 2$	60
Lead sulphide	PbS	238.4

SULPHITES.

Sulphites are usually estimated volumetrically by a standard solution of iodine. (*Vide* p. 745.) Sulphites insoluble in water are diffused in that menstruum, hydrochloric acid added, and the iodine solution then dropped in.

If necessary, sulphites may be estimated gravimetrically by oxidation and precipitation as barium sulphate.

SULPHATES.

These salts are always precipitated and weighed as barium sulphate, the manipulations being identical with those performed in the determination of barium by means of sulphates. (*Vide* p. 664.) The purity of sodium sulphate (*Sodii Sulphas*, U. S. P.), and the presence of not more than a given amount of sulphuric acid in vinegar (*Acetum*, B. P.), are directed, in the British Pharmacopœia, to be ascertained by this process. 10 grains of sodium sulphate yield 7.23 of barium sulphate. 5 ounces of vinegar should yield not more than about $\frac{1}{3}$ of a gramme of barium sulphate.

Sulphates may be estimated volumetrically by a seminormal solution of pure barium chloride, $\text{BaCl}_2 \cdot 2\text{H}_2\text{O} = 244$.

The amount of free sulphuric acid or hydrochloric acid in vinegar, lemon-juice, lime-juice, etc. may also be ascertained volumetrically by adding a known quantity of standard solution of soda, evaporating to dryness, incinerating, dissolving in water, and by standard acid estimating the quantity of soda still remaining free. The soda lost indicates the amount of free mineral acid (Hehner). Thresh first estimates the chlorine in a sample of vinegar, then adds a known additional amount of chlorine, preferably in the form of barium chloride, evaporates, ignites, treats with water, adds sodium bicarbonate to remove excess of barium, filters, and again estimates the chlorine. A loss of 70.8 of chlorine (Cl_2) indicates 98 of free sulphuric acid (H_2SO_4).

The method of estimating free sulphuric, nitric, or hydrochloric acids proposed by Spence and Esilman is founded on their power of decolorizing a standard solution of ferric acetate.

Proportional Weights of Equivalent Quantities of Sulphates.


The sulphuric radical	SO_4	96
Sulphuric acid	H_2SO_4	98
Barium sulphate	BaSO_4	232.8

CARBONATES.

Carbonates are usually estimated by the loss in weight they undergo on the addition of a strong acid.

Process 1.—A small light flask is selected—of such a size that it can be conveniently weighed in a delicate balance. Two narrow glass tubes are fitted to the flask by a cork—the one straight, extending from about two or three centimètres above the cork to the bottom of the flask, the other cut off close to the cork on the inside and curved outward, so as to carry a thin drying-tube horizontally above the flask. (See Fig. 84.) The drying-tube is nearly filled with small pieces of calcium chloride, a plug of cotton-wool preventing escape of any fragments at either end, and is attached by a pierced cork to the free extremity of the curved tube of the flask. A weighed quantity of any pure soluble carbonate is placed in the flask, a little water added, a miniature test-tube containing sulphuric acid lowered into the flask by a thread and supported so that the acid may not flow out, the cork inserted, the outer end of the piece of the straight glass tubing closed by a cap or a fragment of cork, and the whole weighed. The apparatus is then inclined, so that the sulphuric acid and carbonate may slowly react; carbonic acid gas is evolved and escapes through the horizontal tube, any moisture being retained by the calcium chloride. When effervescence has ceased, the gas still remaining in the vessel is sucked out; this is accomplished by adapting a piece of india-rubber tubing to the end of the drying-tube, removing the small plug from the straight tube, and aspirating slowly with the mouth for a few minutes. If the heat produced by the action of the sulphuric acid and solution is considered insufficient to expel all the carbonic acid from the liquid, the plug is again inserted in the tube and the contents of the flask gently boiled for some seconds. When the apparatus is nearly cold more air is again drawn through it, and the whole finally weighed. The loss is due to carbonic acid gas (CO_2), from the weight of which that of any carbonate is ascertained by calculation. Carbonates insoluble in water may be attacked by hydrochloric instead of sulphuric acid; granulated mixtures of carbonates and powdered tartaric or citric acid, by enclosing the preparation in the inner tube and placing water in the flask, or *vice versâ*. The apparatus also may be modified in

Fig. 84.



Estimation of Carbonates.

many ways to suit the requirements, convenience, or practice of the operator.

Process 2.—Carbonates from which carbonic acid gas is evolved by heat may be estimated by the loss they experience on ignition.

Process 3.—Free carbonic acid gas may be absorbed by a solid stick of potash or strong alkaline solution, the loss in volume of the gas or mixture of gases indicating the amount originally present.

Weights of Equivalent Quantities of Carbonic Acid Gas and certain Carbonates.

Carbonic acid gas	CO ₂	44
Carbonic acid	H ₂ CO ₃	62
Anhydrous sodium carbonate .	Na ₂ CO ₃	106
Crystalline sodium carbonate .	Na ₂ CO ₃ , 10H ₂ O .	286
Anhydrous potassium carbonate	K ₂ CO ₃	138
Crystalline potassium carbonate	K ₂ CO ₃ + 16% Aq	164.285
Calcium carbonate	CaCO ₃	100

OXALATES.

Process 1.—The oxalic radical is usually precipitated in the form of calcium oxalate, and weighed as carbonate, the manipulations being identical with those observed in the estimation of calcium. (*Vide* p. 665.) The experiment may be performed on 0.3 or 0.4 of a gramme of pure oxalic acid, 126 parts of which should yield 100 of calcium carbonate.

Process 2.—Oxalates may also be determined by conversion of their acidulous radical into carbonic acid gas, and observation of the weight of the latter. The oxalate, water, and excess of black manganese oxide are placed in the carbonic-acid apparatus (p. 683), a tube containing sulphuric acid lowered into the flask, the whole weighed, and the operation completed as for carbonates. From the following equation it will be seen that every 88 parts of carbonic acid gas evolved indicates the presence of 126 parts of crystallized oxalic acid or an equivalent quantity of other oxalate:



The black manganese oxide used in this experiment must be free from carbonates. The amount of materials employed is regulated by the size of the vessels.

PHOSPHATES.

Process 1.—From phosphates dissolved in water the phosphoric radical may be precipitated and weighed in the form of

magnesium pyrophosphate, the details of manipulation being similar to those observed in estimating magnesium. (*Vide* p. 666.) $\frac{1}{2}$ a gramme or rather more of pure dry crystallized sodium phosphate may be employed in experimental determinations. The official ammonium phosphate (*Ammonii Phosphas*, B. P.) is quantitatively analyzed by this method. "If 20 grains of this salt be dissolved in water, and solution of magnesium ammonio-sulphate be added, a crystalline precipitate falls, which, when well washed upon a filter with solution of ammonia diluted with an equal volume of water, dried, and heated to redness, leaves 16.8 grains." $\frac{1}{2}$ a gramme, or less, is a more convenient quantity if the operations be conducted with care. *Solution of Ammonio-sulphate of Magnesium* (B. P.) is prepared by dissolving 2 parts of magnesium sulphate, 1 of ammonium chloride, and 1 of solution of ammonia (20.6 per cent. NH_4HO) in 18 or 20 of distilled water; such a solution is of considerable use if several phosphoric determinations are about to be made.

Process 2.—Free phosphoric acid is most readily determined as lead phosphate ($\text{Pb}_3\text{2PO}_4$). Of the official (B. P.) diluted phosphoric acid it is stated that "355 grains by weight (or 73.8 of the official 'concentrated' acid) poured upon 180 grains of lead oxide in fine powder leave, by evaporation, a residue (principally lead phosphate) which, after it has been heated to dull redness, weighs 215.5 grains." One-tenth of these quantities may be used for experimental purposes; 1 to 2 grammes will give good results. The lead oxide must be quite pure; it should be prepared by digesting red lead in warm diluted nitric acid, washing, drying, and heating the resulting puce-colored plumbic oxide in a covered porcelain crucible. The increase in weight obtained on evaporating a given amount of solution of phosphoric acid with a known weight of perfectly pure lead oxide (PbO) may be regarded as entirely due to phosphoric anhydride (P_2O_5); $3\text{PbO} + \text{P}_2\text{O}_5 = \text{Pb}_3\text{2PO}_4$, the actual reaction being $3\text{PbO} + 2\text{H}_3\text{PO}_4 = \text{Pb}_3\text{2PO}_4 + 3\text{H}_2\text{O}$. From these equations, and the atomic weights (*vide* Appendix or table on next page), the percentage of phosphoric acid (H_3PO_4) in any specimen of its solution may easily be calculated.

Process 3.—The Strength of Pure Solution of Phosphoric Acid.—This is ascertained by specific gravity and reference to tables.

Process 4.—Bone earth, "superphosphate," the *Calcis Phosphas*, and other forms of calcium phosphate known to be tolerably free from iron or aluminium may be estimated by treating about $\frac{1}{2}$ a gramme with hydrochloric acid somewhat diluted,

filtering if necessary, warming, precipitating with excess of ammonia, collecting the precipitate (Ca_3PO_4), washing, drying, igniting, and weighing. The calcium phosphate of pharmacy (*Calcii Phosphas*, B. P.), if pure, will in this process lose little or no weight.

Process 5.—*Insoluble phosphates* in ashes, manures, etc. are treated as follows: The weighed material (1.0 to 10.0 grammes) is digested in hydrochloric acid diluted with three or four times its bulk of water, filtered (insoluble matter and filter being thoroughly exhausted by water), ammonia added to the filtrate and washings, until, after stirring, a faint cloudy precipitate is perceptible, solution of oxalic acid dropped in until, after agitation for a few minutes, the opalescence is destroyed, ammonium oxalate next added, the whole warmed, calcium oxalate removed by filtration, and the filtrate concentrated if very dilute, the liquid treated with citric acid in such quantity that ammonia when added in excess gives a clear *lemon-yellow* solution (Warington), magnesian mixture poured in (as in Process 1), and the precipitate of ammonio-magnesian phosphate collected, washed, dried, and weighed, as already described in connection with the estimation of magnesium.

Relative Weights of Equivalent Quantities of Phosphoric Compounds.

Phosphoric acid	H_3PO_4	98
Magnesium pyrophosphate	$(\text{Mg}_2\text{P}_2\text{O}_7 = 222) \div 2 =$	111
Lead phosphate	$(\text{Pb}_3\text{P}_2\text{O}_4 = 809.2) \div 2 =$	404.6
Phosphoric anhydride	$(\text{P}_2\text{O}_5 = 142) \div 2 =$	71
Calcium phosphate	$(\text{Ca}_3\text{P}_2\text{O}_4 = 310) \div 2 =$	155
Calcium superphosphate	$(\text{CaH}_4\text{P}_2\text{O}_4 = 234) \div 2 =$	117

QUESTIONS AND EXERCISES.

What quantity of pure rock-salt is equivalent to 4.2 parts of silver chloride? *Ans.* 1.71.—State the percentage of real potassium iodide contained in a sample of which 8 parts yield 10.9 of silver iodide. *Ans.* 96.25.—What is the strength of a solution of hydrocyanic acid 10 parts of which, by weight, yield 0.9 of silver cyanide? *Ans.* 1.81 per cent.—How are nitrates quantitatively estimated?—By what processes may the strength of sulphides be determined?—How much real sodium sulphate is contained in a specimen 10 parts of which yield 14.2 of barium sulphate? *Ans.* 86.61 per cent.—Give details of the operations performed in the quantitative analysis of carbonates.—What amount of carbonic acid gas should be obtained from 10 parts of acid potassium carbonate (or bicarbonate)? *Ans.* 4.4 parts.—To what operation and what proportions of materials does the following equation refer?



Explain the lead process for the estimation of phosphoric acid in the official solution.—State the amount of calcium superphosphate equivalent to 7.6 parts of magnesium pyrophosphate. *Ans.* 8.01 parts.

SILICATES.

Silica (SiO_2) may be separated from alkaline silicates, or from silicates decomposable by hydrochloric acid, by digesting the substance in hydrochloric acid at a temperature of 70° to 80° C. until completely disintegrated, evaporating to dryness, heating in an air-bath, again moistening with acid, diluting with hot water, filtering, washing, drying, igniting, and weighing.

ESTIMATION OF WATER.

Water and other matters readily volatilized are most usually estimated by the loss in weight which a substance undergoes on being heated to a proper temperature. Thus, in the British Pharmacopœia, crystalline gallic acid ($\text{H}_3\text{C}_7\text{H}_3\text{O}_2, \text{H}_2\text{O}$) is stated to lose 9.5 per cent. of its weight at a temperature of 100° C., cerium oxalate ($\text{Ce}_2\text{3C}_2\text{O}_4, 9\text{H}_2\text{O}$) 52 per cent. on incineration, potassium carbonate about 16 per cent. on exposure to a red heat, quinine sulphate ($2\text{C}_{20}\text{H}_{24}\text{N}_2\text{O}_2, \text{H}_2\text{SO}_4, 7\text{H}_2\text{O}$) 15.2 per cent. at 100° C., sodium arsenate ($\text{Na}_2\text{HAsO}_4, 7\text{H}_2\text{O}$) 40.38 per cent. at 149° C., sodium carbonate ($\text{Na}_2\text{CO}_3, 10\text{H}_2\text{O}$) 62.94 per cent., sodium phosphate ($\text{Na}_2\text{HPO}_4, 12\text{H}_2\text{O}$) 63 per cent., and sodium sulphate ($\text{Na}_2\text{SO}_4, 10\text{H}_2\text{O}$) 55.9 per cent. at a low red heat: bismuth oxide heated to incipient redness is not to diminish in weight. Ferric oxyhydrate "yields about 10 per cent. of moisture."

Process.—1 or 2 grammes of substance is sufficient in experiments on desiccation, the material being placed in a watch-glass, covered or uncovered porcelain crucible, or other vessel, according to the temperature to which it is to be exposed.

Rapid desiccation at an exact temperature may be effected by introducing the substance into a tube having somewhat the shape of the letter U, sinking the lower part of the tube into a liquid kept at a definite temperature by aid of a thermometer, and drawing or forcing a current of dry air slowly through the apparatus. Substances liable to oxidation may be desiccated in a current of dried carbonic acid gas. The weights of the U-tube before and after the introduction of the salt, and after desiccation, give the amount of water sought. In all cases the material must be heated until it ceases to lose weight. Occasionally it is desirable to estimate water directly by conveying its vapor in a current of air through a weighed tube containing calcium chloride, and reweighing the tube at the close of the operation; the increase shows the amount of water.

Note.—Highly-dried substances rapidly absorb moisture from the air; they must therefore be weighed quickly, enclosed, if possible,

in tubes (p. 660), a pair of clamped watch-glasses, or a crucible having a tightly-fitting lid.

CARBON, HYDROGEN, OXYGEN, NITROGEN.

The quantitative analysis of animal and vegetable substances is either *proximate* or *ultimate*.

Proximate Quantitative Analysis includes the estimation of water, oil, albumen, starch, cellulose, gum, resin, alkaloids, acids, glucosides, ash. It requires the application of much theoretical knowledge and manipulative skill, and cannot well be studied except under the guidance of a tutor. One of the best works on the subject is by Rochleder, a translation of whose monographs will be found in the *Pharmaceutical Journal*, vol. i. 2d Ser., pp. 562, 610; vol. ii. 2d Ser., pp. 24, 129, 160, 215, 274, 420, 478. Another is by Prescott, "Outlines of Proximate Organic Analysis," Van Nostrand, New York. The fullest is by Dragendorff, translated by H. G. Greenish, "Plant Analysis."

Ultimate Quantitative Organic Analysis can only be successfully accomplished with the appliances of a well-appointed laboratory—a good balance, a gas-furnace (p. 674) giving a smokeless flame (seven or eight centimètres wide and seventy or eighty centimètres long), special forms of glass apparatus, etc. *The theory of the operation* is simple: A weighed quantity of a substance is burnt to carbonic acid gas ($\text{CO}_2 = 44$) and water ($\text{H}_2\text{O} = 18$), and these products collected and weighed; 12 parts in every 44 of carbonic acid gas ($= \frac{3}{11}$) are carbon, 2 in every 18 of water ($= \frac{1}{9}$) are hydrogen; nitrogen, if present, escapes as gas. If nitrogen be a constituent, a second quantity is strongly heated with a mixture of sodium and calcium hydrates; these bodies then split up into oxides, oxygen, and hydrogen; the oxygen burns the carbon of the substance to carbonic acid gas, its hydrogen and nitrogen appearing as water and ammonia respectively; the carbonic acid and water are disregarded, the ammonia collected and weighed in the form of the double platinum and ammonium chloride ($\text{PtCl}_4 \cdot 2\text{NH}_4\text{Cl} = 442.8$), of which 28 parts in every 442.8 are nitrogen. The difference between the sum of the weights of hydrogen and carbon and the weight of substance taken is the proportion of oxygen in the body, supposing nitrogen to be absent. If nitrogen is present, the difference between the sum of the percentages of carbon, hydrogen, and nitrogen and 100 is the percentage of oxygen. Shortly, carbon is estimated in the form of carbonic acid gas, hydrogen as water, nitrogen as ammonia, and oxygen by difference.

The following is the outline of the necessary manipulations: The source of the oxygen for the combustion of carbon and hydrogen is black copper oxide in coarse powder. 200 or 300 grammes of this material are heated in a crucible to low redness for a short time to expel every trace of moisture—then transferred to store-tubes (Fig. 85) resembling test-tubes, half a mètre long and having a slightly narrowed mouth, the tube


being held in a cloth to protect the hand while the hot oxide is being directly introduced into the mouth of the tube by a scooping motion. As soon as the well-corked tube is cool, the oxide is poured, portion by portion, into a similar tube (the

Fig. 85.



combustion-tube), but somewhat longer, drawn out to a quill (bent upward nearly to a right angle) at one end and not constricted at the mouth. Two such tubes are readily made by softening in the blowpipe-flame two or three centimètres of the central part of a tube about a mètre long, and drawing the halves of the tube apart as shown in the following engraving (Fig. 86). The tubes are separated by melting the glass in


Fig. 86.



the middle of the quilled portion. A few décigrammes of fused potassium chlorate should first be dropped into the tube. After 10 or 15 centimètres of oxide have been poured in, about a décigramme of the substance to be analyzed is dropped down the tube, then a few grammes of oxide, then another décigramme of substance, then more oxide, until 3 or 4 décigrammes of the body under examination have been added. The 15 or 20 centimètres of alternate layers are next thoroughly mixed by a long copper wire having a short helix: more oxide is introduced, the wire cleansed by twisting the helix about in the pure oxide, and a plug of dried asbestos finally placed on the top of the oxide at about five centimètres from the mouth of the tube; the tube is then securely corked and set aside. The substance operated on may be pure white sugar, powdered and dried; the tube in which it is contained is weighed before and after the removal of the portions for combustion; the loss is the quantity employed in the experiment. The combustion-furnace may be such as shown on p. 674. If the furnace is very powerful or the combustion-tube not of the hardest glass, the tube should be enclosed in wire-gauze the elasticity of which has been destroyed by heating to redness. If the substance under experiment contain nitrogen, the plug of asbestos

must be displaced by a roll of copper gauze, which serves to reduce any oxides of nitrogen and thus ensure the escape of nitrogen itself—or dry yellow potassium chromate may be used (Perkin). The *water* produced when the prepared tube is heated is collected in a small U-tube containing pieces of calcium chloride or pumice-stone moistened with sulphuric acid (Fig. 87); the *carbonic acid gas* in a series of bulbs (Fig. 87)

Fig. 87.



Calcium Chloride Tube and Potash-bulbs.

containing solution of potash (sp. gr. about 1.27). These bulbs may be purchased at any apparatus-shop. The calcium chloride tube is fitted by a good cork to the combustion-tube, the potash-bulbs by a short piece of india-rubber tubing to the calcium chloride tube. The potash-bulbs may carry a short light tube containing a rod of caustic potash three or four centimetres long: this serves to arrest any moisture that might be carried away from the solution of potash by the dried expanded air which escapes during the operation. The combustion-tube having been placed in the furnace, and the drying-tube and potash-bulbs weighed and attached, the gas is lit under the asbestos, and, when the tube is red hot, the flame slowly extended until nearly the whole tube is at the same temperature, the operation being conducted at such a rate that bubbles of gas escape through the potash-bulbs at about the rate of one per second. When no more gas passes, the extremity of the tube containing the potassium chlorate is gently heated until oxygen ceases to be evolved; perfect combustion of carbon and removal of all carbonic acid gas are thus ensured. The heating of the chlorate must be very carefully conducted, or the evolution of oxygen may become sufficiently rapid to blow some of the fluid out of the potash-bulbs. The drying-tube and bulbs are disconnected and weighed, the increase in weight due to carbonic acid gas and water respectively noted, and the percentages of carbon, hydrogen, and (by loss) oxygen calculated. This method is that of Liebig, with modifications by Bunsen: good combustion-furnaces are those known as Hofmann's, Griffin's, and others.

Lead chromate can be used for combustions in place of copper oxide. The advantages are its less hygroscopic nature and the greater readiness with which it yields its oxygen to organic bodies when heated with them. It must not, however, be used with bodies containing nitrogen, since it would convert so large a proportion of the nitrogen into nitric oxide or higher oxide of nitrogen that it would be necessary to use an inconveniently long layer of the metallic copper to reduce these oxides, and so prevent their absorption in the series of bulbs containing the solution of potash. Organic bodies, however, containing sulphur, bromine, iodine, or chlorine are burnt with advantage by means of lead chromate. If copper oxide were used with bodies containing sulphur, it would be necessary to place an additional tube containing lead peroxide between the calcium chloride tube and the potash-bulbs in order to absorb the sulphurous anhydride formed; this is entirely obviated by using lead chromate, which itself retains the whole of the sulphur. Again, if bodies containing chlorine, iodine, or bromine are burnt by means of copper oxide, then volatile chloride, iodide, or copper bromide is formed, and, collecting in the calcium chloride tube, vitiates the result with regard to the hydrogen; by using lead chromate, however, the chlorine, iodine, and bromine are respectively retained in the combustion-tube as lead chloride, bromide, and iodide.

In order to render the chromate fit for use it is first fused and poured out on a clean iron plate; when cool, it is powdered and heated in a long tube throughout its whole length, while air, dried by passing through calcium chloride or strong sulphuric acid, is drawn over it; when the color of the chromate changes to brown, the heat can be withdrawn and the extremity of the tube farthest from the drying-apparatus closed, so that the air in passing into the tube on cooling may be quite dry; when cool, the drying-tube is removed, and the extremity securely corked; the lead chromate is then ready for direct transference to the combustion-tube.

The general manipulations for substances containing nitrogen resemble the foregoing so far as the use of a combustion-tube and furnace and collection of the ammoniacal gas are concerned. The combustion-tube must be quilled at one end and about a third of a mètre long. The soda-lime is made by slaking quicklime with a solution of soda of such a strength that about 2 parts of quicklime shall be mixed with 1 of sodium hydrate, drying the product, heating to bright redness, and finely powdering; it should be preserved in a well-closed bottle. Some of the soda-lime is introduced into the tube, then layers of sub-

stance and soda-lime, mixture effected by a wire, a good layer of soda-lime added, and a plug of asbestos. Bulbs (Fig. 88),

Fig. 88.



Nitrogen-bulbs.

known as those of Will and Varrentrapp (the originators of the method), containing hydrochloric acid of about 25 per cent., are then fitted by a cork, and the tube heated in the furnace—to a not too bright red heat, or some of the produced ammonia gas may be decomposed. When gas ceases to pass

and combustion is considered to be *quite* complete, the tube is allowed to cool somewhat, the quill is then broken, and aspiration continued slowly until all ammoniacal gas may be considered to have been absorbed by the acid. The bulbs are disconnected, their contents and rinsings poured into a small dish, solution of platinum perchloride added, and the operation completed as in the estimation of potassium and ammonium salts. (*Vide* pp. 661 and 663.) Or the ammonia may be absorbed by a known quantity of standard sulphuric acid, of which the residual excess is estimated by a standard alkali; certain obvious calculations then giving the amount of ammonia produced.

Conversion into ammonia may also be effected by heating the substance with strong sulphuric acid, and, if not then thoroughly attacked, with potassium permanganate (Kjeldahl). Or the nitrogen may be evolved as gas by heating with copper oxide and copper turnings (p. 690), and be collected over alkali, and its volume, and thus indirectly its weight, be determined.

Liquids are analyzed by a similar method to that adopted for solids, volatile liquids being enclosed in small bulbs having a long quill. These are weighed previously to and after the introduction of the liquid; just before being dropped into the combustion-tube the quill is broken. *Solids* are also sometimes similarly burnt from a small boat placed in the tube, a continuous current of purified air being used.

Formulae.—From the percentage composition of an organic substance an empirical formula may be deduced by dividing the weight of each constituent by its atomic weight, and converting the product into the simplest whole numbers; a rational formula by ascertaining the proportion in which the substance unites with a radical or body having a known combining proportion, etc. (*Vide* pp. 386–389.)

Chlorine, Bromine, or Iodine contained in an organic substance may be estimated by heating with fuming nitric acid and silver

nitrate in a sealed tube, or by heating to redness a given weight of the material with ten times as much pure lime in a combustion-tube. Calcium chloride, bromide, or iodide is thus produced. While still hot the tube is plunged into water, the mixture of broken glass and powder treated with pure diluted nitric acid in very slight excess; the filtered liquid precipitated by silver nitrate, and the silver chloride, bromide, or iodide collected, washed, dried, cooled, and weighed.

Sulphur, Phosphorus, and Arsenum in organic salts may be estimated by heating with fuming nitric acid in a sealed tube, or by gradually heating in a combustion-tube 1 part of the substance with a mixture of 10 parts nitre, 2 dried sodium carbonate (in order to moderate deflagration), and 20 sodium chloride. The product is dissolved in water acidulated by nitric acid, the sulphuric radical precipitated and estimated as barium sulphate, the phosphoric and arsenic radicals as ammonio-magnesian phosphate or arsenate.

Limit of Experimental Errors.—Two determinations of carbon may vary to the extent of 0.1 per cent.; of hydrogen, 0.2; of nitrogen, 0.3.

QUININE, ETC.

Process of the British Pharmacopœia for ascertaining the amount of

(1) *Quinine with Cinchonidine*, and (2) *Total Alkaloids in the Succirubra or Red Cinchona Bark* (*Cinchonæ Rubræ Cortex*, B. P.).

1. *For Quinine and Cinchonidine.*—Mix 200 grains of red cinchona-bark, in No. 60 powder, with 60 grains of calcium hydrate; slightly moisten the powder with $\frac{1}{2}$ an ounce of water; mix the whole intimately in a small porcelain dish or mortar; allow the mixture to stand for an hour or two, when it will present the characters of a moist, dark-brown powder, in which there should be no lumps or visible white particles. Transfer this powder to a 6-ounce flask, add 3 fluidounces of a mixture of three volumes of commercial benzol and one volume of amylic alcohol ("benzolated amylic alcohol," B. P.); boil them together for about half an hour, decant and drain off the liquid on to a filter, leaving the powder in the flask; add more of the benzol liquid to the powder, and boil and decant as before; repeat this operation a third time; then turn the contents of the flask on to the filter and wash by percolation with more of the liquid until the bark is exhausted. If, during the boiling, a funnel be placed in the mouth of the flask, and another flask filled with cold water be placed in the funnel, this will form a convenient condenser which will prevent the loss of more than a small quantity of the boiling liquid. Introduce the collected filtrate, while still warm, into a stoppered glass separator; add to it 20 minims of diluted hydrochloric acid, mixed with 2 fluidrachms of water; shake them well together, and when the acid liquid has separated, this may be drawn off, and the process repeated with distilled water slightly acidulated with hydrochloric acid until the whole of the alka-

loids have been removed. The acid liquid thus obtained will contain the alkaloids as hydrochlorates, with excess of hydrochloric acid. It is to be carefully and exactly neutralized with ammonia, and then concentrated to the bulk of 3 fluidrachms. If now about 15 grains of tartarated soda, dissolved in twice its weight of water, be added to the neutral hydrochlorates, and the mixture stirred with a glass rod, insoluble quinine and cinchonidine tartrates will separate completely in about an hour; and these, collected on a filter, washed, and dried, will contain eight-tenths of their weight of the alkaloids quinine and cinchonidine, which, divided by 2, represents the percentage of those alkaloids. The other alkaloids will be left in the mother-liquor.

2. *For Total Alkaloids.*—To the mother-liquor from the preceding process add solution of ammonia in slight excess. Collect, wash, and dry the precipitate, which will contain the other alkaloids. The weight of this precipitate divided by 2, and added to the percentage weight of the quinine and cinchonidine, gives the percentage of total alkaloids.

Note.—If it is desired to obtain each alkaloid separately, the above process may be modified by using dilute sulphuric instead of hydrochloric acid for removing the alkaloids from the benzolated amylic alcohol, and exactly neutralizing with ammonia *while kept hot on a water-bath*; on cooling, the quinine will crystallize out almost completely as neutral sulphate. The cinchonidine may then be precipitated from the filtrate by tartarated soda, and, after its removal, quinidine, if present, by potassium iodide; finally, cinchonine is obtained by precipitating the filtrate from the quinidine with soda or ammonia.

Extractum Cinchonæ Liquidum, B. P.—The greater portion of the alkaloids of red cinchona-bark is dissolved out by water acidulated by hydrochloric acid and containing glycerin. The acid in acting as a solvent probably more or less decomposes the natural compounds of the bark. The glycerin contributes to the permanence of the preparation. The mixture is evaporated to a low bulk, assayed as follows, and further evaporated or diluted until 85 fluid parts contain 5 of total alkaloids. 12½ fluid parts of rectified spirit are then mixed in, and a little water added to make 100 fluid parts. The *assay* is thus conducted: Place 50 fluid grains of the product of the first evaporation (*a*) with ½ an ounce of distilled water in a stoppered glass separator capable of holding 4 fluidounces; add to this 1 fluidounce of benzolated amylic alcohol and ½ a fluidounce of solution of soda; shake them together thoroughly and repeatedly, then allow them to remain at rest until the spirituous solution of the alkaloids shall have separated and formed a distinct stratum over the dark-colored alkaline solution of the other constituents of the extract. Run off the latter by the stopcock, add a little more distilled water to wash away any

adhering alkaline solution from the separator and its contents, and, having run off this as before as completely as possible, decant the spirituous solution into a small porcelain or glass dish the weight of which is known. Evaporate by the heat of a water-bath until a perfectly dry residue is left. The weight now of the dish and its contents, after deducting the known weight of the dish, will give that of the alkaloids, and this multiplied by 2 will give the parts by weight of the alkaloids in 100 fluid parts of the liquid (*a*).

De Vrij's Method for the Separation of the Mixed Alkaloids from Cinchona Barks; De Vrij's Method for the Separation and Quantitative Determination of all the different Cinchona Alkaloids. For these processes the reader is referred to the eleventh edition of this Manual. *Prollius's Method for the Estimation of Total Alkaloids in Cinchona Bark as modified by De Vrij.*—See the thirteenth edition.

Official (B. P.) Methods of Testing Quinine Sulphate for Sulphates of other Alkaloids.

Test for Cinchonidine and Cinchonine.—Heat 100 grains of quinine sulphate in 5 or 6 ounces of boiling water, with 3 or 4 drops of diluted sulphuric acid. [Paul recommends omission of all acid, except what may be required to ensure perfect neutralization.] Set the solution aside until cold. Separate, by filtration, the purified quinine sulphate which has crystallized out. [Evaporate to one-fifth.] To the filtrate, which should nearly fill a bottle or flask, add ether, shaking occasionally, until a distinct layer of ether remains undissolved. Add ammonia in very slight excess, and shake thoroughly, so that the quinine at first precipitated shall be redissolved. Set aside for some hours or during a night. Remove the supernatant clear ethereal fluid, which should occupy the neck of the vessel, by a pipette. Wash the residual aqueous fluid and any separated crystals of alkaloid with a very little more ether, once or twice. Collect the separated alkaloid on a tared filter, wash it with a little ether, dry at 212° F. (100° C.), and weigh. 4 parts of such alkaloid correspond to 5 parts of crystallized cinchonidine sulphate or of cinchonine sulphate.

Test for Quinidine.—Recrystallize 50 grains of the original quinine sulphate as described in the previous paragraph. To the filtrate add solution of potassium iodide, and a little spirit of wine to prevent the precipitation of amorphous hydriodates. Collect any separated quinidine hydriodate, wash with a little water, dry, and weigh. The weight represents about an equal weight of crystallized quinidine sulphate.

Test for Cupreine.—Shake the recrystallized quinine sulphate, obtained in testing the original quinine sulphate for cinchonidine and cinchonine, with 1 fluidounce of ether and a $\frac{1}{4}$ of an ounce of solution of ammonia, and to this ethereal solution, separated, add the

etheral fluid and washings also obtained in testing the original sulphate for the two alkaloids just mentioned. Shake this ethereal liquor with a $\frac{1}{4}$ of a fluidounce of a 10 per cent. solution of caustic soda, adding water if any solid matter separates. Remove the ethereal solution. Wash the aqueous solution with more ether, and remove the ethereal washings. Add diluted sulphuric acid to the aqueous fluid, heated to boiling, until the soda is exactly neutralized. When cold collect any cupreine sulphate that has crystallized out on a tared filter; dry, and weigh.

"Quinine sulphate" should not contain much more than 5 per cent. of sulphates of other cinchona alkaloids.

Quinine sulphate normally contains 15.2 per cent. of water; cinchonidine sulphate, 7.0 per cent.; cinchonine sulphate 6.0 per cent.,—all given off at 100° to 115° C. The drying should therefore be effected at 120° C., and the dried salt weighed in well-fitting weighing tubes. 100 parts of cinchonidine are equivalent to 116 parts of anhydrous cinchonidine sulphate.

Cinchonidine sulphate is almost the only salt likely to be accidentally present in the quinine sulphate of trade, much quinidine being rarely present in bark, and cinchonine sulphate being sufficiently soluble to remain dissolved in the mother-liquors of quinine sulphate. Cinchonidine sulphate may vary from 1 to 18 per cent., but more usually is present to the extent of about 6 per cent.

Quinina, U. S. P.—"If 2 grms. of quinine be mixed, in a small mortar, with 1 grm. of ammonium sulphate and 10 cc. of distilled water, the mixture thoroughly dried on a water bath, the residue (which should be strictly neutral to test-paper) agitated with 20 cc. of water, then allowed to macerate for half an hour at 15° C. (59° F.), with occasional agitation, and filtered through a pellet of glass-wool, 5 cc. of the filtrate, transferred to a test-tube, and gently mixed, without shaking, with 7 cc. of ammonia-water (specific gravity 0.960), should produce a clear liquid. If the temperature during the maceration has been 16° C. (60.8° F.), 7.5 cc. of ammonia-water may be added; if 17° C. (62.8° F.), 8 cc. (In each case a clear liquid indicates the absence of more than small proportions of other cinchona alkaloids.)"

Quininæ Sulphas, U. S. P.—"If 2 grms. of the salt (which must have been previously ascertained to be strictly neutral to litmus-paper, or have been rendered so) be dried, as far as possible, at 100° C. (212° F.), the residue then agitated with 20 cc. of water, and the mixture macerated for half an hour at 15° C. (59° F.), with occasional agitation, upon proceeding further as directed above for Quinine the results there given should be obtained."

De Vrij's Chromate Test for the Purity of Quinine Sulphate.—The purity of quinine sulphate may be tested in the following manner (De Vrij):

Dissolve 1 grm. of the salt in 40 cc. of hot distilled water, and add 6 cc. of a 5 per cent. solution of pure yellow potassium chromate. Set aside for several hours at a temperature *not above* 15° C., when the quinine will be completely precipitated as chromate. Filter, and after adding a few more drops of the chromate to the filtrate, to

make sure that the precipitation has been completed, add 9 or 10 drops of a 5 per cent. solution of soda. If the quinine sulphate was pure, the solution will remain clear, even after a day's standing; but if cinchonidine was present, that alkaloid will be precipitated, and may be filtered off and weighed.

Of the Citrate of Iron and Quinine (Ferri et Quininæ Citras, U. S. P. and B. P.) it is officially (B. P.) stated that "50 grains dissolved in a fluidounce of water and treated with a slight excess of ammonia give a white precipitate, which, when dissolved by successive treatments of the fluid with ether or chloroform, and the latter evaporated and the residue dried (at about 250° F.) until it ceases to lose weight, weighs seven and a half grains. The precipitate is almost entirely soluble in a little pure ether." Metric weights may be used. The British preparation should thus yield 15 per cent., and that of the United States at least 11.5 per cent., of quinine.

A Process for the Determination of the Quinine of the Scaled Compound.—To the residue obtained as stated in the foregoing paragraph is added about 25 cc. of water and enough diluted sulphuric acid to impart a decidedly acid reaction. The mixture is next heated over a water-bath until, the solution remaining acid, the residue has completely dissolved. Dilute soda solution is afterward added with great care until the solution is exactly neutral. The dish is then removed, and the solution allowed to cool and rest over night, when the quinine will have separated in crystals of ordinary sulphate. These should be collected on a filter and the mother-liquor tested with litmus-paper. If it is acid, it must be warmed over a water-bath and dilute soda solution added to exact neutralization, and the solution set aside as before, when some more crystals will probably separate. These are also collected, and with the former ones washed, dried at 120° C., and weighed [$(C_{20}H_{24}N_2O_2)_2 \cdot H_2SO_4 = 746$]. To this weight must be added 1 grm. for 750 cc. of mother-liquor for quinine sulphate which it retains. From this weight of anhydrous quinine sulphate is calculated its equivalent of hydrous quinine ($C_{20}H_{24}N_2O_2)_2 \cdot 2H_2O = 684$, the approximate formula of hydrous quinine dried over a water-bath. The weight thus obtained is compared with the weight of total alkaloid determined, both having been reduced to percentages. The amount of hydrous quinine calculated from the crystals of sulphate should not be much below that weighed directly. In good specimens the difference will be about 1 per cent.

Quinine sulphate should yield, according to De Vrij, at least 91.6 per cent. of quinine tartrate.

(See a paper by Fletcher in the *Pharmaceutical Journal* for September 20, 1869; also in that for September 18, 1880, by De Vrij; and in the *Chemist and Druggist* for October, 1885, by Howard.)

MORPHINE.

The official (U. S. P.) process for the assay of opium is conducted in the following manner: Opium, in any condition to be valued, 10 grms.; ammonia-water, 3.5 cc.; alcohol, ether,

water, each a *sufficient quantity*. "Introduce the opium (which, if fresh, should be in very small pieces, and if dry in very fine powder) into a bottle having a capacity of about 300 cc., add 100 cc. of water, cork it well, and agitate frequently during twelve hours. Then pour the whole as evenly as possible upon a wetted filter having a diameter of 12 cm., and, when the liquid has drained off, wash the residue with water, carefully dropped upon the edges of the filter and the contents, until 150 cc. of filtrate are obtained. Then carefully transfer the moist opium back to the bottle by means of a spatula, add 50 cc. of water, agitate thoroughly and repeatedly during fifteen minutes, and return the whole to the filter. When the liquid has drained off wash the residue as before, until the second filtrate measures 150 cc., and finally collect about 20 cc. more of a third filtrate. Evaporate in a tared capsule, first, the second filtrate to a small volume, then add the first filtrate, rinsing the vessel with the third filtrate, and continue the evaporation until the residue weighs 14 grms. Rotate the concentrated solution about in the capsule until the rings of extract are redissolved, pour the liquid into a tared Erlenmeyer flask having a capacity of about 100 cc., and rinse the capsule with a few drops of water at a time, until the entire solution weighs 20 grms. Then add 10 grms. (or 12.2 cc.) of alcohol, shake well, add 25 cc. of ether, and shake again. Now add the ammonia-water from a graduated pipette or burette, stopper the flask with a sound cork, shake it thoroughly during ten minutes, and then set it aside, in a moderately cool place, for at least six hours or over night."

"Remove the stopper carefully, and, should any crystals adhere to it, brush them into the flask. Place in a small funnel two rapidly-acting filters, of a diameter of 7 cm., plainly folded, one within the other (the triple fold of the inner filter being laid against the single side of the outer filter), wet them well with ether, and decant the ethereal solution as completely as possible upon the inner filter. Add 10 cc. of ether to the contents of the flask, rotate it, and again decant the ethereal layer upon the inner filter. Repeat this operation with another portion of 10 cc. of ether. Then pour into the filter the liquid in the flask, in portions, in such a way as to transfer the greater portion of the crystals to the filter, and, when this has passed through, transfer the remaining crystals to the filter by washing the flask with several portions of water, using not more than about 10 cc. in all. Allow the double filter to drain, then apply water to the crystals, drop by drop, until they are practically free from mother-water, and

afterward wash them, drop by drop from a pipette, with alcohol previously saturated with powdered morphine. When this has passed through displace the remaining alcohol by ether, using about 10 cc. or more if necessary. Allow the filter to dry in a moderately warm place at a temperature not exceeding 60° C. (140° F.), until its weight remains constant, then carefully transfer the crystals to a tared watch-glass and weigh them. The weight found, multiplied by 10, represents the percentage of crystallized morphine obtained from the opium."

Teschemacher and Smith's Method.—Thoroughly exhaust 200 grains of opium with warm distilled water. Concentrate this watery extract to a thin syrup in a shallow dish over a water-bath, the water of which should not quite boil. Transfer this thin syrup to a suitable flask, which permits the use of a soft cork, using a few drops of water successively to wash out the dish. Add to the contents of the flask 50 fluid grains of alcohol, sp. gr. about 0.820, and about 600 fluid grains of ether. Mix gently, but thoroughly, and then add some 50 fluid grains of ammonia, sp. gr. 0.935. Shake the contents of the flask well to precipitate the alkaloids in arenaceous crystals, with occasional agitation during the ensuing eighteen hours. Transfer the contents of the flask to a vacuum filter, and permit all the adherent liquid to be drawn away, washing out the flask with morphinated spirit,* and continue its use till the liquid passes colorless. Then wash with morphinated water† till this also passes colorless. Now dry, slowly at first, finishing at 212° F. Transfer the dried substance to a mortar, reduce it to a very fine powder, and digest it thoroughly in benzene to dissolve the narcotine and such of the opium alkaloids, other than morphine, which may be present. Transfer this mixture to a vacuum filter, wash out the mortar carefully with benzene, which use to wash the powder thoroughly. This, then, will be morphine, free from other opium alkaloids and narcotine, but still containing coloring and possibly other organic matters to the extent of 3 to 10 per cent. Dry and weigh this powder. Now ascertain the percentage by weight of crystallized morphine by titration of this powder with standard hydrochloric acid and litmus as the indicator. (This acid is so made that 1000 grains by weight shall exactly neutralize 100 grains of pure morphine crystallized from water, washed with ether, and gently dried finally at 212° F.)

Of Morphine Hydrochlorate it is officially (in the B. P.) stated that "20 grains of the salt, dissolved in $\frac{1}{2}$ an ounce of warm water, with ammonia added in the slightest possible excess, give on cooling a crystalline precipitate which, when washed with a little cold water and dried in a water-bath, weighs 16 grains;" and respecting the *Injectio Morphine Hypodermica*, B. P., "a fluidrachm of it, rendered slightly alkaline by the addition of solution of ammonia, yields a precipitate of

* *Morphinated Spirit.*—Digest a large excess of morphine in rectified spirit of 80 per cent. for several days, with frequent agitation. Filter for use.

† *Morphinated Water.*—As above, substituting distilled water for spirit.

morphine which, after being washed and dried, should weigh 4.25 grains, corresponding to 6 grains of acetate of morphine." The acetate (*Morphinæ Acetas*, U. S. P.) is liable to lose acetic acid and become basic; hence in the British Pharmacopœia the official morphine acetate must conform to the following requirements: "20 grains of the salt form with 1 drachm of water a slightly turbid solution, which is rendered clear by the addition of 1 grain of acetic acid; and this solution, when mixed with ammonia in slight excess, yields a precipitate which, after washing with a little cold water and drying in a water-bath, weighs 15 grains. If the salt yields a larger proportion of morphine than this, it should be recrystallized from hot water acidulated with acetic acid."

NUX-VOMICA ALKALOIDS.

The British Pharmacopœia directs that its two galenical preparations of nux vomica (*Extractum Nucis Vomicae* and *Tinctura Nucis Vomicae*) shall contain defined proportions of the mixed alkaloids of the drug.—The extract is made by exhausting a pound of powdered nux vomica with a mixture of 64 fluidounces of rectified spirit and 16 of water by percolation. 1 ounce of the percolate is treated as follows: Evaporate almost to dryness over a water-bath, to dispel spirit; dissolve the residue in $\frac{1}{2}$ a fluidounce of diluted sulphuric acid, with an equal bulk of water and add 2 fluidrachms of chloroform; agitate and warm gently. When the liquors have separated, draw off and reject the chloroform; then add to the acid liquor excess of solution of ammonia and $\frac{1}{2}$ a fluidounce of chloroform; well agitate, gently warm, and, after the liquors have completely separated, transfer the chloroform to a weighed dish, evaporate over a water-bath, and dry for one hour at 212° F. (100° C.). Allow the residue of total alkaloid to cool, and then weigh it. Take of the percolated liquid as much as shall contain 131 $\frac{1}{4}$ grains of total alkaloid, distil off the spirit, and evaporate over a water-bath until the extract weighs two ounces. This extract will contain 15 per cent. of total alkaloid.—The tincture is prepared by dissolving 133 grains of the assayed extract in a pint of diluted spirit (made by adding enough rectified spirit to 4 ounces of water to produce a pint of fluid). 1 fluidounce of this tincture will contain 1 grain of the alkaloids of nux vomica.—If properly prepared, 10 grains of the extract when treated in the following manner should yield 1 $\frac{1}{2}$ grains of total alkaloids. Dissolve the extract in $\frac{1}{2}$ a fluidounce of water with the aid of a gentle heat, and add a drachm of sodium carbonate previously dissolved in $\frac{1}{2}$ a fluidounce of water and $\frac{1}{2}$ a fluidounce of chloroform; agitate, warm gently, and sep-

arate the chloroform. To the separated chloroform add $\frac{1}{2}$ a fluidounce of diluted sulphuric acid with an equal bulk of water; again agitate, warm, and separate the acid liquid from the chloroform. To this acid liquor add now an excess of ammonia, and agitate with $\frac{1}{2}$ a fluidounce of chloroform; when the liquors have separated, transfer the chloroform to a weighed dish and evaporate the chloroform over a water-bath. Dry the residue for one hour, and weigh.

SUGAR.

The qualitative test for sugar by means of an alkaline copper solution (*vide* p. 468) may be applied in the estimation of sugar in sacchariferous substances.

Process 1.—34.65 grammes of pure dry crystals of ordinary copper sulphate are dissolved in about 250 cc. of distilled water. 173 grammes of pure crystals of the double potassium and sodium tartrate are dissolved in 480 cc. of solution of caustic soda of sp. gr. 1.14. The solutions are *only mixed when required*, water being then added to form 1 litre, smaller quantities of the fluids being proportionately diluted. 100 cc. of this mixture represent 3.465 grammes of copper sulphate, and correspond to 0.500 of a gramme of pure anhydrous grape-sugar, 0.475 of cane-sugar, 0.807 of maltose, or 0.450 of starch. The solutions must be preserved in well-stoppered bottles to prevent absorption of carbonic acid, and be kept in a dark place. Should the mixture give a precipitate on boiling, a little solution of soda may be added when making experiments. Such a reagent is known as *Fehling's solution*.

Dissolve 0.475 of pure dry powdered cane-sugar in about 50 cc. of water, convert into grape-sugar by acidulating with sulphuric acid and heating for an hour or two on a water-bath, make slightly alkaline with sodium carbonate, and dilute to 100 cc. Place 10 cc. of the copper solution in a small flask, dilute with three or four times its bulk of water, and gently boil. Into the boiling liquid drop the solution of sugar from a burette, 1 cc., or less, at a time, until, after standing for the precipitate to subside, the supernatant liquid has just lost its blue color; 10 cc. of the solution of sugar should be required to produce this effect—equivalent to 0.0475 of cane-sugar, 0.0807 of maltose, or 0.0500 of grape-sugar. Experiments on pure cane-sugar must be practised until accuracy is attained; syrups, diabetic urine, and saccharated substances containing unknown quantities of sugar may then be analyzed.

Starch is converted into grape-sugar by gentle ebullition with diluted acid for eight or ten hours, the solution being finally

diluted so that 1 part of starch, or rather sugar, shall be contained in about 150 of water.

If, instead of Fehling's solution, Pavy's ammoniated solution be used (*Proceedings of the Royal Society of London*, vol. xxviii. p. 260, and vol. xxix. p. 272; or *Lancet*, March 1, 1884, p. 376; or *Pharmaceutical Journal*, 3d Ser., vol. xvii. p. 856), one-fifth more of the copper salt will be required to do the same amount of work.

In cases in which loss of blue color cannot be relied on as indicating the termination of the reaction, copper suboxide should be rapidly filtered out, washed, dried, and, after adding the filter-ash, ignited, and the resulting black copper oxide weighed. 1 gram. of black oxide (or of suboxide or of metallic copper) indicates the subjoined amounts of the respective sugars:

One gramme of	Glucose.	Cane-sugar.	Milk-sugar.	Malt-sugar.
Black oxide . . .	.4535	.4308	.6153	.7314
Suboxide . . .	.5042	.4790	.6843	.8132
Metallic copper . .	.5634	.5395	.7707	.9089

Process 2.—Robert's Method for the Estimation of Sugar in Urine.—About 4 ounces of saccharine urine are put into a 12-ounce bottle, and a lump of German yeast about the size of a cob-nut or small walnut is added. This excess of yeast hastens fermentation and does no harm. The bottle is then covered with a grooved cork (to allow of the escape of carbonic acid gas), and set aside in a warm place to ferment. By the side of it is placed a tightly-corked 4-ounce phial filled with the same urine without any yeast. In about twenty-four hours the fermentation will have ceased and the scum cleared off or subsided. The fermented urine is then decanted and its specific gravity taken. At the same time the specific gravity of the unfermented urine in the companion phial is observed. The *density lost* is thus ascertained. Each degree of density lost represents a grain of glucose per fluidounce.

Sugar is often estimated by the measurement of the carbonic acid gas evolved or of the alcohol produced during fermentation.

Saccharometry.—A generic term for certain quantitative operations undertaken with the view of ascertaining the quantity of sugar present in any matter in which it may be contained.

Saccharometry is frequently performed upon common syrup (*Syrupus*, B. P.) and solutions which are known to contain nothing but cane- (ordinary) sugar, the object being merely to

ascertain the amount present. In such a case it is only necessary to take the specific gravity of the liquid at 60° F., and then refer to a previously prepared table of densities and percentages.

Specific gravity.	Sugar, per cent.	Specific gravity.	Sugar, per cent.	Specific gravity.	Sugar, per cent.
1.007	1.8	1.100	23.7	1.210	46.2
1.014	3.5	1.108	25.6	1.221	48.1
1.022	5.2	1.116	27.6	1.231	50.0
1.029	7.0	1.125	29.4	1.242	52.1
1.036	8.7	1.134	31.5	1.252	54.1
1.044	10.4	1.143	33.4	1.261	56.0
1.052	12.4	1.152	35.2	1.275	58.0
1.060	14.4	1.161	37.0	1.286	60.1
1.067	16.3	1.171	38.8	1.298	62.2
1.075	18.2	1.180	40.6	1.309	64.4
1.083	20.8	1.190	42.4	1.321	66.6 ?
1.091	21.8	1.199	44.3	1.330 (B.P.)	66.6 ?

The spec. grav. may be taken by a hydrometer, technically termed a *saccharometer*. (The above spec. gravs. = 1° to 35° Baumé.)

If a liquid contains other substances besides cane-sugar, the test of specific gravity is of little or no value. Advantage may then be taken of the fact that syrup causes right-handed twisting of a ray of plane-polarized light to an extent proportionate to the amount of sugar in solution. The saccharine fluid is placed in a long tube having opaque sides and transparent ends; and a ray of homogeneous light, polarized by reflection from a black-glass mirror or otherwise, is sent through the liquid and optically examined by a plate of tourmaline, Nicol's prism, or other polarizing eyepiece. Attached to the eyepiece is a short arm which traverses a circle divided into degrees. The eyepiece and arm are previously so adjusted that when the ray is no longer visible the arm points to the zero of the scale of degrees. The saccharine solution, however, so twists the ray as to again render it visible; and the number of degrees which the eyepiece has to be rotated before the ray is once more invisible is proportionate to the strength of the solution. The value of the degrees having been ascertained by direct experiment and the results tabulated, a reference to the table indicates the percentage of sugar in the liquid under examination. Grape-sugar also possesses the property of dextral rotation, but less powerfully than cane-sugar; moreover, the former variety does not, like cane-sugar, suffer inversion of the direction of rotation on the addition of hydrochloric acid to its solution—an operation that furnishes data for ascer-

taining the amount of cane- and of grape-sugar, or of crystallizable and non-crystallizable sugar, present in a mixture. In using the polariscope saccharometer it is convenient to employ tubes of uniform size and always to operate at the same temperature. Various modes are adopted of applying for quantitative purposes this action of syrup on polarized light.

ALCOHOL.

Mulder's process for the determination of the amount of alcohol in wine, beers, tinctures, and other alcoholic liquids containing vegetable matter is as follows: Take the specific gravity and temperature of the liquid and measure off a certain quantity (100 cc.); evaporate to one-half or less, avoiding ebullition in order that particles of the material may not be carried away by the steam. Dilute with water to the original bulk, and take the specific gravity at the same temperature as before. Of the figures representing the latter specific gravity, all over 1000 show to what extent dissolved solid matter affected the original specific gravity of the liquid. Thus, the specific gravity of a sample of wine at 15.5° C. is 0.9951; evaporated till all alcohol is removed and diluted with water to the original bulk, the specific gravity at 15.5° C. is 1.0081. 0.0081 represents the gravitating effect of dissolved solid matter in 0.9951 part of original wine. 0.0081 subtracted from 0.9951 leaves 0.987, which is the specific gravity of the alcohol and water of the wine. Or, divide the sp. gr. of the wine by the sp. gr. of the wine minus alcohol, carrying out the sum to four places of decimals; the quotient shows the sp. gr. of the water and alcohol only of the wine. On referring to a table of the strengths of diluted alcohol of different specific gravities, 0.987 at 15.5° C. is found to indicate a spirit containing 8 per cent. of real alcohol. Mulder's process is that adopted officially (U. S. P.) for ascertaining the strength of white wine (*Vinum Album*) and red wine (*Vinum Rubrum*). If the foregoing operation be conducted in a retort, the liquid being boiled and the steam carefully condensed, the distillate, diluted with water to the original bulk of wine operated on, will still more accurately represent the amount of water and alcohol in the wine—its specific gravity showing the percentage of real alcohol present.

DIALYSIS.

Dialysis (from *διὰ*, *dia*, through, and *λύσις*, *lusis*, a loosing or resolving) is a term applied by Graham to a process of analysis

by diffusion through a septum. The apparatus used in the process is called a *dialyzer*, and is constructed and employed in the following manner. The most convenient septum is the commercial article known as *parchment-paper*, made by immersing unsized paper for a short time in sulphuric acid; it is sold by most dealers in chemical apparatus. A piece of this material is stretched over a gutta-percha hoop, and secured by a second external hoop. Dialyzers of useful size are one or two inches deep and five to ten inches wide. Liquids to be dialyzed are poured into the dialyzer, which is then floated in a flat dish containing distilled water. The portion passing through the septum is termed the *diffusate*, the portion which does not pass through is termed the *dialysate*.

The practical value of dialysis depends upon the fact that certain substances will diffuse through a given septum far more rapidly than others. Uncrystallizable bodies diffuse very slowly. Of such matters as starch, gum, albumen, and gelatin, the last named is perhaps least diffusive; hence substances of this class are termed *colloids*, or bodies like *collin*, which is the soluble form of gelatin. Substances which diffuse rapidly are mostly crystalline; hence bodies of this class are termed *crystalloid*.

Solution of dialyzed iron, Liquor Ferri Dialysatus, B. P., an aqueous solution of about 5 per cent. of highly basic ferric oxychloride, is obtained by saturating 1 fluidounce of the solution of ferric perchloride with fresh ferric hydrate (obtained by precipitating 6 fluidounces of the same solution by ammonia, and washing and draining the product), filtering if necessary, placing on a dialyzer floating in distilled water, and displacing the fluid in the dish by water daily for a week or two, or until the diffusate gives no reaction with silver nitrate, and the fluid on the dialyzer is almost tasteless. The colloid fluid which has thus been subjected to dialysis (hence the only title to the name *dialyzed iron*), and which does not pass through the dialyzer, is the highly basic ferric oxychloride, or so-called "dialyzed iron," or "dialytic iron." This fluid has very little taste of iron. Its value as a medicine has been questioned, its non-diffusibility suggesting that it never passes out of the intestinal canal, and therefore never gets into the blood. Possibly it becomes decomposed in the system and the resulting product becomes assimilated.

It is "a clear dark reddish-brown liquid, free from any marked ferruginous taste." Neutral to test-papers. Specific gravity about 1.047. The solution gives no precipitate with potassium ferrocyanide or with silver nitrate, but after being

heated with hydrochloric acid it yields with potassium ferrocyanide a blue precipitate. 100 grains by weight afford a precipitate with solution of ammonia, which, washed, dried, and ignited, weighs 5 grains.

The phenomena of dialysis show that crystalloids are superior to colloids in affinity for water.

QUESTIONS AND EXERCISES.

Potassium carbonate is said to lose 16 per cent. of water on exposure to a red heat; give the details of manipulation observed in verifying this statement.—Write a few paragraphs descriptive of the process of ultimate organic analysis.—In what forms are carbon, hydrogen, and nitrogen weighed in quantitative analysis?—In the combustion of .41 of a gramme of sugar, what weights of products will be obtained? *Ans.* .632 of carbonic acid gas (CO_2) and .237 of water (H_2O).—How is cinchona assayed for mixed alkaloids?—Give the official method for the estimation of morphine in opium.—Mention the operations necessary for the estimation of the proportion of sugar in saccharated ferrous carbonate or in a specimen of diabetic urine.—What is understood by *saccharometry*?—Give two processes for the estimation of the percentage of alcohol in tinctures, wines, or beer.—Define dialysis.

CONCLUSION.

Detailed instructions for the quantitative analysis of potable water, articles of food, general technical products, special minerals, soils, manures, air, illuminating agents (including solid fats, oils, spirits, petroleum, and gas), dyes, and tanning-materials, would scarcely be in place in this volume.

The course through which the reader has been conducted will, it is hoped, have taught him the principles of the science of chemistry, and given him special knowledge concerning the applications of that science to medicine and pharmacy, as well as have imparted sufficient manipulative skill to meet the requirements of manufacture or analysis. The author would venture to suggest that this knowledge be utilized, not only in the way of personal advantage, but in experimental researches on chemical subjects connected with pharmacognosy, pharmacology, therapeutics, and pharmacy. The discovery and publication of a new truth, great or small, is the best means whereby to aid in advancing the calling in which we may be engaged, increase our own reputation, and contribute to that "ultimate end of knowledge" which Bacon described as "employing the divine gift of reason to the use and benefit of mankind."

APPENDIX.

TABLE OF TESTS FOR IMPURITIES IN PREPARATIONS
OF THE UNITED STATES PHARMACOPŒIA.*

Name of Preparation.	Impurities.	Tests.	Page.
<i>Acacia</i> ,	Starch.	Iodine.	473
	Mineral matter.	Incineration.	102
<i>Acetanilidum</i> ,	Other organic matter.	Sulphuric acid (colored).	
	Aniline, etc.	Ferric chloride.	431
	Copper.	Ammonia in excess.	193
	Lead, copper, etc.	Sulphuretted hydrogen.	258
	Sulphuric acid.	Soluble barium salt.	310
<i>Acidum Aceticum</i> and <i>Acidum Acet-</i> <i>icum Glaciale</i> ,	Hydrochloric acid.	Silver nitrate.	268
	Sulphurous and ferric acid.	Ammonia and silver nitrate (darkened).	
	Empyreumatic substances.	Vol. sol. of potassium permanganate.	652
	Mineral matter.	Evaporation and ignition.	102
	Metallic impurities.	Hydrochloric acid.	221
	Sulphides, etc.	In ammonia (soluble).	
<i>Acidum Arsenosum</i> ,	Mineral matter.	Evaporation and ignition.	102
	Antimony, tin, and cadmium.	Hydrogen sulphide precipitate in Na_2CO_3 (soluble).	
	Cinnamic acid.	Potassium permanganate.	652
	Mineral matter.	Incineration.	102
<i>Acidum Benzoicum</i> ,	Carbonizable organic matter.	Sulphuric acid (color).	
	Chlorine.	Silver nitrate.	268
	Sulphuric acid.	Barium chloride.	310
	Hydrochloric acid.	Silver nitrate.	268
	Pb, Cu, Fe, etc.	Ammonium sulphide.	258
<i>Acidum Boricum</i> ,	Calcium salts.	Ammonium oxalate.	117
	Magnesium salts.	Sodium phosphate.	122
	Ammonium salts.	Heat with potash.	98
	Sodium salts.	Flame test.	89
	Mineral matter.	Incineration.	102
<i>Acidum Carbolicum</i> ,	Creasote and cresylic acid.	Glycerin and dilute (turbidity).	453
<i>Acidum Chromicum</i> ,	Sulphuric acid.	Barium chloride.	310
	Tartaric and oxalic acids.	Potassium acetate.	322
	Calcium salts.	Ammonia and ammonium oxalate.	117
<i>Acidum Citricum</i> ,	Copper, lead, etc.	Sulphuretted hydrogen.	258
	Sulphuric acid.	Barium chloride.	310
	Mineral matter.	Incineration.	102
<i>Acidum Gallicum</i> ,	Tannic acid.	(Isinglass), gelatin.	358
	Mineral matter.	Incineration.	102
<i>Acidum Hydrobromicum Dilutum</i> ,	Iodine.	Cl-water and chloroform.	270
	Sulphuric acid.	Barium chloride.	310

* The manipulations necessary to be observed in testing for impurities will be found described in the paragraphs treating of those substances. The table also includes references to processes for ascertaining deficiency in strength of official articles.

The other characters and tests of pharmacopœial chemical compounds have been given in connection with the respective synthetical and analytical reactions.

TABLE OF TESTS—*Continued.*

Name of Preparation.	Impurities.	Tests.	Page.
<i>Acidum Hydrobromicum Dilutum,</i>	Fixed matter.	Evaporation.	102
	Acids.	Color on standing.	
	General.	Quantitative analysis.	637
	Arsenum.	Bettendorff's test.	175
	Iodine and bromine.	Chloroform and Cl-water.	270
<i>Acidum Hydrochloricum,</i>	Sulphurous acid.	4 drops of I solution (turbidity).	
	Sulphuric acid.	Barium chloride.	310
	Thallium, lead, etc.	Sulphuretted hydrogen.	258
	Copper.	Ammonia in excess.	193
	Arsenum.	Bettendorff's test.	175
<i>Acidum Hydrocyanicum Dilutum,</i>	Fixed matter.	Evaporation.	102
<i>Acidum Hypophosphorosum Dilutum,</i>	Lead, etc.	Sulphuretted hydrogen.	258
	Calcium salts.	Ammonium oxalate.	117
	Sulphuric, tartaric acids, etc.	Barium chloride.	310
	Potassium.	Platinic chloride.	78
	Sulphates.	Barium chloride.	310
<i>Acidum Lacticum,</i>	Chlorides.	Silver nitrate.	268
	Copper, iron, lead, etc.	Ammonium sulphide and ammonia.	224
	Sugars.	Fehling's solution.	468
	Glycerin.	Zinc carbonate and alcohol.	458
	Organic impurities.	Concentrated sulphuric acid (color).	
<i>Acidum Nitricum,</i>	Copper.	Ammonia in excess.	193
	Lead, iron, etc.	Ammonia and ammonium sulphide.	224
	Sulphuric acid.	Barium chloride.	310
	Hydrochloric acid.	Silver nitrate.	268
	Iodine and bromine.	Shake with chloroform.	271
<i>Acidum Oleicum,</i>	Iodic and bromic acids.	Zinc and chloroform (reduction).	460
	Fixed oils.	Mix with alcohol (no oily drops).	460
	Palmitic and stearic acids.	Lead salt, insoluble in ether.	460
	Phosphorous acid.	Mercuric chloride (turbid).	
	Arsenum.	Bettendorff's reaction.	175
<i>Acidum Phosphoricum,</i>	Phosphate.	Add alcohol and ether (turbidity).	
	Sulphuric acid.	Barium chloride.	310
	Hydrochloric acid.	Silver nitrate.	268
	Pyro- and meta-phosphoric acid.	Ferric chloride.	331
	Iron, carbolic acid, and coloring-matter.	Crystallized from alcohol.	492
<i>Acidum Salicylicum,</i>	Organic.	Concentrated sulphuric acid (colored).	
	Hydrochloric acid.	Silver nitrate.	268
	Lead.	Excess of alcohol (ppt.).	
	Nitric and nitrous acid.	Ferrous sulphate and sulphuric acid.	289
	Hydrochloric acid.	Silver nitrate.	268
<i>Acidum Sulphuricum,</i>	Lead, arsenic, copper, etc.	Hydrogen sulphide.	258
	Iron, etc.	Ammonia and ammonium sulphide.	
	Sulphuric acid.	Barium chloride.	310
	Gum and dextrin.	Dilute solution with alcohol (turbidity).	
	Resin.	Dilute with water (turbidity).	

TABLE OF TESTS—*Continued.*

Name of Preparation.	Impurities.	Tests.	Page.
<i>Acidum Tartaricum</i> ,	Oxalic and uric acids.	Calcium sulphate.	316
	Sulphuric acid.	Barium chloride.	310
	Calcium salts.	Ammonium oxalate.	117
	Iron, lead, copper, etc.	Ammonium sulphide.	224
<i>Adeps</i> ,	Alkalies.	Red litmus.	96
	Starch.	Iodine.	473
	Chlorides.	Silver nitrate.	268
	Free fatty acids.	.2 cc. standard KOH solution.	637
<i>Adeps Lanæ Hydro-</i> <i>sus</i> ,	Glycerin.	Evaporation.	461
	Alkalies.	Red litmus.	96
	Free fatty acid.	Drop of KOH solution.	637
	Ash.	Incineration.	102
<i>Æther</i> ,	Residue.	Evaporation.	460
	Aldehyde.	Potassium iodide.	449
	Alcohol and water.	Water.	449
	Amylic alcohol.	Evaporation on blotting-paper (odor).	449
<i>Æther Aceticus</i> ,	Alcohol.	Water.	449
	Organic.	Concentrated sulphuric acid (color).	449
<i>Alcohol</i> ,	Fusel oil.	Evaporation on blotting-paper (odor).	449
	Aldehyde, methyl alcohol, and oak tannin.	Potassium-hydrate solution (color).	449
	Organic.	Silver nitrate (color).	258
	Copper, lead, and zinc.	Hydrogen sulphide.	98
<i>Alumen</i> ,	Ammonium.	Potassium hydrate.	161
	Iron.	Potassium ferro- or ferricyanide.	162
<i>Alumini Hydras</i> ,	Iron.	Potassium ferrocyanide.	310
	Sulphate.	Barium chloride.	258
	Zinc and lead.	Hydrogen sulphide.	655
	Free acid.	Standard sodium hyposulphite.	162
<i>Alumini Sulphas</i> ,	Iron.	Potassium ferrocyanide.	98
	Ammonium salts.	Potassium hydrate.	258
	Copper, lead, or zinc.	Hydrogen sulphide.	310
	Sulphate.	Barium chloride.	268
<i>Ammonii Benzoas</i> ,	Chloride.	Silver nitrate.	99
	Fixed salts.	Non-volatility.	271
	Bromates.	Dilute sulphuric acid.	258
	Metals.	Hydrogen sulphide.	310
<i>Ammonii Bromidum</i> ,	Sulphates.	Barium chloride.	162
	Iron.	Potassium ferrocyanide.	310
	Sulphates.	Barium chloride.	258
	Metals.	Sulphuretted hydrogen.	117
<i>Ammonii Carbonas</i> ,	Calcium.	Ammonium oxalate.	268
	Chloride.	Silver nitrate.	102
	Non-volatile matters.	Evaporation.	258
	Metals.	Hydrogen sulphide.	310
<i>Ammonii Chloridum</i> ,	Sulphate.	Barium chloride.	117
	Calcium.	Ammonium oxalate.	103
	Barium.	Dilute sulphuric acid.	162
	Sulphocyanate.	Ferric chloride.	162
<i>Ammonii Iodidum</i> ,	Iron.	Potassium ferrocyanide.	102
	Non-volatile matter.	Evaporation and ignition.	310
	Sulphate.	Barium chloride.	473
	Iodine (free).	Starch.	641
<i>Ammonii Nitras</i> ,	Chloride or bromide.	Standard silver nitrate.	162
	Iron.	Potassium ferrocyanide.	310
	Sulphate.	Barium chloride.	268
	Chloride.	Silver nitrate.	268
<i>Ammonii Valerianas</i> ,	Acetate.	Ferric chloride (deep color).	310
	Sulphate.	Barium chloride.	310

TABLE OF TESTS—*Continued.*

Name of Preparation.	Impurities.	Tests.	Page.
<i>Ammonii Valerianas,</i>	Chloride.	Silver nitrate.	268
<i>Amyl Nitris,</i>	Aldehyde.	Potassium hydrate (brown).	
	General.	Fractional distillation.	407
	Free acid.	Quantitative analysis.	678
<i>Amylum,</i>	Ash.	Incineration.	102
	Sulphate.	Barium chloride.	310
	Calcium.	Ammonium oxalate.	117
	Chloride.	Silver nitrate.	268
<i>Antimonii et Potas- sii Tartras,</i>	Potassium bitartrate.	Sodium carbonate (no effervescence).	
	Iron and other metals.	Potassium ferrocyanide.	162
	Arsenum.	Bettendorff's test.	175
	Chloride.	Silver nitrate.	268
	Iron and other metals.	Potassium ferrocyanide.	162
<i>Antimonii Oxidum,</i>	Sulphate.	Barium chloride.	310
	Copper and lead.	Ammonium sulphide.	258
	Arsenum.	Bettendorff's test.	175
<i>Antimonii Sulphi- dum,</i>	General.	Quantitative analysis.	671
<i>Antimonium Sul- phuratum,</i>	Chloride.	Silver nitrate.	268
	Calcium.	Ammonium oxalate.	117
	Sulphate.	Barium chloride.	310
	Soluble salts.	Evaporation.	102
	Metallic impurities.	Ammonium sulphide.	224
	Ammonia.	Hydrogen sulphide.	258
	Sulphates.	Mercuric chloride (color).	
<i>Aqua,</i>	Chlorides.	Barium chloride.	310
	Nitrates.	Silver nitrate.	268
	Nitrites.	Diphenylamine and sul- phuric acid (ring).	
	Organic matter.	Zinc, iodide, starch (blue).	
		Potassium permanganate standard solution.	652
	Calcium.	Ammonium oxalate.	117
<i>Aqua Ammonia,</i>	Metallic.	Hydrogen sulphide.	258
	Sulphates.	Barium chloride.	310
	Fixed matter.	Evaporation and ignition.	102
<i>Aqua Chlori,</i>	Hydrochloric acid.	Blue litmus.	96
	Sulphates.	Barium chloride.	310
	Calcium.	Ammonium oxalate.	117
	Chlorides.	Silver nitrate.	268
	Ammonia.	Nessler's reagent.	639
<i>Aqua Destillata,</i>	Carbonic acid.	Lime-water.	314
	Organic matter.	Potassium permanganate.	652
	Nitrates.	Diphenylamine and sul- phuric acid (ring).	
	Nitrites.	Zinc, iodide, starch (blue).	
<i>Aqua Hydrogenii Dioxidii,</i>	Hydrofluoric acid.	Sulphuric acid.	342
	Barium.	Dilute sulphuric acid.	103
<i>Argenti Nitras,</i>	Copper.	Ammonia in excess.	193
	Lead.	Dilute sulphuric acid.	215
	Carbonate.	Effervescence with nitric acid.	219
<i>Argenti Oxidum,</i>	General.	Quantitative analysis.	676
<i>Arseni Iodidum,</i>	Fixed salts.	Incineration.	102
	Organic.	Sulphuric acid to solu- tion (color).	
<i>Atropina,</i>	Morphine.	Sulphuric acid to dry sub- stance (color).	519
<i>Balsamum Peruvi- anum,</i>	Essential oil.	Dilute with water.	414
	Heavy hydrocarbons.	Evaporation.	102
<i>Benzinum,</i>	Sulphur comps.	Silver nitrate and ammo- nia (brown).	
	Benzol.	Nitric and sulphuric acids, warm.	430

TABLE OF TESTS—Continued.

Name of Preparation.	Impurities.	Tests.	Page.
<i>Bismuthi Citras</i> ,	Nitrate.	Sulphuric acid and ferrous sulphate.	289
	Lead.	Dilute sulphuric acid.	215
	Copper.	Ammonia.	193
	Chloride.	Silver nitrate.	268
<i>Bismuthi Subcar- bonas</i> ,	Sulphate.	Barium chloride.	310
	Silver.	Hydrochloric acid.	219
	Ammonium comps.	Potassium hydrate.	98
	Nitrate.	Sulphuric acid and indigo.	291
	Arsenum.	Bettendorff's test.	175
<i>Bismuthi Subnitras</i> ,	Carbonate.	Effervesces with acids.	314
	Arsenum.	Bettendorff's test.	175
	Iodine.	Starch.	473
<i>Bromum</i> ,	General.	Sp. gr. (2.990); boiling-point.	597
<i>Caffeina</i> ,	Organic.	Strong sulphuric acid (color).	
	Other alkaloids.	Mercuric potassium iodide.	630
	Insoluble matter.	Soluble in water.	
	Arsenum, lead, etc.	Hydrogen sulphide.	258
	Iron, aluminium, etc.	Ammonium sulphide.	224
	Iodide.	Starch and chlorine-water.	473
	Sulphate.	Barium chloride.	310
<i>Calcii Bromidum</i> ,	Nitrate and nitrite.	Zinc and potassium hydrate (ammonia).	
	Chloride.	Silver nitrate.	268
	Magnesium and alkalies.	Ammonium oxalate and filtrate (no residue on evaporation).	102
	Sulphate.	Barium chloride.	310
	Chloride.	Silver nitrate.	268
<i>Calcii Carbonas Præcipitatus</i> ,	Iron, aluminium phosphate, etc.	Ammonia.	163
	Arsenum, lead, etc.	Hydrogen sulphide.	258
	Soluble matter.	Shake with water and evaporate.	102
	Iron, aluminium, etc.	Ammonia.	163
<i>Calcii Chloridum</i> ,	Sulphate.	Barium chloride.	310
	Arsenum, lead, etc.	Hydrogen sulphide.	258
	Barium.	Dilute sulphuric acid.	103
<i>Calcii Hypophos- phas</i> ,	Phosphate.	Lead acetate.	332
	Sulphate.	Barium chloride.	310
	Arsenum, etc.	Hydrogen sulphide.	258
	Magnesium.	Sodium phosphate.	122
	Carbonate.	Acid (effervesce).	314
<i>Calcii Phosphas Præcipitatus</i> ,	Sulphate.	Barium chloride.	310
	Barium.	Dilute sulphuric acid.	103
	Arsenum, lead, etc.	Hydrogen sulphide.	258
	Iron.	Potassium ferrocyanide.	161
	General.	Quantitative analysis.	685
<i>Calcii Sulphas Ex- siccatus</i> ,	Carbonate.	Effervescence with acids.	314
	Aluminium.	Ammonia.	163
<i>Calx</i> ,	Barium.	Dilute sulphuric acid.	103
	Carbonate.	Effervescence with acids.	314
<i>Calx Chlorata</i> ,	General.	Quantitative analysis.	657
<i>Calx Sulphurata</i> ,	General.	Quantitative analysis.	115
<i>Cambogia</i> ,	Starch.	Iodine (green).	473
<i>Camphora</i> ,	Fixed salts.	Non-volatility.	421
<i>Carbo Animalis</i> ,	Earthy salts.	Incineration.	102
<i>Carbonei Disulphi- dum</i> ,	Sulphurous acid.	Blue litmus.	96
	Dissolved sulphur.	Evaporated spontaneously.	102
	Hydrogen sulphide.	Lead acetate.	258
<i>Catechu</i> ,	Ash, 6 per cent.	Incineration.	102
<i>Cera Flava</i> ,	Fat acids, Japan wax.	Soda, acid, etc.	450
	Soap.	Water and acid, etc.	450

TABLE OF TESTS—*Continued.*

Name of Preparation.	Impurities.	Tests.	Page.
<i>Cera Flava</i> ,	Paraffin.	Completely soluble in concentrated sulphuric acid.	
	Tallow and other fats.	Ignited, smell of acrolein.	
<i>Cerii Oxalas</i> ,	Carbonate.	Effervescence with acids.	314
	Arsenum, etc.	Hydrogen sulphide.	258
	Aluminium.	Potassium hydrate.	163
	Zinc.	Ammonium sulphide.	136
	Hydrochloric acid and chlorides.	Silver nitrate.	268
<i>Chloral</i> ,	Chloral alcoholate.	Iodoform test.	444
	General.	Quantitative analysis.	487
	Chlorides.	Shake with silver water and test the watery solution.	268
<i>Chloroformum</i> ,	Free chlorine.	Starch and KI.	473
	General.	Specific gravity.	401
	Non-volatile matter.	Residue on evaporation.	401
	Hydrocarbons.	Sulphuric acid.	401
	Organic matter.	Colors with sulphuric acid.	526
<i>Cinchonidine Sulphas</i> ,	Excess of water.	Dried at 100° C.	660
	Cinchonine.	Ammonia after tartrate, etc.	526
	Mineral matter.	Incineration.	102
<i>Cinchonina</i> ,	Organic matter.	Colors with sulphuric acid.	526
	Mineral matter.	Incineration.	102
<i>Cinchoninæ Sulphas</i> ,	Excess of water.	Dried at 100° C.	660
	Cinchonidine.	Insoluble in chloroform.	526
	Organic.	Color sulphuric acid.	526
<i>Cocainæ Hydrochloras</i> ,	Water of crystallization.	Heat at 100° C.	660
	Non-volatile matter.	Incineration.	102
	Coca bases.	1 drop of potassium permanganate gives a color.	652
<i>Coccus</i> ,	Mineral matter.	More than 5 per cent. of ash.	102
<i>Codeina</i> ,	Morphine.	Nitric acid on the solid.	519
	Non-volatile matter.	Incineration.	102
	Fixed oils.	Soft residue on evaporation.	425
<i>Copaiba</i> ,	Turpentine.	Odor during evaporation.	425
	Gurjun balsam.	Carbon disulphide and nitric and sulphuric acids (violet).	
<i>Creasotum</i> ,		Oxidation.	453
	Carbolic acid.	Non-volatility at 100° C.	453
		Crystallization on cooling.	453
<i>Creta Præparata</i> ,	Magnesium.	Sodium phosphate.	122
	Iron.	Potassium ferrocyanide.	161
	Sulphate.	Barium chloride.	310
	Barium.	Dilute sulphuric acid.	103
<i>Crocus</i> ,	Coal-tar colors.	No color given to benzine.	
	Inorganic matter.	Incineration, not more than 7.5 per cent. of ash.	102
			161
<i>Cupri Sulphas</i> ,	Iron, aluminium, etc.	Ammonia.	
	Arsenum, lead, zinc, etc.	Hydrogen sulphide and acetic acid.	258
<i>Elaterinum</i> ,	Non-volatile matter.	Incineration.	102
<i>Eucalyptol</i> ,	Phenols.	Ferric chloride.	454
<i>Ferri Carbonas Saccharatus</i> ,	Sulphate.	Barium chloride.	310
	General.	Quantitative analysis.	650
<i>Ferri et Ammonii Citras</i> ,	Citrates and tartrates of potassium and sodium.	Alkalinity of ash.	96
<i>Ferri et Ammonii Sulphas</i> ,	Aluminium.	Potash, acid, ammonia.	141
<i>Ferri et Ammonii Tartras</i> ,	Citrates and tartrates of fixed alkalies.	Alkalinity of ash.	96

TABLE OF TESTS—*Continued.*

Name of Preparation.	Impurities.	Tests.	Page.
<i>Ferri et Quininae Citras,</i>	Citrates and tartrates of fixed alkalies.	Alkalinity of ash.	96
<i>Ferri et Strychninae Citras,</i>	General.	Quantitative analysis.	697
	Citrates and tartrates of fixed alkalies.	Alkalinity of ash.	96
<i>Ferri Hypophosphis,</i>	Carbonate.	Effervescence with acids.	314
	Calcium.	Ammonium oxalate.	117
	Phosphate.	Magnesium mixture.	331
	General.	Quantitative analysis.	655
<i>Ferri Iodidum Saccharatum,</i>	Iodine.	Starch.	473
	Salts of fixed alkalies.	Soluble after ignition.	102
	Sulphate.	Barium chloride.	310
	Sugar.	Boil with acid, Fehling's solution.	701
	Chloride.	Silver nitrate.	268
<i>Ferri Lactas,</i>	Carbonate.	Effervescence with acids.	314
	General.	Quantitative analysis.	654
	Gum, etc.	Strong sulphuric acid, { Colored brown. Offensive smell.	
	Butyric acid.		
<i>Ferri Phosphas Solubilis,</i>	General.	Quantitative analysis.	651
	Ferric phosphate.	Magnesia mixture and ammonia.	331
<i>Ferri Sulphas, etc.</i>	Copper, zinc, etc.	Ammonia, filter, ammonium sulphide.	223
	General.	Quantitative analysis.	650
<i>Ferri Valerianas,</i>	General.	Quantitative analysis.	650
<i>Ferrum Reductum,</i>	Sulphide.	Lead acetate.	214
	Arsenum.	Bettendorff's test.	175
	General.	Specific gravity.	614
	Fixed matter.	Evaporation and ignition.	102
	Organic matter.	Sulphuric acid (color).	
	Formic, butyric acids, etc.	Odor with dil. sulph. acid.	
<i>Glycerinum,</i>	Sulphuric acid.	Barium chloride.	210
	Hydrochloric acid.	Silver nitrate.	268
	Calcium salts.	Ammonium oxalate.	117
	Oxalic acid.	Calcium chloride.	316
	Sugars.	Fehling's solution.	701
	Metals.	Ammonium sulphide.	161
<i>Glycyrrhizinum Ammoniatum,</i>	Mineral matter.	Incineration.	102
<i>Gossypium Purificatum,</i>	Mineral matter.	More than .8 per cent. of ash.	102
	Acid or alkaline.	Litmus.	96
<i>Hydrarg. Chlor. Corrosivum,</i>	Arsenic.	Hydrogen sulphide: filter, evaporate, and incinerate.	176
	Foreign salts.	Hydrogen sulphide: filter, evaporate, and incinerate.	102
	Mercuric chloride.		
<i>Hydrarg. Chlor. Mite,</i>	Soluble matter.	Potassium hydrate.	98
	Ammonia.	Potassium iodide.	206
<i>Hydrargyri Cyanidum,</i>	Mercuric chloride.		
<i>Hydrarg. Iodidum Flavum,</i>	Fixed salts.	Non-volatility.	209
	Mercuric iodide.	Hydrogen sulphide.	207
	Mercuric chloride.	Litmus.	96
<i>Hydrarg. Iod. Rubrum,</i>	Fixed salts.	Non-volatility.	209
	Chlorides and iodides.	Soluble in water, silver nitrate.	268
<i>Hydrarg. Ox. Flavum,</i>	Chloride.	Silver nitrate.	268
	Fixed salts.	Non-volatility.	209
<i>Hydrarg. Ox. Rubrum,</i>	Nitrate.	Vapor and litmus.	96
	Chloride.	Silver nitrate.	268
	Fixed salts.	Non-volatility.	209

TABLE OF TESTS—*Continued.*

Name of Preparation.	Impurities.	Tests.	Page.
<i>Hydrarg. Subsul.</i> <i>Flavus,</i>	Mercurous and lead salts.	Not soluble in hydrochloric acid.	208
	Fixed salts.	Non-volatility.	209
	Foreign metals.	Yellowish color with $\text{Na}_2\text{S}_2\text{O}_3$.	
<i>Hydrargyrum,</i>	Fixed matter.	Non-volatility.	209
	Carbonate.	Effervescence with acids.	314
	Mercurous salts.	Not soluble in hydrochloric acid.	
	Lead.	Dilute sulphuric acid.	208
<i>Hydrargyrum Ammoniatum,</i>	Fixed salts.	Non-volatility.	215
	General.	Quantitative analysis.	209
	Mercurous oxide.	Hydrochloric acid.	673
	Mercuric oxide.	Hydrogen sulphide.	208
<i>Hydrarg. cum Creta,</i>	Inorganic matter.	Incineration.	207
<i>Hydrastininæ Hydrochloras,</i>			102
<i>Hyoscinæ Hydrobromas,</i>	Non-volatile matter.	Incineration.	
<i>Hyoscyaminæ Hydrobromas,</i>	Other alkaloids.	Platinic chloride (ppt.).	102
<i>Iodoformum,</i>	Fixed matter.	Incineration.	102
	Free acid.	Shake (Litmus.	96
	Coloring matter.	with Colored.	
	Iodide, soluble.	water. Silver nitrate.	275
<i>Iodum,</i>	General.	Quantitative analysis.	656
	Fixed salts.	Non-volatility.	274
	Iodine cyanide.	Physical characters.	274
	Jalap.	Solubility of resin.	505
<i>Jalap.</i>	Ash.	Incineration; .5 per cent.	102
<i>Limonis Succus,</i>	Deficiency of citric acid.	Quantitative analysis.	325
<i>Linum,</i>	Starch.	Iodine.	473
<i>Liquor Ammonii Acetatis,</i>	Non-volatile matter.	Evaporation and incineration.	102
<i>Liquor Calcis,</i>	Deficiency in strength.	Quantitative analysis.	635
<i>Liquor Ferri Acetatis, Liquor Ferri Chloridi, Liquor Ferri Citratis,</i>	Ferrous salts.	Potassium ferricyanide.	161
	Zinc, copper, etc.	Hydrogen sulphide, ammonia.	165
	Nitric acid.	Ferrous sulphate and sulphuric acid.	290
	General impurity or deficiency.	Specific gravity.	614
<i>Liquor Ferri Nitratidis,</i>	General.	Quantitative analysis.	669
<i>Liq. Fer. Subsulphas,</i>	Nitric acid.	Quantitative analysis.	669
<i>Liq. Fer. Tersulphatis,</i>	Ferrous salt.	Ferrous sulphate, etc.	290
<i>Liq. Hydrargyri Nitratidis,</i>	Mercurous salts.	Potassium ferricyanide.	161
<i>Liq. Plumbi Subacetatis,</i>	Fixed salts.	Hydrochloric acid.	208
<i>Liquor Potassæ and Liquor Sodæ,</i>	General.	Non-volatility.	209
<i>Liq. Sodæ Chloratæ,</i>	General.	Quantitative analysis.	631
<i>Lithii Benzoas,</i>	Deficiency.	Specific gravity.	614
	General.	Quantitative analysis.	632
	General.	Quantitative analysis.	659
	Chloride.	Silver nitrate.	268
	Sulphate.	Barium chloride.	310
	Arsenic, lead, iron, etc.	Ammonium sulphide.	165
	Calcium.	Ammonium oxalate.	117
	Potassium.	Sodium cobaltic nitrite.	80
	General.	Quantitative analysis.	634
	Calcium.	Ammonium oxalate.	117
<i>Lithii Bromidum,</i>	Arsenum, lead, etc.	Hydrogen sulphide.	190
	Sulphate.	Barium chloride.	310
	Iron, aluminium, etc.	Ammonium sulphide.	190
	Potassium.	Sodium cobaltic nitrite.	80
	Iodide.	Chlorine-water and starch.	473
	General.	Quantitative analysis.	642

TABLE OF TESTS—*Continued.*

Name of Preparation.	Impurities.	Tests.	Page.
<i>Lithii Bromidum,</i>	Chloride.	Silver nitrate.	268
	Arsenic, lead, etc.	Hydrogen sulphide.	190
	Iron, aluminium, etc.	Ammonium sulphide.	190
<i>Lithii Carbonas</i> and <i>Lithii Citras,</i>	Chloride.	Silver nitrate.	268
	Potassium.	Sodium cobaltic nitrite.	80
	Sulphate.	Barium chloride.	310
	General.	Quantitative analysis.	634
	Calcium.	Ammonium oxalate.	117
	Carbonate.	Effervescence with acids.	314
	Sulphate.	Barium chloride.	310
<i>Lithii Salicylas,</i>	Organic.	Sulphuric acid (color).	
	Chloride.	Silver nitrate.	268
	Calcium.	Ammonium oxalate.	117
	Aluminium, etc.	Ammonium sulphide.	190
	Potassium.	Sodium cobaltic nitrite.	80
	General.	Quantitative analysis.	634
	More than 10 per cent.	Incineration.	427
<i>Lupulinum,</i>	of ash.		
	Sand, etc.	Insoluble in water.	
<i>Lycopodium,</i>	Starch.	Microscope	474
	More than 5 per cent. of mineral matter.	Incineration.	102
<i>Magnesia,</i>	Carbonate.	Effervescence with acids.	314
	Sulphate.	Barium chloride.	310
	Chloride.	Silver nitrate.	268
	Metallic.	Ammonium sulphide.	256
	Water, more than 5 per cent.	Ignition and weigh.	102
	General.	Quantitative analysis.	666
	Calcium.	Ammonium oxalate.	117
<i>Magnesiæ Carbonas,</i>	Sulphate.	Barium chloride.	310
	Chloride.	Silver nitrate.	268
	Metallic.	Ammonium sulphide.	256
	Sodium.	Flame.	90
	Metallic.	Ammonium sulphide.	256
<i>Magnesiæ Sulphas,</i>	Chloride.	Silver nitrate.	268
	Arsenum.	Bettendorff's test.	175
	Antimony.	Ammonium sulphide.	184
	Organic.	No loss on ignition.	102
<i>Mangani Dioxidum,</i>	Sulphides.	Acids.	304
	General.	Quantitative analysis.	667
	Iron.	Potassium sulphocyanate.	162
	Arsenum and copper.	Hydrogen sulphide.	190
	Zinc.	Hydrogen sulphide and acetic acid.	256
	Magnesium and alkalies.	No residue after precipitating with ammonium carbonate.	
<i>Mel,</i>	General.	Quantitative analysis.	666
	Chlorides.	Silver nitrate.	268
	Starch.	Iodine.	473
	Sulphates.	Barium chloride.	310
	Glucose and inorganic matter.	.2 per cent. of ash on incineration.	102
	Thymol.	Sulphuric and nitric acids (blue).	
<i>Menthol,</i>	Wax, paraffin, and inorganic matter.	Non-volatility on water-bath.	112
	Alcohol, chloroform.	Distil.	127
<i>Methyl Salicylas,</i>	Narcotine, etc.	Concentrated sulphuric acid (yellow).	
<i>Morphina, Morphinæ Acetas, and Morphinæ Sulphas,</i>	Strychnine.	Conc. sulph. acid and pot. bichromate (purple).	
	Codeine, etc.	Soluble in ether.	
	Mineral matter.	Incineration.	102
	General.	Quantitative analysis.	699

TABLE OF TESTS—*Continued.*

Name of Preparation.	Impurities.	Tests.	Page.
<i>Naphtalinum</i> ,	Mineral matter.	Incineration.	102
	Coal-tar contaminations.	Sulphuric acid (concentrated) (dark red).	
<i>Naphtol</i> ,	Mineral matter.	Incineration.	102
	Naphtalin.	Insoluble in water.	434
	Organic.	Solution in water.	434
	α -naphtol.	Sulphuric acid.	434
<i>Olea.*</i>			
<i>Opium</i> ,	Deficiency in morphine.	Quantitative analysis.	698
	General.	Melting- and boiling-point.	485
	Sulphate.	Barium chloride.	310
	Chloride.	Silver nitrate.	268
	Amylic alcohol.	Opalescent solution with water.	636
<i>Paraldehydum</i> ,	Free acid.	Standard potassium hydrate.	
	Fixed impurities.	Non-volatility on water-bath.	112
		Litmus.	96
<i>Pepsinum</i> ,	Alkaline.	Quantitative analysis.	550
<i>Petrolatum Liquidum</i> , <i>Petrolatum Molle</i> , and <i>Petrolatum Spissum</i> ,	General.	Sulphuric acid, oily drops.	
	Resins, fats, and fixed oils.		
	Organic.	Concentrated sulphuric acid (black).	
<i>Phosphorus</i> ,	Fixed matter.	Incineration in platinum.	102
	Arsenum.	Hydrogen sulphide.	190
	Sulphur.	Barium chloride.	310
<i>Physostigminæ Salicylas</i> , <i>Physostigminæ Sulphas</i> ,			
	Fixed salts.	Incineration.	102
<i>Picrotoxinum</i> ,	Alkaloids.	Mercuric chloride, platinic chloride, tannin.	571
	Fixed salts.	Incineration.	102
<i>Pilocarpinæ Hydrochloras</i> ,	Fixed salts.	Incineration.	102
	Fixed salts.	Incineration.	102
<i>Piperinum</i> ,	General.	Quantitative analysis.	631
	Carbonate.	Opalescent solution in water.	
<i>Plumbi Acetas</i> ,	Iron and zinc.	Hydrogen sulphide and ammonia.	256
	Insoluble matter.	Not completely soluble in nitric acid.	
<i>Plumbi Carbonas</i> ,	Other elements.	Residue after precipitating Pb with H_2S .	
<i>Plumbi Iodidum</i> ,	Nitrate.	Sulphuric acid and indigo.	291
	Acetate.	Ferric chloride.	300
	Soluble salts.	Solution evaporated, residue left.	102
<i>Plumbi Nitras</i> ,	Iron and copper.	Potassium ferrocyanide (colored ppt.).	
	Zinc, etc.	Ammonium sulphide.	162
	Carbonate.	Effervescence with acid.	314
<i>Plumbi Oxidum</i> ,	More than 2 per cent. of moisture.	Heat strongly and weigh.	102
	Copper.	Ammonia.	193
	Iron.	Ammonia.	161
<i>Potassa</i> ,	Silicate, barium sulphate.	Not completely soluble in HNO_3 .	355
	Organic matter.	Not clear solution in water.	
	Arsenum, lead, etc.	Hydrogen sulphides.	222
	Iron, aluminium, etc.	Ammonia.	165
	Calcium.	Ammonium oxalate.	117

* The tests for the respective oils do not admit of arrangement in tabular form. The student is therefore referred to the text of this Manual or to the Pharmacopœia itself.

TABLE OF TESTS—*Continued.*

Name of Preparation.	Impurities.	Tests.	Page.
<i>Potassa,</i>	Chloride.	Silver nitrate.	268
	Sulphate.	Barium chloride.	310
	General.	Quantitative analysis.	630
	Carbonate.	Effervescence with acids.	314
	Nitrate.	Sulphuric acid and indigo.	291
	Silica.	Add an acid.	355
	Arsenum.	Odor on heating.	168
<i>Potassii Acetas,</i>	Lead, etc.	Hydrogen sulphide.	222
	Sulphate.	Barium chloride.	310
	Chloride.	Silver nitrate.	268
	Iron, etc.	Ammonium sulphide.	165
	Carbonates.	Effervescence with acids.	314
	Organic.	Sulphuric acid (colored).	
	Carbonate.	Phenolphthalein.	87
<i>Potassii Bicarbonas,</i>	Metallic.	Hydrogen sulphide.	258
	Iron.	Potassium ferrocyanide.	162
	Chloride.	Silver nitrate.	268
	Chloride.	Silver nitrate.	268
	Sulphate.	Barium chloride.	310
	Insoluble matter.	Not soluble in ammonia.	
	Calcium.	Ammonium oxalate.	117
<i>Potassii Bitartras,</i>	Copper, lead, etc.	Ammonium sulphide.	258
	Sodium.	Flame test.	90
	Bromate.	Color with dilute acids.	296
	Iodine.	Chlorine-water and starch.	473
	Aluminium, etc.	Ammonium sulphide.	258
	Arsenum, etc.	Hydrogen sulphide.	222
	General.	Quantitative analysis.	642
<i>Potassii Bromidum,</i>	Calcium.	Ammonium oxalate.	117
	Barium.	Potassium sulphate.	103
	Sulphate.	Barium chloride.	310
	Iron.	Potassium ferrocyanide.	162
	Chloride.	Silver nitrate.	268
	Insoluble residue.	Not completely soluble in water.	
	Sodium.	Flame test.	90
<i>Potassii Carbonas,</i>	Metallic.	Hydrogen sulphide.	258
	Hyposulphite.	Dilute acids.	347
	Nitrate.	Ferrous sulphate, etc.	289
	Sulphate.	Barium chloride.	310
	Chloride.	Silver nitrate.	268
	Sulphide.	Add dilute acid.	304
	Cyanide.	Ferrous sulphate, ferric chloride, and HCl (blue color).	
<i>Potassii Chloras,</i>	General.	Quantitative analysis.	631
	Sodium.	Flame test.	90
	Sulphate.	Barium chloride.	310
	Calcium.	Ammonium oxalate.	117
	Chloride.	Silver nitrate.	268
	Metals.	Hydrogen sulphide.	258
	Nitrate and nitrite.	Zinc and potassium hydrate.	
<i>Potassii Citras,</i>	Carbonate.	Effervescence with acids.	314
	Sulphate.	Barium chloride.	310
	Chloride.	Silver nitrate.	268
	Tartrate.	A ppt. with acetic acid.	
<i>Potassii Cyanidum,</i>	Carbonate.	Effervescence with acids.	314
	Ferrocyanide.	Ferric chloride.	162
	Sulphocyanate.	Ferric chloride.	162
	General.	Quantitative analysis.	641
<i>Potassii et Sodii Tartras,</i>	Calcium.	Ammonium oxalate.	117
	Arsenum, lead, etc.	Hydrogen sulphide.	258
	Ammonium salts.	Potassium hydrate.	98
	Sulphate.	Barium chloride.	310
	Chloride.	Silver nitrate.	268

TABLE OF TESTS—*Continued.*

Name of Preparation.	Impurities.	Tests.	Page.
<i>Potassii et Sodii Tartras,</i>	General.	Quantitative analysis.	632
<i>Potassii Ferrocyanidum,</i>	Carbonate.	Effervescence with acids.	314
	Sulphate.	Barium chloride.	310
	Chloride.	Silver nitrate.	268
	Ferri cyanide.	Silver nitrate, red ppt.	
	Calcium.	Ammonium oxalate.	117
<i>Potassii Hypophosphis,</i>	Carbonates.	Effervescence with acids.	314
	Chloride.	Silver nitrate.	268
	Sulphate.	Barium chloride.	310
	Phosphate.	Magnesia mixture.	331
	General.	Quantitative analysis.	655
	Iodate.	Starch and sulphuric acid.	76
	Arsenum, lead, etc.	Hydrogen sulphide.	258
<i>Potassii Iodidum,</i>	Sulphate.	Barium chloride.	310
	Nitrate and nitrite.	Zinc and potassium hydrate (ammonia).	
	Iron.	Potassium ferrocyanide.	162
	General.	Quantitative analysis.	643
	Cyanide.	Ferrous sulphate and KOH (blue color).	
<i>Potassii Nitras,</i>	Iron.	Potassium ferrocyanide.	162
	Sulphate.	Barium chloride.	310
	Chloride.	Silver nitrate.	268
	Calcium.	Ammonium oxalate.	117
	Metallic.	Hydrogen sulphide.	258
	Iodine.	Starch.	473
	Chlorate.	Concentrated sulphuric acid (yellow).	
<i>Potassii Permannas,</i>	Sulphate.	Barium chloride.	310
	Nitrate and chlorate.	Sulphuric acid and diphenylamine (color).	
	General.	Quantitative analysis.	635
	Chloride.	Silver nitrate.	268
	Sodium.	Flame test.	90
<i>Potassii Sulphas,</i>	Arsenum, lead, etc.	Hydrogen sulphide.	258
	Iron, etc.	Ammonium sulphide.	165
	Magnesium.	Sodium phosphate.	122
	Chloride.	Silver nitrate.	268
	Calcium.	Ammonium oxalate.	117
<i>Pyrogallol,</i>	Fixed matter.	Incineration.	102
	Organic matter.	Concentrated sulphuric acid (color).	
<i>Quinidinæ Sulphas,</i>	Morphine.	Nitric acid.	519
	Other cinchona alkaloids.	Solubility in ether.	
	Organic.	Sulphuric acid (color).	
<i>Quinina,</i>	Cinchonine, cinchonidine, or quinidine.	Sulphate of ammonium and ammonia (insoluble).	
	Fixed matter.	Incineration.	102
<i>Quininæ Bisulphas,</i>	Organic.	Sulphuric acid (color).	
	General.	Quantitative analysis.	695
	Fixed matter.	Incineration.	102
<i>Quininæ Hydrobromas and Quininæ Sulphas,</i>	Fixed matter.	Incineration.	102
	Free water.	Dry on water-bath and weigh.	112
	Organic.	Sulphuric acid (color).	
<i>Quininæ Hydrochloras,</i>	General.	Quantitative analysis.	695
	Sulphate.	Barium chloride.	310
	Fixed salts.	Incineration.	102
	Barium.	Dilute sulphuric acid.	103
	Organic.	Sulphuric acid (color).	
<i>Quininæ Valerianas,</i>	Free water.	Dry on water-bath.	112
	General.	Quantitative analysis.	617
	Fixed salts.	Incineration.	102

TABLE OF TESTS—*Continued.*

Name of Preparation.	Impurities.	Tests.	Page.
<i>Quininæ Valerianas,</i>	Sulphate.	Barium chloride.	310
	Organic.	Sulphuric acid (color).	
	Water.	Heat on water-bath.	112
<i>Resina Jalapæ,</i>	Soluble matter.	Partially soluble in water.	
	Common resin.	Ammonia solution, gelatinized on cooling.	
<i>Resorcinum,</i>	Fixed salt.	Incineration.	102
	Empyreumatic bodies.	Colored solution in water.	
<i>Saccharum,</i>	Grape-sugar.	Silver nitrate and ammonia.	467
	Insoluble matter.	Deposit on standing from solution.	
<i>Saccharum Lactis,</i>	Cane-sugar.	Cold sulphuric acid (color).	
<i>Salicinum,</i>	Mineral matter.	Incineration.	102
	Alkaloids.	Mercuric-potassium iodide.	571
<i>Salol,</i>	Sulphate.	Barium chloride.	310
	Chloride.	Silver nitrate.	268
	Mineral matter.	Incineration.	102
<i>Santoninum,</i>	Carbolic or salicylic acid.	Ferric chloride (color).	
	Mineral matter.	Incineration.	102
	Organic.	Sulphuric acid (colored).	
	Alkaloids.	Mercuric-potassium iodide.	571
<i>Sapo,</i>	Water in excess.	Dry at 110°; not more than 36 per cent. loss.	102
	Animal fats.	Alcoholic 4 per cent. solution; gelatinize on cooling.	
	Metallic.	Ammonium sulphide.	163
<i>Scammonium,</i> <i>Sinapis Alba</i> and <i>Sinapis Nigra,</i>	Sodium carbonate, etc.	Quantitative analysis and solubility in alcohol.	663
	Silica, etc.	Solubility in water.	
	Alkalinity.	Vol. est. of alkalinity.	635
	Starch.	Iodine.	473
<i>Soda,</i>	Starch.	Iodine.	473
	Organic matter.	Not clear aqueous solution.	
	Potassium.	Sodium cobaltic nitrite.	80
	Arsenum, etc.	Hydrogen sulphide.	258
	Iron, etc.	Ammonium sulphide.	165
	Calcium.	Ammonium oxalate.	117
	Chloride.	Silver nitrate.	268
	Sulphate.	Barium chloride.	310
	Silicate.	Alcohol to aqueous solution (ppt.).	
	Carbonate.	Effervescence with acids.	314
<i>Sodii Acetas,</i>	Nitrate.	Sulphuric acid and indigo.	291
	General.	Quantitative analysis.	631
	Sulphate.	Barium chloride.	310
	Calcium.	Ammonium oxalate.	117
	Chloride.	Silver nitrate.	268
<i>Sodii Arsenas,</i>	Arsenum, etc.	Hydrogen sulphide.	258
	Iron, etc.	Ammonium sulphide.	165
	Potassium.	Sodium cobaltic nitrite.	80
	Arsenite.	Reduce AgNO ₃ on boiling.	
<i>Sodii Benzoas,</i>	Metallic.	Ammonium sulphide.	258
	Chloride.	Silver nitrate.	268
	Potassium.	Sodium cobaltic nitrite.	80
	Metallic.	Ammonium sulphide.	258
<i>Sodii Bicarbonas,</i>	General.	Quantitative analysis.	630
	Sulphocyanate.	Ferric chloride.	162
	Potassium.	Sodium cobaltic nitrite.	80
	Normal carbonate.	Phenolphthalein.	87
	Arsenum, etc.	Hydrogen sulphide.	258
	Iron, etc.	Ammonium sulphide.	165

TABLE OF TESTS—*Continued.*

Name of Preparation.	Impurities.	Tests.	Page.
<i>Sodii Bicarbonas,</i>	Calcium.	Ammonium oxalate.	117
	Chloride.	Silver nitrate.	268
	Sulphate, etc.	Barium chloride.	310
	Ammonium.	Heat with potassium hydrate.	98
<i>Sodii Bisulphis,</i>	General.	Quantitative analysis.	631
	Chloride.	Silver nitrate.	268
	Hyposulphite.	Not clear solution in dilute HCl.	
	Sulphate.	Barium chloride.	310
<i>Sodii Boras,</i>	Arsenum, etc.	Hydrogen sulphide.	258
	Carbonate.	Effervescence with acids.	314
	Arsenum, etc.	Hydrogen sulphide.	258
	Calcium.	Ammonium oxalate.	117
	Phosphate.	Magnesia mixture.	331
	Chloride.	Silver nitrate.	268
	Sulphate.	Barium chloride.	310
	Nitrate.	Sulphuric acid and indigo.	314
	Potassium.	Sodium cobaltic nitrite.	80
	Calcium.	Ammonium oxalate.	117
<i>Sodii Bromidum,</i>	Sulphate.	Barium chloride.	310
	Iodine.	Chlorine-water and starch.	473
	Bromate.	Dilute sulphuric acid on salt (yellow).	
	Arsenum, etc.	Hydrogen sulphide.	258
<i>Sodii Carbonas,</i>	Excess of moisture.	Heat on water-bath and weigh.	112
	Iron, etc.	Ammonium sulphide.	165
	Calcium.	Ammonium oxalate.	117
	Sulphocyanate.	Ferric chloride.	162
	Metallic.	Ammonium sulphide.	258
	Sulphate, etc.	Barium chloride.	310
	Potassium.	Sodium cobaltic nitrite.	80
	Chlorides.	Silver nitrate.	268
	General.	Quantitative analysis.	631
	Tartrate.	Alkalinity of ash.	96
<i>Sodii Chloras,</i>	Potassium.	Sodium cobaltic nitrite.	80
	Arsenum, etc.	Hydrogen sulphide.	258
	Iron, etc.	Ammonium sulphide.	165
	Magnesium.	Sodium phosphate.	122
	Calcium.	Ammonium oxalate.	117
	Sulphate.	Barium chloride.	310
	Chloride.	Silver nitrate.	268
	Potassium.	Sodium cobaltic nitrite.	80
	Calcium.	Ammonium oxalate.	117
	Sulphate.	Barium chloride.	310
<i>Sodii Chloridum,</i>	Metallic.	Hydrogen sulphide and ammonia.	258
	Magnesium.	Sodium phosphate.	122
	Iodide or bromide.	Starch and chlorine-water.	473
	Carbonate, etc.	Effervescence with acids.	314
	Potassium.	Sodium cobaltic nitrite.	80
	Calcium.	Ammonium oxalate.	117
	Arsenum, etc.	Hydrogen sulphide.	258
	Chloride.	Silver nitrate.	268
	Phosphate.	Magnesia mixture.	331
	Sulphate.	Barium chloride.	310
<i>Sodii Hypophosphis,</i>	Iron.	Potassium ferrocyanide.	161
	General.	Quantitative analysis.	654
	Calcium.	Ammonium oxalate.	117
	Metallic.	Hydrogen sulphide and ammonia.	258
	Carbonates, etc.	Effervescence with acids.	314
	Sulphide.	Silver nitrate.	307
	Sulphate.	Barium chloride.	310
	General.	Quantitative analysis.	647

TABLE OF TESTS—Continued.

Name of Preparation.	Impurities.	Tests.	Page.
<i>Sodii Iodidum,</i>	Excess of water.	Water-bath and weigh.	112
	Potassium.	Sodium bitartrate.	79
	Calcium.	Ammonium oxalate.	117
	Arsenum, etc.	Hydrogen sulphide.	258
	Zinc, etc.	Ammonium sulphide.	165
	Iron.	Potassium ferrocyanide.	162
	Iodine.	Starch.	473
	Iodate.	Acid and starch.	296
	Sulphate.	Barium chloride.	310
	Cyanide.	Ferrous sulphate and KHO (blue).	
<i>Sodii Nitras,</i>	Nitrate, etc.	Iron, zinc, and NaHO (ammonia).	
	General.	Quantitative analysis.	643
	Metallic.	Ammonium sulphide.	258
	Magnesium, etc.	Ammonia and sodium phosphate.	126
	Potassium.	Sodium bitartrate.	79
<i>Sodii Nitris,</i>	Iodide, iodate.	Starch, H ₂ S, and chlorine-water (blue).	
	Sulphate.	Barium chloride.	310
	Chloride.	Silver nitrate.	268
	Insoluble matter.	Not soluble in water.	
<i>Sodii Nitris,</i>	Iodide.	Starch and chlorine-water.	473
	Lead, etc.	Hydrogen sulphide.	258
	Calcium.	Not completely soluble in water.	
		Hydrogen sulphide.	258
<i>Sodii Phosphas and Sodii Pyrophosphas,</i>	Potassium.	Sodium bitartrate.	79
	Carbonate.	Effervescence with acids.	314
	Arsenum.	Bettendorff's test.	175
	Chloride.	Silver nitrate and HNO ₃ .	268
	Sulphate.	Barium chloride.	310
	Hypophosphite, etc.	Silver nitrate ppt., boiled (brown).	
		Effervescence with acids.	314
<i>Sodii Salicylas,</i>	Metallic.	Hydrogen sulphide.	258
	Sulphate.	Barium chloride.	310
	Chloride.	Silver nitrate.	268
	Organic.	Sulphuric acid chars.	492
	Magnesium.	Ammonium oxalate.	117
<i>Sodii Sulphas,</i>	Carbonate.	Effervescence with acids.	314
	Metallic.	Hydrogen sulphide.	258
	Chloride.	Silver nitrate.	268
	General.	Quantitative analysis.	682
<i>Sodii Sulphis,</i>	Sulphate.	Barium chloride.	310
	Chloride.	Silver nitrate.	268
	Metallic.	Hydrogen sulphide.	258
<i>Sodii Sulphocarbolas,</i>	Hyposulphite.	Acids liberate sulphur.	347
	Sulphate.	Barium chloride.	310
	Chloride.	Silver nitrate.	268
	Metallic.	Ammonium sulphide.	257
<i>Sparteinae Sulphas,</i>	Fixed salts.	Incineration.	102
	Ammonium salts.	Heat with potassium hydrate.	98
<i>Spiritus Ætheris Nitrosi,</i>	General.	Specific gravity.	615
	Aldehyde.	Potas. hydrate (brown).	
	Deficiency of nitrous compound.	Quantitative analysis.	404
<i>Spiritus Frumenti and Spiritus Vini Gallici,</i>	Fusel oil.	Disagreeable odor after evaporation.	
	Glycerin.	Sweet taste after evaporation.	
	Tannin.	Ferric chloride.	358
	Free acid.	Potassium hydrate V. S., about 1.1 cc.	636

TABLE OF TESTS—*Continued.*

Name of Preparation.	Impurities.	Tests.	Page.
<i>Strontii Bromidum</i> ,	Arsenum, etc.	Hydrogen sulphide.	258
	Iron, etc.	Ammonium sulphide.	62
	Iodine.	Starch.	473
	General.	Quantitative analysis.	643
	Barium.	Dilute sulphuric acid.	102
<i>Strontii Iodidum</i> ,	Arsenum, etc.	Hydrogen sulphide.	258
	Iron, etc.	Ammonium sulphide.	62
	Barium.	Dilute sulphuric acid.	102
	General.	Quantitative analysis.	643
	Metallic.	Hydrogen sulphide and ammonia.	258
<i>Strontii Lactas</i> ,	Carbonate, etc.	Effervescence with acids.	314
	Barium.	Dilute sulphuric acid.	102
	Chloride.	Silver nitrate.	268
	Butyrate, etc.	Penetrating odor on heating gently the acid solution.	
<i>Strychnina</i> and <i>Strychninæ</i> Sul- <i>phas</i> ,	Organic.	Sulphuric acid (color).	
	General.	Quantitative analysis.	633
	Brucine.	Nitric acid.	529
	Mineral matter.	Incineration.	102
<i>Sulphur Lotum</i> and <i>Sulphur Præcipi- tatum</i> ,	Metallic, etc.	Completely soluble in NaHO.	
	Arsenum.	Hydrogen sulphide.	257
	Acid.	Litmus.	96
	Ammonia.	Litmus.	96
	Selenium,	Hydrochloric acid and KCN, boil (red).	
<i>Sulphur Sublima- tum</i> ,	Mineral matter.	Incineration; not more than .5 per cent.	102
<i>Syrupus Acidi Hy- driodici</i> ,	Iodine.	Starch.	473
<i>Tamarindus</i> ,	Traces of copper.	Iron.	161
<i>Terebenum</i> ,	Acids.	Litmus.	96
	Unaltered oil of turpentine.	Action on polarized light.	
<i>Terpeni Hydras</i> ,	Resinous matter.	Not completely volatile.	71
	Free acid.	Litmus.	96
	Fixed salts.	Incineration.	102
	Thymín.	Solution in caustic potash (oily drops).	
<i>Thymol</i> ,	Paraffin, etc.	Non-volatile on water-bath.	122
	Metallic.	Hydrogen sulphide and ammonia.	258
<i>Tinctura Ferri Chlo- ridi</i> ,	Fixed alkalies.	No residue after precipitating the iron.	
	Nitrate.	Ferrous sulphate, etc.	289
	Oxychloride.	Opalescent on diluting and boiling.	
<i>Veratrina</i> ,	Mineral matter.	Incineration.	102
<i>Vinum Album</i> ,	Free acid.	Volumetric potassium hydrate.	636
	Tannic acid.	Ferric chloride.	358
	Aniline colors.	Heated with chloroform and NaHO (isonitrile).	
<i>Vinum Rubrum</i> ,	Fuchsin.	Coloring silk.	
	Free acid.	Volumetric potassium hydrate.	636
<i>Zinci Acetas</i> ,	Tannic acid.	Ferric chloride.	358
	Arsenum, etc.	Hydrogen sulphide.	258
	Aluminium, etc.	Filtrate from zinc hydrate; contain solid matter.	
	Sulphate.	Barium chloride.	310
	Chloride.	Silver nitrate.	268

TABLE OF TESTS—*Continued.*

Name of Preparation.	Impurities.	Tests.	Page.
<i>Zinci Bromidum,</i>	Iodine.	Starch and chlorine-water.	473
	Arsenum, etc.	Hydrogen sulphide.	258
	Iron, calcium, etc.	Insoluble in excess of ammonium carbonate.	
	Magnesium.	Sodium phosphate.	122
	General.	Quantitative analysis.	643
<i>Zinci Carbonas Præcipitatus,</i>	Arsenum, etc.	Hydrogen sulphide.	258
	Iron, calcium, etc.	Insoluble in excess of ammonium carbonate.	
	Lead.	Insoluble in dilute sulphuric acid.	251
	Alkali.	Volumetric oxalic acid.	634
	Oxychloride.	Opalescent when diluted with alcohol.	
<i>Zinci Chloridum,</i>	Arsenum, etc.	Hydrogen sulphide.	258
	Iron, calcium, etc.	Insoluble in excess of ammonium carbonate.	
	Sulphate.	Barium chloride.	310
	General.	Quantitative analysis.	643
	Arsenum, etc.	Hydrogen sulphide.	258
<i>Zinci Iodidum,</i>	Iron, calcium, etc.	Insoluble in excess of ammonium carbonate.	
	Sulphate.	Barium chloride.	310
	Magnesium.	Sodium phosphate.	122
	General.	Quantitative analysis.	643
	Arsenum, etc.	Hydrogen sulphide.	258
<i>Zinci Oxidum,</i>	Iron, calcium, etc.	Insoluble in excess of ammonium carbonate.	
	Chloride.	Silver nitrate.	268
	Magnesium.	Sodium phosphate.	122
	Carbonate.	Effervescence with acids.	314
	Sulphate.	Barium chloride.	310
<i>Zinci Phosphidum,</i>	Lead, silicate, etc.	Insoluble in sulphuric acid.	257
	Insoluble matter.	Insoluble in dilute hydrochloric acid.	
	Iron, etc.	Potassium ferrocyanide.	162
	Lead, etc.	Ammonium sulphide.	258
	Arsenum, etc.	Hydrogen sulphide.	258
<i>Zinci Sulphas,</i>	Insoluble matter.	Insoluble in water.	
	Arsenum, etc.	Hydrogen sulphide.	258
	Iron, calcium, etc.	Insoluble in excess of ammonium carbonate.	
	Magnesium.	Sodium phosphate.	122
	Chloride.	Silver nitrate.	268
<i>Zinci Valerianas,</i>	Free acid.	Litmus.	96
	Metallic.	Ammonium sulphide.	258
	Magnesium.	Sodium phosphate.	122
	Acetate.	Ferric chloride.	300
	Butyrate.	Turbid solution with copper acetate.	364
<i>Zincum,</i>	Sulphur.	Dissolve in acids; H_2S given off.	304
	Arsenum, etc.	Hydrogen sulphide.	258
	Iron, etc.	Ammonium sulphide.	165

SATURATION TABLES.

Equivalent weights of Citric Acid, Tartaric Acid, Carbonate of Potassium, Bicarbonate of Potassium, Carbonate of Sodium, Bicarbonate of Sodium, Carbonate of Ammonium, and Carbonate of Magnesium; repeated (in black) for 20 parts of each, and incidentally (in roman) for other proportions. (Exact to two places of decimals.)

Citric Acid ($\text{H}_3\text{C}_6\text{H}_5\text{O}_7, \text{H}_2\text{O}$) $\div 3 \times 2 = 140$	20.00	18.66	16.96	14.00	9.78	16.66	26.74	29.31
Tartaric Acid $\text{H}_2\text{C}_4\text{H}_4\text{O}_6 = 150$	21.43	20.00	18.26	15.00	10.49	17.85	28.65	31.41
Carbonate of Potassium $\text{K}_2\text{C}_3\text{O} + 16\% \text{ Aq.} = 164.285$	23.47	21.90	20.00	16.43	11.48	19.52	31.38	34.40
Bicarbonate of Potassium $2(\text{KHCO}_3) = 200$	28.57	26.66	24.34	20.00	13.98	23.81	38.20	41.90
Carbonate of Sodium $\text{Na}_2\text{CO}_3, 10\text{H}_2\text{O} = 286$	40.86	38.13	34.81	28.60	20.00	34.04	54.63	59.98
Bicarbonate of Sodium $2(\text{NaHCO}_3) = 168$	24.00	22.45	20.45	16.80	11.74	20.00	32.09	35.18
Carbonate of Ammonium $(\text{N}_2\text{H}_5\text{C}_2\text{O}_5) \div 3 \times 2 = 104.7$	14.96	13.95	12.68	10.47	7.31	12.46	20.00	21.86
Carb. of Magnes. $(\text{MgCO}_3, \text{Mg}_2\text{HO}_4\text{H}_2\text{O}) \div 4 = 95.5$	13.64	12.73	11.62	9.55	6.68	11.37	18.24	20.00

The amount of acid given in any column will saturate the amount of carbonate in the same column, and *vice versa*.

The amounts of carbonate in any column are equal to each other in chemical power.

Lemon-juice (sp. gr. 1.040) contains, on an average, 9 per cent. by weight of citric acid.

The same Table in round numbers, for purposes of prescribing and dispensing.

Citric Acid	20	19	17	14	10	17	27	30
Tartaric Acid	22	20	18	15	11	18	29	32
Carbonate of Potassium (Potassii Carbonas, U. S. P.)	24	22	20	16	12	20	31	35
Bicarbonate of Potassium (Potassii Bicarbonas, U. S. P.)	29	27	24	20	14	24	38	42
Carbonate of Sodium (cryst.) (Sodii Carbonas, U. S. P.)	41	38	34	28	20	34	55	60
Bicarbonate of Sodium (Sodii Bicarbonas, U. S. P.)	24	22	20	17	12	20	32	36
Carbonate of Ammonium (Ammonii Carbonas, U. S. P.)	15	14	13	11	7	13	20	22
Carbonate of Magnesium (Magnesii Carbonas, U. S. P.)	13	13	11	9	7	11	18	20

The table is read thus: 20 grains of Citric Acid will saturate 29 grains of Bicarbonate of Potassium; 20 grains of Bicarbonate of Sodium will saturate, or be saturated by, 18 grains of Tartaric Acid; 11 grains of Tartaric Acid = 7 grains of Carbonate of Ammonium; 20 grains of Bicarbonate of Sodium are equivalent to, or will do as much work as, 34 grains of Carbonate of Sodium; 14 grains of Citric Acid are as strong as 15 of Tartaric Acid. It is occasionally convenient to double the numbers, halve them, or take some other proportion; also to employ them in weights other than grains.

Lemon-juice contains, on an average, 41 grains of Citric Acid in 1 fluidounce, or 5 grains per fluidrachm.

THE ELEMENTS.

	Symbol and atomic value.	Atomic weight. B. P.	Atomic weight, U. S. P.
Aluminium (Al_2^{VI})	Al^{IV}	27	27.04
Antimony (Sb^{III})	Sb^{V}	120	119.6
Arsenum (As^{III}) (74.9, Kessler)	As^{V}	75	74.9
Barium ($^{136.84}$, Marignac; $^{137.12}$, if $\text{O} = 16$)	Ba^{II}	136.8	136.9
Beryllium (or Glucinum) ($^{9.3}$, Awdziejew; Klatzo)	Be^{II}	9.1	9.03
Bismuth (Bi^{III}) (210 , Dumas; $^{207.5}$, if $\text{O} = 15.96$)	Bi^{V}	208	208.9
Boron ($^{10.9}$, Berzelius; $^{10.97}$, Clarke)	B^{III}	11	10.9
Bromine ($^{79.75}$, Stas; $^{79.96}$, if $\text{O} = 16$)	Br^{I}	80	79.76
Cadmium ($^{111.7}$, Lenssen)	Cd^{II}	112	111.5
Cæsium ($^{132.7}$, Johnson & Allen, Bunsen; 133 , if $\text{O} = 16$)	Cs^{I}	133	132.7
Calcium ($^{39.9}$, Erdmann and Marchand; 40 if $\text{O} = 16$)	Ca^{II}	40	39.91
Carbon (C^{II})	C^{IV}	12	11.97
Cerium (Ce^{III}) (138 , Mendelejeff)	Ce^{VI}	141.5	139.9
Chlorine ($^{35.368}$, Stas) $^{35.4}$, if $\text{O} = 16$	Cl^{I}	35.5	35.37
Chromium (Cr_2^{VI}) ($^{52.08}$, Siewert)	Cr^{VI}	52.5	52
Cobalt (Co^{II})	Co^{VI}	58.6	58.6
Copper ($^{63.12}$, Millon & Commaille; $^{63.3}$, if $\text{O} = 16$)	Cu^{II}	63.3	63.18
Didymium (?)	D^{II}	145	142
Erbium (?)	Eb^{III}	166	166
Fluorine ($^{18.96}$, Luca, Louyet)	F^{I}	19	19
Gallium	Ga	70	69.9
Germanium ($^{72.3}$, Boisbaudran)			72.3
Glucinum, see Beryllium			9.03
Gold (Au^{I}) ($^{196.2}$, Berzelius; $^{196.64}$, Kruss; $^{196.85}$, Thorpe and Laurie)	Au^{III}	197	196.7
Hydrogen ($^{1.0025}$, if $\text{O} = 16$)	H^{I}	1	1
Indium	In^{VI}	113.7	113.6
Iodine ($^{126.533}$, Stas; $^{126.86}$, if $\text{O} = 16$)	I^{I}	127	126.53
Iridium	Ir^{IV}	193	192.5
Iron (Fe^{II} and Fe_2^{VI}) ($^{55.9}$, Dumas)	Fe^{VI}	56	55.88
Lanthanium ($^{138.85}$, Clarke, Meyer, and Seubert)	La^{III}	138	138.2
Lead (Pb^{II}) ($^{206.9}$, Stas)	Pb^{IV}	206.4	206.4
Lithium	L^{I}	7	7.01
Magnesium ($^{24.4}$, Van de Plaats, Ostwald)	Mg^{II}	24	24.3
Manganese (Mn^{II} and Mn^{IV}) ($^{64.8}$, Dewar & Scott)	Mn^{VI}	55	54.8
Mercury ($^{199.8}$, Erdmann and Marchand; $^{200.1}$ to $^{200.4}$, others)	Hg^{II}	200	199.98
Molybdenum ($^{95.9}$, Dumas; Debray)	Mo^{VI}	96	95.9
Nickel (Ni^{II})	Ni^{VI}	58.6	58.6
Niobium (Columbium)	Nb^{V}	94	93.7
Nitrogen (N^{I} and N^{III}) ($^{14.009}$, Stas; $^{14.04}$, if $\text{O} = 16$)	N^{V}	14	14.01

THE ELEMENTS (*continued*).

	Symbol and atomic value.	Atomic weight, B. P.	Atomic weight, U. S. P.
Osmium (200, Ostwald).....	Os ^{IV}	185	193
Oxygen (15.96, Stas, Nilson)	O ^{II}	16	15.96
Palladium	Pd ^{IV}	106.2	106.35
Phosphorus (P ^{III}) (30.96, Schrotter)	P ^V	31	30.96
Platinum (194.9, if O = 16)	Pt ^{IV}	194.4	194.3
Potassium (39.04, Stas; 39.1, if O = 16)	K ^I	39	39.03
Rhodium (104.1, Berzelius)	Rh ^{IV}	104.3	102.9
Rubidium (85.2, Bunsen)	Rb ^I	85.4	85.2
Ruthenium (103.5, Berzelius)	Ru ^{IV}	104.2	101.4
Samarium	Sm	149.6	149.62
Scandium	Sc	44	43.97
Selenium (or Selenion) (79, various observers)	Se ^{IV}	78.9	78.87
Silicon (28.3, Clarke)	Si ^{IV}	28	28.3
Silver (107.66, Stas; 107.93, if O = 16)	Ag ^I	108	107.66
Sodium (22.98, Stas; 23.05, if O = 16)	Na ^I	23	23
Strontium (87.3, if H = 1)	Sr ^{II}	87.5	87.3
Sulphur (S ^{II} and S ^{IV})	S ^{VI}	32	31.98
Tantalum (182.8, Van der Plaats, Ostwald)	Ta ^V	182	182
Tellurium (128, Meyer, and Seubert)	Te ^{VI}	125	125
Terbium	Tb	159	159.1
Thallium (203.5, Crookes)	Tl ^{III}	204	203.7
Thorium (or Thorium) (234, Clarke)	Th ^{II}	233	231.9
Tin (Sn ^{II}) (117.4, Dumas; 117.97, Clarke)	Sn ^{IV}	118	118.8
Titanium (50, Clarke)	Ti ^{IV}	48	48
Tungsten (183.6, if H = 1)	W ^{VI}	184	183.6
Uranium (239.8, Ebelmen)	U ^{VI}	240	238.8
Vanadium (51.37, Clarke)	V ^V	51	51.1
Ytterbium	Yb	173	172.6
Yttrium (89, Ostwald)	Y ^{III}	90	88.9
Zinc (64.7, Axel Erdmann; 65.4, Ostwald)	Zn ^{II}	64.9	65.1
Zirconium (89.57, Clarke; 90.7, Ostwald)	Zr ^{IV}	90.4	90.4

The quantivalence or atomic value (for combination or exchange) of some elements is, apparently, variable: in the above table the full coefficients are given in the column of symbols, other common values in brackets.

Atomic weights were somewhat obscurely termed equivalents by Wollaston.

Other elements than the above exist. They are very rare. Some of the rarer so-called elements may not be truly elementary.

Note.—Students must expect the figures representing the atomic weights of elements to vary slightly from time to time, in accordance with the advancement of knowledge in the direction of purity of materials and improvements in manipulation, and as regards modes of research and realization of chemical and physical analogies amongst elements. Atomic weights have also been founded, at different times, on different units or starting-points—namely, H = 1 and O = 16; H = 1 and O = 15.96; and O = 16 and H = 1.0025. In the foregoing table the atomic weights in large figures are those which, as a rule, at present will be found more useful by English and American medical and pharmaceutical workers. (See a suggestive paper, with lists, by Venables, in the *Chemical News*, vol. lix. pp. 77 and 89, from the *Journal of the Elisha Mitchell Scientific Society*, vol. v. part 2.)

INDEX.

- Abies balsamea*, 412, 427
excelsa, 425.
Abrī radix, 504.
Abrin, 504.
Abrus precatorius, 504.
Absinthe, 500.
Absinthin, 500.
Absinthol, 417.
Absolute alcohol, 442, 597, 615.
Absorption spectra, 561.
Acacia catechu, 359.
suma, 359.
Acaciæ gummi, 116.
impurities in, 707.
Acetal, 486.
Acetamide, 431.
Acetanilide, 431, 544 (table).
Acetanilidum, 431.
impurities in, 707.
Acetate of ammonium, 93.
solution of, 93.
amyl, 406.
copper, 192.
ethyl, 300, 406.
iron, 153, 300.
lead, 211.
morphine, 517.
potassium, 70.
sodium, 83.
zinc, 134.
Acetates, 297.
analytical reactions of, 300.
decomposition of aqueous solutions of, 298.
Acetic acid, 297, 298, 398.
anhydrous, 299.
constitution of, 387, 398.
glacial, 299.
volumetric estimation of
free, 637.
aldehyde, 484.
anhydride, 299.
ether, 300, 406.
series of acids, 484.
relations of, 490.
Acetone, 300, 498.
Acetonitrate of barium, 127.
iron, 158.
Acetonitrile, 483.
Acetophenone, 498.
Acetoximes, 515.
Acetum, 398.
cantharidis, 298.
ipecacuanhæ, 298, 537.
opii, 298.
scillæ, 298.
Acetyl, 298.
chloride, 396.
Acetylene, 411.
series of hydrocarbons, 411.
Acetylenes, relations to paraffins
and olefines, 410
Acid carbonate of potassium, 71.
of sodium, 83.
salts, 73, 74, 302.
solution of arsenic, 168
tartrate of potassium, 62, 74,
79, 319.
Acidimetry, 639.
Acids, of acetic series, 484.
acrylic series, 491.
analytical detection of, 369.
antidotes to, 264.
benzoic or aromatic series, 491.
cinnamic series, 494.
definition of, 263.
dibasic, 493.
free, estimated, 636.
glyoxylic series, 490.
hexabasic, 498.
hydroxybenzoic series, 492.
lactic series, 489.
malic series, 496.
of chlorine, 295.
of phosphorus, 352.
organic, 482.
phthalic series, 496.
polybasic, 498.
quantitative estimation of, 637.
succinic series, 495.
sulphonic, 445.
table showing the relations of
acetic, lactic, and glyoxylic,
490.
acetic and dibasic
series, 497.

- Acids**, table showing the relations of benzoic and hydroxybenzoic, 493.
 tartaric series, 496.
 tetrabasic, 498.
 tribasic, 497.
 trihydroxybenzoic series, 494.
Acidulous radicals, formulæ and quantivalence of, 67, 124, 263.
 qualitative detection of, 367.
 quantitative estimation of salts of, 677.
 tables to aid in the detection of, 368, 369.
 volumetric estimation of, 639.
- Acidum aceticum*, 298.
 impurities in, 707.
dilutum, 299.
glaciale, 299.
 impurities in, 707.
arsenosum, 168.
benzoicum, 335.
boricum, 333, 334.
 impurities in, 707.
carbolicum, 453.
liquefactum, 453.
chromicum, 238.
 impurities in, 707.
citricum, 323.
 impurities in, 707.
gallicum, 360.
 impurities in, 707.
hydrobromicum dilutum, 270.
 impurities in, 707.
hydrochloricum, 30, 265.
 impurities in, 708.
dilutum, 265.
hydrocyanicum dilutum, 281.
 impurities in, 708.
hypophosphorum dilutum, 344.
 impurities in, 708.
lacticum, 347.
dilutum, 347.
 impurities in, 708.
meconicum, 349.
nitricum, 288.
 impurities in, 708.
dilutum, 288.
nitro-hydrochloricum, 288.
dilutum, 187, 289.
oleicum, 460.
 impurities in, 708.
phosphoricum, 328.
dilutum, 329.
 impurities in, 708.
salicylicum, 492.
 impurities in, 708.
- Acidum sulphuricum*, 310.
aromaticum, 310.
dilutum, 310.
 impurities in, 708.
sulphurosum, 305.
 impurities in, 708.
tannicum, 357.
 impurities in, 708.
tartaricum, 319.
 impurities in, 709.
- Acipenser*, 549.
Acokanthera, 506.
Aconine, 530.
Aconiti ferocis radix, 530.
folia, 529.
heterophylli radix, 530.
radix, 529.
Aconitia, 529.
Aconitic acid, 324.
Aconitina, 529.
Aconitine, 529.
Aconitum ferox, 530.
heterophyllum, 530.
napellus, 530.
- Acorin*, 420.
Acorus calamus, 420.
Acrinyl sulphocyanate, 451.
Acrolein, 458.
Acrylic acid, 489.
 aldehyde, 458.
Actea racemosa, 509.
Adeps, 462.
benzoatus, 462.
lanæ, 460.
 impurities in, 709.
hydrosus, 460.
 impurities in, 709.
- Adhesion**, 57.
Adipocere, 548.
Adraganthin, 479.
Advice to Students, xvii., 381.
Ægle Marmelos, 359.
Ærated bread, 470.
 water, 86.
Æsculin, 537.
Æther, 444, 448.
 impurities in, 709.
aceticus, 406.
 impurities in, 709.
purus, 448.
- Affinity**, chemical, 38.
 units of, 56, 123.
- Agate**, 354.
- Air**, composition of, 27.
 gas-burner, 23.
 influence of animals and plants on, 20.
 nitrogen in the, 25.
 oxygen in the, 16.

- Air**, relative weights of the, 27.
weight of 1 cubic cent., 620.
of 100 cubic inches, 621.
- Ajowan oil**, 415.
- Ajwain oil**, 415.
flowers, 415.
- Alwainka-puhl**, 415.
- Alabaster**, 106.
- Albumen**, 544
detection of, in urine, 573.
vegetable, 548.
- Albumen ovi*, 544.
- Albumenoids**, 549, 647.
- Albumens**, 549.
- Albumins**, 549.
- Albumoses**, 549.
- Alchemy**, 13, 191
- Alcohol**, absolute, 442, 597, 615.
allylic, 451.
amyllic, 363, 449.
benzylic, 456.
butylic, 364, 449.
cetylic, 450.
cerylic, 450.
cinnamic, 495.
decylene, 452.
deodorized, 443.
ethylic, 439.
from sugar, 439.
hydroxybenzylic, 456.
impurities in, 709.
melissic, 450.
methylic, 437.
pentylic, 449.
phenic, 452.
propargyl, 411.
propenyl, 457.
propylic, 447.
purity of, 444.
quantitative estimation of, 704.
real, 442.
salicylic, 456.
test for, 444.
tolyl, 455.
- Alcoholates of bromal**, 488.
of chloral, 488.
- Alcoholic drinks**, 441.
- Alcoholometer**, 616.
- Alcohols**, 436.
allylic series, 451.
benzylic, 455.
dihydric, 409.
ethylic series, 456.
hexhydric, 465.
monhydric, 436.
penthydric, 465.
primary, 437.
saligenin, 455.
secondary, 437.
- Alcohols**, tertiary, 437.
tetrahydric, 465.
trihydric, 457, 465.
- Aldehyde**, 483.
acrylic, 457.
benzoic, 336, 432, 491.
cinnamic, 417.
cumic, 417.
euodic, 419.
formic, 484.
glycolic, 409.
lauric, 419.
orthohydroxybenzoic, 494.
oxalic, 409.
parahydroxybenzoic, 494.
rutic, 419.
salicylic, 407, 457, 494.
test for, 485.
- Aldehydes**, 482.
general reactions, 483.
general formation, 482.
- Aldose**, 467.
- Aldoximes**, 515.
- Ale**, 478.
- Alexandrian senna**, 501.
- Alizarate of potassium**, 434.
- Alizarin**, 434, 554.
- Alkalies**, analytical separation of
the, 101.
antidotes to, 264.
quantitative estimation of the,
628.
- Alkalimetry**, 628.
- Alkaline carbonates**, volumetric
estimation of the, 631.
earths, 127.
solution of arsenic, 166.
- Alkaloids**, 510.
animal, 512.
antidotes to the, 516.
nomenclature of, 515.
poisonous, examination for, 566,
et seq.
reagents for, 570.
vegetal, 512.
vegeto-animal, 515.
- Alkanet**, 554.
- Alkanna tinctoria*, 554.
- Alkyl salts**, 407, 484.
- Allium*, 452.
- Allotropes**, 482.
- Allotropic bodies**, 482.
- Allotropy**, 480.
- Alloxan**, 362.
- Alloy**, 196.
- Allyl cyanide**, 451.
sulphide, 452.
sulphocyanate, 451.
- Allylene**, 411.

- Allylic series of alcohols*, 451.
 alcohol, 451.
Almond-oil, 463.
Almonds, oil of bitter, 491, 499.
 test for nitroben-
 zol in, 500.
Aloe barbadensis, 434.
 purificata, 434.
 socotrina, 434.
Aloes, 434.
Aloins, 434.
 formulæ of, 435.
Alstonia constricta, 537.
 scholaris, 537.
Alstonicine, 537.
Alstonine, 537.
Althæa officinalis, 479.
Alum, 138.
 cake, 139.
 chrome, 139, 338.
 dried, 140.
 flour, 139.
 impurities in, 709.
 iron, 139.
 potash, 139.
 roche or rock, 140.
 soda, 139.
Alumen, 138.
 exsiccatum, 140.
 impurities in, 709.
Alumina, 140.
Aluminium, 139.
 • analytical reactions of, 140.
 and ammonium sulphate, 138.
 and sodium, double chloride of,
 138.
 bronze, 138.
 chloride of, 138.
 derivation of word, 33.
 detection of, in presence of iron
 and zinc, 163.
 hydrate, 140.
 oxide, 140.
 quantitative estimation of, 668.
 separation of, from chromium
 and iron, 240.
 silicate, 138.
 steel, 138.
 sulphate, 140.
Amalgam, 194.
 ammonium, 91.
Amber, 356.
 oil of, 356.
American pennyroyal, 418.
 turpentine, 412.
Amianth, 354.
Amidacetic acid, 552.
Amide-bases, 431, 510, 515.
Amides, 206, 431.
Amido-acet-phenetidin, 431.
Amido-benzene, 430, 513.
 -chloride of mercury, 206.
Amido-succinamic acid, 496.
Amidogen, 206, 431.
Amines, constitution of, 431.
Ammonia, 91, 92.
 acetate, 93.
 benzoate, 95, 337.
 carbonate, 93.
 citrate, 95.
 detected by Nessler's test, 630.
 fetid spirit of, 94.
 gas, composition of, 91, 130.
 in drinking-water, 630.
 nitrate, 94.
 oxalate, 95.
 phosphate, 95.
 preparation of, 92.
 solution of, 92.
 sulphate, 91.
 type, 379.
 volcanic, 91.
 volumetric estimation of solu-
 tions of, 629.
Ammoniacal liquor, 92.
 salts, sources of, 91.
Ammoniacum, 427.
Ammonix benzoas, 95, 337.
 carbonas, 93.
 liquor, 92.
 fortior, 92.
 nitras, 94.
 phosphas, 95.
 sesquicarbonas (see *Ammonii Car-*
 bonas).
 spiritus aromaticus, 94.
 fetidus, 94.
Ammoniated mercury, 206.
 varieties of, 206.
Ammonii acetatis, liquor, 93.
 benzoas, 95, 337.
 impurities in, 709.
 bromidum, 95, 271.
 impurities in, 709.
 carbonas, 93.
 impurities in, 34, 709.
 chloridum, 91.
 impurities in, 709.
 citratiss, liquor, 95.
 nitras, 94.
 impurities in, 709.
 phosphas, 95.
 sulphas, 91.
Ammonio-chloride of mercury, 206,
 207.
 -citrate of iron, 154.
 -magnesian phosphate, 122, 330.
 -nitrate of silver, 178, 207.

- Ammonio-sulphate of copper, 177.
 magnesium, 685.
 -tartrate of iron, 154.
- Ammonium**, 90.
 acetate, 93.
 amalgam, 91.
 analytical reactions of, 98.
 and bismuth, citrate of, 254.
 and magnesium, arsenate, 122.
 phosphate, 122, 330.
 and platinum double chloride,
 99, 249.
 arsenate, 169.
 aspartate, 349.
 benzoate, 95, 337.
 bicarbonate, 94.
 bromide, 95, 271.
 vol. estimation of, 630.
 carbamate, 93.
 carbonate, 93.
 solution of, 94.
 chloride, 91.
 chromate, 238.
 citrate, 95.
 cyanate, 339.
 derivation of word, 32.
 derivatives, 207.
 hydrate, 92.
 iodides, 273, 274.
 molybdate, 331.
 nitrate, 94, 288.
 oxalate, 95.
 periodide, 273.
 phosphate, 95.
 potassium and sodium, separa-
 tion of, 101.
 quantitative estimation of, 663.
 salts, source of, 91.
 volatility of, 99.
 sulphate, 91.
 and ferrous sulphate, 652.
 sulphide, 96.
 sulphydrate, 96.
 tartrate, 99.
 urate, 362.
 volumetric estimation of car-
 bonate of, 630.
- Amomum melegueta*, 418.
- Amorphous alkaloid, 526.
 meaning of, 78.
 phosphorus, 328.
- Amphicreatinine, 513.
- Amrad, 479.
- Amygdala amara*, 463, 499.
 dulcis, 463, 499.
- Amygdalin, 499.
- Amyl**, acetate, 406.
 nitrite, 351, 407.
 valerianate, 363, 407.
- Amyl nitris*, 351, 407.
 impurities in, 710.
- Amylamine, 511.
- Amylene, 408.
 hydrate, 450.
- Amylic alcohols, 363, 449.
- Amyloids, 472.
- Amyloses, 472.
- Amylum*, 472.
 impurities in, 710.
- Amyric acid, 426.
- Amyrin, 426.
- Anacyclus pyrethrum*, 425.
- Analogies between chlorine, bro-
 mine, and iodine, 272, 278.
- Analogy of carbon, boron, and sil-
 icon, 356.
 of oxygen, sulphur, selenium,
 and tellurium, 302.
 of sodium and potassium salts,
 89.
 of nitrogen, phosphorus, arse-
 num, and antimony, 166, 332,
 512.
- Analysis**, blowpipe, 372, 373.
 gas, 361, 560.
 gravimetric, 592, 659.
 meaning of word, 62.
 of gases and vapors, 560.
 of insoluble salts, 375, *et seq.*
 of medicines, 557.
 of salts, 376.
 of substances having unknown
 properties, 557.
 organic, 384.
 practical, 100, 257.
 proximate, 688.
 quantitative, 592.
 spectral, 262, 560.
 systematic, for the detection
 and separation of the metals,
 101, 125, 163, 186, 220, 257.
 ultimate, 688.
 volumetric, 625.
- Analytical chemists, 14.
 detection of the acidulous radi-
 cals of salts soluble in water,
 365.
 memoranda, 222, 259.
- Anamirta paniculata*, 506.
- Anamirtin, 506.
- Anchusa tinctoria*, 554.
- Anchusin, 554.
- Andira Araroba*, 338.
- Andrographis paniculata*, 509.
- Andropogon citratus*, 420.
 nardus, 417.
 schænanthus, 418.
- Aneroid barometer, 594.

- Anethene, 415.
 Anethol, 415.
 Angelate of potassium, 415.
 Angelic acid, 415.
 powder, 181.
Angelica, 428.
 Angostura bark, false, 527.
 true, 536.
 Angosturin, 536.
 Anhydride, acetic, 299.
 antimonic, 181.
 antimonious, 181.
 arsenous, 168.
 boric, 338.
 carbonic, 312.
 chlorochromic, 240, 278.
 chromic, 239.
 molybdic, 332.
 nitric, 288, 290.
 nitrous, 289, 290.
 phosphoric, 26, 313, 349.
 phthalic, 591.
 silicic, 355.
 sulphocarbonic, 314.
 sulphuric, 310.
 sulphurous, 305.
 Anhydrides, 85, 300.
 Anhydrochromate of potassium, 238.
 Anhydrous bodies, 85.
 arsenous acid, 168.
 Anhydrous chromic acid, 239.
 ferrous chloride, 149.
 nitric acid, 288.
 perchloride of iron, 148.
 stannic sulphide, 244.
 sulphate of copper, 192.
 Aniline, 431, 513.
 blue, 557.
 colors, 557.
 green, 557.
 red, 557.
 yellow, 557.
 Animal alkaloid, 512.
 charcoal, 112.
 decolorizing power of, 113.
 rouge, 337.
 starch, 472.
 Animals and plants, complementary
 action of air, 19.
 Aniseed oil, 415.
 Anise-fruit, 415.
 Anisum, 415.
 Annatto, 554.
 Anode, 245.
 Anthemem, 415.
Anthemis, 415, 509.
Anthemis nobilis, 415.
 Anthracene, 434, 554.
 Anthracite, 241.
 Anthraquinone, 434.
 Antichlor, 306.
Antidotes to acids, 264.
 alkalies, 264.
 alkaloids, 516.
 antimony, 186.
 arsenic, 152, 168.
 barium, 104.
 carbolic acid, 454.
 copper, 194.
 cyanides, 283.
 hydrochloric acid, 268.
 hydrocyanic acid, 283.
 lead, 215.
 mercurey, 209.
 nitric acid, 292.
 oxalic acid, 317.
 silver, 220.
 sulphuric acid, 311.
 tin, 244.
 zinc, 137.
 Antifebrin, 431.
 Antimonial wine, 182.
 Antimonic anhydride, 181.
 chloride, 180.
 oxide, 181.
Antimonii chloridi, liquor, 180.
 oxidum, 181.
 impurities in, 710.
 Antimonious anhydride, 181.
 chloride, 181.
 oxide, 181.
 oxychloride, 181.
Antimonii et potassii tartras, 182.
 impurities in, 710.
 sulphidum, 180.
 impurities in, 710.
 sulphuratum, 182.
 impurities in, 710.
Antimonium tartaratum, 182, 319.
 impurities in, 710.
 quantitative estimation of
 antimony in, 671.
 Antimoniuretted hydrogen, 185.
Antimony, 180.
 analytical reactions of, 184.
 and arsenic, analytical separa-
 tion of, 187.
 and potassium, tartrate of, 182,
 319.
 antidote to, 186.
 black, 180.
 bromide, 180.
 butter of, 180.
 chloride, 180.
 crocus, 180.
 crude, 180.
 derivation of word, 33.

- Antimony** from arsenic, to distinguish, 187.
 glass, 180.
 hydride, 184.
 in organic mixtures, detection of, 562.
 iodide, 180.
 oxide, 181.
 oxychloride, 181, 184.
 oxysulphide, 182.
 pentachloride, 180.
 potassio-tartrate, 182.
 quantitative estimation of, 647, 671.
 solution of chloride of, 180.
 sulphide, 177, 184.
 sulphur, salts of, 183.
 sulphurated, 182.
 tartarated, 182.
 volumetric estimation, 647.
- Antipyrine**, 431.
Antiseptic, 454, 457, 492.
Antozone, 585.
Apatite, 332.
Apocynum, 509.
Apomorphinæ hydrochloras, 520.
Apomorphine, 519.
Aporetine, 338.
Apothecaries, 14.
Apparatus, xii., xiii., 16.
 for experiments, xiii.
 for volumetric analysis, 626.
 lists of, xiii.
- Apple essence**, 407.
 oil, 407.
 wine, 441.
- Aqua**, 129.
 impurities in, 710.
 ammoniacæ, 92.
 fortior, 92.
 impurities in, 710.
 amygdala amara, 499.
 anethi, 414.
 anisi, 414.
 aurantii florum fortior, 416.
 camphoræ, 422.
 carui, 414.
 impurities in, 710.
 chlori, 29, 267.
 chloroformi, 414.
 cinnamomi, 414.
 creasoti, 453.
 destillata, 129.
 impurities in, 710.
 feniculi, 414.
 fortis, 288.
 duplex, 288.
 simplex, 288.
 hydrogenii oxidum, 103.
- Aqua hyd. oxidum**, impurities in, 710.
 laurocerasi, 282, 500, 641.
 menthæ piperitæ, 414.
 viridis, 414.
 pimentæ, 414.
 regiæ, 187, 288.
 rosæ, 414, 419.
 sambuci, 414.
- Arabic acid**, 478.
Arabin, 116.
Arabinose, 466.
Arachidic acid, 490.
Arachin, 464.
Arachis, 464.
 hypogæa, 464.
 oil, 464.
- Araroba powder**, 338.
Arbor Dianæ, 220.
Arbutin, 359, 501.
Archil, 555.
Arctium lappa, 509.
Arctostaphylos uva ursi, 501.
- Are**, 604.
Areca catechu, 360.
 nuts, 360.
- Arecaïne**, 360.
Arecoline, 360.
Arekane, 360.
Areometers, 616.
- Argal. Vide Acid Tartrate of Potassium.**
- Argent-ammon-ammonium, nitrate of**, 207.
- Argenti et potassii nitras**, 218.
 cyanidum, 220.
 iodidum, 220.
 nitras, 216.
 impurities in, 710.
 nitras dilutus, 218.
 fuscus, 218.
 oxidum, 219.
 impurities in, 710.
- Argentie chloride, sulphide, etc. Vide Salts of Silver.**
- Argentiferous galena**, 216.
Argentum, 33.
 purificatum, 218.
- Argol. Vide Acid Tartrate of Potassium.**
- Armenian bole**, 555.
Armoraciz radix, 415.
Arnatto, 554.
Arnica flores, 423.
Arnica rhizoma, 423.
Arnicin, 423.
Arnotto, 554.
- Aromatic acids**, 491.
 glycols, 456.
 series of hydrocarbons, 429.

- Arrow-poison, 506.
 Arrowroot starch (fig.), 475.
 Arsenate of ammonium, 169.
 barium, 105, 178.
 calcium, 178.
 copper, 178.
 iron, 146, 170, 178.
 magnesium and ammonium, 122.
 silver, 178, 220.
 sodium, 170.
 volumetric estimation of, 646.
 zinc, 178.
 Arsenates, 169.
 Arsenic, 33, 167, 168.
 acid, 169.
 solution of, 169.
 and arsenical solutions, volumetric estimation of official, 646.
 anhydride, 169.
 antidotes to, 152, 168.
 derivation of word, 33.
 in carbonate of potassium, solution of, 167.
 in hydrochloric acid, solution of, 168.
 odor of, 169.
 white, 167.
 Arsenical ores, 167.
 sulphur, 176.
 Arsenicum, 33, 168.
 Arsenide of cobalt, 234.
Arseni iodidum, 167.
 Arsenio-sulphide of iron, 167.
 nickel, 236.
 Arsenous acid, 168.
 anhydride, 168.
 oxide, 168.
 sulphide, 176.
 Arsenite of copper, 178, 194.
 potassium, 168.
 silver, 177.
 sodium, 168.
 Arsenites, 168.
 Arsenum, 167, 168.
 analytical reactions of, 171.
 and antimony, analytical separation of, 187.
 Bettendorff's test, 177.
 bromide, 166.
 chloride, 166.
 detection of, in metallic copper, 173.
 in organic mixtures, 562.
 Fleitmann's test for, 175.
 from antimony, to distinguish, 187.
 Arsenum, hydride, 174.
 iodide, 167.
 Marsh's test for, 173.
 molecular weight of, 166.
 quantitative estimation of, 670.
 red native sulphide of, 167.
 reduction of arsenates to arsenites, 169.
 Reinsch's test for, 172.
 sources of, 167.
 sulphide, 167, 176.
 yellow native sulphide of, 167.
 Arseniuretted hydrogen, 174.
 Arsines, 512.
 Art of Chemistry, 13.
 Artemisia absinthium, 499.
 pauciflora, 507.
 Artificial alkaloids, 510.
 gastric juice, 550.
 Arum-root, 360.
Asafetida, 427.
Asagrea officinalis, 543.
 Asbestos, 354.
 Asclepidin, 509.
Asclepias tuberosa, 509.
 Aselline, 463.
 Aseptol, 445.
 Ash, 102.
 black, 88.
 bone, 112.
 soda, 89.
 Asparagine, 349, 496.
 Aspartate of ammonium, 349.
 Aspidospermine, 530.
 Aspirator, water, 309.
 Asphalte, 427.
 Asymmetrical atoms, 320.
Assafetida, 427.
 -ate, meaning of, 73, 76.
 Atees, 530.
 Ateesine, 530.
 Atis, 530.
 Atmosphere, aqueous vapor in, 27.
 carbonic acid in, 313.
 composition of, 27.
 minor constituents, 27.
 nitrogen in, 25.
 oxygen in, 25.
 Atmospheric pressure, quantitative determination of, 593.
 Atom, definition of, 57.
 weights, definition of, 59.
 Atomic attraction, 139.
 Atomic weights as indicated by—
 chemical "periodicity," 380.
 properties, 387, 392.
 density of gases and vapors, 53, 619.
 electricity, 623.

- Atomic weights** as indicated by—
 isomorphism, 55.
 specific heat, 622.
 substitution, 193.
- Atomic proportions**, 52, 198.
 theory, 51, 54, 198, 380.
 weights, 52.
- Atomicity**, 56.
- Atoms**, 36, *et seq.*
 conception of, 389.
 quantivalence of, 56.
 definition of, 59.
- Atropa belladonna*, 531.
- Atropia*, 530.
- Atropina*, 532.
 impurities in, 710.
- Atropinæ sulphas*, 533.
- Atropine*, 531.
- Attar of rose*, 419.
- Attraction**, atomic, 139.
 molecular, 139.
- Aurantii cortex*, 415, 509.
fructus, 415.
- Auri et sodii chloridum*, 246.
- Auric chloride**, 246.
 sulphide, 246.
- Aurous salts**, 246.
- Aurum*, 34.
- Australian kino**, 359.
- Avenæ farina*, 473.
- Avignon grains**, 553.
- Avogadro's and Ampère's "law,"**
 53.
- Azadiracta indica*, 509.
- Azedarach*, 509.
- Azo**, 513.
- Azobenzene**, 436.
- Azoimide**, 515.
- Azoxybenzene**, 430.
- BACTERIA**, 440.
- Bacterium aceti**, 484.
mycoderma, 297.
- Bael-fruit**, 359.
mucilage, 479.
- Bahia powder**, 338.
- Baking-powder**, 470.
- Balance**, 600.
- Balloons**, coal-gas for, 24.
 hydrogen for, 24.
- Balm-of-Gilead fir**, 427.
- Balsam**, Canada, 412, 427.
copaiva, 426.
Gurjun, 426.
 of Peru, 422, 495.
 of storax, 422, 495.
 of Tolu, 422, 495.
- Balsamodendron myrrha*, 428.
- Balsams**, 422.
- Balsamum peruvianum*, 422, 495.
 impurities in, 710.
tolutanum, 422, 495.
- Baptin**, 532.
- Baptisia tinctoria*, 532.
- Baptism**, 532.
- Baptitoxine**, 532.
- Barbadoes aloes**, 435.
- Barbaloin**, 435.
- Barberry**, 533.
- Baric chloride**, nitrate, etc. *Vide*
 Salts of Barium.
- Barii dioxidum*, 103.
 vol. est. of, 663.
- Barium**, 103.
 acetonitrate, 127.
 analytical reactions of, 104.
 and calcium, separation of,
 from magnesium, 126.
 antidotes to, 104.
 arsenate, 105.
 carbonate, 103.
 native, 103.
 chloride, 103.
 chromate, 105.
 neutral, 105.
 derivation of word, 32.
 detection of, in presence of cal-
 cium and magnesium, 117,
 126.
 dioxide, 103.
 vol. est. of, 663.
 flame, 105.
 hydrate, 103.
 nitrate, 103.
 oxalate, 105, 317.
 oxide, 103.
 peroxide, 103.
 phosphate, 105, 332.
 quantitative estimation of, 664.
 salts, antidote to, 104.
 sulphate, 103.
 sulphide, 103.
 sulphite, 307.
 sulphocarbonate, 454.
- Barley starch** (fig.), 475.
 sugar, 471.
- Barometer**, 593.
- Barwood**, 420, 555.
- Baryta**, 103.
 nitrate, 103.
 water, 103.
- Barytes**, nitrate of, 103.
- Basalt**, 138.
- Base**, meaning of, 262.
 organic, 392.
- Basic**, meaning of, 262.
- Bassorin**, 116, 479.
- Bastard saffron**, 555.

- Basyulous hydrocarbons, 392.
 radicals, 60, 124.
 Bath brick, 354.
 Bauxite, 138.
 Bay rum, 441.
 salt, 81.
 Bearberry, 359.
 Beaver tree, 509.
 Beberia, or Beberine, 532.
 Bebeeru, 532.
Beberinæ sulphas, 532.
 Bebirine, 532.
 Beer, 441, 478.
 Beeswax, 450.
 Beetroot, 469.
 Behenic acid, 490.
Bela fructus, 359.
Belladonnæ folia, 531.
 radix, 531.
 Bell-metal, 242.
 Bend glass tubes, to, 17.
 Benné oil, 464.
 Benzaldehyde, 492, 500.
 artificial, 500.
Benzene, 429.
 disulphonic acid, 455.
 nitro, 430.
 series of hydrocarbons, 429.
 constitution of, 432.
 sulphonic acid, 445.
 Benzin, 398, 430.
 Benzine-Collas, 430.
Benzinum, 398.
 impurities in, 710.
 Benzoate of ammonium, 95.
 of iron, 337.
 of sodium, 337.
 Benzoated lard, 337.
 Benzoates, 335.
 analytical reactions of, 337.
 Benzodichloride, 337.
 Benzoëne, 429, 431.
 Benzoic acid, 335, 431, 491.
 aldehyde, 336, 431, 491.
 Benzoin, 336, 422.
Benzoinum, 335, 422.
 Benzol, 429.
 Benzoline, 398.
 Benzotrichloride, 336.
 Benzoyl chloride, 492.
 ecgonine, 534.
 hydrate, 492, 534.
 hydride, 492.
 sulphonic imide, 445.
 Benzyl benzoate, 495.
 cinnamate, 495.
 derivatives, 431.
 hydrate, 492, 495.
 Benzylic alcohol, 456, 495.
 Benzylic alcohols, 452.
 Berbamine, 533.
 Berberine or Berberia, 533.
Berberis cortex, 533.
 Bergamot oil, 416.
 Berlin blue, 555.
 red, 555.
 Berthollet's laws, 379.
 Beryllium, 725.
 Betaine, 515.
 Beta-naphtol, 434.
 Betel-nuts, 360.
 Bettendorff's test for arsenum, 175.
 Betula lenta, 407.
 Bhang, 424.
 Bi-, the prefix, 73.
 Bibasic. *See* Dibasic.
 Bibirine, 532.
 Bibiru, 532.
 Biborate of sodium, 333.
 Bibulous paper, 109.
 Bicarbonate of ammonium, 94.
 potassium, 71.
 sodium, 83, 313.
 chemically pure, 628.
 lozenges, 87.
 Bicarbonates, test for, 315.
 Bichloride of mercury, 201.
 Bichromate of potassium, 238.
 Bikh, 530.
 Bile, 552.
 detection of, in urine, 575.
 test for presence of, 552.
 Biliary calculi, 589.
 Bimeconate of morphine, 517.
 Binary hypotheses, 286.
 Birch oil, 407.
 Bish, 533.
Bismuth, 251.
 analytical reactions of, 254.
 and ammonium, solution of
 citrate of, 254.
 carbonate, 253.
 citrate, 253.
 derivation of word, 34.
 hydrate, 254.
 iodides, double, 253, 571.
 lozenge, 252.
 nitrate, 251.
 oxide, 253.
 quantitative estimation of, 772.
 salts, composition of, 252, 253.
 subcarbonate or oxycarbonate,
 253.
 subnitrate or oxynitrate, 252.
 sulphate, 254.
 sulphide, 254.
Bismuthi citras, 253.
 impurities in, 711.

- Bismuthi et ammonii citras*, 254.
oxidum, 253.
subcarbonas, 253.
 estimation of bismuth in, 772.
 impurities in, 711.
subnitras, 252.
 impurities in, 711.
Bismuthum, 251.
purificatum, 251.
 Bisulphide of carbon, 314.
 Bisulphite of lime, 306.
 of sodium, 306.
 Bitartrate of potash, 79.
 Bitter almonds, oil of, 491, 499.
 cassava, 472.
 orange-rind oil, 415.
 principles, 509.
 Bittern, 269.
 Bitter-sweet, 541.
 Bituminous coal, 241.
 Bivalence, 56.
 Bivalent radicals, 56, 67, 123.
Bixa orellana, 554.
 Bixin, 554.
Black alder-bark, 338.
 antimony, 179.
 ash, 88, 313.
 band, 142.
 bone-, 112.
 cherry-bark, 500.
 coloring-matters, 556.
 dyes, 556.
 flux, 169.
 haw, 510.
 hellebore, 505.
 ink, 162, 556.
 lead, 30.
 oxide of copper, 191.
 of iron, 142.
 of manganese, 231.
 of mercury, 505.
 pepper, 541.
 snake-root, 509.
 Bladder-green, 556.
Blanc de perle, 253, 556.
 Bleaching by chlorine, 29.
 liquor, 115.
 powder, 114.
 Blende, 131.
 Blistering collodion, 480.
 Block tin, 241.
 Blood, 545.
 composition of, 546.
 detection of, in organic matter, 584.
 hydrocyanic acid in the, 284.
 root, 541.
 stains, 584.
 Blowpipe analysis, 372.
 flame, 135.
Blue coloring-matters, 555.
 cohosh, 509.
 copperas, 144, 192.
 flag, 509.
 gum tree, 359.
 indigo, 291.
 ointment, 195.
 pill, 195.
 Prussian, 341.
 stone, 192.
 Turnbull's, 161, 342.
 vitriol, 144, 192.
 "Boiled oil," 463.
 Boiling-point, definition of, 597.
 Boiling-points of various substances, 597.
 Boldine, 416.
 Boldo, 416.
 Bonduc-seeds, 509.
 Bone-ash, 112.
 -black, 112.
 -earth, 112, 327.
 Bone oil, 513.
 Bones, composition of, 112, 327.
 Boneset, 509.
 Boracic acid, 333.
 Borate of glyceryl, 335.
 Borates, 333.
 analytical reactions of, 334.
 Borax, 333.
 bead, 232.
 Bordeaux turpentine, 412.
 Boric acid, 333.
 as an antiseptic, 547.
 anhydride, 333.
 Borneène, 421.
 Borneo camphor, 451.
 Borneol, 451.
Boron, 333.
 chloride, 333.
 derivation of word, 34.
 flame, 334.
 fluoride, 333.
 molecular weight of, 333.
 Borotartrate of potassium, 334.
Bos taurus, 552.
Boswellia, 428.
 Botany Bay kino, 359.
 Bourdon barometer, 594.
 Boyle's law, 53, 619.
 Brandy, 443.
 Brass, 131.
Brassica alba, 451.
juncea, 451.
nigra, 451.
 Brayera, 506.
 Brazil powder, 338.

- Brazil wood, 556.
 Bread, 471.
 aërated, 471.
 Bread-crumbs, 471.
 Bread-making, 471.
 Breidin, 426.
 Brezilín, 554.
 Bricks, 355.
 Bright's disease, 474.
 Britannia metal, 180, 210, 241.
 British gum, 477.
 Bromal, 488.
 alcoholates, 488.
 hydrate, 488.
 Bromate of potassium, 77.
 Bromates, 77, 271, 296.
 Bromic acid, 77, 296.
 Bromide of—
 ammonium, 95, 270.
 antimony, 180.
 arsenum, 166.
 ethyl, 402.
 iron, 148.
 phosphorus, 329.
 potassium, 77, 270.
 volumetric estimation of, 643.
 silver, 220, 271.
 sodium, 270.
 sulphur, 304.
 Bromides, 270.
 analytical reactions of, 271.
 quantitative analysis of, 678.
 separation of, from chlorides and iodides, 276.
Bromine, 270.
 analytical separation of, 276.
 derivation of word, 33.
 its analogy to chlorine and iodine, 273, 278.
 solution of, 271.
 volumetric estimation of, free, 678.
 solution of, 658.
Bromum, 269.
 impurities in, 711.
 Bronze, 241.
 aluminium, 138.
 coinage, 191.
 Bronzing-powder, 244.
 Broom-tops, 542.
 Brown coloring-matters, 556.
 hæmatite, 142, 153.
 rosin, 423.
 sugar, 469.
 Brucia, 528.
 Brucine, 528.
 distinction from morphine, 529.
 Brunswick green, 177.
 Bryoidin, 426.
 Bryonin, 501.
Bryonia alba, 501.
 dioica, 501.
Buchu, 416.
 oil of, 416.
 Buckthorn green, 556.
 juice, 502.
 "Bumping," 282.
 Bunsen's gas-burners, 23.
 valve, 652.
 Burdock, 509.
 Burette, Muhr's, 627.
 Burgundy pitch, 425.
 Burners, gas, 17, 23.
 Burnett's disinfecting fluid, 132.
 Burnt ochre, 555.
 sugar, 471.
 umber, 556.
 Butane, 397.
Butea frondosa, 359.
 Butter, 463, 546.
 of antimony, 180.
 of cacao, 462.
 of cocoa, 462.
 of kokum, 462.
 of orris, 418.
 Butyl-chloral, 488.
 hydras, 488.
 hydrate, 488.
 sulphocyanate, 415.
 Butylene, 408.
 Butylic alcohol, 364, 449.
 Butyrate of ethyl, 407.
 Butyrates, 364.
 Butyric acid, 364, 488.
 aldehyde, 488.
 chlorinated, 488.
 Butyrone, 498.
 Buxine, 533.
Buxus sempervirens, 533.
 By-products, 201.
CABBAGE-ROSE petals, 555.
 oil of, 419.
 Cacao-butter, 462.
 Cadaverine, 513.
Cadmium, 249.
 analytical reactions of, 250.
 derivation of word, 34.
 hydrate, 250.
 iodide, 250.
 oxide of, 250.
 sulphide, 250.
 sulphate, 250.
Cæsalpinia bonducella, 509.
 brasiliensis, 554.
 Cæsium, 726.
Caffeina, 242.

- Caffeina*, impurities in, 711.
citrata, 542.
effervescens, 543.
- Caffeine, 542.
citrate, 542.
- Cajuput oil, 416.
- Cajuputene, 416.
- Cajuputol, 416.
- Caking coal, 241.
- Calabar bean, 540.
- Calamina præparata*, 134.
- Calamine, 131, 420.
prepared, 134.
- Calamus, 420.
- Calamus draco*, 424.
- Calcic sulphate, phosphate, etc.
Vide Salts of Calcium.
- Calcii bromidum*, 271.
volumetric estimation of, 643.
impurities in, 711.
- carbonas præcipitatus*, 108.
impurities in, 711.
- chloridi, liquor*, 106.
- chloridum*, 106.
impurities in, 711.
- chlorinata, liquor*, 115.
- hydras*, 108.
- hypophosphis*, 343.
impurities in, 711.
- phosphas præcipitatus*, 113.
impurities in, 711.
- sulphas exsiccatus*, 106.
impurities in, 711.
- Calcined magnesia, 122.
- Calcium**, 106.
analytical reactions of, 116.
and barium, separation from
magnesium, 126.
bisulphite, 306.
bromide, 271.
volumetric estimation of, 643.
- carbonate, 106, 108.
prepared, 113.
- chloride, 106.
- chromate, 117.
- citrate, 324.
- derivation of word, 32.
- flame, 117.
- fluoride, 106.
in bones, 114.
- gummate, 116.
- hydrate, 108.
- hypochlorite, 114.
- hypophosphite, 343.
volumetric estimation of, 654.
- hyposulphite, 303.
- Calcium**, in presence of barium and
magnesium, detection of, 126.
oxalate, 117, 316.
oxide, 108.
phosphate, 106, 114, 327.
polysulphide, 303.
quantitative estimation of, 665.
silicate, 106.
sulphate, 106, 115.
sulphide, 115.
sulphite, 305.
tartrate, 325.
- Calc-spar, 106.
- Calcis, liquor, 108.
syrupus, 108.
- Calculi, urinary, 572.
examination of, 587.
- Calendula officinalis*, 509.
- Calendulin, 509.
- Caliche, 286.
- Calomel, 203, 208.
test for corrosive sublimate in, 203.
- Calotropis*, 509.
- Calumbæ*, 533.
- Calx*, 108.
impurities in, 711.
- chlorata*, 114.
impurities in, 711.
- sulphurata*, 115.
impurities in, 711.
- Cambogia*, 427.
impurities in, 711.
- Camphene, 412.
- Camphor laurel, 421.
mixture, 422.
oil, 421.
water, 422.
- Camphora*, 421.
impurities in, 711.
- officinarum*, 421.
- Camphoric acid, 422.
- Camphoronic acid, 422.
- Camphors, 421.
- Cam-wood, 554.
- Canada balsam, 412, 427.
- Canadian hemp, 509.
turpentine, 412.
- Candle-flame, composition of, 23.
- Canellæ cortex*, 509.
- Cane-sugar, 469.
- Cannabene, 423.
hydride of, 423.
- Cannabin, 423.
- Cannabinine, 423.
- Cannabis indica*, 423.
sativa, 423.
- Cantharides, vinegar of, 298.
- Cantharidic acid, 422.

- Cantharidin, 422.
Cantharis, 422.
 Caoutchin, 421.
 Caoutchouc, 420.
 Capacity, unit, 603.
 Capillary, 595.
 Capric acid, 490.
 Caproate of glyceryl, 462.
 Caproic acid, 462, 490.
 Caprylate of glyceryl, 462.
 Caprylic acid, 462.
 Capsaicin, 425, 534.
 Capsicin, 425.
 Capsicine, 425, 534.
Capsicum, 508, 638.
 Capsicum-fruit, oleoresin of, 425.
 oil, 425.
 Caramel, 471.
 Caraway-fruit, 416.
 oil, 416.
 Carbamate of ammonium, 93.
 ethyl, 489.
 Carbamic acid, 489.
 Carbamide, 489.
 Carbamines, 483, 544 (table).
 Carbazotic acid, 454, 554.
 Carbinols, 436.
Carbo animalis, 112.
 purificatus, 112.
 impurities in, 711.
 ligni, 113.
 Carbohydrates, 466.
 Carbolates, 454.
 Carbolic acid, 452.
 antidote to, 454.
Carbon, 30.
 bisulphide, 314.
 combustion of, 30.
 compounds, chemistry of, 383.
 derivation of word, 32.
 disulphide, 314.
 quantitative estimation of, in
 organic compounds, 688, *et*
 seq.
 tetrachloride, 401.
 Carbonate of ammonium, 93.
 solution of 94.
 barium, 102, 104.
 bismuth, 254.
 calcium, 106, 108.
 prepared, 113.
 iron, 145.
 saccharated, 145.
 lead, 210, 215.
 lithium, 227.
 magnesium, 119, 122.
 potassium, 62.
 acid, 71.
 sodium, 85.
 Carbonate of sodium, acid, 83.
 chemically pure, 628.
 manufacture of, 88, 312.
 strontium, 229.
 zinc, 134, 137.
Carbonates, 312.
 acidulous radical in, 66, 312.
 analytical reactions of, 314.
 gravimetric estimation of, 683.
 volumetric estimation of alkali-
 line, 631.
 Carbonic acid, 31, 312, 489.
 gas, generation of, 72.
 solubility of, in water,
 86.
 anhydride, 312.
 oxide, 341.
Carbonei disulphidum, 314.
 impurities in, 711.
 Carbonization, 102.
 Carbonyl, 437.
 Carboxyl group, 437, 489.
 Carburetted hydrogen, light, 396.
 heavy, 409.
 Cardamom oil, 416.
 greater, 418.
 lesser, 416.
Cardamomum, 416.
Carica papaya, 534, 551.
 Carolina jasmine, 537.
 Carmine, 337.
 Carminic acid, 337.
 Carnallite, 61.
 Carnine, 515.
 Carpaine, 534.
 Carrageen moss, 479.
 Carrotin, 554.
 Carthamin, 555.
Carthamus tinctorius, 555.
Carum ajowan, 415.
 Carvene, 416.
 Carvol, 415, 416.
 Caryophylene, 412.
 Caryophyllin, 416.
Cascara sagrada, 338.
 Cascarilla oil, 417.
Cascarillæ, 417.
 Cascarillin, 509.
 Casein, 546.
 vegetable, 548.
 Cassava, bitter, 473.
Cassia acutifolia, 501.
 elongata, 501.
 oil, 417.
Cassia pulpa, 469.
 Cast iron, 142.
Castanea, 359.
 Castile soap, 461.
Castilleja elastica, 420.

- Castor, 424.
 fiber, 424.
 oil, 464.
Castoreum, 424.
 Castorin, 424.
Cataplasma sinapis, 451.
 Catechin, 359.
Catechu, 359, 556.
 impurities in, 711.
 nigrum, 359.
 Catechuic acid, 359.
 Cathartic acid, 501.
 Cathartogenic acid, 501.
 Cathode, 245.
Caulophyllum, 509.
 Caustic, 218.
 alcohol, 443.
 lime, 107.
 lunar, 218.
 mitigated, 218.
 points, 218.
 potash, 62.
 soda, 82.
 toughened, 218.
 Cayenne pepper, 534.
 Cedra oil, 416.
 Cedrene, 419.
 Celandine, 534, 541.
 Celestine, 229.
 Cellulin, 479.
 Celluloid, 480.
 Cellulose, 479.
 of starch, 474.
 Celsius's thermometer, 596.
 Cements, 355.
 Centesimal composition, 385.
 Centiare, 604.
 Centigrade thermometer, 596.
Cephaelis ipecacuanha, 437.
Cera alba, 451.
 flava, 451.
 impurities in, 711.
 Cerasin, 479.
Cerasus virginiana, 500.
 Cerates, 589.
 Cerebrin, 546.
 Ceresine, 450.
Cerevisiæ fermentum, 439.
Cerii oxalas, 236.
 impurities in, 712.
 Cerite, 230.
 Cerium, 230.
 derivation of word, 34.
 oxalate, 230.
 Ceroleine, 450.
 Cerotic acid, 399, 489, 490.
 Ceryl cerotate, 450.
 Cerylic alcohol, 450.
Cetaceum, 450.
 Cetine, 450.
Cetraria, 338, 476.
 Cetraric acid, 338.
 Cetyl hydrate, 450.
 palmitate, 450.
 Cetylic alcohol, 450.
 Cevadilla, 543.
 Cevadilline, 543.
 Cevadine, 539, 543.
 Ceylon "moss," 479.
 Chalcedony, 354.
 Chalk, 106.
 French, 557.
 precipitated, 109.
 prepared, 111.
 stones, 589.
 Chalybeate water, 142.
 Chameleon, mineral, 232.
 Chamomile oil, 415.
 Char, 102.
 Charas, 424.
 Charcoal, 30.
 animal, 112.
 decolorizing power of, 112.
 wood, 113.
 Charles's law, 53.
Charta potassii nitratis, 286.
 sinapis, 451.
 Chartreuse, 441.
 Chaulmugra, 510.
Chavica officinarum, 541.
 Chavicic acid, 541.
 Cheese, 546.
 poison, 513, 571.
 Chelerythrine, 534.
 Chelidonic acid, 534.
 Chelidonine, 534.
 Chelidonium, 534, 541.
Chemical action, illustration of, by
 symbols, 41.
 definition of, 35, 57.
 affinity, 38.
 combination, laws of, 59.
 by volume, laws of, 53, *et seq.*
 by weight, laws of, 47, *et seq.*
 different from mechanical,
 31, 36.
 compound, 31.
 definition of, 58.
 diagrams, 46, 47, 64, 65.
 equations, 47, 63.
 definition of, 58.
 force, 38, *et seq.*
 conditions for the manifesta-
 tion of, 40.
 its relations to heat and
 electricity, 623.
 formula, 58.

- Chemical formulæ**, 42.
 notation, 41, 46.
 philosophy, principles of, 35, *et seq.*
 preparations of the British Pharmacopœia, 589.
 reagents, xiii.
 symbol, definition of, 58.
 symbols, 41.
 toxicology, 561.
- Chemicals**, lists of, xiv.
- Chemism**, 38.
- Chemist and Druggist**, 14.
- Chemistry**, art of, 13.
 definition of, 57.
 derivation of the word, 14.
 inorganic, 383.
 object of, 14.
 of carbon compounds, 383.
 organic, 383.
 science of, 14.
- Chemists**, analytical, 14.
 manufacturing, 14.
 pharmaceutical, 14.
- Chenopodium***, 420.
- Cherry-laurel water**, 500.
 sugar in, 467.
 tree gum, 479.
 wild black, 500.
- Chestnut brown**, 556.
- Chian turpentine**, 413.
- Chicory**, 476.
- Chili saltpetre**, 286.
 nitre, 286.
- Chimaphila umbellata***, 501.
- China clay**, 355.
- Chinese green**, 556.
 red, 554.
 wax, 450.
 white, 557.
 yellow, 553.
- Chinoidin**, 526.
- Chinoline**, 514.
- Chirata***, 351.
- Chiratin**, 351.
- Chiratogenin**, 351.
- Chiretta**, 351.
- Chloral**, 400, 485.
 alcoholates, 486, 488.
 butyl, 488.
 croton, 488.
 impurities in, 712.
 hydrate, 486.
 estimation of, 487.
- Chlorate of potassium**, 293.
 preparation of oxygen from, 16.
 of sodium, 294.
- Chlorates**, 293.
- Chlorates**, analytical reaction of, 295.
- Chloric acid**, 114, 292.
- Chloride of acetyl**, 396.
 aluminium, 138.
 ammonium, 91.
 antimony, 180.
 arsenum, 166.
 barium, 103.
 boron, 333.
 calcium, 106.
 removal of iron from, 106.
 chromium, 238.
 ethylene, 411.
 ethylidene, 411.
 gold, 246.
 iridium, 571.
 iron, 148, 149.
 lead, 214.
 lime, 114.
 magnesium, 118.
 manganese, 231.
 mercuric-ammonium, 206.
 mercurous-ammonium, 207, 208.
 mercury, 201.
 methyl, 399.
 palladium, 571.
 platinum and ammonium, 99, 247.
 and lithium, 228.
 and potassium, 79, 247.
 and sodium, 247.
 silicon, 355.
 silver, 217.
 sulphur, 304.
 tin, 242.
 solution of, 242.
 zinc, 132.
- Chlorides**, 265.
 estimation of, 677.
 separation of, from bromides and iodides, 276.
 tests for, 268.
- Chlorinated butyric aldehyde**, 488.
 lime, 114.
 volumetric estimation of, 657.
 soda, solution of, 87.
 volumetric estimation of, 657.
- Chlorine**, 27, 267.
 acids, 295.
 as a disinfectant, 29.
 bleaching by, 29.
 collection of, 28.
 derivation of word, 32.
 inhalation of, 29.
 its analogy to bromine and iodine, 272, 278.
 liquid, 268.

- Chlorine**, preparation of, 27.
 properties of, 28.
 relative weight of, 29.
 solubility in water, 28.
 the active agent in bleaching-
 powder, 29.
 volumetric estimation of, 656.
 water, 28, 267.
- Chlorochromic anhydride**, 240, 277.
- Chloroform**, 399.
 impurities in, 712.
 -water, 401.
- Chloronitric gas**, 289.
- Chloronitrous gas**, 289.
- Chlorophyll**, 553, 556.
- Chlorous acid**, 295.
- Chocolate**, 462.
- Cholesterin**, 460, 589.
- Choline**, 513, 552.
- Chondrine**, 549.
- Chondrus crispus*, 479.
- Christmas rose**, 505.
- Chromate of ammonium**, 238.
 barium, 105.
 calcium, 117.
 lead, 214.
 potassium, red, 238.
 standard solution of,
 648.
 yellow, 105, 238.
 and ammonium, 105.
 of silver, 220.
- Chromates**, 238.
 analytical reactions of, 240.
- Chromes**, 215.
- Chrome-alum**, 139, 238.
 ironstone, 238.
 -orange, 215.
 -red, 215, 554.
 -yellow, 215, 553.
- Chromic acid**, 238.
 anhydrous, 239.
 anhydride, 239.
 hydrate, 240.
 salts, 239.
- Chromium**, 238.
 analytical reactions of, 240.
 chloride, 238.
 derivation of word, 34.
 oxide of, 238.
 separation of, from aluminium
 and iron, 241.
 sulphate, 238.
- Chromogens in urine**, 578.
- Chromous salts**, 238.
- Chromule**, 556.
- Chrysarobin**, 338.
- Chrysarobinum*, 338.
- Chrysotropic acid**, 532.
- Chrysophan**, 338.
- Chrysophanic acid**, 338, 434.
- Churras**, 424.
- Chyme**, 551.
- Cicuta virosa*, 417.
- Cicutine**, 536.
- Cider**, 441.
- Cimicifuga racemosa*, 509.
- Cimicifugæ rhizoma*, 509.
- Cimicifugin**, 509.
- Cinchamidine**, 526.
- Cinchona calisaya*, 521.
lancifolia, 521.
officinalis, 521.
- Cinchona succirubra*, 521.
- Cinchonæ rubræ cortex*, 521.
- Cinchonicine**, 526.
- Cinchonidinæ**, 526.
 impurities in, 712.
 sulphates, 526.
 impurities in, 712.
- Cinchonidine*, 526.
 impurities in, 712.
- Cinchonina*, 525.
- Cinchoninæ sulphas*, 526.
 impurities in, 712.
- Cinchonine**, 525.
- Cinchovatine**, 526.
- Cinnabar**, 195, 554.
- Cinnamon**, 495.
- Cinnamene**, 495.
- Cinnamic acid**, 337, 494.
 alcohol, 495.
 aldehyde, 417.
 series of acids, 494.
- Cinnamol**, 495.
- Cinnamon-bark**, 417.
- Cinnamon oil**, 417.
- Cinnamyl cinnamate**, 495.
- Cissampeline**, 533.
- Cissampelos pareira*, 533.
- Citrate of ammonium**, 95.
 bismuth, 254.
 ammonium, 254.
 caffeine, 542.
 calcium, 325.
 iron and quinine, 155, 522.
 lithium, 227.
 magnesium, 122.
 nicotine, 539.
 potassium, 73.
 volumetric estimation of,
 632.
 quinine, 523.
 silver, 326.
 strychnine, 527.
- Citrate**, 323.
 analytical reactions of, 325.
- Citrenes**, 412.

- Citric acid, 323, 498.
 action of heat on, 324.
 saturating power of, 325.
 volumetric estimation of, 637.
 Citron oil, 416.
 Citronella oil, 417.
 Citronellol, 417.
 Citro-tartrate of sodium, 88.
 Citrus, 416.
 bergamia, 323.
Classification, 102, 123, 257, 393.
 Clausius's theory, 24.
Claviceps purpurea, 424.
 Clay, 138, 354.
 fire-, 202.
 ironstone, 142.
 Cloves, oil of, 416.
 Club-moss, 464.
 Coal, 241.
 brasses, 142.
 gas, 408, 454.
 for balloons, 24.
 Coal-tar, products of, 453, 557.
 colors, 557.
Cobalt, 234.
 analytical reactions of, 235.
 arsenide, 235.
 blue, 555.
 derivation of word, 34.
 glance, 235.
 hydrate, 235.
 oxide, 235, 555.
 separation of, from nickel, 237.
 sulphate, 235.
 sulphide, 235.
 Cobaltic ultramarine, 555.
 Cobalticyanide of potassium, 235.
 nickel, 237.
 Coca, 534.
 Cocaidine, 534.
Cocainæ hydrochloras, 534.
 impurities in, 712.
 Cocaine, 534.
 Cocamine, 534.
 Coccerin, 238.
Cocculus indicus, 506.
Coccus, 337.
 impurities in, 712.
 cacti, 337.
 ilicis, 182.
 Cochineal, 337, 555.
 Cocoa, 462, 543.
 nibs, 462.
 nut, 462.
 oil, 462.
Cocos nucifera, 462.
 Codamine, 518.
Codeina, 517.
Codeina, impurities in, 712.
 Codeine, 517, 520.
 Cod-liver oil, 463.
 Coffee, 542.
 Cohesion, 57.
 Coinage, copper, 191, 617.
 gold, 245, 617.
 silver, 216, 617.
 Coke, 30.
Colchici cormus, 535.
 semini, 535.
 Colchicein, 535.
 Colchicin, 535.
 Colchicine, 535.
Colchicum autumnale, 535.
 Coleothar, 153, 555.
 Collection of gases, 16, 17.
 Collidine, 513, 532.
 Collin, 705.
 Collodion, 480.
 blistering, 480.
 flexible, 480.
Collodium, 480.
 cantharidatum, 480.
 flexile, 480.
 vesicans, 480.
 Colloid bodies, 705.
Colocynthis pulpa, 502.
 Colocynthin, 502.
 Colophene, 413.
 Colopholic acid, 423.
 Colophononic acid, 423.
 hydrate, 423.
 Colophonine, 423.
 Coloring-matters, 553.
Combination, chemical, by weight, 47.
 by volume, 52, *et seq.*
 Combining proportions, 50, 198, 592, *et seq.*
 Combustible, 23.
 Combustion, 23.
 analysis for carbon and hydrogen, 688, *et seq.*
 for nitrogen, 691.
 definition of, 58.
 spontaneous, 158.
 supporters of, 23.
 Composition of atmosphere, 27.
 bismuth salts, 252, 254.
 centesimal, 385.
 empirical, 385.
 molecular, 386.
 oils and fats, 459.
 organic compounds, 384.
 Compounds, chemical, 31.
 definition of, 58.
 different from mechanical, 36.

- Compounds**, 36.
 of the elements, 60.
 Conchinine, 524.
 Concrete oil of mangosteen, 463.
 Condensation, 128.
 Condenser, 128.
 Condensing-tub, 128.
 worm, 128.
 Confections, 470.
 Conhydrine, 536.
 Conia, 536.
 Conicine, 536.
Conii folia, 536.
 Conine, 532, 536.
Conium, 536.
maculatum, 536.
 Conquinine, 524.
 Constant proportions, law of, 48.
 Constant white, 557.
Constitution. See also "Structure."
Constitution as indicated by—
 chemical "periodicity," 380.
 properties, 387, 395.
 density of gases and vapors,
 53, 55, 619.
 electricity, etc., 623.
 isomorphism, 55.
 specific heat, 622.
 substitution, 394.
Constitution of alkaloids, 510,
 513.
 benzene series, 432.
 bleaching powder, 114.
 magnesium carbonate, 121.
 matter, 42, 130.
 molecular, 130, 138, 139, 193,
 386, 390.
 organic compounds, 387, 393,
 396.
 of salts, 66, 87, 124, 286, 299,
 378, 387, 390, 396, 513.
 Constitutional formula, 386.
 Construction of formulæ, 46, 389,
 392, 396.
 Convolvulin, 505.
 Convolvulinol, 505.
Convolvulus scammonia, 508.
 Conylia, 536.
Copaiba, 426.
 impurities in, 712.
 Copaiva, 426.
 oil, 417.
 Copaivaöl, 426.
 Copaivic acid, 426.
 Copal, 424.
Copper, 190.
 acetate, 192.
 ammonio-sulphate, 178, 193,
 207.
Copper, analytical reactions of, 192.
 antidotes to, 194.
 arsenate, 177.
 arsenite, 177, 194.
 black oxide of, 192.
 blue, 555.
 carbonate, 190.
 coinage, 191.
 derivation of word, 33.
 detection of arsenum in, 173.
 ferrocyanide, 194.
 flame, 194.
 foil, 192.
 hydrate, 194.
 hydride, 345.
 in organic mixtures, detection
 of, 564.
 iodide, 276.
 melting-point of, 599.
 metallic, 190.
 nitrate, 192.
 oxide, 191.
 oxyacetate, 192.
 pyrites, 190.
 quantitative estimation of, 671.
 quantivalence of, 191.
 subacetate, 192.
 sulphate, 192.
 anhydrous, 192.
 sulphide, 192.
 zinc couple, 679.
 Copperas, blue, 144, 192.
 green, 144.
 Coptis-root, 533.
 Coriander oil, 417.
 Cork, specific gravity of, 618.
 borers, 16.
 Cornic acid, 338.
 Cornin, 338.
 Cornutene, 424.
 Corpse fat, 548.
 Correction of the volume of a gas
 for pressure, 619.
 for temperature, 619.
 Correlation of force, 38.
 matter, 38.
 Corrosive sublimate, 201.
 test for, in calomel, 203.
Cortex pruni serotinæ, 500.
 Corydaline, 536.
Corydalis, 536.
 Cotarnine, 518.
 Coto-bark, 502.
 Cotoin, 502.
 Cotton-seed oil in olive oil, 459.
 cake, 515.
 Cotton-wool, 479.
 Couch-grass, 511.
 Coumarin, 494.

- Cowbane, 417.
 Cowhage, 242.
 Cow's milk, 546.
 Cranesbill, 360.
 Cream, 547.
 of tartar, 62, 80, 318.
 soluble, 334.
 Creasol, 453.
 Creasote, 453.
Creasotum, 453.
 impurities in, 712.
 Creatine, 512.
 Creatinine, 512.
 Cremnitz white, 557.
 Cresol, 453, 455.
 Cresotic acid, 493.
 Cresylic acid, 453.
Creta, 111.
 præparata, 111.
 impurities in, 712.
Crini radix, 508.
Crinum asiaticum, 508.
 Crith, 621.
 Crocetin, 554.
 Crocin, 554.
 Crocus (mineral), 153.
 (vegetable), 554.
 impurities in, 712.
 of antimony, 180.
Crocus sativus, 554.
 Croton chloral, 488.
 hydrate, 488.
 oil, 463.
 Crotonic acid, 491.
 Crotonylene, 411.
 Crucibles, 70, 355.
 Crude antimony, 180.
 potashes, 62.
 Crum's test for manganese, 234.
 Cruso-creatinine, 513.
 Cryohydrates, 81.
 Cryolite, 376.
 Cryptopine, 518.
 Crystallization, water of, 86.
 fractional, 78, 378.
 Crystalloid bodies, 705.
 Cubeb pepper, 541.
Cubeba, 541.
 Cubebene, 417.
 Cubebin, 541.
 Cubebs, oil of, 417.
 oleo-resin, 428.
 Cubic inches in a gallon, 621.
 nitre, 286.
 Cuca. See Coca.
Cucurbita pepo, 510.
 Cudbear, 555.
 Culver's root, 510.
 Cumin, 417.
 Cuminic acid, 417.
Cuminum cyminum, 417.
 Cummin, 417.
 Cupel, 676.
 Cupellation, estimation of silver
 by, 676.
 Cupr-diammon - diammonium, sul-
 phate of, 207.
 Cuprea-bark, 521.
 Cupreine, 526.
Cupri sulphas, 192.
 impurities, 712.
 nitras, 192.
Cupric arsenite, 177, 194.
 compounds, 191.
 ferrocyanide, 194.
 hydrate, 194.
 nitrate, 192.
 oxide, 191.
 sulphate, 192.
 sulphide, 192.
 Cuprous hydride, 345.
 iodide, 191.
 oxide, 191, 468.
Cuprum, 190.
 Curacao, 441.
 Curari, 528.
 Curarine, 528.
Curcuma longa, 420, 554.
 Curcumin, 554.
 Curd soap, 461.
 Curds, 546.
 and whey, 471, 546.
 Currants, sugar in, 467.
 Curry powder, odor and flavor of,
 420.
Cuspariæ cortex, 536.
 Cusparidine, 536.
 Cusparine, 536.
Cusso, 424.
 Cutch, 359.
 Cyanates, 338.
 Cyanic acid, 338.
 Cyanide of allyl, 451.
 mercury, 280.
 nickel, 237.
 potassium, 280.
 silver, 220, 283.
 Cyanides, 280.
 analytical reactions of metallic,
 283.
 antidote to, 283.
 double, 280.
 quantitative estimation of, 641,
 678.
Cyanogen, 162, 279.
 Cyanurets. *Vide* Cyanides.
 Cyder. See Cider.
 Cymene, 415, 417, 429, 432.

- Cymol, 415.
 Cypripedin, 510.
Cypripedium pubescens, 510.
 Cystin, 581.
 Cytisine, 436.
- DAHLIA**, 476.
 Dalleiochin, 523.
 Dalton's atomic theory, 51.
 laws, 47, 48, 49.
 Dambose, 469.
 Dandelion, 476.
Daphne laureola, 425.
 mezereum, 425, 502.
 Daphnetin, 502.
 Daphnin, 502.
Datura alba, 538.
 Daturia, or Daturin, 538.
 Daughlish's bread, 470.
 Davy safety-lamp, 24.
 Deadly nightshade, 531.
 Decane, 398.
 Decantation, 110.
 Decimal coinage, 605.
 weights, 602.
 Decoction, 589.
 Decolorizing power of animal charcoal, 113.
 Decrepitation, 372.
 Decylene alcohol, 452.
- Definition** of an atom, 57.
 atomic weights, 59.
 a chemical compound, 58.
 equation or diagram, 58.
 formula, 58.
 symbol, 58.
 a gas, 58.
 a liquid, 58.
 a mixture, 57.
 a molecule, 57.
 a solid, 58.
 an element, 57.
 chemical action, 57.
 force, 57.
 chemistry, 57.
 combustion, 58.
 law of diffusion, 58.
 molecular weights, 59.
 quantivalence of atoms, 59.
- Deflagrating flux, 377.
 Deflagration, 75.
 Deliquescence, 89.
 Delphine, Delphinine, or Delphia, 536.
Delphinium staphysagri, 536.
 Density, 614.
 of vapors, 619.
 Dentifrices, action of, 242.
 Deodorizers, 29.
- Deodorizing liquid, 132.
 power of animal charcoal, 113.
 Deoxidizers, 645.
 Deposits, urinary, 579.
 Derivations of names of elements, 32, *et seq.*
 Desiccation, 661, 687.
 Desiccators, 661.
 Destructive distillation, 129.
 Detonation, 75.
 De Valangin's solution, 168.
 Dextrin, 476.
 Dextrogyrate, 469.
 Dextroracemic acid, 320.
 Dextrose, 467.
 Dextrotartaric acid, 320.
 Dhak tree, 359.
 Dhatura, 538.
 Diabetic urine, 575.
 Diacid alcohols, 456.
 Diamide, 515.
 Dicarbonyl benzene, 521.
 Diagram, chemical definition of, 63.
Diagrams, chemical, 42, 63.
 Dialysate, 705.
 Dialysis, 705.
 Dialytic iron, 705.
 Dialyzed iron, 160, 705.
 Diamines, 512.
 Diamond, 30.
 Diaphragms, 71.
 Diastase, 477.
 action of, upon starch, 477.
 Diazo, 513.
 Diazobenzene, 513.
 Dibasic acids, 264, 495.
 Dibasyloous radicals, 264.
 Dibromethane, 409.
Dicentra formosa, 536.
 Dichloromethane, 399.
 Dichloromethylbenzene, 432.
Dichopsis gutta, 420.
 Didymium, 230.
 Dietetics, 15.
 Diethyl, 397.
 Diethylamine, 511.
 Diethylsulphon - dimethylmethane, 446.
 Diffusate, 705.
 Diffusion of gases, 24.
 law of, definition of, 24, 58.
 Digallic acid, 357.
 Digitalein, 503.
 Digitalin, 502.
 Digitaliretin, 502.
Digitalis, 502.
 folia, 502.
 Digitin, 503.
 Digitogenin, 503.

- Digitonin, 503.
 Digitoxin, 503.
 Dihydric alcohols, 409, 456.
 Dihydroxyacetic acid, 490.
 Dihydroxybenzoic acid, 493.
 Dihydroxybutyric acid, 490.
 Dihydroxyl benzenes, 455.
 derivatives of the hydrocarbons, 456.
 Dihydroxypropionic acid, 490.
 Dihydroxysuccinic acid, 496.
 Di-iodo-paraphenol-sulphonic acid, 445.
 Di-iodosalicylic acid, 493.
 Diketone, 521.
 Dill oil, 415.
 Dipentene, 412.
 Dimeric-ammmonium, 207.
 Dimethyl, 397.
 Dimethyl-ethyl-carbonol, 450.
 Dinitrocellulin, 479.
 Diosphenol, 416.
Diospyri fructus, 360.
Diospyros embryopteris, 360.
Dipterocarpi balsamum, 426.
Dipterocarpus lævis, 426.
 turbinatus, 426.
 Disinfectant, chlorine as a, 29.
Disinfectants, 29.
 Disinfecting fluid, Burnett's, 132.
 carbolic acid, 453.
 Condy's, 78, 232.
 powder, 114.
 solution, 114.
 Dissociation, 622.
 Distillation, 127.
 destructive, 129.
 dry, 129.
 fractional, 378, 439.
 Distilled water, 129.
 Disulphide of carbon, 314.
 Dita, 537.
 Dитай, 537.
 Ditamine, 537.
 Dithionic acid, 346.
 Dock, 338.
 Dolomite, 118.
 Donovan's solution, 167.
Dorema ammoniacum, 427.
 Double chloride of aluminium and sodium, 139.
 cyanides, 280.
 salts, 79, 139.
 Doundaké bark, 424.
 Dover's powder, 537.
 Draconyl, 495.
 Dragon's blood, 424.
 Dried alum, 140.
 Drops, 608.
 Dry distillation, 129.
 Drying apparatus, 112.
 in vacuo, 112.
 oils, 463.
 precipitates, 112.
Dryobalanops aromatica, 421.
Duboisia myoporoides, 538.
 Duboisine, 538.
Dulcamara, 541.
 Dulcamarin, 541.
 Dulcin, 431.
 Dulcite, 465.
 Dulong and Petit's law, 623.
 Dutch camphor, 421.
 liquid, 411.
 Dyads, 123.
 Dyeing by mordants, 140.
 Dyer's saffron, 555.
 Dynamic electricity, production of, 132.
 Dynamicity, 56.
 Dynamite, 458.

 EARTH, bone-, 112, 327.
 Earthenware, 354.
 Earth-nut oil, 464.
 Earth pitch, 427.
 Earths, alkaline, 127.
 Eau de Cologne, 405.
 Eau de Javelle, 88.
 Ebonite, 420.
 Ebullition, 282.
Ecballii fructus, 503.
 Ecboline, 424.
 Ecgonine, 534.
Echites scholaris, 537.
 Effervescing citrate of magnesia, 88, 121.
 citro-tartrate of sodium, 88.
 soda-water, 86.
 solution of potash, 72.
 sulphate of magnesium, 119.
 tart. soda powder, 321.
 Efflorescence, 89.
 Egg, oil of yelk of, 545.
 white of, 544.
 Elæometer, 616.
 Elæoptens, 413.
 Elaidic acid, 491.
Elastica, 421.
 Elaterin, 503.
Elaterium, 503.
 impurities in, 503, 712.
 Elder-flower oil, 419.
 Elecampane, 418.
 Electric amalgam, 196.
 Electricity, production of dynamic, 132.
 related to chemical action, 623.

- Electrolysis, 22, 623.
 Element, definition of, 57.
Elements, 13, 15, 32, 380.
 and their compounds, 60.
 classification of, according to analogy, 102.
 etymology of names of, 32, *et seq.*
 metallic, 15.
 non-metallic, 16.
 of medical or pharmaceutical interest, 15.
 quantivalence of, 123.
 symbols of, 32, 58.
 atomic values, and weights of the, 725.
 of, and derivation of name of, 32, *et seq.*
- Elemi, 426.
 Elutriation, 134.
 fractional, 378.
 Emerald green, 536.
 Emetine, 537.
 Emodin, 338.
 Empirical formulæ, 385.
 composition, 385.
Emplastra, 213, 423.
Emplastrum menthol, 418.
plumbi, 213.
iodidi, 213.
saponis fuscum, 298.
 Emulsin, 500.
 Emulsions, 428.
 Enemas, 589.
 English red, 555.
 blue, 555.
 Enzymes, 441.
 Eosin, 434.
 Epsom salt, 118.
 Equation, chemical definition of, 58.
 Equations, 46, 64.
 Equisetic acid, 324.
 Equivalence, 56.
 Equivalents, 726.
 Erbium, 725.
 Ergosterin, 424.
 Ergot, 424.
Ergota, 424.
 Ergotin, 424.
 Ergotine, 424.
 Ergotinic acid, 424.
 Ergotinine, 424.
 Ergotinum, 424.
 Ericolin, 501.
 Erlangen blue, 556.
 Error in experiment, 693.
 Erucic acid, 464.
 Erythrite, 465, 496.
 Erythrorretine, 338.
Erythroxyton coca, 534.
 Esculin. See *Æsculin*.
 Eseridine, 540.
 Eserine, 540.
 Essence of aniseed, 414.
 apple, 407.
 ginger, 420.
 greengage, 407.
 melon, 407.
 mirbane, 430.
 mulberry, 407.
 peppermint, 414.
 pineapple, 407.
 quince, 407.
 Essences, 415.
Essentia anisi, 414.
menthæ piperitæ, 414.
 Essential oils. See *Oils*.
 Ester, 484.
 Etching, 343.
 Ethal, 450.
 Ethane, 397.
 substitution products of, 402.
 Ether, 446.
 acetic, 300, 406.
 hydrobromic, 402.
 nitrous, 351.
 petroleum, 398.
 Etheral salts, 407, 483.
 Ethers, 444, 484.
 mixed, 447.
 sulphur, 447.
 Ethiop's mineral, 208.
Ethyl, acetate, 300, 406
 bromide, 402.
 butyrate, 407.
 carbamate, 489.
 formic acid, 488.
 group, 392.
 hydrate, 436.
 hydride, 397.
 hydrogen sulphate, 439, 447.
 hydroxylamine, 515.
 iodide, 402.
 nitrite, 403.
 oenanthylate, 407.
 oxide, 445.
 pelargonate, 407.
 sebacate, 407.
 suberate, 407.
 sulphuric acid, 447.
 Ethylamine, 511.
 Ethylate of sodium, 443.
 Ethylene, 408.
 chloride, 411.
 dibromide, 409.
 hydrate, 436.
 sulphate, 409.
 Ethylic acid, 490.

- Ethylic alcohol, 439.
 bromide, 402.
 iodide, 402.
 series of alcohols, 436.
 Ethylidene, chloride of, 411.
 compounds, 489.
 lactic acid, 489.
 Ethylsulphonic acid, 445.
 Ethylsulphuric acid, 447.
 Etymology of names of elements,
 32.
 Eucalypti gummi, 418.
 rostrata, 418.
Eucalyptol, 418.
 impurities in, 712.
 Eucalyptus oil, 417.
 var., 359.
Eucalyptus globulus, 417.
 Euchlorine, 296.
 Eudiometry, 568.
 Eugeniac acid, 416.
 Eugenin, 416.
 Eugenol, 416.
 Euodic aldehyde, 419.
 Euonymi cortex, 509.
 Euonymin, 510.
Euonymus atropurpureus, 510.
Eupatorium perfoliatum, 509.
 Euphorbium, 427.
 Euphorbon, 427.
 Euxanthate of magnesium, 554.
 Evaporation, 71.
 Everett's yellow salt, 282.
 Ewe's milk, 548.
Examinations of the Pharmaceuti-
 cal Society of Great Britain,
 14. *Vide* prefatory matter,
 xi.
 Experimental error, 693.
 Explosions of gas, 22.
 Extract of malt, 478.
 Extracts, 589.
Extractum cinchonæ liquidum, 694.
 euonymi siccum, 509.
 glycyrrhizæ purum, 504.
 fluidum, 504.
 hamamelidis liquidum, 509.
 nucis vomicæ, 700.
 saturni, 212.
FACE-ROUGE, 337.
 Fæces, 573.
 Fahrenheit's thermometer, 596.
Farina tritici, 473.
Fats and oils, composition of, 459.
 Fats, etc., analysis of, 706.
 solid, 462.
 to determine the melting-point
 of, 597.
 Fatty acids, 462.
 bodies, 458.
 series of, 395.
 Fehling's solution, 701.
Fel bovinum purificatum, 532.
Fel bovis, 532.
 Felspar, 355, 376.
 Fenchene, 412.
 Fennel oil, 418.
 Fenugreek, 443.
Fer réduit, 158.
Fermentation, 440.
 acetic, 440.
 alcoholic, 440.
 ammoniacal, 440.
 butyric, 440.
 by soluble albumenoids, 440.
 lactic, 440.
 mannitic, 440.
 putrefactive, 440.
 viscous, 440.
 Ferments, 441.
 Ferrate of potassium, 143.
Ferri acetatis, liquor, 154.
 fortior, 154.
 tinctura, 154.
 arsenas, 146, 170.
 carbonas, 145, 153.
 saccharatus, 145.
 impurities in, 712.
 citras, 154.
 liquor, 154.
et ammonii citras, 45.
 impurities in, 712.
 quantitative estima-
 tion of iron in, 669.
et ammonii sulphas, 140.
 impurities in, 712.
et ammonii tartras, 157.
et potassii tartras, 156.
 impurities in, 712.
et quininæ citras, 156, 522.
 impurities in, 712.
 quantitative estima-
 tion of quinine in,
 697.
et quinnina citras solubilis, 155.
 impurities in, 713.
et strychninæ citras, 154.
 impurities in, 713.
hypophosphis, 344.
 impurities in, 713.
iodidum saccharatum, 148.
 impurities in, 713.
lactas, 348.
 impurities in, 713.
oxidum hydratum, 151.
perchloridi, liquor, 150.
pernitratiss, liquor, 158.

- Ferri peroxidum hydratum*, 151.
persulphatis, liquor, 151.
phosphas solubilis, 146, 157.
 impurities in, 712.
pulvis, 158.
subcarbonas, 145, 153.
sulphas granulatus, 144.
 impurities in, 713.
 exsiccatus, 144.
 granulata, 144.
 impurities in, 713.
valerianas, 363.
 impurities in, 713.
- Ferric** acetate, 154, 300.
 aceto-nitrate, 158.
 benzoate, 337.
 chloride, 148.
 hydrous, 149.
 citrate, 155.
 hydrate, 151.
 hypophosphite, 344.
 vol. est. of, 654.
 iodate, 296.
 nitrate, 157.
 oxide, 152, 154.
 separated from phosphates
 and oxalates, 375.
 oxyhydrate, 157.
 oxyiodate, 296.
 oxysulphate, 145.
 phosphate, 157, 331.
 salts, 143, *et seq.*
 analytical reactions of, 161.
 sulphate, 150.
 sulphocyanate, 162, 356.
 valerianate, 363.
- Ferricyanide of potassium, 341.
 Ferricyanides, 341.
 Ferricyanogen, 162, 342.
 Ferrocyanide of potassium, 280,
 341.
 of zinc, 137.
 Ferrocyanides, 341.
 Ferrocyanogen, 162, 341.
- Ferrous** arsenate, 146, 170.
 ammonium sulphate, 649.
 arsenate, vol. est. of, 650.
 bromide, 148.
 carbonate, 145.
 vol. est. of, 650.
 chloride, 149.
 anhydrous, 149.
 citrate, 157.
 hydrate, 161.
 iodide, 147.
 lactate, 348.
 phosphate, 146.
 vol. est. of, 651.
 salts, 144, *et seq.*
- Ferrous** salts, analytical reactions
 of, 160.
 sulphate, 144.
 vol. est. of, 650, 653.
 sulphide, 31, 148, 160, 162.
 tartrate, 157.
- Ferrum*, 142.
 reductum, 158, 669.
 impurities in, 713.
 tartaratum, 154, 156.
 est. of iron in, 669.
- Ferula Narthex*, 427.
 Ferulaic acid, 427.
- Fibrin, 545.
 vegetable, 548.
- Ficus*, 467.
 elastica, 420.
- Fig, 467.
- Filicic acid, 464.
- Filix mas*, 464.
- Filter, to dry, 660.
- Filtering-paper, 109.
- Filters, 109.
- Filtrate, 125.
- Fine gold, 245.
- Fire-clay, 202.
- Fire-damp, 396.
- Fir wool, 413.
 oil, 413.
- Fish-poison, 513.
- Fixed oils, 463.
 and volatile oils, difference
 between, 463.
- Flag, blue, 509.
- Flake white, 557.
- Flame, oxidizing, 135.
 reducing, 136.
 structure of, 23.
- Flare, 462.
- Flashing-point, 413.
- Flaxseed, 479.
- Fleitmunn's test for arsenum, 175.
- Flexible collodion, 480.
- Flint, 354.
- Flores zinci*, 135.
- Flour, 472.
- Flowers of sulphur, 301.
- "Fluid magnesia," 120.
- Fluorescein, 434.
- Fluoric acid, 342.
- Fluoride of boron, 333.
 calcium, 106.
 in bones, 112.
 lithium, 227.
 silicon, 342.
- Fluorides, 342.
- Fluorine, 343.
 derivation of word, 33.
- Fluor-spar, 106, 342.

- Fœniculum*, 418.
 Foil, copper, 191.
 tin, 241.
Food, analysis of, 706.
 elements of, 548.
 how disposed of in the bodies
 of animals, 548.
 Force, chemical, 37, *et seq.*
 origin of, 38.
 Formates, 339.
 Formic acid, 339, 484, 490.
Formica rufa, 339.
 Formula, chemical, definition of,
 58.
 official, 29.
 officinal, 29.
Formulæ, 42, 723.
 constitutional, 386.
 construction of, 56.
 empirical, 386.
 graphic, 138.
 molecular, 55, 386.
 rational, 386.
 structural, 138, 386, 390, 430.
 two-volume, 55, 386.
 typical, 379.
 why used at all, 391.
 Fousel oil, 449.
 Fowler's solution, 168.
 Foxglove, 502.
 Fractional distillation, 378, 439.
 crystallization, 78, 378.
 elutriation, 134, 378.
 fusion, 378.
 lixiviation, 89, 378.
 operations, 378.
 precipitation, 79, 378.
 sifting, 372, 378.
 solution, 89.
 sublimation, 93, 378.
 Frangula, 502.
 bark, 338.
 Frangulin, 338, 503.
Fructus pychotis, 415.
 Frankincense, Arabian, 428.
 common, 428.
Fraxinus ornus, 465.
 Free acids, 367.
 estimated, 636.
 Freezing-mixture, 305.
 French chalk, 557.
 turpentine, 412.
 Fruit-essences, 407.
 Fuchsine, 557.
 Fulminating silver, 219.
 mercury, 219.
 Fume-cupboard, 97.
 Fumerolles, 333.
 Fuming sulphuric acid, 310.
 Funnel-tubes, 21.
 "Fur" in water-vessels, 315.
 Furniture of a laboratory, xiii.
 Furze, 536.
 Fusel oil, 449.
 Fusibility of metals, table of the,
 599.
 Fusible white precipitate, 206.
 Fusing-points of fats, 598.
 Fusion, fractional, 378.
 Fustic, 553.
 GAB tree, 360.
 Gadinine, 513.
 Galactose, 471.
Galbanum, 427.
 Galena, 210.
 argentiferous, 216.
 Galenical preparations of the
 British Pharmacopœia, 589.
Galipea cusparia, 536.
 Galipine, 536.
 Galipot, 423.
 Gall of the ox, 552.
Galla, 357.
 Gallic acid, 460, 493, 494.
 Gallium, 725.
 Gallon, 668.
 Gallotannic acid, 357, 494.
 Galls, Aleppo, 357.
 English, 357.
 Gall-stones, 589.
 Galvanic test for mercury, 208.
 Galvanized iron, 131.
 Gambier, 359.
 Gamboge, 427.
 Gambogic acid, 427.
 Ganja, 423.
 Garancin, 554.
Garcinia indica, 463.
 Hanburii, 427.
 oil, 462.
 pictoria, 427.
 purpurea, 463.
Garciniæ purpurea oleum, 463.
 Garden thyme, 417.
 Garlic, essential oil of, 452.
 Gas analysis, 361, 550.
 -burners, 17, 23.
 coal, 408, 454, 557.
 definition of, 59.
 for balloons, coal-, 24.
 lamp, 17.
Gases and vapors, analysis of, 361.
 collection of, 16, 17.
 correction of the volume of, 619.
 for pressure, 619.
 for temperature, 619.
 diffusion of, 24.

- Gases**, law of solubility of, in water, 86.
 relation of, to liquids and solids, 42.
 specific gravity of, 619.
Gastric juice, 550.
 artificial, 550.
Gaultheria procumbens, 407.
Gaultheric acid, 407.
Gay-Lussac's law, 52.
Gelatinogenous substances, 549.
Gelatin, 549.
 sugar, 552.
 tests for, 550.
 vegetable, 479.
Gelatinum, 555.
Gelsemine, 537.
Gelseminic acid, 537.
Gelseminine, 537.
Gelsemium, 537.
Gentian bitter, 503.
Gentiana lutea, 503.
Gentianæ radix, 503.
Gentianic acid, 504.
Gentiogenin, 504.
Gentiopierin, 503.
Gentisic acid, 504.
Gentisin, 504.
Geraniol, 418.
Geranium maculatum, 360.
 oil, 418.
German silver, 131, 236.
Ghatti, 479.
Gin, 441.
Gingelly oil, 464.
Ginger, essence of, 420.
 -grass oil, 418.
 oil, 420.
Gingerol, 534.
Girdwood and Roger's method for detecting strychnine, 567.
Glacial acetic acid, 299.
 phosphoric acid, 330.
Glass, 355.
 liquid, 355.
 of antimony, 180.
 rods, 70.
 soluble, 355.
 tubes, to bend, 17.
 to cut, 17.
 to draw out, 109.
Globulin, 546.
Globulins, 549.
Glonoine, 458.
Glucinum, 725.
Glucose, 467.
Glucoses, 467.
Glucosides, 499.
Glucosimide, 445.
Glue, 549.
Gluside, 445.
Glusidum, 445.
Glutanic acid, 497.
Gluten and Glutin, 473.
Glyceric acid, 490.
 hydrate, 436.
Glycerin, 213, 436, 457.
 test for, 457.
Glycerinum, 457.
 impurities in, 713.
 acidi carbolici, 453, 458.
 gallici, 458.
 tannici, 357, 458.
 aluminis, 458.
 amylī, 458.
 boracis, 333, 458.
Glyceritum boroglycerini, 334.
 plumbia subacetatis, 212.
Glycerols, 457.
Glycyl, 457.
Glyceryl, 457.
 borate, 335.
 caproate, 462.
 caprylate, 462.
 hydrate, 459.
 laurate, 462.
 margarate, 462.
 myristate, 462.
 oleate, 459.
 palmitate, 462.
 ricinoleate, 464.
 rutate, 462.
 tristearate, 459.
Glycocholates, 552.
Glycocine, 552.
Glycocoll, 552.
Glycogen, 476.
Glycol, 409, 456.
 trichlorethylidene, 486.
Glycols, 409, 456.
Glycols, aromatic, 456.
Glycollic acid, 409, 489.
 aldehyde, 409.
Glycyrrhetin, 504.
Glycyrrhizæ radix, 471, 504.
Glycyrrhizic acid, 504.
Glycyrrhizin, 504.
Glycyrrhizinum ammoniatum, 504.
 impurities in, 713.
Glyoxal, 409.
Glyoxylic acid, 490.
 series, 490.
Guoscopine, 518.
Goa powder, 338.
Gold, 245.
 analytical reactions of, 246.
 and sodium chloride, 246.
 coin, 245.

- Gold**, derivation of word, 34.
 earth, 553.
 fine, 245.
 jewellers', 245.
 leaf, 245.
 ochre, 553.
 perchloride, 246.
 sulphide, 246.
 yellow, 553.
Golden seal, 533.
 syrup, 471.
Gossypium purificatum, 479.
 impurities in, 713.
Gothite, 153.
Goulard's cerate, 212.
 extract, 212.
 water, 212.
Gracillaria, 479.
Graham's dialytic process, 704.
 law of diffusion, 24, 58.
Grain, 608.
Grains of paradise, 418.
Gramme, 603.
 relation of, to grains, 610.
Granati cortex, 359.
 radicis cortex, 359.
Granatum, 359.
Granite, 138.
Granulated phosphorus, 328.
 sulphate of iron, 144.
 tin, 242.
 zinc, 20.
Granulose, 473.
Grape-sugar, 318, 467.
Grapes, dried, 318.
 sugar in, 467.
Graphic formulæ, 138, 389, 392, 397.
Graphite, 30.
Grass oils, 417, 418, 420.
Gravel, 579.
Gravimetric quantitative analysis, 659.
Gravitation, 600.
Gravity, 600.
Gray powder, 195.
Green chloride of iron, 149.
 Chinese, 556.
 copperas, 144.
 emerald, 556.
 iodide of iron, 147.
 mercury, 197.
 pigments, 556.
 sulphate of iron, 144.
 ultramarine, 556.
 vitriol, 144.
Greengage essence, 407.
Griffith's mixture, 146.
Grindelia robusta, 537.
Grindeline, 537.
Ground-nut oil, 464.
Group tests, 258.
Guaiaci lignum, 504.
 resini, 504.
Guaiacin, 504.
Guaiacol, 453.
Guaiaconic acid, 504.
Guaiacum, resin of, 504.
Guaiaretic acid, 504.
Guaiaretin, 504.
Guaiaretinic acid, 504.
Guanine, 515.
Guano, 362.
Guarana, 542.
Guaranine, 542.
Guaza, 424.
Guilandina bonducella, 509.
Guinea grains, 418.
Gulancho, 509.
Gum, 116, 478.
 acacia, 116, 478.
 Arabic, 116, 478.
 benjamin, 335.
 British, 477.
 cherry-tree, 479.
 resins, 422, 427.
 tragacanth, 116, 479.
Gummate of calcium, 116.
 lead, 116.
Gummic acid, 479.
Gunj, 504.
Gunjah, 423.
Gun-cotton, 479.
 metal, 241.
Gunpowder, 291.
Gurjun balsam, 426.
Gutta percha, 420.
Guttæ, 608.
Gynocardia odorata, 509.
Gypsum, 106, 116.
HAMAMELIDIS cortex, 509.
 folia, 509.
Hæmatein, 555.
Hæmatin, 546.
Hæmatite, brown, 142, 153.
 red, 142, 153.
Hæmatoxylin, 555.
Hæmatoxylin, 359, 555.
Half-sovereign, weight of the, 246.
Haloid salts, 287.
Hamamelis virginiana, 509.
Hambro' blue, 555.
Hard soap, 461.
Hardness of water, 315.
Hashish, 424.
Heat, latent, 86, 129.
 related to chemical action, 622.
 source of, 17.

- Heat**, specific, 130.
Heavy carbonate of magnesium,
 119.
 magnesia, 122.
 spar, 103.
 white, 103.
Heberden's ink, 162.
Hectare, 604.
Hedeoma pulegioides, 418.
Hedeomol, 418.
Helenin, 418.
Heliotrope, 507.
Hellebore, black, 505.
 green, 505, 539.
 white, 539.
 American, 539.
Helleborein, 505.
Helleborin, 505.
Helleborus niger, 505.
 viridis, 505.
Hemialbumose, 551.
Hemidesmi radix, 340.
Hemidesmic acid, 340.
Hemlock, 417.
Hemp, Canadian, 509.
 Indian, 423.
Hempseed calculi, 589.
Henbane, 538.
Henry and Dalton's laws, 86.
Heptane, 398.
Heptoic aldehyde, 417.
Heptylene, 408.
Heptylic acid, 490.
Herapathite, 524.
Hesperidene, 416.
Hesperidin, 509.
Hevea (Siphonia) Brasiliensis, 420.
Hexabasic acids, 498.
Hexabromobenzene, 430.
Hexachlorobenzene, 430.
Hexads, 124.
Hexahydric alcohols, 465.
Hexahydropyridine, 541.
Hexahydrodipyridyl, 539.
Hexane, 398.
Hexylene, 408.
Hexylic acid, 490.
Hibisci capsulæ, 479.
Hibiscus esculentus, 479.
Hippuric acid, 340, 580.
Hips, 469.
Hoffner's blue, 556.
Hollway's smelting process, 196.
Homatropinæ hydrochloras, 531.
 hydrobromas, 531.
Homatropine, 531.
Homoche lidine, 541.
Homologous series, 394.
Homology, 394.
Homopterocarpin, 555.
Homoquinine, 527.
Homotartaric acid, 497.
Honey, 469.
Honeydew, 471.
Hop, 427, 515, 539.
 essential oil of, 427.
Hordeum decorticatum, 473.
 starch of (fig.), 475.
Horehound, 510.
Horse-chestnut, 537.
Horseradish oil, 415.
Huile de Cade, 427.
Humulus, 427.
 lupulus, 427, 539.
Hydrargyri chloridum corrosivum,
 201.
 impurities in, 713.
 mite, 203.
 impurities in, 713.
 cyanidum, 280.
 impurities in, 713.
 iodidum flavum, 197.
 impurities in, 713.
 rubrum, 199.
 impurities in, 713.
 nitratis acidus, liquor, 200.
 oxidum flavum, 204.
 impurities in, 713.
 rubrum, 204.
 impurities in, 713.
 perchloridum, 202.
 subchloridum, 203.
 subsulphas flavus, 228.
 impurities in, 714.
 sulphuretum cum sulphure, 208.
Hydrargyrum, 33, 195.
 impurities in, 714.
 ammoniatum, 206.
 impurities in, 714.
 cum creta, 195.
 impurities in, 714.
Hydrastine, 535.
Hydrastininæ hydrochloras, 538.
 impurities in, 714.
Hydrastinine, 538.
Hydrastis canadensis, 533.
 rhizoma, 533.
Hydrate of aluminium, 140.
 ammonium, 92.
 barium, 103.
 benzoyl, 493, 534.
 bismuth, 254.
 bromal, 488.
 butyl chloral, 488.
 cadmium, 250.
 calcium, 107.
 cetyl, 450.
 chloral, 486.

- Hydrate** of chromium, 240.
cobalt, 235.
copper, 193, 194.
lime, 107.
manganese, 234.
nickel, 237.
potassium, 62.
sodium, 82.
zinc, 136.
- Hydrated oxide of iron**, 151, 152.
substances, 85.
- Hydrates**, composition of, 66.
identified, 377.
of the hydrocarbons, 436.
- Hydraulic cement**, 355.
- Hydrazine**, 515.
- Hydrazobenzene**, 430.
- Hydric acetate**, chloride, nitrate, sulphate, etc. See the respective acids, Acetic, Hydrochloric, etc.
- Hydride of antimony**, 185.
arsenum, 174.
benzoyl, 492.
copper, 345.
ethyl, 397.
methyl, 396.
phosphorus, 343.
silicon, 355.
- Hydrides**, 124.
- Hydriodic acid**, 272.
volumetric estimation of, 644.
- Hydrium*, 21.
- Hydrobromic acid**, 269.
ether, 402.
- Hydrocarbons**, 391.
acetylene series, 411.
anthracene series, 434.
basylous, 391.
benzene series, 429.
dihydroxyl derivatives, 456.
monohydroxyl derivatives, 436.
naphthalene series, 433.
neutral, 391.
normal, 391.
olefine series, 408.
paraffin series, 396.
polyhydroxyl derivatives, 465.
saturated, 392.
series of, 394.
terpene series, 412.
trihydroxyl derivatives, 457.
unsaturated, 393.
- Hydrochlorate of apomorphine**, 519.
morphine, 516.
quinine, 522.
- Hydrochloric acid**, 30, 265.
analytical reactions of, 268.
- Hydrochlorides**, 266.
- Hydrocotarnine**, 518.
- Hydrocotoin**, 502.
- Hydrocotyle asiatica*, 509.
- Hydrocyanic acid**, 279.
analytical reactions of, 283.
antidotes to, 283.
diluted, 281.
from bitter almond and cherry-laurel, 500.
inhalation of, 282.
in organic mixtures, detection of, 565.
in the blood, 284.
Schönbein's test for, 284.
volumetric estimation of, 640.
- Hydroferricyanic acid**, 341.
- Hydroferrocyanic acid**, 340.
- Hydrofluoric acid**, 342.
- Hydrogen**, 21.
antimoniuretted, 185.
arseniuretted, 174.
benzoate, borate, etc. *Vide* the respective acids, Benzoic, Boracic, etc.
combustion of, 22.
dioxide, 103.
volumetric estimation of, 653.
derivation of word, 32.
explosion of, 22.
functions of, 124.
heavy carburetted, 408.
in artificial light-producers, 22.
light carburetted, 396.
lightness of, 22.
peroxide, 103.
persulphide, 303.
phosphoretted, 343.
preparation of, 21.
properties of, 21, 24.
quantitative estimation of, in organic compounds, 638, *et seq.*
salts of, 263.
siliciuretted, 355.
sulphuretted, 97, 301, 303.
type, 379.
used for balloons, 24.
weight compared with air, 24.
weight of 1 litre, 621.

- Hydrogen**, weight of 100 cubic inches, 621.
Hydrogenium, 21.
 Hydrolysis, 460.
 Hydrolysts, 441.
 Hydrometers, 616.
 Hydroquinine, 527.
 Hydroquinone, 455, 501, 521.
 Hydrosulphuric acid, 301.
 Hydrosulphurous acid, 346.
 Hydrosulphyl, 303.
 Hydrous butyl chloral, 488.
 compounds, 85, 121.
 ferric chloride, 150.
 Hydroxyacetic acid, 409, 492.
 Hydroxybenzoic acid, 493.
 aldehyde, 493.
 Hydroxybenzyl alcohol, 456.
 Hydroxybutyric acid, 490.
 Hydroxycaproic acid, 490.
 Hydroxydodecyl alcohol, 490.
 Hydroxyformic acid, 489, 490.
 Hydroxyheptilic acid, 490.
 Hydroxyl, 66, 303.
 Hydroxylamine, 516.
 Hydroxyoctilic acid, 490.
 Hydroxypentilic acid, 490.
 Hydroxy - propane - tricarboxylic acid, 498.
 Hydroxypropionic acid, 490.
 Hydroxysuccinic acid, 496.
 Hydroxytoluic acid, 493.
 Hygiene, 15.
Hyoscinæ hydrobromas, 538.
 impurities in, 714.
 Hyoscine, 538.
 Hyoscyamia, 538.
Hyoscyaminæ hydrobromas, 538.
 impurities in, 714.
 Hyoscyamine, 538.
Hyoscyamus, 538.
 Hyper-, meaning of, 149.
 Hypnone, 498.
 Hypo-, meaning of, 344.
 Hypobromites, 271.
 Hypochloride of sulphur, 304.
 Hypochlorite of calcium, 114.
 sodium, 88.
 Hypochlorites, 292.
 Hypochlorous acid, 292.
 Hypogæin, 464.
 Hypophosphite of calcium, 343.
 sodium, 344.
 Hypophosphites, 344.
 Hypophosphoric acid, 353.
 Hypophosphorous acid, 344.
 estimation of, 638, 654.
 Hyposulphite of calcium, 303.
 sodium, 345.
 Hyposulphite of sodium, standard solution of, 655.
 Hyposulphites, 345.
 Hyposulphurous acid, 345.
 -IC, meaning of, 73, 143.
 Icacin, 426.
 Iceland moss, 338, 476.
Ichthyocolla, 549.
 -ide, meaning of, 76.
 Igasurine, 528.
 Ignition, 102.
Illicium, 415.
Illicium anisatum, 415.
 Illuminating agents, analysis of, 706.
 Imidazoic acid, 515.
 Imidogen bases, 431.
 Incense, 428.
 Inch, 608.
 Incineration, 102.
 of filters in quantitative analysis, 659.
 Indelible ink, 218.
 Indestructibility of force, 38.
 of matter, 13, 38.
Indian, barberry, 533.
 gamboge, 427.
 hemp, 423.
 ink, 556.
 ipacacuanha, 537.
 liquorice, 504.
 melissa oil, 420.
 mustard, 451.
 pennywort, 509.
 red, 555.
 yellow, 553.
 India-rubber, 420.
 vulcanized, 420.
 Indican, 291.
 Indicator, 626.
 Indiglucin, 291.
 Indigo, 291.
 blue, 291.
 sulphate of, 291.
 white, 291.
 wild, 532.
 Indigogen, 291.
 Indigotin, 291.
 disulphonic acid, 291.
 Indium, 725.
 Infusible white precipitate, 207.
 Infusions, 589.
Infusum cinchonæ acidum, 522.
 Inhalation of chlorine, 29.
 conine, 536.
 hydrocyanic acid, 282.
 Inhalations, 589.
Injectio apomorphinæ hypodermica, 520.

- Injectio morphinæ hypodermica*, 517.
- Ink**, black, 162, 358, 556.
 Heberden's, 162.
 indelible, 218.
 Indian, 556.
 invisible, 236.
 marking, 218.
 printer's, 556.
 sympathetic, 236.
- Inorganic chemistry, 383.
 compounds, 383.
- Inosite, 469.
- Insecticide, 425.
- Introduction, 13.
- Inula*, 418.
- Inula Helenium*, 418, 476.
- Inulic anhydride, 418.
- Inulin, 476.
- Inulol, 418.
- Inverted sugar, 468.
- Invisible ink, 236.
- Iodal, 488.
- Iodate of potassium, 75, 396.
- Iodates, 396.
- Iodic acid, 396.
- Iodide of ammonium, 274.
 antimony, 180.
 arsenum, 168.
 bismuth and potassium, 571.
 cadmium, 250.
 copper, 276.
 ethyl, 402.
 hydrogen, 273.
 iron, 31, 147.
 lead, 213.
 potassium, 75, 274.
 detection of iodate in, 76.
 silver, 220, 275.
 starch, 275, 473.
 sulphur, 274.
- Iodides**, 272.
 analytical reactions of, 275.
 of mercury, 197.
 quantitative estimation of, 677.
 separation of, from bromides
 and chlorides, 276.
- Iodine**, 30, 272.
 its analogy to chlorine and bro-
 mine, 272, 278.
 chloride, 273.
 derivation of word, 32.
 moisture in, 678.
 molecular weight of, 198, 273.
 solution of, 274.
 standard solution of, 645.
 tincture of, 274.
 volumetric estimation of, 656.
 water, 274.
- Iodoformum*, 401, 444.
- Iodoformum*, impurities in, 714.
- Iodo-salicylic acid, 493.
- Iodol, 513.
- Iodopyrrol, 513.
- Iodum*, 272.-
 impurities in, 714.
- Ipecacuanha*, 537.
- Ipomœa orizabensis*, 505.
 purga, 505.
 simulans, 505.
 turpethum, 201.
- Iridin, 509.
- Iridium, 247, 725.
 chloride, 571.
- Iris florentina*, 418.
- Iris versicolor*, 509.
- Irish moss, 479.
- Irisin, 509.
- Iron**, 142.
 acetate, 153, 300.
 acetonitrate, 158.
 alum, 139.
 ammonio-citrate, 154.
 ammonio-tartrate, 154.
 analytical reactions of, 160.
 arsenate, 146, 170, 179.
 bromide, 148.
 carbonate, 145.
 saccharated, 145.
 volumetric estimation of,
 650.
 cast, 142.
 chloride, green, 149.
 citrate, 157.
 and quinine, 155, 523.
 compounds, nomenclature of,
 143.
 derivation of word, 32.
 detection of, in presence of
 aluminium and zinc, 163.
 dialyzed, 160.
 ferrocyanide of, 341.
 galvanized, 131.
 hydrate, 151.
 hydrated oxide, 151.
 iodate, 396.
 iodide, 31, 147.
 magnetic oxide of, 142.
 estimation of iron in,
 651.
 meconate, 349.
 nitrate (pernitrate) of, 157.
 in compounds, estimation of,
 669.
 ore, magnetic, 142.
 needle, 153.
 spathic, 142.
 specular, 142.
 oxide, 152, 153.

- Iron oxyhydrate**, 153.
 oxysulphate, 145.
 perchloride, anhydrous, 148.
 perhydrate, 151.
 pernitrate, 157.
 peroxide, 152.
 separation of, from phosphates and oxalates, 375.
 peroxhydrate, 151.
 persulphate, 150.
 phosphate, 146, 161, 331.
 volumetric estimation of, 651.
 potassio-citrate, 154.
 potassio-tartrate, 154.
 pyrites, 142.
 quantitative estimation of, 669.
 red oxide of, 153.
 reduced, 158.
 rust of, 143.
 saccharated carbonate of, 145.
 volumetric estimation of, 650.
 salts, nomenclature of, 143.
 scale, compounds of, 154.
 separation of, from aluminium and chromium, 240.
 sodio-citrate, 157.
 sodio-tartrate, 157.
 sulphate, 144.
 sulphide, 31, 147, 161, 162.
 sulphocyanate, 162, 356.
 tartrate, 156.
 wine of, 157.
 wrought, 142.
Ironstone, clay, 142.
 chrome, 238.
Isaconitine, 530.
Isatropyl-ecgonine, 534.
Isinglass, 504.
Iso-, meaning of, 55, 480.
Isoamylic hydride, 398.
Isobutane, 398.
Isoheptoic acid, 418.
Isomeric bodies, 480.
Isomerides, 480.
 physical, 482.
Isomerism, 480.
Isomers, 480.
Isomorphism, the doctrine of, 55.
Isomorphous bodies, 55.
Isonandra gutta, 420.
Isonitriles, 483.
Isophthalic acid, 497.
Isoprophylactic acid, 488.
Isorottlerin, 425.
Ispaghul, 479.
 -ite, meaning of, 73.
Ivory-black, 556.
- JABORANDI**, 540.
Jaboridine, 541.
Jaborine, 540.
Jalap, Mexican male, 505.
 resin, 505.
 Tampico, 505.
 true, 505.
Jalapa, 505.
 impurities in, 714.
Jalapic acid, 505.
Jalapin, 505.
Jalapinol, 505.
 "James's powder," 184.
Japaconitine, 530.
Jaune brillant, 250.
Jelly, vegetable, 479.
Jequeritin, 504.
Jequerity, 504.
Jequerity-zymase, 504.
Jervine, 539.
Juglandine, 539.
Juices, 589.
Juniper oil, 418.
Juniper-tar oil, 427.
Juniperus oxycedrus, 427.
Juniperus sabina, 419.
- KAINITE**, 61.
Kairine, 514, 526.
Kairoline, 526.
Kaladana resin, 505.
Kali, 32, 88.
Kalium, 32.
Kamala, 425.
Kaolin, 355.
Kariyat, 509.
Kelp, 272.
Kermes, mineral, 184.
Ketone-chloroform, 402.
Ketones, 498.
Ketose, 467.
Kieselguhr, 548.
Kieserite, 118.
Kilbride, mineral, 153.
Kiln, 107.
Kilogramme, 604.
Kilolitre, 604.
Kilomètre, 604.
Kinates. See **Quinates**.
Kinetic theory, 24.
King's blue, 555.
Kino, 359.
Kinone. See **Quinone**.
Kiwach, 242.
Kjeldahl's process, 692.
Kokum butter, 462.
Kola-nut, 543.
Kosin, 424.
Koussin, 424.

- Koussou, 424.
Krameria radix, 359.
 Kunch, 504.

 "LABBARAQUE'S SOLUTION," 88.
 Laboratory furniture, xiii.
 Laburnum, 536.
 Lac, 546.
 Lac, 555.
 dye, 555.
 seed, 555.
 shell, 555.
 stick, 555.
 Lactams, 507.
 Lactates, 347.
 Lactic acid, 347, 489, 490.
 volumetric estimation of, 637.
 series of acids, 489.
 relations to acetic and glyoxylic series, 490.
 Lactims, 507.
 Lactoglucose. See Glucose.
 Lactometer, 547.
 Lactone, 507.
 Lactones, 507.
 Lactose, 471.
Lactuca, 509.
 Lactucarium, 509.
 Lactucin, 509.
 Ladies' slipper, 510.
 Lævogyrate, 467.
 Lævoracemic acid, 320.
 Lævorotation, 467.
 Lævotartaric acid, 320.
 Lævulose, 468, 469.
 Lakes, 554.
Lamellæ atropinæ, 531.
 physostigminæ, 540.
 Lampblack, 556.
 Lamps, gas-, 17.
Lana philosophica, 135.
 Lanolin, 460.
 Lanthanum, 726.
 Lanthopine, 518.
Lapis lazuli, 556.
 Lappa, 509.
Lappa officinalis, 509.
 Larch-bark, 359.
 Lard, 462.
 benzoated, 462.
 oil, 462.
 prepared, 462.
 purified, 462.
Laricis cortex, 359.
Larix europæa, 359.
 Larixin, 360.
 Larixinic acid, 360.
 Latent heat, 86, 129.

 Laudanine, 518.
 Laudanosine, 518.
 Laughing-gas, 95.
 Laurate of glyceryl, 462.
 Laurel camphor, 421.
 Lauric acid, 462, 490.
 aldehyde, 419.
Laurocerasi folia, 500.
 Lavender oil, 418.
 -water, 415.
 "Law," Avogadro's and Ampère's, 53.
 Berthollet's, 379.
 Boyle's, 53, 619.
 Charles's, 53, 619.
 Dalton's, 48, 49.
 Dulong and Petit's, 623.
 Gay-Lussac's, 54.
 Graham's, 24, 58.
 Henry and Dalton's, 86.
 Malaguti's, 379.
 Mariotte's, 54, 619.
 Law concerning molecular weight, 55.
 of constant proportions, 47.
 diffusion, definition of, 58.
 indestructibility of matter, 13.
 multiple proportions, 49, 198.
 solubility of gases in liquids, 86.
 The Periodic, 380.
 Laws of chemical combination by weight, 47, 52, 59, 198, *et seq.*
 chemical combination by volume, 49, 59, *et seq.*
 Lead, 210.
 acetate, 211.
 estimation of, 631.
 analytical reactions of, 214.
 antidotes to, 215.
 carbonate, 211, 215.
 chloride, 214.
 chromate, 215.
 derivation of word, 33.
 detection of, in organic mixtures, 563.
 gummate of, 116.
 hydrato-carbonate, 210.
 iodide, 213.
 nitrate, 212.
 oleate, 213.
 oxide, 210.
 oxyacetate, 211.
 oxychromate, 215.
 perchloride, 213.
 peroxide, 212.
 plaster, 213.
 puce-colored oxide or peroxide of, 212.

- Lead**, pyrophorous, 159.
 quantitative estimation of, 675.
 quantivalence of, 211.
 red, 212.
 shot, 210.
 subacetate, 211.
 sugar of, 211.
 sulphate, 215.
 sulphide, 214.
 native, 210.
 test for, in water, 214.
 tree, 215.
 volumetric estimation of solu-
 tions of acetate of, 631.
 white, 210.
- Leadstone**, 142.
- Leaf-green**, 556.
- Lecanora**, 556.
- Lees**, 318.
- Legumin**, 548.
- Lemon and kali**, 88.
- Lemon-chrome**, 215.
 -grass oil, 420.
 -juice, 325.
 estimation of mineral acids
 in, 682.
 oil, 416.
- Length**, unit of, 603.
- Lentisk tree**, 424.
- Lepidolite**, 227.
- Leptandra**, 510.
- Leptandrin**, 510.
- Leucic acid**, 490.
- Leucine**, 513.
- Leucomaines**, 513.
- Levisticum**, 428.
- Levulose**, 468.
- Lichen blue**, 555.
 sugar, 465.
- Lichenin**, 476.
- Lichestearic acid**, 338.
- Light magnesium carbonate**, 119.
 carburetted hydrogen, 395.
 magnesia, 122.
- Lignin**, 479.
- Lime**, bisulphite, 306.
 carbonate, 108.
 precipitated, 108.
 caustic, 107.
 chloride of, 114.
 hydrate, 107.
 juice, 325.
 estimation of mineral acids
 in, 682.
 kiln, 107.
 oil, 416.
 phosphate, 113.
 quick, 107.
 slaked, 107.
- Lime**, sulphurated, 115.
- Lime-water**, 108.
- Limestone**, 106.
 magnesian, 118.
 mountain, 118.
- Limonenes**, 412.
- Limonis cortex**, 416.
 succus, 324.
 impurities in, 714.
- Limonite**, 153.
- Linamarin**, 500.
- Liniment of mercury**, 195.
- Linimentum ammoniæ**, 461.
 calcis, 461.
- Linkage of atoms**, 389, 430.
- Linoleine**, 463.
- Linoxyn**, 463.
- Linseed**, 463, 479.
 cake, 463.
 meal, 463.
 oil, 463.
 tea, 479.
- Linum usitatissimum**, 479.
 impurities in, 714.
- Liqueurs**, 441.
- Liquid**, definition of, 58.
 camphor, 421.
- Liquidambar orientale**, 495.
- Liquids**, specific gravity of, 614.
 official, specific gravity of, 614.
- Liquor acidi arsenosi**, 168.
 vol. est. of, 648.
- acidi chromici**, 239.
 arsenosi, 168.
- ammonii acetatis**, 93.
 impurities in, 714.
- citratis**, 95.
 fortior, 95.
- antimonii chloridi**, 180.
 estimation of anti-
 mony in, 671.
- arseni et hydrargyri iodidum**, 167.
- bismuthi**, 254.
 et ammonii citratis, 254.
 estimation of bis-
 muth in, 672.
- calcis**, 108.
 impurities in, 714.
 vol. est. of, 635.
 chlorinatæ, 115.
- cocainæ hydrochloratis**, 534.
- ferri acetatis**, 150.
 estimation of iron
 in, 669.
 impurities in, 714.
- citratis**, 154.
 impurities in, 714.
- chloridi**, 150.
 impurities in, 714.

- Liquor ferri dialysatus**, 160, 705.
 est. of iron in, 669.
nitratis, 158.
 estimation of iron in, 669.
 impurities in, 714.
persulphatis, 151.
 estimation of iron in, 669.
 impurities in, 714.
subsulphatis, 151.
 impurities in, 714.
gutta-percha, 421.
glonoini, 458.
hydrargyri nitratis, 200.
 impurities in, 714.
perchloridi, 202.
iodi compositus, 317.
lithiæ effervescens, 228.
magnesi carbonatis, 120.
 citratis, 122.
morphinæ acetatis, 517.
bimeconatis, 518.
hydrochloratis, 517.
sulphatis, 518.
nitroglycerini, 458.
plumbi subacetatis, 212.
 dilutus, 212.
 impurities in, 714.
potassæ, 63, 67.
 impurities in, 714.
effervescens, 72.
 neutralizing power of, 68.
 specific gravity of, 614.
 to prepare, 68.
potassii arsenitis, 168.
permanganatis, 232.
sodæ, 82.
 impurities in, 714.
chloratæ, 87.
 impurities in, 714.
effervescens, 86.
sodii arsenatis, 170.
ethylatis, 443.
silicatis, 355.
strychninæ hydrochloratis, 528.
zinci chloridi, 133.
Liquorice, 471, 504.
 sugar, 471, 504.
List of apparatus, xii.
 chemical substances, xiv.
 reagents, xiii.
Litharge, 210.
Lithates, 362.
Lithic acid, 362.
Lithii benzoas, 227.
 impurities in, 714.
bromidum, 227.
 impurities in, 714.
Lithii carbonas, 227.
 impurities in, 715.
citras, 227.
 impurities in, 715.
salicylas, 227.
 impurities in, 715.
Lithium, 227.
 analytical reactions of, 228.
 and platinum chloride, 228.
 benzoate, 227.
 bromide, 227.
 volumetric estimation of, 634.
 carbonate, 227.
 vol. est. of, 634.
 citrate, 227.
 vol. est. of, 634.
 derivation of word, 31.
 flame, 228.
 fluoride, 227.
 salicylate, 227.
 volumetric estimation of, 634.
 silicate, 227.
 sulphate, 228.
 urate, 228.
Litmus, 96, 555.
 paper, 96.
 solution of, 96, 636.
 tincture of, 96.
Litre, 603.
 relation of, to pints, 604.
Liver of sulphur, 69.
Lixiviation, 89.
 fractional, 378.
Loadstone or lodestone, 142.
Lobelia, 539.
Lobelina, 539.
Lobeline, 539.
Lodestone, 142.
Loganetin, 506.
Loganin, 506.
Logwood, 359, 555.
 solution of, bleached by chlorine, 29.
Lokas, 556.
Long pepper, 541.
Looking-glasses, 241.
Lotio hydrargyri flava, 204.
nigra, 205.
Louisa blue, 556.
Lozenges, 303, 589.
 bicarbonate of sodium, 87.
 bismuth, 252.
 chlorate of potassium, 295.
 morphine, 517.
 and ipecacuanha, 517.
 reduced iron, 158.
 sulphur, 303.

- Lucifers, 25.
 Lump-sugar, 469.
 Lunar caustic, 218.
Lupulin, 427.
 oleo-resin of, 427.
 Lupuline, 539.
 Lupulinic acid, 427.
 Lupulinum, 427.
 impurities in, 715.
Lupulus, 427, 539.
 Luteolin, 554.
 Lutidine, 513.
 Luting, 98.
 fire-clay, 202.
 linseed meal, 98.
 Lycopodium, 664.
 impurities in, 715.
Lycopodium clavatum, 664.

 MACE, fixed oil of, 462.
 volatile oil of, 419.
 Macleyine, 541.
 Madder, 434, 554.
 Magenta, 557.
 Magnesia, 122.
 calcined, 122.
 carbonate of, 119, 120.
 effervescing citrate of, 122.
 fluid, 120.
 heavy, 122.
 hydrous carbonate of, 119.
 impurities in, 715.
 light, 122.
 sulphate, 118.
Magnesia carbonas levis, 129.
 levis, 122.
 ponderosa, 122.
 Magnesian limestone, 108.
Magnesii carbonas, 119.
 impurities in, 715.
 ponderosa, 119.
 carbonatis, liquor, 120.
 citratis effervescens, 122.
 liquor, 122.
 sulphas, 118.
 impurities in, 715.
 Magnesite, 118.
Magnesium, 118.
 analytical reactions of, 122.
 and ammonium arsenate, 122.
 phosphate, 122, 330.
 sulphate, 685.
 carbonate, 119.
 chloride, 118.
 citrate, 122.
 derivation of word, 32.
 detection of, in presence of
 barium and calcium, 126.
 euxanthate, 554.

Magnesium, for analytical purposes,
 174.
 oxide, 122.
 phosphates in bones, 112.
 purrate, 553.
 quantitative estimation of, 666.
 separation from barium and
 calcium, 126.
 silicate, 354.
 sulphas effervescens, 119.
 sulphate, 118.
 Magnetic iron ore, 142.
 vol. est. of, 651.
 Magnolia, 509.
 Magpie test for mercury, 209.
 Maize starch (fig.), 475.
 Malachite, 190.
 Malaguti's law, 379.
 Malate of atropine, 531.
 nicotine, 539.
 Malates, 348.
 Male fern, oil of, 464.
 Malic acid, 348, 496.
 series of acids, 496.
 Mallow tea, 479.
 Malonic acid, 497.
 Malt, 478.
 extract, 478.
 substitutes, 468.
 Maltose, 468, 470, 471.
 Manganate of potassium, 77, 231.
 sodium, 232.
Manganese, 231.
 analytical reactions of, 233.
 black oxide of, 231.
 Crum's test for, 234.
 derivation of word, 34.
 dioxide, 231.
 quantitative analysis of black
 oxide of, 667.
Mangesii oxidum nigrum, 231.
Mangani dioxidum, 231.
 impurities in, 715.
 sulphas, 234.
 impurities in, 715.
 Manganous chloride, 231.
 hydrate, 234.
 sulphide, 233.
 Mangosteen oil, 463.
Manihot, starch of (fig.), 475.
 Manna, 465.
 Mannite, 465.
 Manufacturing chemists, 14.
 Manures, analysis of, 706.
Maranta, starch of (fig.), 475.
 Maraschino, 441.
 Marble, 106.
 Margarine, 462.
 Margosa-bark, 509.

- Marigold, 509.
 Marine soap, 462.
 Mariotte's law, 53, 619.
 Marking ink, 218.
 Marl, 138.
Marmor album, 106.
Marrubium, 510.
 Marseilles soap, 461.
 Marsh gas, 396.
 Marsh-gas series, 395.
 Marshmallow, 479.
 Marsh's test for arsenum, 173.
Massa ferri carbonatis, 145.
 hydrargyri, 195.
 Massicot, 210.
 Mastic, 424.
Mastiche, 424.
 Mastichic acid, 424.
 Masticin, 426.
 Maté, 542.
Maticæ folia, 510.
 Matico, 510.
Matricaria, 415
 chamomilla, 415.
Matter, indestructible, 13, 38.
 origin of, 38.
 Mauve, 501, 757.
 May-apple, 425.
 Mayer's reagent, 630.
 Meadow-sweet, 493, 507.
 oil of, 407, 493, 507.
 Measures, 605, *et seq.*
 Mechanical and chemical combina-
 tion, 31, 36.
 medicines, 242.
 Meconate of iron, 349.
 morphine, 517.
 Meconic acid, 349, 567.
 Meconidine, 518.
 Meconin, 518.
 Meconoisin, 518.
 Meerschau, 354.
Mel, 471.
 impurities in, 715.
 boracis, 333.
 despumatum, 559.
 Melam, 357.
Melassa, 419.
 Melasses, 471.
 Melegueta pepper, 418.
 Meletizose, 471.
Melia azedarach, 510.
 Melissa oil, 420.
 Melissic acid, 489, 490.
 alcohol, 450.
 Melissyl, palmitate of, 450.
 Melitose, 471.
 Melitic acid, 498.
 Mellon, 357.
 Melon essence, 407.
 Melting-points, table of, 598, 599.
 of fats, etc., 548.
 to determine, 598.
 of metals, 599.
 Memoranda, analytical, 100, 224,
 259, 365.
Menispermum canadense, 533.
Mentha arvensis, 419.
 pulegium, 419.
 Menthene, 419.
Menthol, 419, 452.
 impurities in, 715.
 Mercaptans, 445.
 Mercurialine, 512.
Mercuric ammonium, chloride of,
 205.
 chloride, 201.
 cyanide, 280.
 hexiodide, 273.
 iodide, 197.
 nitrate, 199.
 oxide, 204.
 oxynitrates, 200.
 oxysulphate, 201.
 phenylate, 454.
 potassium iodide, volumetric
 solution of, 658.
 salts, 196.
 analytical reactions of, 205.
 sulphate, 200.
 sulphide, 208.
 sulphocyanate, 357.
Mercurius vitæ, 181.
Mercurous ammonium, chloride of,
 206, 208.
 chloride, 203, 209.
 chromate, 209.
 compounds, 196.
 iodide, 197.
 nitrate, 199.
 oxide, 205.
 salts, 196.
 analytical reactions of, 205,
 208, 209.
 sulphate, 201.
 sulphide, 208.
Mercury, 195.
 amido-chloride, 206.
 ammoniated, 206.
 ammonio-chloride, 206.
 analytical reactions of, 205.
 antidotes to, 209.
 basic sulphate, 201.
 bichloride, 201.
 black oxide, 204.
 carbonates, 209.
 chlorides, 201.
 derivation of word, 33.

- Mercury**, detection of, in organic mixtures, 562.
 formula of, 195.
 fulminate, 219.
 galvanic test for, 209.
 hexiodide, 273.
 iodides, 197.
 magpie test for, 209.
 molecular weight of, 195.
 native sulphide, 195.
 nitrates, 199.
 nomenclature of salts of, 196.
 of life, 181.
 oleate of, 460.
 oxides, 204.
 oxynitrates, 200.
 oxysulphate, 201.
 oxysulphide, 208.
 perchloride, 201.
 persulphate, 200.
 phenylate, 454.
 quantitative estimation of, 673.
 subchloride, 203, 209.
 sulphates, 200.
 sulphide, 195.
 yellow oxide, 204.
- Mesitylene**, 429.
Mesoxalic acid, 493.
Meta-, meaning of, 349, 455.
Metaboric acid, 333.
Metachloral, 486.
Metacinnaméin, 495.
Metagummic acid, 479.
Metaldehyde, 485.
Metallic elements, 15.
Metalloids, 16.
Metals, 16.
 of minor pharmaceutical importance, 226.
 quantitative estimation of, 659.
 table of the fusibility of, 599.
- Metamerides**, 481.
Metamerism, 481.
Metamers, 481.
Metantimonic acid, 181.
Metaphosphates, 349.
Metaphosphoric acid, 330, 349.
Metastannates, 243.
Metastannic acid, 243.
Metastyrol, 495.
Metathesis, 59, 79, 107.
Metavanadates, 332.
Methacrylic acid, 491.
Methane, 396.
 series, 395.
 substitution-products of, 399.
Methoxycatechol, 336.
Methylal, 486.
Methylamine, 512.
- Methylated spirit**, 437.
 sweet spirit of nitre, 438.
- Methyl benzene**, 431.
 arbutin, 501.
 carbinol, 439.
 chloride of, 399.
 conine, 536.
 dichlorobenzene, 432.
 ethyl, 397.
 formic acid, 484.
 group, 431.
 hydride of, 396.
 monochlorobenzene, 432.
 morphine, 520.
 nonyl-ketone, 419, 498.
 phenoène, 429, 431.
 prophyl-phenoène, 429, 432.
 protocatechuic aldehyde, 364.
 salicylate, 407, 492.
 impurities in, 715.
 theobromine, 543.
 trichlorobenzene, 432.
- Methyl salicylas**, 492.
 impurities in, 715.
- Methylic acid**, 490.
 alcohol, 437.
 detected in presence of ethylic alcohol, 438.
- Metre**, 604.
 relation of, to inches, 604.
- Metric system**, 601, *et seq.*
- Metrical system**, weights and measures of, 604, *et seq.*
- Metrical system of weights and measures**, its relation to the British, 608, *et seq.*
- Meum**, 428.
Mezereum, 425.
Mica, 138.
Mica panis, 471.
- Microcosmic salts**, 373.
- Microscopic examination of urinary sediments**, 581.
- Microscopy of starches**, 474.
- Micro-spectroscope**, 560.
- Milk**, 546.
 poison, 513, 571.
 curdling ferment, 546.
 of sulphur, 303.
 sugar, 471.
- Mimetesite**, 332.
Mimotannic acid, 359.
- Mineral acids**, detection of, in organic mixtures, 564.
- Mineral**, chameleon, 232.
 kermes, 182.
 Kilbride, 153.
 purple, 555.
 rouge, 153, 555.

- Minerals, general analysis of, 370,
et seq.
 special analysis of, 687.
- Minim, 608.
- Minium, 210.
- Mint, 419.
- Mirbane, essence of, 430.
- Mishmi bitter, 533.
- Mispickel, 168.
- Mistura ferri aromatica*, 162.
composita, 146.
- Mitigated caustic, 218.
- Mixed ethers, 447.
- Mixture, different from chemical
 combination, 31, 36.
 definition of, 57.
- Mixtures, 589.
- Mohr's burette, 627.
- Moist sugar, 469.
- Molasses, 471.
- Molecular** arsenum, 168.
 attraction, 138.
 composition, 386.
 constitution or structure, 130,
 138, 195, 386, 393, 432.
 density of gases and vapors, 53,
 55, 387, 619.
 electricity, etc., 623.
 ferric chloride, 149.
 formulæ as indicated by chem-
 ical "periodicity," 380.
 properties, 387, 395.
 isomorphism, 55.
 phosphorus, 328.
 specific heat, 622.
 substitution, 194.
 sulphur, 302.
 theory, vii.
 volume, 55.
 weight, 55, 621.
 weights, 149, 198.
 definition of, 59.
- Molecule, definition of, 57.
- Molecules, 38.
 conception of, 391, 393.
- Molybdates, 332.
- Molybdenum, 331, 726.
 oxide, 331.
 sulphide, 331.
- Molybdic anhydride, 331.
- Monads, 123.
- Mouamines, 512.
- Mon-iodoethane, 403.
- Monobasic acids, 264.
- Monobasylous radicals, 264.
- Monobrom-acetanilide, 431.
- Monobrom-camphor, 421.
- Monobrom-ethane, 403.
- Monobromobenzene, 430.
- Monochlorobenzene, 430.
- Monochloromethane, 399.
- Monochloromethylbenzene, 432.
- Monoformin, 451.
- Monohydroxyl derivatives of the
 paraffins, 436.
- Mononitrocellulin, 479.
- Monoxynaphthalenes, 434.
- Morbid urine, 572.
- Mordants, 140.
- Mori succus*, 554.
- Morphina*, 516.
- Morphinæ acetat*, 517.
 impurities in, 715.
hydrochloras, 517.
sulphas, 517.
 impurities in, 715.
- Morphine**, or Morphia, 516.
 acetate, 517.
 analytical reactions of, 518.
 bimeconate, 517.
 distinction from brucine, 529.
 hydrochlorate, 517.
 in organic mixtures, detection
 of, 567.
 quantitative estimation of,
 698.
 sulphate, 517.
 tartrate, 517.
- Morrhaine, 463.
- Mortar, 355.
- Mosaic gold, 244.
- Moschus*, 549.
moschiferus, 549.
- "Mother-liquor," 109.
- Motion from heat, 86.
- Mottled soap, 461.
- Mountain-blue, 555.
 limestone, 118.
- Mucic acid, 472.
- Mucilage of bael, 479.
 gum acacia, 116.
 linseed, 479.
 marshmallow, 479.
 squill, 479.
 starch, 473.
 tragacanth, 116.
- Mucilago acaciæ*, 116.
amyl, 473.
tragacanthæ, 116.
- Mucuna pruriens*, 242.
- Mucus in urine, 580.
- Mudar tree, 509.
- Mulberry calculus, 588.
 essence, 407.
 juice, 554.
 sugar in, 467.
- Mulder's process for estimating al-
 cohol, 704.

- Multiple proportions, law of, second law of combination, 49, 198.
- Murex, 362.
- Murexid, 362.
- Muscarine, 513.
- Musk, 549.
artificial, 549.
deer, 549.
- Mustard, 451.
artificial oil of, 451.
essential oil of, 420.
fixed oil of, 464.
"plaster," 451.
- Mycoderma aceti*, 297.
- Mydriatics (table), 544.
- Mylabris cichorii*, 422.
- Myotics (table), 544.
- Myrcia acris*, 441.
- Myristate of glyceryl, 462.
- Myristic acid, 462, 490.
- Myristicene, 419.
- Myristicin, 419.
- Myristicol, 419.
- Myristin, 462.
- Myronate of potassium, 451.
- Myrosin, 451.
- Myroxylon Pereiræ*, 495.
toluifera, 495.
- Myrrh, 428.
- Myrrha*, 428.
- Myrrhic acid, 428.
- Myrtus communis*, 419.
- "Mystery gold," 245.
- Mytiloxine, 513.
- NAPELLINE, 530.
- Naphthalene, 434, 514.
series of hydrocarbons, 433.
- Naphthalic acid, 336.
- Naphthols, 434.
- Naphtalinum*, 434.
impurities in, 716.
- Naphthyl alcohols, 434.
- Naphtol*, 434.
impurities in, 716.
- Narceine, 518.
- Narcotina, 518.
- Narcotine, 518.
- Narthex*. See *Ferula*.
- Nascent state, 40.
- Natal aloes, 435.
- Nataloin, 435.
- Natrium*, 32.
- Natron, 32, 286.
- Natural philosophy, 43.
- Nectandra Rodiei*, 532.
- Nectandræ cortex*, 532.
- Nectandrine, 533.
- Needle iron ore, 153.
- Negative pole, 245.
- Neroli oil, 416.
- Nessler's test, 744.
- Neuridine, 513.
- Neurine, 513.
- Neutral chromate, 105.
hydrocarbons, 391.
- Neutralization, 96.
- Nickel, 236.
analytical reactions of, 236.
arseno-sulphide, 236.
cobalticyanide, 237.
cyanide, 237.
derivation of word, 34.
hydrate, 237.
separation of, from cobalt, 237.
silver, 236.
sulphide, 236.
- Nicker-nuts, 509.
- Nicotiana tabacum*, 539.
- Nicotine, Nicotia, Nicotina, or Nicotylia, 539.
malate, 539.
- Nihilum album*, 135.
- Nim, 509.
- Niobium, 725.
- Nitrate of ammonium, 94, 288.
argent-ammon-ammonium, 207.
barium, 102.
bismuth, 251.
copper, 192.
iron, 157.
lead, 212.
mercury, 199.
pilocarpine, 540.
potassium, 73, 285.
silver, 216, 217.
and potassium, 217.
standard solution of, 639.
toughened, 218.
sodium, 81, 285.
strontium, 229.
- Nitrates, 285.
analytical reactions of, 289.
quantitative estimation of, 678,
et seq.
- Nitre, 285.
Chili, 286.
cubic, 286.
prismatic, 285.
sweet spirit of, 351, 403, 438.
- Nitric acid, 285, 287.
anhydrous, 288.
antidotes to, 292.
dil., 288.
in organic mixtures, detection of, 564.
volumetric estimation of, 638.

- Nitric anhydride, 288, 290.
 oxide, preparation of, 289.
 peroxide, 289, 290.
 Nitrification, 286.
 Nitrile bases, 431, 510, 515.
 Nitriles, 483.
 Nitrite of amyl, 351, 407.
 ethyl, 351, 402.
 potassium, 350.
 sodium, 351.
Nitrites, 350.
 analytical reactions, 350.
 in water, test for, 350.
 Nitrobenzene, 430.
 Nitrobenzol, 430, 500.
 in oil of bitter almonds, test
 for, 500.
 Nitrocellulin, 470.
 Nitroethane, 405.
Nitrogen, 25, 289.
 derivation of word, 32.
 in the atmosphere, 25.
 oxides, 289, 290.
 peroxide, 289, 290.
 preparation of, 25.
 properties of, 26.
 quantitative estimation of, in
 organic compounds, 691, *et seq.*
 relative weight of, 27.
 Nitroglycerin, 458.
 Nitrohydrochloric acid, 187, 289.
 Nitromannite, 465.
 Nitrometer, 351, 405.
 Nitropentane, 408.
 Nitrosulphonic acid, 308.
 Nitrosyl-sulphonic acid, 309.
 Nitrous acid, 289, 290, 350.
 anhydride, 289, 290.
 ether, 403.
 oxide, 95.
Nomenclature of alkaloids, 515.
 anhydrides, 85.
 anhydrous bodies, 85.
 -ate, 73, 76.
 carbonization, evaporation, ig-
 nitition, incineration, 102.
 double salts, 79.
 glucosides, 499.
 hydrates, 66.
 hydric, -ous, 76.
 hydrous bodies, 85.
 -ide, -ite, 76.
 iron salts, 143.
 mercury compounds, 196.
 notes on, 66, 73, 76, 79, 85, 102,
 143, 196, 515.
 Nonane, 398.
 Non-drying oils, 463.
 Non-metallic elements, 15.
 Non-metals, 16.
 Nonylic acid, 490.
 Nordhausen sulphuric acid, 310.
 Normal hydrocarbons, 391.
 solutions, 626.
 Notation, 41.
 of organic compounds, 389, 396.
 Notes, analytical, 224, 259.
 Nutmeg, expressed oil of, 462.
 oil of, 419.
 Nutrition, plastic elements of, 549.
 Nux vomica, 527.
 assay of, 700.
 OATMEAL, 473.
 Occlusion, 248, 249.
 Ochre, 553.
 Octahedron, 172.
 Octylic acid, 490.
 CEnanthylate of ethyl, 407.
 CEnanthylic acid, 490.
 Official formula, 28.
 liquids, specific gravity of, 614.
 substances, volumetric estima-
 tion of, 627, *et seq.*
Oil of ajowan, 415.
 ajwain, 415.
 almond, 463.
 amber, 356.
 American pennyroyal, 418.
 aniseed, 415.
 apple, 407.
 arachis, 464.
 benne, 464.
 bergamot, 416.
 birch, 407.
 bitter almond, 491, 500.
 artificial, 430, 500.
 "boiled," 463.
 boldo, 416.
 buchu, 416.
 cacao, 462.
 cake, 463.
 cajuput, 416.
 camphor, 421.
 cannabis indica, 423.
 capsicum, 421.
 caraway, 416.
 cardamoms, 416.
 cascarilla, 417.
 cassia, 417.
 castor, 463.
 cedra, 416.
 chamomile, 415.
 cinnamon, 417.
 citron, 416.
 citronella, 417.
 cloves, 416.
 cocoa-nut, 462.

Oil, cod-liver, 463.
 copaiva, 417.
 coriander, 417.
 cowbane, 417.
 croton, 463.
 cubeb, 417.
 cummin, 417.
 dill, 415.
 earth-nut, 464.
 eggs, 545.
 elder-flower, 419.
 eucalyptus, 412, 417.
 fennel, 418.
 fir wool, 413.
 garcinia, 462.
 garlic, 452.
 geranium, 418.
 gingelly, 464.
 ginger, 420.
 ginger-grass, 418.
 grains of paradise, 418.
 grass, 417.
 ground-nut, 464.
 hop, 427.
 horseradish, 415.
 Indian hemp, 423.
 jaborandi, 540.
 juniper, 418.
 lard, 462.
 lavender, 418.
 foreign, 418.
 lemon, 416.
 lemon-grass, 420.
 lime, 416.
 linseed, 463.
 lycopodium, 464.
 mace, fixed, 462.
 volatile, 419.
 male fern, 464.
 mangosteen, 462.
 meadow-sweet, 407, 493, 507.
 melissa, 420.
 mustard, artificial, 451.
 essential, 420, 451.
 fixed, 464.
 myrtle, 419.
 neroli, 416.
 nutmeg, fixed, 462.
 volatile, 419.
 of vitriol, 309.
 olibanum, 428.
 olive, 459, 464.
 omum, 415.
 orange-flower, 416.
 orange-rind, 416.
 orris, 418.
 palm, 462.
 paraffin, 398.
 pennyroyal, 419.

Oil, pepper, 541.
 peppermint, 418.
 petit-grain, 416.
 phellandrium, 412.
 phosphorated, 328.
 pilocarpus, 540.
 pimento, 416.
 pine wool, 413.
 ptychotis, 415.
 resin, 423.
 rose, 419.
 rosemary, 419.
 rue, 419, 498.
 saffron, 554.
 sage, 419.
 sandal-wood, 419.
 sassafras, 420.
 savin, 419.
 sesame, 464.
 shark-liver, 465.
 spearmint, 419.
 sperm, 450.
 spike, 418.
 star-anise, 415.
 sweet-birch, 407.
 sweet-flag, 420.
 teal, 464.
 theobroma, 462.
 thyme, 420.
 turmeric, 420.
 turpentine, 412.
 valerian, 420.
 verbenæ, 420.
 water-hemlock, 417.
 wine, 409.
 winter-green, 407.
 wood, 426.
 Oils, analysis of, 706.
 and fats, composition of, 459.
 drying, 463.
 essential, 413.
 tested for alcohol, 414.
 fixed, 463.
 non-drying, 463.
 volatile, 413, *et seq.*
 process for, 414.
 Ointments, 589.
 Okra, 479.
 Ol, meaning of, 457.
 Oleate of glyceryl, 459.
 lead, 213.
 mercury, 460.
 potassium, 459.
 veratrine, 460.
 zinc, 460.
 Oleates, 459.
 Oleatum hydrargyri, 460.
 veratrinæ, 460.
 zinci, 136.

- Olefiant gas, 408.
 Olefine series of hydrocarbons, 408.
 Olefines, relation to paraffins and acetylenes, 410.
 Oleic acid, 459, 491.
 Oleine, 459.
Oleo-resina cubebæ, 426.
 Oleo-resins, 422, 425.
Oleum amygdalæ, 463.
 amara, 415.
 andropogi citratis, 420.
 anethi, 415.
 anisi, 415.
 anthemidis, 415.
 arachis, 464.
 aurantii corticis, 416.
 florum, 416.
 bergamottæ, 416.
 betulæ volatile, 407.
 cadinum, 427.
 cajaputi, 416.
 cari, 416.
 carui, 416.
 caryophylli, 416.
 chenopodii, 421.
 cinnamomi, 417.
 copaibæ, 417.
 coriandrum, 417.
 crotonis, 463.
 cubebæ, 417.
 erigerontis, 417.
 eucalypti, 417.
 fœniculi, 418.
 juniperi, 418.
 lavandulæ, 418.
 florum, 418.
 limonis, 416.
 lini, 463.
 macis, 419.
 menthæ piperitæ, 418.
 viridis, 419.
 morruæ, 464.
 myrcia, 418.
 myristicæ, 419.
 expressum, 462.
 myrti, 419.
 olivæ, 459, 464.
 phosphoratum, 328.
 picis liquidæ, 426.
 pimentæ, 416.
 pini sylvestris, 413.
 ricini, 464.
 rosæ, 419.
 rosmarini, 419.
 rutæ, 419, 498.
 sabinæ, 419.
 santali, 419.
 sassafras, 420.
 sinapis volatile, 419.
 succini, 356.
 terebinthinæ, 412, 413.
 theobromatis, 462.
 thymi, 420.
 Olibanum, 428.
 Olive oil, 459, 464.
 Omentum, 462.
 Omum oil, 415.
 Opal, 354.
Ophelia chirata, 351.
 Ophelic acid, 351.
 Opianic acid, 518.
 Opianine, 518.
Opii pulvis, 516.
Opium, 516.
 alkaloids, 516.
 deodoratum, 516.
 detection of, in organic mixtures, 567.
 estimation of morphine in, 619.
 impurities in, 716.
 Orange chrome, 215.
 Orange-flower oil, 416.
 water, 416.
 Orange-rind oil, 416.
 wine, 441.
 Orchil, 555.
 Orchis tuber, 479.
 Orcin, 456.
 Ordeal-poison, 540.
 Orellin, 554.
 Organic analysis, 557.
Organic bases, 510.
 chemistry, 383.
 Advice to Students, 381.
 compounds, composition of, 384.
 constitution of, 387.
 notation of, 389, 396.
 radicals, 392.
 Orpiment, 168, 553.
 Orris, butter of, 418.
 camphor of, 418.
 oil of, 418.
 Ortho-, meaning of, etc., 350.
 Orthodihydroxyl benzene, 455.
 Orthohydroxybenzoic aldehyde, 494.
 Orthophenolsulphonic acid, 454.
 Orthophosphates, 332, 350.
 Orthophosphoric acid, 350.
 Orthovanadates, 332.
Oryza, 473.
 sativa, 472, 473.
 starch of (fig.), 475.
Oryzæ farina, 473.
Os ustum, 111.
 Osmium, 726.
 Otto of rose, 419.
 Ouabain, 506.

- Ounce, 608.
 Ourari, 528.
 -ous, meaning of, 76, 143.
Ovi albumen, 544.
 Ox-bile, 552.
 Ox-gall, 552.
 Oxalate of ammonium, 95.
 barium, 105, 317.
 calcium, 117, 316.
 cerium, 230.
 silver, 317.
 sodium, 316.
 strontium, 230.
Oxalates, 316.
 analytical reactions of, 316.
 from phosphates and ferric
 oxide, separation of, 375.
 quantitative estimation of, 684.
 Oxalic acid, 316, 410, 496.
 antidotes to, 317.
 chemically pure, 316.
 in organic mixtures, detec-
 tion of, 564.
 standard solution of, 634.
 Oxamide, 496.
Oxide of aluminium, 140.
 antimony, 181.
 barium, 103.
 bismuth, 253.
 cadmium, 251.
 calcium, 107.
 chromium, 239.
 cobalt, 235, 555.
 copper, 92.
 iron, 153.
 black, 142.
 magnetic, 142.
 lead, 210.
 magnesium, 122.
 manganese, 231.
 mercury, 204.
 molybdenum, 331.
 silicon, 354.
 silver, 218.
 tin, 243.
 zinc, 135.
 Oxides of nitrogen, 290.
 identified, 377.
 Oxidizing flame, 135.
 Oxyacanthine, 533.
 Oxyacetate of copper, 192.
 lead, 211.
 Oxyacid salts, 287.
 Oxyacids of sulphur, 346.
 Oxycarbonate of bismuth, 253.
 Oxychloride of antimony, 181, 184.
 Oxychromate of lead, 215.
 Oxydizing flame, 135.
Oxygen, 16.
 Oxygen, analogies of, 173.
 derivation of word, 32.
 from ozone and autozone, 275.
 in the air, 16, 25.
 its relation to animal and
 vegetable life, 19.
 preparation of, 16, 103.
 properties of, 19.
 quantitative estimation, of, in
 organic compounds, 688, *et*
 seq.
 solubility in water, 19.
 specific gravity of, 24.
 weight of 100 cubic inches, 621.
 Oxygenated water, 103.
 Oxyhydrates of iron, 152.
 Oxyiodide of iron, 147.
 Oxymalonic acid, 497.
 Oxymel, 470.
 of squill, 470.
Oxymel scillæ, 470.
 Oxynitrates of bismuth, 252.
 mercury, 200.
 Oxysalts, 287.
 Oxysuccinic acid, 497.
 Oxysulphate of iron, 145.
 mercury, 201.
 Oxysulphide of antimony, 182.
 mercury, 208.
 Ozokerite, 450.
 Ozone, 275.

 PALAS TREE, 359.
 Palladium, 726.
 chloride, 571.
 Palm oil, 462.
 Palmitate of cetyl, 450.
 glyceryl, 462.
 melissyl, 450.
 Palmitic acid, 462, 490.
 Palmitine, 462.
 Pancreatin, 551.
Pancreatinum, 552.
 Papain, 551.
Papaver rhæas, 555.
 somniferum, 516.
 Papaverine, 518.
Papaveris capsulæ, 516.
 Papaw, 551.
 Paper, bibulous, 109.
 filtering, 109.
 litmus-, blue, 96.
 red, 96.
 turmeric, 96.
 Papers, test-, 96.
 Para-, meaning of, etc., 320, 455.
 Para-acetphenetidin, 431.
 Paracotin, 502.
 Paracyanogen, 280, 282.

- Paradol, 418.
 Paraffin, 398.
 oil, 398.
 series of hydrocarbons, 394, 396.
 wax, 398.
 Paraffinic acid, 399.
 Paraffins, gaseous, 396.
 hard, 399.
 monohydroxyl derivatives of, 436.
 relations to olefines and acet-
 lenes, 410.
 soft, 398.
 solid, 399.
Paraffinum durum, 398.
liquidum, 398.
molle, 398.
 Paraguay tea, 542.
 Parahydroxybenzoic aldehyde, 494.
 Paraldehyde, 485.
 Paraldehydum, 485.
 impurities in, 716.
 Parallin, 507.
 Parapeptone, 551.
 Para-phenetol-carbamide, 431.
 Paratartaric acid, 320.
Pareiræ radix, 533.
 Paricine, 533.
 Parietinic acid, 338.
 Parigenin, 507.
 Parilla, 533.
 Paris blue, 556.
 red, 554.
 Particles, elementary, 36.
 Patent sugar, 469.
 Pearlash, 62.
 Pearl barley, 473.
 sago starch (fig.), 475.
 white, 252.
 Pear wine, 441.
 Pectin, 479.
 Pelargonate of ethyl, 407.
 Pelargonic acid, 490.
 Pelargonium, 418.
 Pelletierine, 359.
 Pellitory-root, 425.
 Pelosine, 533.
 Pennyroyal, American oil, 418.
 oil, 419.
 Pennywort, Indian, 509.
 Pentachloride of antimony, 181.
 Pentads, 124.
 Pentane, 398.
 Pentathionic acid, 346.
 Penthidric alcohols, 465.
 Pentylic acid, 490.
 alcohol, 449.
 Pepper, black, 541.
 Pepper, Cayenne, 534.
 cubeb, 541.
 long, 541.
 Melegueta, 418.
 oil of, 541.
 resin of, 541.
 white, 541.
 Peppermint oil, 418.
Pepo, 510.
 Pepsin, 550.
 Pepsinum, 550.
 impurities in, 716.
 saccharatum, 551.
 vegetable, 551.
 Peptone, 550, 551.
 Peptones in urine, 574.
 Per-, meaning of, 148.
 Perbromates, 271.
 Percha tree, 420.
 Perchlorate of potassium, 295.
 Perchloric acid, 295.
 Perchloride of gold, 246.
 iron, 148.
 mercury, 201.
 platinum, 248.
 tin, 242.
 Perfumes, 415.
 Perhydrate of iron, 150.
Periodic law, 380.
 Periodide of ammonium, 273.
 mercury, 273.
 potassium, 273.
 Permanganate of potassium, 77, 232.
 standard solution of, 652.
 vol. est. of, 635.
 Pernitrate of iron, 157.
 Peroxide of barium, 103.
 hydrogen, 103, 353.
 iron, 152.
 hydrated, 153.
 lead, 212.
 nitrogen, 289, 290.
 Perry, 441.
 Persian berries, 553.
 Personne's solution, 644.
 Persulphate of iron, 151.
 mercury, 200.
 Persulphide of hydrogen, 303.
 Peru, balsam of, 422, 495.
 Peruvine, 495.
 Petalite, 227.
 Petit-grain oil, 416.
Petrolatum, 398.
liquidum, 398.
 impurities in, 716.
 molle, 398.
 impurities in, 716.
 spissum, 398.
 impurities in, 716.

- Pétroléine**, 398.
Petroleum benzin, 398.
 ether, 398.
 gas, 397.
 light, 398.
 soft, 398.
 spirit, 398, 430.
 testing, 413.
Pettenkofer's test for presence of bile, 553.
Peumus boldus, 416.
Pewter, 185, 210, 241.
Phæoretine, 338.
Pharaoh's serpents, 357.
Pharbitis nil, 505.
Pharbitisin, 505.
Pharmaceutical Society of Great Britain, examinations of, 14.
Pharmacy, 15.
Pharmacognosy, 15.
Pharmacology, 15.
Phellandrene, 412.
Phellandrium aquaticum, 412.
Phenacetin, 431.
Phenacetinum, 431 (table) 544.
Phenazonum, 431 (table) 544.
Phenic acid, 452.
 alcohol, 452.
Phenocoll, 431.
Phenoëne, 429.
Phenol, 452.
 carbamine (table), 544.
 constitution of, 454.
 isonitrile (table), 544.
 salicylic, 492.
Phenol-mercury, 454.
Phenolphthalein, 434, 497, 635.
Phenols, 452.
Phenol-sulphonic acid, 445, 454.
Phenylacetamide, 431.
Phenylamine, 431, 514.
Phenylates, 454.
Phenylcarbinol, 456.
Phenyl-dimethyl-pyrazolon, 431.
Phenyl group, 431.
Phenylmethyl ketone, 498.
Phenyl salicylate, 492.
Phosphate of ammonium, 95.
 barium, 105, 332.
 calcium, 106, 112, 357.
 iron, 146, 156, 331.
 magnesium and ammonium, 122, 331.
 from oxalates and ferric oxide, separation of, 375.
 in bones, 112, 327.
 silver, 219.
 sodium, 88.
Phosphate of sodium, how prepared from phosphate of calcium, 113.
Phosphates, 327.
 analytical reactions of, 330.
 quantitative estimation of, 684.
Phosphites, 352.
 test for, 352.
Phospho-antimonic acid, 571.
Phosphomolybdic acid, 331, 571.
Phosphorated oil, 326.
Phosphoretted hydrogen, 343.
Phosphoric acid, 25, 327, 352.
 diluted, 329.
 glacial, 330.
 meta-, 330.
 ortho-, 330.
 pyro-, 330.
 quantitative estimation of free, 684.
 vol. est. of, 638.
 anhydride, 25, 330.
Phosphorous acid, 351, 352.
Phosphorus, 25, 327.
 acids of, 352.
 bromide of, 329.
 combustion of, 25.
 derivation of word, 32.
 detection of, in organic mixtures, 566.
 impurities in, 716.
 iodide, 329.
 granulated, 328.
 molecular weight of, 328.
 pill, 328.
 properties of, 25, 328.
 red or amorphous, 328.
 trihydride, 343.
Phosphotungstic acid, 571.
Phthaleins, 432.
Phthalic acid, 336, 432, 496.
 anhydride, 497.
 series of acids, 496.
Phyllocyanin, 457.
Phylloxanthin, 457.
Physical isomerides, 482.
Physics, 43.
Physostigma, 540.
 impurities in, 716.
Physostigmina, 540.
Physostigminæ salicylas, 540.
 impurities in, 716.
 sulphas, 540.
 impurities in, 716.
Physostigmine, 540.
 salicylate, 540.
Phytolacca fructus et radix, 510.
Phytolaccin, 510.
Picoline, 513.

- Picric acid, 454, 554.
 Picrotin, 506.
 Picrotoxin, 506, 544.
Picrotoxinum, 506.
 impurities in, 716.
 Pigment, 553.
Pigmentum nigrum, 557.
 Pills, 589.
 Pilocarpidine, 540.
 impurities in, 716.
Pilocarpinæ nitras, 540.
 hydrochloras, 540.
Pilocarpus, 540.
Pilocarpus pennatifolius, 540.
Pilula aloes et ferri, 144.
 ferri carbonatis, 145.
 compositæ, 146.
 iodidi, 31.
 phosphori, 328.
 plumbi cum opii, 211.
 Pimaric acid, 423.
 Pimento oil, 416.
Pimpinella anisum, 413.
 Pine-apple, essence of, 407.
 wool, 413.
 Pinene, 412.
 Pinic acid, 423.
 Pink saucers, 555.
 the common, 507.
 Pins, 242.
 Pint, 608.
Pinus, 412, 423.
 larix, 359, 412.
 Pipe-clay, 139.
 Piperazine, 512.
Piper nigrum, 541.
 Piperia, 541.
 Piperic acid, 541.
 Piperidia, 541.
 Piperidine, 541.
 Piperine, 541.
Piperinum, 541.
 impurities in, 716.
 Pipette, 272.
Pistachia terebinthus, 412.
 Pitch, 427.
 Burgundy, 425.
 Pituri, 540.
Pix burgundica, 425.
 liquida, 413, 426.
Plantago ispaghula, 479.
 Plants and animals, complementary
 action of, on air, 19.
 Plaster of ammoniacum and mer-
 cury, 195.
 mercury, 195.
 Paris, 106.
 Plasters, 213, 423.
 Plastic elements of nutrition, 548.
 Plastic sulphur, 301.
 Platinic chloride, 248.
 salts, 248.
 sulphide, 248.
 Platinous chloride, 248.
 salts, 248.
Platinum, 247.
 analytical reactions of, 248.
 and ammonium, chloride of, 99,
 249.
 and lithium, chloride of, 228.
 and potassium, chloride of, 79,
 249.
 and sodium, chloride of, 249.
 black, 248.
 derivation of word, 34.
 -foil, 247.
 perchloride, 248.
 residues, to recover, 249.
 spongy, 249.
 Pleurisy-root, 509.
 Plumbago, 30.
Plumbi acetas, 211.
 impurities in, 716.
 carbonas, 211.
 impurities in, 716.
 emplastrum, 213.
 iodidum, 213.
 impurities in, 716.
 nitras, 212.
 impurities in, 716.
 oxidum, 210.
 impurities in, 716.
 subacetatis, liquor, 212.
 Plumbic acetate, sulphate, etc.
 Vide Salts of Lead.
 peroxide, 212.
Plumbum, 30.
 "Plummer's pill," 203.
Pocula emetica, 180.
 Podophyllin, 425.
 Podophyllotoxin, 425.
Podophyllum, 425.
 resina, 425.
 Podophyllum resin, 425.
 Poisonous alkaloids, 566.
 Poisons of cheese, milk, fish, etc.,
 513, 571.
 antidotes to. See Antidotes.
 detection of, in organic mix-
 tures, 562, *et seq.*
 Pokeberry and root, 510.
 Polybasic acids, 264.
 Polybasyous radicals, 264.
 Polychroite, 554.
Polygala senega, 508.
 Polygalic acid, 508.
 Polyhydroxyl derivatives of hydro-
 carbons, 465.

- Polymerides, 481.
 Polymerism, 480.
 Polymers, 481.
 Polymorphism, 482.
 Polymorphous bodies, 482.
 Polysulphide of calcium, 303.
 Pomegranate-rind, 359.
 Pomegranate-root bark, 359.
 Poppy capsules, 516.
 Porcelain, 355.
 Porridge, 473.
 Porter, 441.
 Portland cement, 355.
 Port wine, 441.
 Positive pole, 245.
Potash, acetate, 70.
 alum, 139.
 bicarbonate, 71.
 bichromate, 238.
 bitartrate, 79.
 carbonate, 62.
 caustic, 63, 67.
 to prepare pure, 68.
 solution of, 63.
 chlorate, 293.
 citrate, 73.
 effervescing solution of, 73.
 iodate, solution of, 296.
 nitrate, 73, 285.
 permanganate, 77, 231.
 prussiate, red, 343.
 yellow, 280, 340.
 sulphate, 75.
 sulphurated, 68.
 tartrate, 74.
 acid, 79.
 and soda, 87.
 volumetric estimation of solu-
 tions of, 631.
 volumetric solutions of, 636.
 Potash-water, 72.
 Potashes, 62.
 Potassa, 68.
 impurities in, 717.
 cum calce, 68.
 sulphurata, 68.
Potassæ effervescens, liquor, 72.
 liquor, 63, 67.
 to prepare pure, 67.
 sulphuratæ unguentum, 69.
 Potassic hydrate, carbonate, etc.
 Vide Salts of Potassium.
Potassii acetat, 70, 242.
 impurities in, 717.
 bicarbonas, 71.
 impurities in, 717.
 bichromas, 238.
 bitartras, 74, 79, 320.
 impurities in, 717.
Potassii bromidum, 77.
 impurities in, 717.
 carbonas, 62.
 impurities in, 717.
 chloras, 293.
 impurities in, 717.
 citras, 73.
 impurities in, 717.
 cyanidum, 280.
 impurities in, 717.
 et sodii tartras, 74, 87, 320.
 impurities in, 717.
 ferricyanidum, impurities in,
 718.
 ferrocyanidum, 280, 340.
 hypophosphis, 344.
 impurities in, 717.
 iodidum, 74.
 impurities in, 718.
 nitras, 73, 285.
 impurities in, 718.
 permanganas, 79, 231.
 impurities in, 718.
 sulphas, 73, 288.
 impurities in, 718.
 Potassio-citrate of iron, 154.
 Potassio-tartrate of antimony, 181.
 iron, 154.
Potassium, 59.
 acetate, 70.
 acid carbonate. *Vide* Bicar-
 bonate.
 alizarate, 434.
 analytical reactions of, 78.
 and platinum chloride, 79, 249.
 and sodium tartrate, 87, 320.
 angelate, 413.
 anhydrochromate, 238.
 bicarbonate, 71.
 bichromate, 238.
 borotartrate, 334.
 bromate, 79.
 bromide, 79, 270.
 volumetric estimation of,
 642.
 carbonate, 62.
 chlorate, 16, 291.
 chloride, 78.
 chromate, 105.
 citrate, 73.
 cobalticyanide, 235.
 cyanate, 338.
 cyanide, 280.
 volumetric estimation of,
 641.
 derivation of word, 32.
 ferrate, 143.
 ferricyanide, 341.
 ferrocyanide, 280, 341.

- Potassium flame, 80, 101.
 hypophosphite, 344.
 volumetric estimation of, 655.
 hydrate, 63.
 to prepare pure solution, 68.
 standard solution, 636.
 iodate, 74, 296.
 iodide, 74, 274.
 volumetric estimation of, 643.
 manganate, 77, 231.
 myronate, 451.
 nitrate, 73, 285.
 nitrite, 350.
 oleate, 459.
 perchlorate, 295.
 periodide, 273.
 permanganate, 77, 231.
 volumetric estimation of, 652.
 preparation of, 63.
 properties of, 63.
 quantitative estimation of, 659.
 quantivalence of, 63.
 red chromate, 238.
 volumetric solution of, 648.
 red prussiate, 342.
 salts, analogy of, to sodium salts, 89.
 sodium and ammonium, separation of, 101.
 sources, 61.
 succinate, 356.
 sulphate, 73, 288.
 sulphide, 68.
 sulphocyanate, 356.
 volumetric solution of, 658.
 tartrate, 74, 320.
 acid, 62, 74, 79, 318.
 tri-iodide, 273.
 yellow chromate of, 105, 238.
 prussiate of, 280, 341.
- Potato, 472, 475.
 oil, 449.
 starch (fig.), 475.
- Poultices, 589.
- Pound, 608.
- Powder, bleaching, 114.
- Powders, 589.
 soda, 88.
 specific gravity of, 617.
- Practical analysis, 100.
- Precipitant, 79.
- Precipitate, 79.
- Precipitated carbonate of lime, 108.
 chalk, 19.
 sulphur, 303.
- Precipitates soluble in solutions of salts, 224.
 to wash, 109.
 to weigh, 681.
- Precipitation, 79.
 fractional, 378.
- Preparations of the pharmacopœias, chemical, 589.
 galenical, 589.
- Prepared calamine, 132.
 carbonate of calcium, 113.
 chalk, 143.
 lard, 462.
 suet, 462.
- Pressure, correction of volume of gas for, 619.
 gauges, 594.
- Prickly ash, 533.
- Primary alcohols, 436.
- Principles**, 35, *et seq.* See, also,
 "Analogy," "Atomic,"
 "Chemical Action, Affinity, and Combination," "Classification," "Constitution," "Formulae," "Fractional," "Homology," "Indestructibility," "Laws," "Molecular," "Notation," "Nomenclature," "Solution," "Substitution," "Structure," "Valency," etc.
- Printer's ink, 556.
- Prismatic nitre, 285.
- Proof spirit, 442.
- Propane, 397.
- Propanetricarboxylic acid, 498.
- Propargyl alcohol, 411.
- Propenyl, 457.
 alcohol, 457.
 hydrate, 457.
- Propepsin, 550.
- Propeptone, 551, 574.
- Prophetin, 503.
- Propione, 498.
- Propionic acid, 488.
- Proportions**, atomic, 51, 198.
 constant, 48.
 multiple, 49, 198.
 reciprocal, 50.
- Propylamine, 512.
- Propylene, 408.
- Propylformic acid, 488.
- Propylic acid, 488.
 alcohol, 457.
- Propylmethylbenzene, 432.
- Proteid principles, 544.
- Protocatechuic aldehyde, 364.
Protococcus vulgaris, 465.
- Protopine, 518, 541.

- Proximate analysis, 688.
 Prune, 469.
Prunum, 469.
Prunus serotina, 500.
 virginiana, 500.
 Prussian blue, 162, 342.
 Prussiate of potash, red, 342.
 yellow, 280, 341.
 Prussic acid, 280.
 Pseudaconitine, 530.
 Pseudojervine, 539.
 Pseudomorphine, 518.
 Pseudoxanthine, 513.
Pterocarpī lignum, 555.
 Pterocarpin, 555.
Pterocarpus marsupium, 359.
 santalinus, 419, 555.
 Ptomaines, 513, 571.
 Ptyalin, 586.
Ptychotis ajowan, 415.
 Puce-colored oxide of lead, 212.
 Puddling iron, 142.
 Pulezone, 419.
Pulvis algarothi, 181.
 angelicus, 181.
 antimonialis, 184.
 effervescens, 88, 321.
 effervescentes compositus, 321.
 ipecacuanhæ et opii, 537.
 morphinæ compositus, 517.
 sodæ tartaratæ effervescens, 321.
 Purified ox-bile, 552.
 Purple of Cassius, 247.
 foxglove, active principle in,
 502.
 pigment, 362.
 Purpurin, 579.
 Purrate of magnesium, 553.
 Purree, 553.
 Pusch's test for citric and tartaric
 acids, 326.
 Pus in urine, 585.
 Putrescine, 513.
 Putty-powder, 243.
 Pyrethric acid, 425.
 Pyrethrin, 425, 541.
Pyrethrum, 425.
 carneum, 425.
 Pyridine, 513.
 derivatives, 513, 536, 541.
 Pyrites, copper, 190.
 iron, 142.
 Pyroarsenate of sodium, 169.
 Pyroarsenates, 169.
 Pyroborate of sodium, 333.
 Pyrocatechin, 455.
 Pyrochromate of potassium, 238.
 Pyrogallol, 360, 465, 494.
 impurities in, 718.
 Pyroligneous acid, 297.
 Pyrolusite, 231.
 Pyromellitic acid, 498.
 Pyrometers, 598.
 Pyromorphite, 333.
 Pyrophorus, 159.
 Pyrophosphates, 353.
 Pyrophosphoric acid, 330, 353.
 Pyrosulphuric acid, 310.
 Pyrotartaric acid, 497.
 Pyrovanadates, 332.
 Pyroxylic spirit, 437.
Pyroxylin, 479.
 Pyrrol, 513.
Pyrus cydonia, 500.
- QUADRIVALENCE, 56.
 Qualitative analysis, 125, 370, 557.
Quantitative analysis, 598, *et seq.*
 determination of atmospheric
 pressure, 593.
 of temperature, 594.
 of weight, 600.
Quantivalence, 56, 123.
 of acidulous radicals, 67, 124,
 264.
 of atoms, definition of, 57.
 variation in, 139.
- Quartz, 354.
Quassia lignum, 506.
 Quassin, 506.
 Quebrachine, 530.
 Quebracho bark, 530.
 Queen's root, 542.
 Quercitrin, 554.
 Quercitron, 554.
Quercus cortex, 358.
 tinctoria, 554.
 Quevenne's iron, 158.
 Quicklime, 107.
 Quillaic acid, 508.
Quillaja, 508.
 saponaria, 508.
 Quinamine, 526.
 Quinate of quinine, 521.
 Quince-seed, 479, 500.
 Quinia. See Quinine, 521.
 Quinic acid, 521.
 Quinicine, 526.
 Quinidine, 524.
 sulphas, 525.
Quinina, 696.
 impurities in, 718.
Quininæ hydrochloras, 522.
 impurities in, 718.
 hydrobromas, 523.
 impurities in, 718.

- Quininæ bisulphas*, 522.
 impurities in, 718.
sulphas, 522.
 impurities in, 718.
valerianas, 523.
 impurities in, 719.
- Quinine**, 521.
 amorphous, 526.
 analytical reactions of, 523.
 bisulphate, 522.
 citrate, 523.
 of iron and, 55, 522.
 hydrobromate, 523.
 hydrochlorate, 522.
 iodosulphate, 523..
 kinate, 521.
 quantitative estimation of, 693.
 quininate, 521.
 sulphates, 522.
 valerianate, 523.
 wine, 522.
- Quiniretin**, 526.
- Quinlan's test** for presence of bile, 553.
- Quinoidine**, 526.
- Quinoline**, 514.
- Quinone**, 501, 521.
- Quinquivalence**, 56.
 of atoms, definition of, 59.
- RACEMIC acid**, 320.
- Radicals**, acidulous, 67, 123.
 formulae of, 67.
 basylous, 60, 123.
 definition of, 67.
 organic, 392.
- Rai**, 451.
- Raisins**, 318, 467.
- Raoult's experiments**, 623.
- Raspberry**, sugar in, 467.
- Ratafia**, 441.
- Rational formula**, 390.
- Rattan palm**, 424.
- Ratti**, 504.
- Reactions**, analytical, 62.
 synthetical, 62.
- Reagents**, list of, xiii.
- Real alcohol**, 442.
- Realgar**, 168.
- Réaumur's thermometer**, 596.
- Reciprocal proportions**, law of, 50.
- Rectification**, 129.
- Rectified oil** of turpentine, 412.
 spirit, 129, 442.
- Red**, Chinese, 554.
 chromate of potassium, 238.
 chrome, 554.
 coloring-matter, 554.
 corpuscles in blood, 546.
- Red earth**, 555.
 enamel colors, 555.
 gravel, 579.
 gum, 418.
 hæmatite, 142.
 iodide of mercury, 198.
 lead, 212.
 litmus-paper, 96.
 ochre, 555.
 oxide of iron, 142, 153, 555.
 mercury, 204.
 Paris, 554.
 phosphorus, 328.
 poppy-petals, 555.
 precipitate, 204.
 prussiate of potash, 341.
 rose-petals, 555.
 sandal-wood, 419, 555.
 sanders-wood, 420, 555.
 sulphide of mercury, 204.
 Venetian, 153.
- Reduced indigo**, 291.
 iron, 153.
 volumetric estimation of, 654.
- Reducing flame**, 135.
- Reinsch's test** for arsenum, 172.
- Relative weight** of hydrogen and oxygen, 24.
- Remijia bark**, 526.
- Rennet**, 546.
- Reseda luteola***, 554.
- Resin**, 412, 423.
 arnica, 423.
 cannabis, 423.
 capsicum, 424.
 castor, 424.
 ergot, 424.
 guaiacum, 504.
 Indian hemp, 423.
 jalap, 505.
 kaladana, 505.
 kamala, 425.
 koussou, 424.
 mastic, 424.
 mezereon, 425.
 oils, 422.
 pepper, 425, 541.
 podophyllum, 425.
 pyrethrum, 425.
 roottlera, 425.
 scammony, 508.
 soap, 462.
- Resina***, 423.
 copaibæ, 425.
 jalapæ, 505.
 impurities in, 719.
- Resins**, 423.
- Resorcin**, 423.

- Resorcinum*, 456.
 impurities in, 719.
 Respiratory materials of food, 548.
 Retort, 128.
Rhamni frangula cortex, 338.
 Purshiani cortex, 338.
 succus, 502, 556.
 Rhamnin, 554.
 Rhamnose, 554.
Rhamnus catharticus, 556.
 frangula, 503.
 Rhaponticin, 338.
 Rhatany-root, 359.
 Rheic acid, 338.
 Rhein, 338.
 Rheum, 338.
 Rheumin, 338.
 Rhodium, 726.
Rhæadas petala, 555.
 Rhœadine, 518.
 Rhubarb, oxalate of calcium from,
 316.
 Rhubarbaric acid, 338.
 Rhubarbarin, 338.
 coriaria, 360.
Rhus cotinus, 553.
 glabra, 360.
 toxicodendron, 361.
 Rice, 473.
 starch (fig.), 475.
 Ricin, 464.
 Ricinine, 464.
 Ricinoleate of glyceryl, 464.
 Ricinoleine, 464.
 Ringworm powder, 338.
Roccella tinctoria, 465.
 Roché alum, 140.
 Rochelle salt, 74, 87, 320.
 Rock alum, 140.
 salt, 81.
 Rohun-bark, 510.
 Roll sulphur, 301.
 Roman cement, 355.
 Rosa canina, 364.
Rosæ caninæ fructus, 469.
 centifoliæ petala, 419, 555.
 gallicæ petala, 555.
 Rose, 419, 469, 555.
 oil, 419.
 Roseaniline, 431, 557.
 Rosemary oil, 419.
 Rose-petals, 419, 555.
 Rose-water, 414, 419.
 Rosin, 412, 423.
 Rotang palm, 424.
 Rotten-stone, 138.
Rottlera tinctoria, 425.
 Rottlerin, 425.
 Rouge, animal, 337.
 Rouge, mineral, 153, 555.
 vegetable, 555.
Rubia tinctoria, 554.
 Rubianic acid, 554.
 Rubidium, 726.
 Rubijervine, 439.
Rubus, 360.
 Ruby, 138.
 Rue oil, 419.
 Rum, 441.
 Rumex, 338.
 Rumicin, 338.
 Rust of iron, 143.
 Rutate of glyceryl, 462.
 Ruthenium, 726.
 Rutic acid, 462.
 aldehyde, 419.
 radical, 462.
 SABADILLA, 543.
 Sabadilline, 543.
 Sabadine, 543.
 Sabadinine, 543.
 Sabatrine, 543.
Subina, 419.
 Saccharated carbonate of iron, 145.
 volumetric estimation
 of, 653.
 Saccharic acid, 472.
 Saccharin, 446.
 soluble, 446.
 Saccharine, 469.
 Saccharometer, 703.
 Saccharometry, 702.
Saccharomyces cerevisiæ, 440.
 Saccharons, 469.
 Saccharoses, 467, 469.
Saccharum lactis, 471.
 impurities in, 719.
 purificatum, 469.
 ustum, 471.
 Sacred bark, 338.
 Safety lamp, 24.
 tube, 266.
 Safflower, 555.
 Safranin, 554.
 Saffron, 554.
 bastard, 555.
 dyer's, 555.
 oil of, 554.
 Safrol, 420.
 Sage, oil of, 419.
 Sago, 473.
 starch (fig.), 475.
 Sal ammoniac, 91.
 prænella, 286.
 volatile, 94.
 Salep, 479.
 Salicin, 456, 506.

- Salicinum*, 506.
 impurities in, 719.
 Salicylate of methyl, 407, 492.
 phenyl, 492.
 sodium, 493.
 Salicylic acid, 407, 457, 492.
 aldehyde, 407, 457, 493.
 Salicylol, 407.
 Salicylous acid, 407.
 Saligenin, 492, 506.
 alcohols, 457.
 Saligenol, 453, 506.
 Saliretin, 506.
 Saliva, 586.
 Salol, 492.
 Salol, 492.
 impurities in, 719.
 Salseparin, 507.
 Salt cake, 312.
 common, 81.
 definition of a, 60.
 Epsom, 118.
 of sorrel, 316.
 Rochelle, 74, 87, 320.
 rock, 81.
 Saltpetre, 286.
 Chili, 286.
Salts, acids, 73, 74, 302.
 action of the blowpipe on, 373.
 of heat on, 372.
 of sulphuric acid on, 372.
 alkyl, 484.
 analogies of, 90.
 analysis of insoluble, 376.
 constitution of, 61, 124, 262, 286,
 299, 378.
 résumé, 378.
 double, 79.
 formation of, 79.
 nomenclature of, 73, 76.
 of ammonium, volatility of, 99.
 of iron, nomenclature of, 143.
 physical properties of, 371.
 substitution of, for each other,
 89.
 table of the solubility or insol-
 ubility of, in water, 368.
 Salviol, 419.
 Sambucene, 419.
Sambuci flores, 419.
 Sand, 355.
 Sandal-wood, oil of, 419.
 red, 555.
 white, 419.
 yellow, 419.
 Sand-bath, 29.
 Sand-stone, 354.
 Sand-tray, 29.
 Sanders-wood, red, 555.
Sanguinaria canadensis, 541.
 Sanguinarine, 541.
 Santalin, 555.
Santalum album, 419.
 Santonic acid, 507.
Santonica, 507.
 Santonin, 507.
Santoninum, 507.
 impurities in, 719.
 Santoniretin, 507.
 Sap-green, 556.
 Sapan-wood, 554.
Sapo, 461.
 impurities in, 719.
 animalis, 461.
 durus, 461.
 kalinus venalis, 461.
 mollis, 461.
 viridis, 461.
 Sapogenol, 507.
 Saponetin, 507.
 Saponification, 462.
 Saponin, 507.
 Sapotoxin, 508.
 Sapphire, 138.
 Saprine, 513.
Sarcina ventriculi in urine, 587.
Sarcocephalus esculentus, 424.
 Sarcolactic acid, 348, 489.
 Sarkine, 515.
 Sarkosine, 512.
Sarracenia purpurea, 543.
Sarsaparilla, 507.
 Sassafras oil, 420.
 radix, 420.
 swamp, 509.
 Sassafrol, 420.
 Saturated hydrocarbons, 391.
 solutions, boiling-points of, 596.
 Saturating power of citric acid,
 325.
 of tartaric acid, 321.
 Saturation, 71.
 tables, 724.
 Saturn, 211.
 Saturnine colic, 211.
 Savin oil, 419.
 Saxon blue, 555.
 Saxony blue, 555.
 Scale compounds of iron, 154, *et seq.*
Scammonie radix, 508.
 resina, 508.
 Scammonin, 508.
 Scammonioid, 508.
Scammonium, 508.
 impurities in, 719.
 Scammony, resin of, 508.
 Scandium, 726.
 Scents, 415.

- Scheele's green, 178.
 Schist, 138.
Schœnocaulon officinale, 543.
 Schonbein's test for hydrocyanic acid, 284.
 Schweinfurth green, 177.
 Science of Chemistry, 14.
Scilla, 470, 508.
 Scillin, 508.
 Scillipierin, 508.
 Scillitin, 508.
 Scillitoxin, 508.
 Sclerotic acid, 424.
 Sclerotinic acid, 424.
 Scoparin, 542.
Scoparius, 542.
 Scopoleine, 532.
 Scopoletin, 532.
Scopola Japonica, 515, 532.
 Scutellaria, 510.
 Scyllite, 469.
 Sea-salt, 81.
 Sebacate of ethyl, 407.
Secale cereale, 424.
 Secondary alcohols, 436.
 Sediments, urinary, 579.
 microscopic examinations of, 581.
 Seed-lac, 555.
 Seidlitz powder, 321.
 Selenic acid, 302.
 Selenion, 726.
 Selenious acid, 320.
 Selenium, 726.
Semina cardamomi majoris, 418.
Senegæ radix, 508.
Senna, 501.
Senna of Alexandria, 501.
 Sepia, 556.
 Sepiadæ, 556.
 Serolin, 546.
Serpentaria, 510.
 Serpent's excrement, 580.
 Sesamé oil, 464.
Sesamum indicum, 464.
 Sesquiterpene, 412.
Sevum præparatum, 462.
 Sexivalence, 57.
 Shale, 138.
 Shark-liver oil, 465.
 Shellac, 555.
 Shell-fish poison, 513.
 Sherry wine, 441.
 Shot, 210.
 Siam benzoin, 364.
 Sidee, 424.
 Sienna, 556.
 Sifting an aid to analysis, 372, 378.
 Silica, 354.
 Silicate of aluminium, 354.
 calcium, 354, 355.
 lithium, 227.
 magnesium, 354.
Silicates, 354.
 quantitative estimation of, 687.
 tests for, 355.
 Silicic acid, 354, 355.
 anhydride, 354.
 Siliciuretted hydrogen, 355.
 Silicon, chloride, 356.
 derivation of word, 34.
 fluoride, 342, 356.
 hydride, 355.
 oxide, 355.
Silver, 216.
 ammonio-nitrate, 177, 207.
 analytical reactions of, 219.
 antidotes, 220.
 arsenate, 177, 219.
 arsenite, 177.
 bromide, 220, 271.
 chloride, 217.
 chromate, 220.
 citrate, 326.
 coinage, 216, 617.
 cyanide, 220, 283.
 derivation of word, 35.
 estimation of, by cupellation, 676.
 extraction of, 216.
 fulminate, 219.
 German, 171, 236.
 iodide, 228, 275.
 nickel, 236.
 nitrate, impure, 216.
 pure, 217.
 toughened, 218.
 volumetric solution of, 639.
 oxalate, 317.
 oxide, 219.
 phosphate, 219.
 pure, 217.
 quantitative estimation of, 675.
 standard solution of nitrate of, 639.
 sulphate, 216.
 sulphide, 219.
 native, 216.
 sulphite, 307.
 tartrate, 322.
 tree, 220.
 Sinalbin, 451.
Sinapis, 451.
 albæ, 451.
 impurities in, 719.
 juncea. See *Brassica*.
 nigræ, 451.
 impurities in, 719.

Sinapisine, 451.
 Sinigrin, 451.
 Sinistrin, 507.
 Siphon; 110.
 Size, 549.
 Skullcap, 510.
 Slaked lime, 107.
 Slate, 138.
 Smalt, 235, 555.
 Smilacin, 507.
 Snake-root, black, 509.
 Virginia, 510.
Soap, ammonium, calcium, Castile,
 green, hard, Marseilles, mot-
 tled, potassium, sodium, soft,
 461.
 curd, 461.
 marine, 462.
 resin, 462.
 yellow, 462.
 Soap-bark, 508.
 Soap-stone, 557.
 Soap-wort, 507.
 Socaloin, 435.
 Socatrine aloes, 434.
Soda, 82, 312.
 acetate, 83.
 alum, 139.
 arsenate, 169.
 ash, 89, 312.
 benzoate, 337.
 bicarbonate, 83, 313.
 carbonate, 84, 89, 312.
 caustic, 82.
 chlorinated, 87.
 citro-tartrate, 88.
 hydrate, 82.
 hypochlorite, 87.
 hypophosphite, 344.
 hyposulphite, 345.
 lime, 691.
 nitrate, 82, 286.
 phosphate, 88, 113.
 powders, 88.
 salicylate, 493.
 solution of chlorinated, 87.
 standard solution of, 639.
 sulphate, 267.
 sulphite, 306.
 sulphocarbonate, 454.
 valerianate, 362.
 volumetric estimation of, 632.
 solution of, 639.
Soda, 82.
 impurities in, 719.
 tartarata, 74, 87, 319.
 Soda-water, 86.
 Sodic carbonate, etc. *Vide* Salts of
 Sodium.

Sodii acetat, 83.
 impurities in, 719.
arsenas, 169.
 impurities in, 719.
benzoas, 337.
 impurities in, 719.
bicarbonas, 83, 313.
 impurities in, 719.
bisulphis, 306.
 impurities in, 720.
boras, 333.
 impurities in, 720.
bromidum, 88.
 impurities in, 720.
carbonas, 84, 313.
 impurities in, 720.
 exsiccatus, 85.
chloras, 294.
 impurities in, 720.
chloratæ, liquor, 87.
chloridum, 81.
 impurities in, 720.
citro-tartras effervescens, 88.
hypophosphis, 344.
 impurities in, 720.
hyposulphis, 345.
 impurities in, 720.
iodidum, 88.
 impurities in, 720.
nitras, 286.
 impurities in, 720.
nitris, 351.
 impurities in, 720.
phosphas, 113.
 impurities in, 721.
 effervescens, 88.
salicylas, 493.
 impurities in, 721.
silicatis, liquor, 355.
sulphas, 267.
 impurities in, 721.
 effervescens, 88.
sulphis, 306.
 impurity in, 721.
sulphocarbonas, 454.
 impurity in, 721.
valerianas, 362.
 Sodio-citrate of iron, 156.
Sodium, 81.
 acetate, 83.
 acid carbonate, 83.
 sulphate, 265.
 analytical reactions of, 89.
 and aluminium double chloride,
 138.
 and platinum chloride, 249.
 and potassium, tartrate of, 87,
 318.
 arsenate, 169.

- Sodium arsenite, 168, 169.
 benzoate, 337.
 biboate, 333.
 bicarbonate, 83, 88, 313.
 chemically pure, 628.
 lozenges, 87.
 bisulphite, 306.
 volumetric estimation of, 648.
 bromate, 77, 271.
 bromide, 88.
 carbonate, 81, 85, 313.
 chemically pure, 628.
 manufacture of, 88, 312.
 chlorate, 294.
 chloride, 81.
 pure, preparation of, 658.
 volumetric solution of, 657.
 cholate, 552.
 citro-tartrate, 88.
 derivation of word, 32.
 ethylate, 444.
 flame, 89, 101.
 glycocholate, 552.
 gravimetric estimation, 663.
 hydrate, 82.
 hypochlorite, 87.
 hypophosphite, 344.
 volumetric estimation of, 654.
 hyposulphite, 345.
 iodide, 87.
 manganate, 232.
 metaborate, 333.
 nitrate, 81, 286.
 nitrite, 351.
 oxalate, 316.
 permanganate, 232.
 phosphate, 113.
 effervescent, 88.
 how prepared from phosphate of calcium, 113.
 potassium, and ammonium, separation of, 111.
 pyroarsenate, 169.
 pyroborate, 333.
 pyrophosphate, 353.
 quantitative estimation of, 663.
 salicylate, 493.
 salts, analogy of, to potassium salts, 89.
 sources of, 81.
 stannate, 243.
 sulphate, 267.
 effervescent, 88.
 sulphite, 306.
 volumetric estimation of, 647.
 sulphycarbolate, 454.
 Sodium tartrate, 321.
 taurocholate, 552.
 valerianate, 362.
 Soft soap, 461.
 Soils, analysis of, 706.
 Solania, 541.
 Solanidine, 542.
 Solanine, 541.
 Solanum dulcamara, 541.
 tuberosum, 541.
 starch of (fig.), 475.
 Solazzi juice, 504.
 Soldier, 210, 241.
 Solid, definition of, 58.
 fats, 462.
 potash, 68.
 Solids, to take the specific gravity of, 617, *et seq.*
 heavier than water, to take the specific gravity of, 617.
 Solubility of carbonic-acid gas in water, 87.
 of gases in water, 87.
 or insolubility of salts in water, table of, 368.
 Soluble cream of tartar, 334.
 glass, 355.
 saccharin, 445.
 starch, 477.
 substances, to take the specific gravity of, 617.
 tartar, 74.
 Solution of albumen, 544.
 ammonia, 92.
 ammonio-nitrate of silver, 177, 207.
 ammonio-sulphate of copper, 177.
 magnesium, 685.
 ammonium acetate, 92.
 chloride, 91.
 citrate, 95.
 oxalate, 95.
 antimony chloride, 180.
 arsenic, acid, 169.
 alkaline, 167.
 barium chloride, 103.
 bismuth and ammonium citrate, 254.
 boric acid, 334.
 bromine, 271.
 calcium chloride, 103.
 carbonate of ammonium, 94.
 chlorinated lime, 115.
 soda, 87.
 chlorine, 28, 268.
 copper acetate, 192.
 ferric chloride, 149.
 fractional, 86, 89, 378.

- Solution** of gold chloride, 246.
 perchloride, 246.
 gutta-percha, 421.
 iodine, 274.
 iron acetate, 154.
 perchloride, 149.
 pernitrate, 158.
 persulphate, 151.
 isinglass, 549.
 lead oxyacetate, 116, 211.
 lime, 108.
 litmus, 96, 636.
 mercury nitrate, 200.
 perchloride, 202.
 nature of, 139, 623.
 phosphoric acid, 228.
 platinum perchloride, 248.
 potash, 63.
 effervescing, 72.
 potassio-mercuric iodide, 658.
 potassium acetate, 70.
 ferricyanide, 341.
 ferrocyanide, 341.
 iodate, 296.
 red prussiate of potash, 341.
 soda, 82.
 sodium acetate, 83.
 ethylate, 443.
 phosphate, 113.
 silicate, 355.
 stannous chloride, 242.
 sulphate of calcium, 116.
 indigo, 291.
 iron, 151, 155.
 sulphhydrate of ammonium, 96.
 tartaric acid, 320.
 tin chloride, 242.
 yellow chromate of potassium, 105.
 zinc chloride, 133.
Sonnenschein's process for poisonous
 alkaloids, 569.
Soporine, 536.
Sorbin, -ose, 469.
Sorrel salt, 316.
Soot, 30.
Soubresauts, 282.
Source of heat, 18.
Sovereign, weight of the, 246.
Soymida febrifuga, 510.
Sozoiodol, 445.
Sozolic acid, 445.
Spanish liquorice, 504.
Spar, fluor-, 106, 342.
 heavy, 102.
Sparteine sulphas, 542.
 impurities in, 721.
Sparteine, or **Sparteia**, 542.
Spathic iron ore, 142.
Spearmint oil, 419.
Specific gravity, 613, *et seq.*
 bottles, 614.
 of gases, 619.
 of liquids, 614.
 of official liquids, 614.
 of oxygen, 24.
 of powders, 617.
 of solids, 616.
 lighter than water, 618.
 of soluble substances, 618.
 heat, 130, 622.
 weight, 613.
Spectral analysis, 262, 560.
Spectroscope, 262, 560.
Specular iron ore, 142.
Speculum metal, 241.
Speiss, 236.
Sperm oil, 450.
Spermaceti, 450.
Spermatozoa in urine, 586.
Spigeline, 542.
Spiræa ulmaria, 407, 493, 507.
Spirit, methylated, 437.
 of French wine, 443.
 of myrcia, 441.
 of nitrous ether, 403.
 adulterated, 438.
 of turpentine, 412.
 of wine, 442.
 petroleum, 398, 430.
 proof, 442.
 pyroxylic, 437.
 rectified, 129, 442.
 wood-, 437.
Spirits, 441.
 analysis of, 704.
Spiritus ætheris, 449.
 compositus, 409.
 nitrosi, 351, 403.
 impurities in, 721.
 ammoniacæ, 93.
 aromaticæ, 94, 615.
 fœtidus, 94.
 armoraciæ compositus, 415.
 cajuputi, 415.
 chloroformi, 401.
 cinnamomi, 415.
 frumenti, 441.
 impurities in, 721.
 juniperi, 415.
 lavandulæ, 415.
 menthæ piperitæ, 415.
 myrciæ, 441.
 myristicæ, 415.
 rectificatus, 442.
 rosmarina, 415.
 tennior, 442.
 vini gallici, 443.

- Spiritus vini gallici*, impurities in, 721.
 Spodumene, 227.
 Spogel-seeds, 479.
 Sponge, 548.
 Spongy platinum, 249.
 Spontaneous combustion, 159.
 Spotted cranesbill, 360.
 Spruce fir, 425.
 Spurge laurel, 425.
Squalus carcharias, 465.
 Squill, 470, 479, 508.
 bulb, 508.
 vinegar of, 297.
Standard solution of hyposulphite
 of sodium, 655.
 iodine, 645.
 nitrate of silver, 639.
 oxalic acid, 634.
 potash, 636.
 potassium permanganate,
 652.
 red chromate of potassium,
 648.
 soda, 639.
 sulphuric acid, 628.
 Stannate of sodium, 243.
 Stannates, 243.
 Stannic acid, 243.
 anhydride, 243.
 chloride, 242.
 oxide, 241, 243.
 sulphide, 244.
 anhydrous, 244.
 Stannous chloride, 242.
 solid, 242.
 hydrate, 244.
 oxide, 244.
 sulphide, 243.
Stannum, 34.
Staphisagriæ semina, 536.
 Star-anise oil, 415.
Starch, 472.
 action of diastase upon, 477.
 of dilute acids upon, 477.
 animal, 476.
 blue, 472.
 cellulose, 474.
 granules, composition of, 473.
 iodide of, 275, 473.
 mucilage of, 473.
 potato, 472.
 quantitative estimation of,
 702.
 soluble, 476.
 wheat, 472.
 white, 472.
 Starches, microscopy of, 474.
 Stas's process for poisonous alka-
 loids, 568.
 Stavesacre, 536.
 Steam-bath, 112.
 Stearic acid, 457, 488, 490.
 Stearine, 459.
 Stearoptens, 413.
 Steatite, 557.
 Steel, 142.
 wine, 157.
 Stereochemical conceptions, 390,
 393.
 Stibines, 512.
Stibium, 35.
 Stick lac, 555.
 liquorice, 504.
 Still, 127.
Stillingia, 542.
 Stone coal, 241.
 red, 555.
 ware, 355.
 Storax, 422, 495.
 Stout, 441.
Stramonii folia, 538.
 semina, 538.
 Strasburg turpentine, 412.
 Strawberry, sugar in, 467.
 Stream-tin, 241.
 Strontianite, 229.
Strontii bromidum, 229.
 impurities in, 721.
 iodidum, 229.
 impurities in, 721.
 lactas, 229.
 impurities in, 721.
Strontium, 229.
 analytical reactions of, 229.
 bromide, 229.
 volumetric estimation of,
 643.
 carbonate, 229.
 derivation of word, 34.
 flame, 230.
 nitrate, 229.
 sulphate, 229.
 Strophanthidin, 509.
 Strophanthin, 509.
 Strophanthus, 509.
 Structural formulæ, 386.
Structure, as indicated by—
 chemical "periodicity," 380.
 properties, 388, 396.
 density of gases and vapors, 53,
 55, 619.
 electricity, etc., 623.
 isomorphism, 55.
 specific heat, 622.
 substitution, 193.
 Structure of flame, 23.
 of molecules, 138, 139, 193, 387,
 392.

- Strychnine**, or *Strychnia*, 527.
 impurities in, 721.
 analytical reactions of, 528.
 citrate, 527.
 in organic mixtures, detection of, 566.
 sulphate, 527.
 with brucine, estimation of, 700.
Strychnos ignatus, 527.
 nux vomica, 506, 527.
Styracin, 495.
Styrax præparatus, 495.
Styrol, 495.
Styrone, 495.
Subacetate of copper, 192.
 lead, 211.
Subcarbonate of iron, 145.
 bismuth, 253.
Subchloride of iron, 149.
 mercury, 203.
Suberate of ethyl, 407.
Sublimation, 94.
Sublimed sulphur, 301.
Subnitrate of bismuth, 252.
Substances readily deoxidized,
 quantitative estimation of, 652.
 oxidized, quantitative estimation of, 645.
Substitution, 172, 193, 394.
 products, 399.
Succi, 589.
Succinate of potassium, 336.
Succinates, 336.
Succinic acid, 336, 496.
Succinum, 336.
Succus limonum, 324.
Sucrose, 469.
Suet, 462.
 prepared, 462.
Suffioni, 333.
Sugar, action of alkali upon, 470.
 amount in various fruits, 468.
 barley, 471.
 brown, 469.
 candy, 469.
 detection of, in urine, 574.
 from caoutchouc, 469.
 eucalyptus, 471.
 fish, 469.
 gum arabic, 469.
 larch, 471.
 melitose, 471.
 mountain ash, 469.
 muscles, 469.
 starch, 469.
 Turkish manna, 471.
 grape-, 467.
 inverted, 468.
Sugar, lichen, 465.
 lump, 469.
 maple, 471.
 milk, 471.
 moist, 469.
 of gelatin, 550.
 of lead, 211.
 patent, 469.
 quantitative estimation of, 702.
 sand, 471.
 syrup, 469.
 test for, 468.
Sugar-cane, 468, 469.
Suint, 460.
Sulphamido-benzoates, 446.
Sulphate of aluminium, 139.
 and ammonium, 139.
 ammonium, 91.
 barium, 103.
 beberine, 532.
 bismuth, 253.
 calcium, 106, 115.
 chromium, 238.
 cinchonidine, 525.
 cinchonine, 525.
 cobalt, 235.
 copper, 192.
 anhydrous, 192.
 cupr - diammon - diammonium, 207.
 indigo, 291.
 iron, 144.
 and ammonium, 138, 144.
 solution of, 144.
 lead, 215.
 lithium, 228.
 magnesium, 118.
 mercury, 200, 201.
 morphine, 517.
 potassium, 73, 288.
 quinine, 522.
 sodium, 267.
 strontium, 229.
 strychnine, 527.
 zinc, 131.
Sulphates, 307.
 analytical reactions of, 310.
 quantitative estimation of, 682.
Sulphethylic acid, 447.
Sulphide of allyl, 452.
 ammonium, 96.
 antimony, 179, 184.
 arsenum, 168, 176.
 native, 168.
 barium, 103.
 bismuth, 254.
 cadmium, 250.
 calcium, 115.
 cobalt, 235.

- Sulphide of iron**, 31, 144, 161, 162.
 lead, 214.
 native, 210.
 manganese, 233.
 mercury, 207.
 native, 195.
 molybdenum, 331.
 nickel, 237.
 potassium, 68.
 silver, 219.
 native, 216.
 zinc, 136.
 native, 131.
- Sulphides**, 301.
 analytical reactions of, 304.
 quantitative estimation of, 681.
- Sulphindigotic acid**, 291.
- Sulphindyllic acid**, 291.
- Sulphite of barium**, 307.
 calcium, 307.
 silver, 307.
 sodium, 306.
 zinc, 136.
- Sulphites**, 305.
 analytical reactions of, 306.
 quantitative estimation of, 682.
- Sulphocarbonates**, 454.
- Sulphocarbolic acid**, 454.
- Sulphocarbonates**, 313.
- Sulphocarbonic anhydride**, 314.
- Sulphocyanate of acrinyl**, 451.
 allyl, 451.
 butyl, 415.
 iron, 162, 336.
 mercury, 357.
 tetryl, 415.
- Sulphocyanates**, 336.
- Sulphocyanic acid**, 336.
- Sulphocyanides**, 357.
- Sulphocyanogen**, 357.
- Sulphonol**, 446; (table) 544.
- Sulphonic acids**, 445.
- Sulphophenates**, 454.
- Sulphophenic acid**, 454.
- Sulphostannates**, 244.
- Sulphovinic acid**, 310, 447.
- Sulphur**, 30, 301.
 adulteration of, 304.
 alcohols, 445.
 allotropy of, 301.
 analogies, 177, 302.
 analytical reactions of, 304.
 arsenic in, 176.
 bromide, 304.
 chloride, 304.
 derivation of word, 32.
 estimation of, 681.
 ethers, 445, 447.
 flowers of, 301.
- Sulphur hypochloride**, 304.
 iodide, 274.
 liver of, 68.
 lozenges, 303.
 milk of, 302.
 molecular weight of, 302.
 oxyacids, 346.
 plastic, 301.
 precipitated, 303.
 roll, 301.
 sublimed, 301.
- Sulphur lotum**, 301.
 impurities in, 721.
- præcipitatum**, 303.
 impurities in, 721.
- sublimatum**, 301.
 impurities in, 721.
- Sulphurated antimony**, 182.
 lime, 115.
 potash, 68.
- Sulphurets**. *Vide* Sulphides.
- Sulphuretted hydrogen**, 97, 144, 301.
- Sulphuric acid**, 307.
 antidotes to, 311.
 aromatic, 310.
 diluted, 310.
 estimation of, 638.
 in vinegar, 682.
 fuming, 310.
 Nordhausen, 310.
 organic mixtures, detection
 of, in, 564.
 purification of, 309.
 standard solution of, 628,
 634.
 volumetric estimation of,
 638.
 anhydride, 310.
- Sulphuris iodidum**, 274.
- Sulphurous acid**, 30, 305.
 volumetric estimation of,
 645.
 anhydride, 305.
- Sulphydrate of ammonium**, solu-
 tion of, 96.
- Sulphydric acid**, 301.
- Sumatra camphor**, 421.
- Sumbul**, 427.
 radix, 427.
 root, 427.
- Supporters of combustion**, 23.
- Suppositoria acidi tannici**, 358.
cum sapone, 358
morphinæ, 517.
plumbi composita, 211.
- Suppositories**, 589.
- Surface unit**, 603.
- Surgery**, 15.
- Swamp sassafras**, 509.

- Sweet birch, oil of, 407.
 flag, oil of, 420.
 spirit of nitre, 351, 403, 438.
 adulterated, 438.
- Sweetbread, 551.
- Sylvestrine, 412.
- Sylvic acid, 423.
- Symbol, function of a, 58.
- Symbols** of elements, 32, *et seq.*, 725.
 illustration of chemical action
 by, 46.
- Sympathetic inks, 236.
- Symmetrical compounds, 433.
- Synaptase, 500.
- Synthesis, 62.
- Syphon. See Siphon.
- Syrup, golden, 471.
- Syrups, 589.
 specific gravities of, 615.
- Syrupus*, 469, 702.
 acidi hydriodici, 274.
 impurities in, 721.
 aurantii, 416.
 floris, 416.
 calci lacto-phosphatis, 173.
 calcis, 108.
 impurities in, 721.
 ferri iodidum, 148.
 phosphatis, 146.
 subchloridi, 149.
- TABACUM**, 539.
- Tabellæ nitroglycerina*, 458.
- Tables, various. See Appendix.
- Talc, 138.
- Tamarindus*, 323.
 impurities in, 721.
- Tampico jalap, 505.
- Tanacetie acid, 507.
- Tanacetum*, 507.
- Tannic acid, or Tannin, 357, 495.
- Tanning, 358.
- Tantalum, 726.
- Tapioca, 473.
 starch (fig.), 475.
- Tar, 413, 426.
- Taraxaci radix*, 510.
- Taraxacin, 510.
- Tartar, cream of, 62, 79, 319.
 emetie, 189, 319.
 estimation of antimony in,
 671.
 meaning of, 319.
 soluble, 74.
 cream of, 334.
- Tartarated antimony, 181.
- Tartaric acid, 318, 496.
 saturating power of, 321.
 solution of, 320.
- Tartaric acid, volumetric estimation
 of, 639.
 series of acids, 496.
- Tartarus boraxatus*, 334.
- Tartrate** of ammonium, 99, 181.
 antimony and potassium, 181,
 319.
 calcium, 322.
 morphine, 517.
 potassium acid, 62, 74, 79, 318.
 neutral, 74, 320.
 and sodium, 87, 320.
 silver, 322.
 sodium, 321.
- Tartrates**, 318.
 analytical reactions of, 322.
 volumetric estimation of, 632.
- Tartronic acid, 497.
- Taurine, 512, 552.
- Taurocholates, 552.
- Taxine, 542.
- Tea, 542.
- Teal oil, 464.
- Telini fly, 422.
- Tellurium, 726.
- Temperature, correction of volume
 of gas for, 619.
 quantitative determination of,
 594.
- Terebenthene, 412.
- Terebene, 413.
- Terebenum*, 413.
 impurities in, 721.
- Terebinthinæ canadensis*, 412, 427.
- Terephthalic acid, 497.
- Terpene series of hydrocarbons, 412.
- Terpenes, 412.
- Terpin hydrate, 413.
- Terpini hydras*, 413.
 impurities in, 721.
- Terpinene, 412.
- Terpinol, 413.
- Terpinolene, 412.
- Terra di Sienna, 555.
- Tertiary alcohols, 436.
 amylic alcohol, 450.
- Terra japonica*, 359.
- Test-papers, 96.
- Test-tubes, 16.
- Tetanine, 513.
- Tetano-caunabin, 423.
- Tetrabasic acid, 498.
- Tetrachloride of carbon, 401.
- Tetrachloromethane, 401.
- Tetrads, 124.
- Tetrahydric alcohols, 465.
- Tetra-iodo-pyrrol, 513.
- Tetramines, 512.
- Tetrane, 397.

- Tetrathionic acid, 346.
 Tetrylic acid, 488, 497.
 sulphocyanate, 415.
 Thalleioquin, 523.
 Thalline, 526.
 Thallium, 726.
 Thebaine, 518.
 Theia, 542.
 Theine, 542.
 relation to theobromine, 543.
 Thenard's blue, 555.
Theobroma cacao, 462, 543.
 oil, 462.
 Theobromine, relation of, to theine, 543.
 or Theobromia, 543.
 Theophylline, 543.
 Therapeutics, definition and derivation of, 15.
Theriaca, 471.
 Thermolysis, 622.
 Thermometers, 595.
 Celsius's, 595.
 Centigrade, 595.
 Fahrenheit's, 595.
 Réaumur's, 595.
 Thermometric scales, conversion of degrees of, 596.
 Thio-acids, 302.
 Thio-alcohols, 445.
 Thio-ethers, 445, 447.
 Thionic acids, 346.
 Thiosulphates, 345.
 Thorinum, 726.
 Thorium, 726.
 Thorn-apple, 538.
 Thoroughwort, 509.
Thus americanum, 427.
 masculum, 428.
 Thyme, 420.
 Thymene, 420.
Thymol, 420.
 impurities in, 721.
Thymus vulgaris, 417, 420.
 Tiglic acid, 464.
 Tiles, 355.
Tin, 241.
 amalgam, 242.
 analytical reactions of, 243.
 antidotes to, 244.
 block, 241.
 chloride, 242.
 derivation of word, 24.
 dropped, or grain, 241.
 granulated, 241.
 oxide, 243.
 perchloride, 242.
 prepare liquor, 243.
 tacks, 242.
 34 *
 Tin-foil, 241.
 Tin-plate, 242.
 Tin-stone, 241.
 Tin-white cobalt, 235.
 Tincal, 333.
Tinctura ferri acetatis, 154.
 chloridi, 150.
 impurities in, 721.
 iodi, 274.
 nucis vomicæ, 700.
 quininæ, 523.
 ammoniata, 522.
Tincturæ, 150.
 Tincture of phenolphthalein, 639.
 Tinctures, 150, 589.
Tinospora cordifolia, 509.
 Titanium, 726.
 Tobacco, 539.
Toddalix radix, 510.
 Toluene, 495.
 Tolu, balsam, 422, 429, 495.
 Toluene, 429, 431.
 dihydric alcohols, 456.
 sulphonic acid, 445.
 amide, 445.
 chloride, 445.
 Toluol, 431.
 Toluyl alcohol, 455.
 derivatives, 432.
 Tonic substances, 509.
 Tonka bean, 494.
 Toughened caustic, 218.
 nitrate of silver, 218.
 Tourmalines, 354.
 Toxicodendric acid, 361.
 Toxicology, 561.
 Tragacanth, 116, 479.
Tragacantha, 116.
 Tragacanthin, 479.
 Treacle, 471.
 Tree, lead, 215.
 silver, 220.
 Triads, 122.
 Triamines, 512.
 Triangle, wire, 100.
 Tribasic acids, 498.
 Tribasylous radicals, 124.
 Tribromophenol, 454.
 Tricarballic acid, 498.
 Trichloroacetal, 486.
 Trichloraldehyde, 485.
 Trichlorobutylidene glycol, 488.
 Trichlorethylidene ethyl ether, 488.
 glycol, 486.
 Trichloromethane, 399.
 Trichloromethylbenzene, 336, 432.
 Triethylamine, 511.
 Triethylia, 511.
 Trigoneline, 543.

- Trihydric alcohols, 457.
 Trihydroxybenzene, 465.
 Trihydroxybenzoic acid, 493, 494.
 Trihydroxyl derivatives of the hydrocarbons, 457.
 Tri-iodomethane, 401.
 Trimethyl methane, 397.
 phenoëne, 429.
 Trimethylamine, 512.
 Trinitrine, 458.
 Trinitrocellulin, 480.
 Trinitro-phenol, 454.
 Triphane, 227.
 Triple phosphate, 581.
 Tripoli, 354.
 Trithionic acid, 346.
Triticum repens, 510.
 sativum, 473.
 starch, fig. of, 475.
Trituratio elaterina, 503.
 Tritylia, 512.
 Trivalence, 56.
 Trivalent radicals, 56, 67, 123.
Trochisci acidi tannici, 358.
 bismuthi, 252.
 ferri redacti, 158.
 morphinæ, 517.
 et ipecacuanhæ, 517.
 potassii chloratis, 295.
 sodii bicarbonatis, 87.
 sulphuris, 303.
 Tropate of tropine, 532.
 Tropic acid, 532.
 Tropidine, 532.
 Tropine, 532.
 Tube-funnels, 22.
 Tubes for collecting gases, 68.
 glass. See Glass Tubes.
 Tungsten, 726.
 Tunicin, 480.
 Turgite, 153.
 Turkey corn, 536.
 Turmeric, 420, 554.
 oil, 420.
 paper, 96.
 Turmerol, 420.
 Turnbull's blue, 161, 342.
 Turnsole, 555.
Turpentine, 412, 427.
 American, 412.
 Bordeaux, 412.
 Canadian, 412.
 Chian, 412.
 French, 412.
 Russian, 412.
 impure, 412.
 rectified oil of, 412.
 spirit of, 412.
 Strasburg, 412.
Turpentine, Venice, 412.
 Turpeth mineral, 201.
 Turps, 412.
Tylophora asthmatica, 537.
 Type-metal, 180, 210, 251.
 Types, chemical, 379.
 Typical formulæ, 379.
 Tyrosine, 513.
 Tyrotoxicon, 513, 571.

ULEXINE, 536.
 Ultimate analysis, 688.
 Ultramarine blue, 555.
 green, 556.
 Ultraquinine, 526.
 Umbelliferone, 428.
 UMBER, 556.
Uncaria gambier, 359.
Unguentum aconitinæ, 530.
 antimonii tartarati, 182.
 cerussæ, 211.
 glycerini plumbi subacetatis, 212.
 hydrargyri, 195.
 ammoniaci, 207.
 iodidi rubri, 199.
 nitratæ, 200.
 dilutæ, 200.
 oxidi rubri, 204.
 subchloridi, 203.
 iodi, 274.
 paraffini, 398.
 plumbi acetatis, 211.
 carbonatis, 211.
 iodidi, 213.
 potassæ sulphuratæ, 69.
 staphisagriæ, 537.
 veratrinæ, 544.
 zinci oxidi, 135.
 Unitary hypothesis, 286.
 Units of length, surface, capacity, and weight, 603.
 Univalence, 56.
 Univalent radical, 56, 123.
 Unsaturated hydrocarbons, 393.
 Unsymmetrical compounds, 433.
 Uranium, 726.
 Urari, 528.
 Urate of lithium, 228.
 Urates, 361.
Urceola elastica, 420.
 Urea, 339, 489, 512, 575.
 artificial, 339, 578.
 nitrate of, 576.
 test for excess of, in urine, 576.
 Urethane, 489.
 Uric acid, 361, 575, 579, 581, 587.
 Urinary calculi, 572.
 examination of, 587.

- Urinary deposits or sediments,
 plates of, 582.
 sediments, 579.
 microscopical examination
 of, 581.
- Urine**, 361, 572.
 diabetic, 575.
 estimation of sugar in, 702.
 of urea in, 576.
 morbid examination of, 572.
- Urinometer, 575.
 Urobilin, 573.
 Uroerythrin, 582.
 Urostealith, 587.
Uvæ, 318, 469.
Uva ursi, 359.
- Valency**, 56.
 variation in, 139.
 conception of, 391.
- Valerine, 420.
 Valerian oil, 420.
Valerianæ, 420
 Valerianate of amyl, 333, 401.
 sodium, 362.
 zinc, 136, 363.
 Valerianates, 362.
 Valerianic or valeric acid, 362, 420,
 488.
 Valerol, 420, 427.
 Valerone, 498.
 Valylene, 412.
 Vanadates, 332.
 Vanadinite, 332.
 Vanadium, 332, 726.
 relationship to nitrogen, phos-
 phorus, and arsenum, 332.
- Vanilla, 364.
Vanilla planifolia, 364.
 Vanillic acid, 364.
 Vanillin, 364.
Vapor acidi hydrocyanici, 282.
 chlori, 29.
 coninæ, 536.
 iodi, 273.
 olei pini sylvestris, 413.
- Vapor-density, 619.
Variolaria, 556.
- Vegetable** albumen, 548.
 alkaloids, 512.
 and animal life, relation of, 20.
 casein, 548.
 crocus, 554.
 fibrin, 548.
 gelatin, 479.
 green, 556.
 jelly, 479.
 oil, 458.
 rouge, 555.
- Vegeto-animal alkaloids, 512.
 Venetian red, 153.
 Venice turpentine, 412.
 Veratralbine, 539.
Veratri viridis rhizoma, 539.
 Veratridine, 544.
Veratrina, 721.
 impurities in, 853.
 Veratrine, or Veratria, 539, 543.
 oleate of, 460.
 Veratroidine, 539.
Veratrum album, 539.
 officinale, 543.
 viride, 539.
 Verbena oil, 420.
 Verdigris, 192.
 Vermilion, 208, 554.
Veronica virginica, 510.
 Viburnin, 510.
Viburnum, 510.
 Vinegar, 297.
 estimation of mineral acids in,
 682.
 of cantharides, 298.
 of squill, 298.
- Vinum album*, 441.
 impurities in, 721.
 antimonii, 182.
 aurantii, 441.
 ferri, 157.
 citratis, 157.
 ipecacuanhæ, 537.
 quininæ, 522.
 rubrum, 441.
 impurities in, 721.
 xericum, 441.
- Violet, salicylic acid in, 492.
 Virginia snakeroot, 510.
 Vitelline, 545.
Vitellus, 545.
 Vitriol, blue, 144, 191.
 green, 144, 309.
 oil of, 309, 310.
 white, 144.
- Volatile oils. See Oils.
 Volatility of salts of ammonium, 99.
 Volatilization, 99.
 Volcanic ammonia, 91.
 Volume, combination by, 52.
 of gas, corrections of, 619, 620.
 molecular, 55.
- Volumetric** quantitative analysis,
 625.
- Volumetric estimation of—
 acetic acid, 637.
 acid, phosphoric, 639.
 hypophosphorous, 639.
 acids, 636.
 acidum arsenosi liq., 646.

- Volumetric estimation of acidum
 hydriodici syrupus, 644.
 ammonia, solutions of, 629.
 ammonium bromide, 641.
 carbonate, 630.
 antimony, 646.
 arsenic, 647.
 barium dioxide, 653.
 calcium bromide, 642.
 chlorine, solution of, 656.
 citric acid, 637.
 ferrous arsenate, 650.
 carbonate, 653.
 phosphate, 651.
 sulphate, 653.
 hydrochloric acid, 637.
 hydrocyanic acid, 640.
 hydrogen dioxide, 653.
 hyposulphurous acid, 654.
 iodine, 656.
 compound tincture, 657.
 tincture, 657.
 iron, arsenate, 650.
 iodide, 644.
 magnetic oxide, 651.
 phosphate, 651.
 reduced, 653.
 saccharated carbonate, 650.
 sulphate, 653.
 lactic acid, 637.
 lead acetate, 631.
 lime, chlorinated, solution of, 657.
 lime-water, 635.
 lithium bromide, 642.
 benzoate, 634.
 carbonate, 634.
 citrate, 634.
 salicylate, 634.
 nitric acid, 638.
 potash, 632.
 potassæ liq., 632.
 potassium, 631.
 bicarbonate, 631.
 bromide, 642.
 carbonate, 631.
 citrate, 632.
 cyanide, 641.
 iodide, 643.
 permanganate, 635.
 tartrate, 632.
 soda, 632.
 chlorinated solution of, 657.
 sodæ liq., 632.
 sodium, 631.
 arsenate, 646.
 bicarbonate, 631.
 bisulphite, 648.
 bromide, 642.
- Volumetric estimation of sodium
 carbonate, 631.
 citrate, 632.
 hyposulphite, 647.
 sulphite, 647.
 tartrate, 632.
spiritus ammoniæ aromaticus, 630.
 strontinum bromide, 643.
 lactate, 633.
 sulphuric acid, 639.
 sulphurous acid, 646.
 tartar emetic, 647.
 tartaric acid, 639.
 zinc bromide, 643.
 chloride, 643.
 iodide, 643.
- Volumetric solutions, 628, 534, 636,
 639, 645, 648, 652, 655.
- Vulcanite, 420.
 Vulcanized india-rubber, 420.
- WAHOO BARK, 510.
 Wakhuma, 530.
 Warmth of animals, how kept up,
 20.
 Washing bottles, 109.
 flasks, 109.
 precipitates, 110.
 Wasp-sting, 339.
- Water, 127.
 aërated, 88.
 ammonia in potable, 630.
 baryta-, 103.
 boiling-point of, 596.
 chalybeate, 142.
 chlorine-, 28, 267.
 chloroform-, 401.
 composition of, 22.
 cubic inches of, in a gallon, 621.
 distilled, 128.
 estimation of, 687.
 evaporation of, 71.
 formation of, expressed by sym-
 bols, 46.
 hardness of, 315.
 hemlock-, 417.
 lead in, 214.
 lime-, 108.
 nitrites in, 350.
 of crystallization, 85, 139.
 oxygenated, 103.
 potash-, 72.
 preparation of, 22.
 purification of, 127.
 quantitative estimation of, 687,
 soda-, 86.
 softness of, 315.
 weight of a cubic inch of, 608,
 621.


- Water**, weight of minim, drachm, ounce, pint, and gallon, 608.
 Water-aspirator, 309.
 Water-bath, 109, 112.
 Water-oven, 660.
 Water-type, 379.
 Wax, 450.
 Chinese, 450.
 paraffin, 399.
 Wedgewood ware, 355.
 Weighing-tubes, 660.
Weight, 600.
 estimation of, 600.
 molecular, 55.
 definition of, 59.
 of air, 620.
 of hydrogen, 620.
 of water, 621.
 specific, 600, 613.
Weights, atomic, 52.
 balance, 600.
 and measures, 600.
 and measures of the metric decimal system, *et seq.*, 601.
 of the British Pharmacopœia of 1885, 608.
 relative, 52.
 of litres at different temperatures, 620.
 Weld, 554.
 Welding, 142, 249.
 Wheaten flour, 473.
 Wheat starch (fig.), 475.
 Whey, 411, 546.
 Whiskey, 441.
 White arsenic, 168.
 Castile soap, 461.
 indigo, 201.
 lead, 210, 557.
 marble, 106.
 pepper, 541.
 pigments, 557.
 precipitate, 206.
 fusible, 206.
 infusible, 207.
 resin, 423.
 vitriol, 144.
 wax, 450.
 Whiting, 121.
 Whortleberry, sugar in, 467.
 Wild indigo, 530.
 Wild black-cherry bark, 500.
 Willow-bark, 456, 506.
 Wine, 441.
 antimonial, 182.
 apple, 441.
 heavy oil of, 409.
 ipêcacuanha, 537.
 iron, 157.
 Wine, oil of, 409.
 orange, 441.
 pear, 441.
 quinine, 522.
 sherry, 441.
 steel, 157.
 Winter-green, oil of, 407.
 Wire triangle, 100.
 Wire-gauze tray, 29.
 Witch-hazel, 509.
 Witherite, 103.
 Wood, specific gravity of, 618.
 Wood-charcoal, 113.
 Wood-creasote, 453.
 Wood-naphtha, 437.
 Wood-oil, 429.
 Wood-spirit, 437.
 Wood-tar, 413, 426.
 Woody night-shade, 541.
 Wool fat, 461.
 Woorara, 628.
 Wormwood, 417, 479.
 Wourali, 628.
 Wrought iron, 142.

XANTHINE, 543, 589.
 Xanthocreatinine, 513.
 Xanthorrhiza *apiifolia*, 533.
 Xanthoxylon *fraxineum*, 533.
 Xanthoxylum, 533.
 Xylene, 429.
 Xylenes, 455.
 Xyloidin, 473.
 Xylonite, 480.

YARD, 608.
 Yeast, 439.
 Yelk of egg, 545.
 Yellow chromate of potassium, 105.
 coloring-matters, 553.
 dock, 338.
 ochre, 553.
 oxide of mercury, 204.
 parilla, 533.
 prussiate of potash, 280.
 sienna, 553.
 soap, 462.
 wax, 450.
 wood, 553.
 Yolk of egg, 545.
 Ytterbium, 726.
 Yttrium, 726.

ZAFFRE, 235.
 Zanaloin, 435.
 Zanzibar aloes, 435.
 Zea *mays*, 472.
 starch of (fig.), 475.
Zinc, 131.

- Zinc** acetate, 134.
 analytical reactions of, 136.
 antidotes to, 137.
 bromide, 134.
 volumetric estimation of, 643.
 carbonate, 131, 134.
 chloride, 132.
 volumetric estimation of, 643.
 derivation of word, 33.
 detection of, in presence of aluminium and iron, 163.
 ethide, 397.
 ferrocyanide of, 137.
 granulated, 21.
 hydrate, 136.
 in organic mixtures, detection of, 564.
 iodide, 134.
 volumetric estimation of, 643.
 molecular weight of, 195.
 oleate of, 136, 460.
 oxide, 135.
 phosphide, 328.
 quantitative estimation of, 667.
 sulphate, 131.
 sulphide, 136.
 native, 131.
 sulphite, 136.
 sulphocarbolate, 454.
- Zinc**, valerianate, 136, 363.
 white, 134.
Zinci acetas, 135.
 impurities in, 721.
bromidum, 134.
 impurities in, 721.
carbonas, 134.
precipitatus, 134.
 impurities in, 721.
chloridi liquor, 133.
chloridum, 131.
 impurities in, 721.
iodidum, 134.
 impurities in, 721.
oleatum, 136.
oxidum, 135.
 impurities in, 721.
phosphidum, 387.
 impurities in, 721.
sulphas, 131.
 impurities in, 721.
sulphocarbolas, 454.
unguentum, 135.
valerianas, 136, 363.
 impurities in, 721.
Zincum, 21, 131.
 impurities in, 721.
Zingiber, 420.
 Zirconium, 726.
 Zymotic alkaloids, 513.
 Zymosis, 441.





Catalogue of Books

PUBLISHED BY

Lea Brothers & Company,

706, 708 & 710 Sansom St., Philadelphia.

The books in the annexed list will be sent by mail, post-paid, to any Post Office in the United States, on receipt of the printed prices. No risks of the mail, however, are assumed, either on money or books. Gentlemen will therefore in most cases find it more convenient to deal with the nearest bookseller.

PERIODICALS 1894.

The Medical News,

The Leading Medical Weekly of America,

Combines most advantageously for the practitioner the features of the newspaper and the weekly magazine. Its frequent issues keep the reader posted on all matters of current interest and in touch with the incessant progress in all lines of medical knowledge. Close adaptation to the needs of the active practitioner is shown by a list of subscribers large enough to justify the reduction in price to **\$4.00 per annum**, so that it is now the cheapest as well as the best large medical weekly of America. It contains from twenty-eight to thirty-two quarto pages of reading matter in each issue.

The American Journal^{of} the Medical Sciences.

Containing 112 to 128 octavo pages each month, THE AMERICAN JOURNAL accommodates elaborate Original Articles from the leading minds of the profession, careful Reviews and classified Summaries of Medical Progress. According to the highest literary authority in medicine, "from this file alone, were all other publications of the press for the last fifty years destroyed, it would be possible to reproduce the great majority of the real contributions of the world to medical science during that period."

Price, \$4.00 Per Annum.

Combinations at Reduced Rates.

THE AMERICAN JOURNAL OF THE MEDICAL SCIENCES,	} Together	\$7.50
\$4.00		
THE MEDICAL NEWS, \$4.00	} Together	\$8.50
THE MEDICAL NEWS VISITING LIST for 1894 (see below and on page 16), \$1.25		
With either or both above periodicals, 75 cents.		
THE YEAR-BOOK OF TREATMENT for 1894 (see page 16), \$1.50.	}	
With either or both above periodicals, 75 cents.		

The Medical News Visiting List.

This LIST, which is by far the most handsome and convenient now attainable, has been thoroughly revised for 1894. A full description will be found on page 16. It is issued in four styles. Price, each, \$1.25. Thumb-letter Index for quick use 25 cents extra. For Special Combination Rates with periodicals and the Year-Book of Treatment see above.

(S. 1. 4.)

ABBOTT (A. C.). PRINCIPLES OF BACTERIOLOGY: a Practical Manual for Students and Physicians. In one 12mo. volume of 259 pages, with 32 illustrations. Cloth, \$2.

ALLEN (HARRISON). A SYSTEM OF HUMAN ANATOMY; WITH AN INTRODUCTORY SECTION ON HISTOLOGY, by E. O. SHAKESPEARE, M. D. Comprising 813 double-columned quarto pages, with 380 engravings on stone on 109 plates, and 241 woodcuts in the text. In six sections, each in a portfolio. Price per section, \$3.50. Also, bound in one volume, cloth, \$23; half Russia, \$25. *Sold by subscription only.*

AMERICAN SYSTEM OF DENTISTRY. In treatises by various authors. Edited by WILBUR F. LITCH, M.D., D.D.S. In three very handsome super-royal octavo volumes, containing 3180 pages, with 1863 illustrations and 9 full-page plates. Per volume, cloth, \$6; leather, \$7; half Morocco, \$8. *For sale by subscription only.* Prospectus free on application to the Publishers.

AMERICAN SYSTEMS OF GYNECOLOGY AND OBSTETRICS. In treatises by the most eminent American specialists. Gynecology edited by MATTHEW D. MANN, A. M., M. D., and Obstetrics edited by BARTON C. HIRST, M. D. In four large octavo volumes comprising 3612 pages, with 1092 engravings, and 8 colored plates. Per volume, cloth, \$5; leather, \$6; half Russia, \$7. *For sale by subscription only.* Prospectus free on application to the Publishers.

ASHHURST (JOHN, JR.). THE PRINCIPLES AND PRACTICE OF SURGERY. For the use of Students and Practitioners. New (6th) and revised edition. In one large and handsome octavo volume of 1161 pages, with 656 engravings. Cloth, \$6; leather, \$7. *Just ready.*

ASHWELL (SAMUEL). A PRACTICAL TREATISE ON THE DISEASES OF WOMEN. Third edition. 520 pages. Cloth, \$3.50.

A SYSTEM OF PRACTICAL MEDICINE BY AMERICAN AUTHORS. Edited by WILLIAM PEPPER, M. D., LL. D. In five large octavo volumes, containing 5573 pages and 198 illustrations. Price per volume, cloth, \$5; leather \$6; half Russia, \$7. *Sold by subscription only.* Prospectus free on application to the Publishers.

ATTFIELD (JOHN). CHEMISTRY; GENERAL, MEDICAL AND PHARMACEUTICAL. Twelfth edition, specially revised by the Author for America. In one handsome 12mo. volume of 782 pages, with 88 illustrations. Cloth, \$2.75; leather, \$3.25.

BARNES (ROBERT). A PRACTICAL TREATISE ON THE DISEASES OF WOMEN. Third American from third English edition. In one 8vo. vol. of about 800 pages, with about 200 illus.

BARNES (ROBERT AND FANCOURT). A SYSTEM OF OBSTETRIC MEDICINE AND SURGERY, THEORETICAL AND CLINICAL. The Section on Embryology by PROF. MILNES MARSHALL. In one large octavo volume of 872 pages, with 231 illustrations. Cloth, \$5; leather, \$6.

BARTHOLOW (ROBERTS). CHOLERA; ITS CAUSATION, PREVENTION AND TREATMENT. In one 12mo. volume of 127 pages, with 9 illustrations. Cloth, \$1.25.

— **MEDICAL ELECTRICITY. A PRACTICAL TREATISE ON THE APPLICATIONS OF ELECTRICITY TO MEDICINE AND SURGERY.** Third edition. In one octavo volume of 308 pages, with 110 illustrations.

BASHAM (W. R.). RENAL DISEASES; A CLINICAL GUIDE TO THEIR DIAGNOSIS AND TREATMENT. In one 12mo. volume of 304 pages, with 21 illustrations. Cloth, \$2.

BELL (F. JEFFREY). COMPARATIVE ANATOMY AND PHYSIOLOGY. In one 12mo. volume of 561 pages, with 229 engravings. Cloth, \$2. See *Students' Series of Manuals*, p. 14.

BELLAMY (EDWARD). A MANUAL OF SURGICAL ANATOMY.

In one 12mo. vol. of 300 pages, with 50 illustrations. Cloth, \$2.25.

BERRY (GEORGE A.). DISEASES OF THE EYE; A PRACTICAL TREATISE FOR STUDENTS OF OPHTHALMOLOGY. New (2d) edition. Very handsome octavo vol. of 745 pages, with 197 original illus. in the text, of which 87 are exquisitely colored. Cloth, \$8.**BILLINGS (JOHN S.).** THE NATIONAL MEDICAL DICTIONARY.

Including in one alphabet English, French, German, Italian and Latin Technical Terms used in Medicine and the Collateral Sciences. In two very handsome imperial octavo volumes, containing 1574 pages and two colored plates. Per volume, cloth, \$6; leather, \$7; half Morocco, \$8.50. *For sale by subscription only.* Specimen pages on application to publishers.

BLOXAM (C. L.). CHEMISTRY, INORGANIC AND ORGANIC.

With Experiments. New American from the fifth London edition. In one handsome octavo volume of 727 pages, with 292 illustrations. Cloth, \$2; leather, \$3.

BRISTOWE (J. S.). A TREATISE ON THE SCIENCE AND PRACTICE OF MEDICINE. Eighth edition. In one octavo volume of 1325 pages, with 114 illustrations.**BROADBENT (W. H.).** THE PULSE. In one 12mo. vol. of 317 pages, with 59 engravings. Cloth, \$1.75. See *Series of Clinical Manuals*, p. 13.**BROWNE (LENNOX).** THE THROAT AND NOSE AND THEIR DISEASES. New (4th) and enlarged edition. In one imperial octavo volume of 751 pages, with 235 engravings and 120 illustrations in color. Cloth, \$6.50.

— **KOCH'S REMEDY IN RELATION ESPECIALLY TO THROAT CONSUMPTION.** In one octavo volume of 121 pages, with 45 illustrations, 4 of which are colored, and 17 charts. Cloth, \$1.50.

BRUCE (J. MITCHELL). MATERIA MEDICA AND THERAPEUTICS. Fifth edition. In one 12mo. volume of 591 pages. Cloth, \$1.50. See *Student's Series of Manuals*, p. 14.**BRUNTON (T. LAUDER).** A MANUAL OF PHARMACOLOGY, THERAPEUTICS AND MATERIA MEDICA; including the Pharmacy, the Physiological Action and the Therapeutical Uses of Drugs. Third and revised edition, in one octavo vol. of 1305 pp., with 230 illus.**BRYANT (THOMAS).** THE PRACTICE OF SURGERY. Fourth American from the fourth English edition. In one imperial octavo vol. of 1040 pages, with 727 illustrations. Cloth, \$6.50; leather, \$7.50.**BUMSTEAD (F. J.) AND TAYLOR (R. W.).** THE PATHOLOGY AND TREATMENT OF VENEREAL DISEASES. New edition. See *Taylor on Venereal Diseases*.**BURNETT (CHARLES H.).** THE EAR: ITS ANATOMY, PHYSIOLOGY AND DISEASES. A Practical Treatise for the Use of Students and Practitioners. Second edition. In one 8vo. volume of 580 pages, with 107 illustrations. Cloth, \$4; leather, \$5.**BUTLIN (HENRY T.).** DISEASES OF THE TONGUE. In one pocket-size 12mo. volume of 456 pages, with 8 colored plates and 3 engravings. Limp cloth, \$3.50. See *Series of Clinical Manuals*, page 13.**CARPENTER (WM. B.).** PRIZE ESSAY ON THE USE OF ALCOHOLIC LIQUORS IN HEALTH AND DISEASE. New edition, with a Preface by D. F. CONDIE, M.D. One 12mo. volume of 178 pages. Cloth, 60 cents.

— **PRINCIPLES OF HUMAN PHYSIOLOGY.** In one large octavo volume.

- CARTER (R. BRUDENELL) AND FROST (W. ADAMS).** OPTHALMIC SURGERY. In one pocket-size 12mo. volume of 559 pages, with 91 engravings and one plate. Cloth, \$2.25. See *Series of Clinical Manuals*, p. 13.
- CASPARI (CHARLES JR.).** A TEXT-BOOK ON PHARMACY. For Students and Pharmacists. In one handsome octavo volume, thoroughly illustrated. *In press for early publication.*
- CHAMBERS (T. K.).** A MANUAL OF DIET IN HEALTH AND DISEASE. In one handsome 8vo. vol. of 302 pages. Cloth, \$2.75.
- CHAPMAN (HENRY C.).** A TREATISE ON HUMAN PHYSIOLOGY. In one octavo volume of 925 pages, with 605 illustrations. Cloth, \$5.50; leather, \$6.50.
- CHARLES (T. CRANSTOUN).** THE ELEMENTS OF PHYSIOLOGICAL AND PATHOLOGICAL CHEMISTRY. In one handsome octavo volume of 451 pages, with 38 engravings and 1 colored plate. Cloth, \$3.50.
- CHURCHILL (FLEETWOOD).** ESSAYS ON THE PUERPERAL FEVER. In one octavo volume of 464 pages. Cloth, \$2.50.
- CLARKE (W. B.) AND LOCKWOOD (C. B.).** THE DISSECTOR'S MANUAL. In one 12mo. volume of 396 pages, with 49 engravings. Cloth, \$1.50. See *Students' Series of Manuals*, p. 14.
- CLASSEN'S** QUANTITATIVE ANALYSIS. Translated by EDGAR F. SMITH, Ph. D. In one 12mo. vol. of 324 pp., with 36 illus. Cloth, \$2.
- CLELAND (JOHN).** A DIRECTORY FOR THE DISSECTION OF THE HUMAN BODY. In one 12mo. vol. of 178 pages. Cloth, \$1.25.
- CLOUSTON (THOMAS S.).** CLINICAL LECTURES ON MENTAL DISEASES. With an Abstract of Laws of U. S. on Custody of the Insane, by C. F. FOLSOM, M. D. In one handsome octavo volume of 541 pages, illustrated with engravings and 8 lithographic plates. Cloth, \$4. DR. FOLSOM'S *Abstract* is also furnished separately in one octavo volume of 108 pages. Cloth, \$1.50.
- CLOWES (FRANK).** AN ELEMENTARY TREATISE ON PRACTICAL CHEMISTRY AND QUALITATIVE INORGANIC ANALYSIS. From the fourth English edition. In one handsome 12mo. volume of 387 pages, with 55 engravings. Cloth, \$2.50.
- COATS (JOSEPH).** A TREATISE ON PATHOLOGY. In one vol. of 829 pages, with 339 engravings. Cloth, \$5.50; leather, \$6.50.
- COHEN (S. SOLIS).** A HANDBOOK OF APPLIED THERAPEUTICS. One large 12mo. volume, with illustrations. *Preparing.*
- COLEMAN (ALFRED).** A MANUAL OF DENTAL SURGERY AND PATHOLOGY. With Notes and Additions to adapt it to American Practice. By THOS. C. STELLWAGEN, M.A., M.D., D.D.S. In one handsome octavo vol. of 412 pages, with 331 engravings. Cloth, \$3.25.
- CONDIE (D. FRANCIS).** A PRACTICAL TREATISE ON THE DISEASES OF CHILDREN. Sixth edition, revised and enlarged. In one large 8vo. volume of 719 pages. Cloth, \$5.25; leather, \$6.25.
- CORNIL (V.).** SYPHILIS: ITS MORBID ANATOMY, DIAGNOSIS AND TREATMENT. Translated, with Notes and Additions, by J. HENRY C. SIMES, M.D. and J. WILLIAM WHITE, M.D. In one 8vo. volume of 461 pages, with 84 illustrations. Cloth, \$3.75.
- CULVER (E. M.) AND HAYDEN (J. R.).** MANUAL OF VENEREAL DISEASES. In one 12mo. volume of 289 pages, with 33 engravings. Cloth, \$1.75.

- DALTON (JOHN C.).** A TREATISE ON HUMAN PHYSIOLOGY. Seventh edition, thoroughly revised and greatly improved. In one very handsome octavo volume of 722 pages, with 252 engravings. Cloth, \$5; leather, \$6.
- DOCTRINES OF THE CIRCULATION OF THE BLOOD. In one handsome 12mo. volume of 293 pages. Cloth, \$2.
- DANA (JAMES D.).** THE STRUCTURE AND CLASSIFICATION OF ZOÖPHYTES. With illustrations on wood. In one imperial 4to. volume. Cloth, \$4.
- DAVENPORT (F. H.).** DISEASES OF WOMEN. A Manual of Non-Surgical Gynecology. For the use of Students and General Practitioners. Second edition. In one handsome 12mo. volume of 314 pages, with 107 engravings. Cloth, \$1.75.
- DAVIS (F. H.).** LECTURES ON CLINICAL MEDICINE. Second edition. In one 12mo. volume of 287 pages. Cloth, \$1.75.
- DE LA BECHE'S** GEOLOGICAL OBSERVER. In one large octavo volume of 700 pages, with 300 engravings. Cloth, \$4.
- DENNIS (FREDERIC S.) AND BILLINGS (JOHN S.)** A SYSTEM OF SURGERY. In Treatises by various Authors; In very handsome octavo volumes, richly illustrated. *Preparing.*
- DRAPER (JOHN C.).** MEDICAL PHYSICS. A Text-book for Students and Practitioners of Medicine. In one handsome octavo volume of 734 pages, with 376 engravings. Cloth, \$4.
- DRUITT (ROBERT).** THE PRINCIPLES AND PRACTICE OF MODERN SURGERY. A new American, from the twelfth London edition, edited by STANLEY BOYD, F.R.C.S. In one large octavo volume of 965 pages, with 373 engravings. Cloth, \$4; leather, \$5.
- DUANE (ALEXANDER.)** THE STUDENT'S DICTIONARY OF MEDICINE AND THE ALLIED SCIENCES. Comprising the Pronunciation, Derivation and Full Explanation of Medical Terms. Together with much Collateral Descriptive Matter, Numerous Tables, etc. In one square octavo volume of 658 pages. Cloth, \$4.25; half leather, \$4.50; full sheep, \$5. Thumb-letter Index for quick use, 50 cents extra. *Just ready.*
- DUNCAN (J. MATTHEWS).** CLINICAL LECTURES ON THE DISEASES OF WOMEN. Delivered in St. Bartholomew's Hospital. In one octavo volume of 175 pages. Cloth, \$1.50.
- DUNGLISON (ROBLEY).** A DICTIONARY OF MEDICAL SCIENCE. Containing a full explanation of the various subjects and terms of Anatomy, Physiology, Medical Chemistry, Pharmacy, Pharmacology, Therapeutics, Medicine, Hygiene, Dietetics, Pathology, Surgery, Ophthalmology, Otology, Laryngology, Dermatology, Gynecology, Obstetrics, Pediatrics, Medical Jurisprudence, Dentistry, etc., etc. By ROBLEY DUNGLISON, M.D., LL.D., late Professor of Institutes of Medicine in the Jefferson Medical College of Philadelphia. Edited by RICHARD J. DUNGLISON, A.M., M.D. New (21st) edition, thoroughly revised and greatly enlarged and improved, with the Pronunciation, Accentuation and Derivation of the Terms. In one magnificent imperial octavo volume of 1191 pages. Cloth, \$7; leather, \$8. Thumb-letter Index for quick use, 75 cents extra. *Just ready.*
- EDES (ROBERT T.).** TEXT-BOOK OF THERAPEUTICS AND MATERIA MEDICA. In one 8vo. volume of 544 pages. Cloth, \$3.50; leather, \$4.50.
- EDIS (ARTHUR W.).** DISEASES OF WOMEN. A Manual for Students and Practitioners. In one handsome 8vo. volume of 576 pages, with 148 engravings. Cloth, \$3; leather, \$4.
- ELLIS (GEORGE VINER).** DEMONSTRATIONS IN ANATOMY. Being a Guide to the Knowledge of the Human Body by Dissection. From the eighth and revised English edition. In one octavo volume of 716 pages, with 249 engravings. Cloth, \$4.25; leather, \$5.25.

EMMET (THOMAS ADDIS). THE PRINCIPLES AND PRACTICE OF GYNÆCOLOGY, for the use of Students and Practitioners. Third edition, enlarged and revised. In one large 8vo. volume of 880 pages, with 150 original engravings. Cloth, \$5; leather, \$6.

ERICHSEN (JOHN E.). THE SCIENCE AND ART OF SURGERY. A new American from the eighth enlarged and revised London edition. In two large octavo volumes containing 2316 pages, with 984 engravings. Cloth, \$9; leather, \$11.

FARQUHARSON (ROBERT). A GUIDE TO THERAPEUTICS. Fourth American from fourth English edition, revised by FRANK WOODBURY, M. D. In one 12mo. volume of 581 pages. Cloth, \$2.50.

FIELD (GEORGE P.). A MANUAL OF DISEASES OF THE EAR. Fourth edition. In one octavo volume of 391 pages, with 73 engravings and 21 colored plates. Cloth, \$3.75. *Just ready.*

FLINT (AUSTIN). A TREATISE ON THE PRINCIPLES AND PRACTICE OF MEDICINE. New (7th) edition, thoroughly revised by FREDERICK P. HENRY, M. D. In one large 8vo. volume of 1143 pages, with engravings. Cloth, \$5.00; leather, \$6.00. *Just ready.*

——— A MANUAL OF AUSCULTATION AND PERCUSSION; of the Physical Diagnosis of Diseases of the Lungs and Heart, and of Thoracic Aneurism. Fifth edition, revised by JAMES C. WILSON, M. D. In one handsome 12mo. volume of 274 pages, with 12 engravings. Cloth, \$1.75.

——— A PRACTICAL TREATISE ON THE DIAGNOSIS AND TREATMENT OF DISEASES OF THE HEART. Second edition, enlarged. In one octavo volume of 550 pages. Cloth, \$4.

——— A PRACTICAL TREATISE ON THE PHYSICAL EXPLORATION OF THE CHEST, AND THE DIAGNOSIS OF DISEASES AFFECTING THE RESPIRATORY ORGANS. Second and revised edition. In one octavo volume of 591 pages. Cloth, \$4.50.

——— MEDICAL ESSAYS. In one 12mo. vol. of 210 pages. Cloth, \$1.38.

——— ON PHTHISIS: ITS MORBID ANATOMY, ETIOLOGY, ETC. A Series of Clinical Lectures. In one 8vo. volume of 442 pages. Cloth, \$3.50.

FOLSOM (C. F.). AN ABSTRACT OF STATUTES OF U. S. ON CUSTODY OF THE INSANE. In one 8vo. vol. of 108 pages. Cloth, \$1.50. Also bound with *Clouston on Insanity*.

FOSTER (MICHAEL). A TEXT-BOOK OF PHYSIOLOGY. Fifth and revised American from the fifth English edition. In one large octavo volume of 1064 pages, with 316 engravings. Cloth, \$4.50; leather, \$5.50. *Just ready.*

FOTHERGILL (J. MILNER). THE PRACTITIONER'S HANDBOOK OF TREATMENT. Third edition. In one handsome octavo volume of 664 pages. Cloth, \$3.75; leather, \$4 75.

FOWNES (GEORGE). A MANUAL OF ELEMENTARY CHEMISTRY (INORGANIC AND ORGANIC). Twelfth edition. Embodying WATTS' *Physical and Inorganic Chemistry*. In one royal 12mo. volume of 1061 pages, with 168 engravings, and 1 colored plate. Cloth, \$2.75; leather, \$3.25.

FOX (TILBURY AND T. COLCOTT). EPITOME OF SKIN DISEASES, with Formulæ. For Students and Practitioners. Third American edition, revised by T. C. FOX. In one small 12mo. volume of 238 pages. Cloth, \$1.25.

- FRANKLAND (E.) AND JAPP (F.R.).** INORGANIC CHEMISTRY. In one handsome octavo volume of 677 pages, with 51 engravings and 2 plates. Cloth, \$3.75; leather, \$4.75.
- FULLER (HENRY).** ON DISEASES OF THE LUNGS AND AIR PASSAGES. Their Pathology, Physical Diagnosis, Symptoms and Treatment. From second English edition. In one 8vo. volume of 475 pages. Cloth, \$3.50.
- GANT (FREDERICK JAMES).** THE STUDENT'S SURGERY. A Multum in Parvo. In one square octavo volume of 845 pages, with 159 engravings. Cloth, \$3.75.
- GIBBES (HENEAGE).** PRACTICAL PATHOLOGY AND MORBID HISTOLOGY. In one very handsome octavo volume of 314 pages, with 60 illustrations, mostly photographic. Cloth, \$2.75.
- GIBNEY (V. P.).** ORTHOPEDIC SURGERY. For the use of Practitioners and Students. In one 8vo. vol. profusely illus. *Preparing.*
- GOULD (A. PEARCE).** SURGICAL DIAGNOSIS. In one 12mo. vol. of 589 pages. Cloth, \$2. See *Student's Series of Manuals*, p. 14.
- GRAY (HENRY).** ANATOMY, DESCRIPTIVE AND SURGICAL. Edited by T. PICKERING PICK, F.R.C.S. A new American, from the thirteenth English edition, thoroughly revised. In one imperial octavo volume of 1118 pages, with 636 large and elaborate engravings. Price, with illustrations in colors, cloth, \$7; leather, \$8. Price, with illustrations in black, cloth, \$6; leather, \$7. *Just ready.*
- GRAY (LONDON CARTER).** A TREATISE ON NERVOUS AND MENTAL DISEASES. For Students and Practitioners of Medicine. In one handsome octavo volume of 681 pages, with 168 engravings. Cloth, \$4.50; leather, \$5.50.
- GREEN (T. HENRY).** AN INTRODUCTION TO PATHOLOGY AND MORBID ANATOMY. Sixth American from the seventh London edition. In one handsome octavo volume of 540 pages, with 167 engravings. Cloth, \$2.75.
- GREENE (WILLIAM H.).** A MANUAL OF MEDICAL CHEMISTRY. For the Use of Students. Based upon BOWMAN'S *Medical Chemistry*. In one 12mo. vol. of 310 pages, with 74 illus. Cloth, \$1.75.
- GROSS (SAMUEL W.).** A PRACTICAL TREATISE ON IMPO-TENCE, STERILITY, AND ALLIED DISORDERS OF THE MALE SEXUAL ORGANS. Fourth edition. Edited by F. R. STURGIS, M.D. In one handsome octavo vol. of 165 pages, with 18 illus. Cloth, \$1.50.
- GROSS (SAMUEL D.).** A PRACTICAL TREATISE ON THE DISEASES, INJURIES AND MALFORMATIONS OF THE URINARY BLADDER, THE PROSTATE GLAND AND THE URETHRA. Third edition, thoroughly revised and edited by SAMUEL W. GROSS, M. D. In one octavo vol. of 574 pages, with 170 illus. Cloth, \$4.50.
- HABERSHON (S. O.).** ON THE DISEASES OF THE ABDOMEN, comprising those of the Stomach, Œsophagus, Caecum, Intestines and Peritoneum. Second American from the third English edition. In one octavo volume of 554 pages, with 11 engravings. Cloth, \$3.50.
- HAMILTON (ALLAN MCLANE).** NERVOUS DISEASES, THEIR DESCRIPTION AND TREATMENT. Second and revised edition. In one octavo volume of 598 pages, with 72 engravings. Cloth, \$4.
- HAMILTON (FRANK H.).** A PRACTICAL TREATISE ON FRACTURES AND DISLOCATIONS. Eighth edition, revised and edited by STEPHEN SMITH, A. M., M. D. In one handsome octavo volume of 832 pages, with 507 engravings. Cloth, \$5.50; leather, \$6.50.
- HARDAWAY (W. A.).** MANUAL OF SKIN DISEASES. In one 12mo. volume of 440 pages. Cloth, \$3.

HARE (HOBART AMORY). A TEXT-BOOK OF PRACTICAL THERAPEUTICS, with Special Reference to the Application of Remedial Measures to Disease and their Employment upon a Rational Basis. With articles on various subjects by well-known specialists. New (4th) edition, revised and enlarged. In one octavo volume of 740 pages. Cloth, \$3.75; leather, \$4.75. *Just ready.*

HARE (HOBART AMORY), EDITOR. A SYSTEM OF PRACTICAL THERAPEUTICS. By American and Foreign Authors. In a series of contributions by 78 eminent Physicians. Three large octavo volumes comprising 3544 pages, with 434 engravings; Price per volume, cloth, \$5; leather, \$6; half Russia, \$7. *For sale by subscription only.* Address the publishers.

HARTSHORNE (HENRY). ESSENTIALS OF THE PRINCIPLES AND PRACTICE OF MEDICINE. Fifth edition. In one 12mo. volume, 669 pages, with 144 engravings. Cloth, \$2.75; half bound, \$3.

— A HANDBOOK OF ANATOMY AND PHYSIOLOGY. In one 12mo. volume of 310 pages, with 220 engravings. Cloth, \$1.75.

— A CONSPECTUS OF THE MEDICAL SCIENCES. Comprising Manuals of Anatomy, Physiology, Chemistry, Materia Medica, Practice of Medicine, Surgery and Obstetrics. Second edition. In one royal 12mo. vol. of 1028 pages, with 477 illus. Cloth, \$4.25; leather, \$5.

HERMAN (G. ERNEST). FIRST LINES IN MIDWIFERY. In one 12mo. vol. of 198 pages, with 80 engravings. Cloth, \$1.25. See *Student's Series of Manuals*, p. 14.

HERMANN (L.). EXPERIMENTAL PHARMACOLOGY. A Handbook of the Methods for Determining the Physiological Actions of Drugs. Translated by ROBERT MEADE SMITH, M. D. In one 12mo. volume of 199 pages, with 32 engravings. Cloth, \$1.50.

HILL (BERKELEY). SYPHILIS AND LOCAL CONTAGIOUS DISORDERS. In one 8vo. volume of 479 pages. Cloth, \$3.25.

HILLIER (THOMAS). A HANDBOOK OF SKIN DISEASES. Second edition. In one royal 12mo. volume of 353 pages, with two plates. Cloth, \$2.25.

HIRST (BARTON C.) AND PIERSOL (GEORGE A.). HUMAN MONSTROSITIES. Magnificent folio, containing 220 pages of text and illustrated with 123 engravings and 39 large photographic plates from nature. In four parts, price each, \$5. *Limited edition. For sale by subscription only.*

HOBLYN (RICHARD D.). A DICTIONARY OF THE TERMS USED IN MEDICINE AND THE COLLATERAL SCIENCES. In one 12mo. volume of 520 double-columned pages. Cloth, \$1.50; leather, \$2.

HODGE (HUGH L.). ON DISEASES PECULIAR TO WOMEN, INCLUDING DISPLACEMENTS OF THE UTERUS. Second and revised edition. In one 8vo. vol. of 519 pp., with illus. Cloth, \$4.50.

HOFFMANN (FREDERICK) AND POWER (FREDERICK B.). A MANUAL OF CHEMICAL ANALYSIS, as Applied to the Examination of Medicinal Chemicals and their Preparations. Third edition, entirely rewritten and much enlarged. In one handsome octavo volume of 621 pages, with 179 engravings. Cloth, \$4.25.

HOLDEN (LUTHER). LANDMARKS, MEDICAL AND SURGICAL. From the third English edition. With additions by W. W. KEEN, M. D. In one royal 12mo. volume of 148 pages. Cloth, \$1.

HOLLAND (SIR HENRY). MEDICAL NOTES AND REFLECTIONS. From third English edition. In one 8vo. volume of 493 pages. Cloth, \$3.50.

HOLMES (TIMOTHY). A TREATISE ON SURGERY. Its Principles and Practice. A new American from the fifth English edition. Edited by T. PICKERING PICK, F.R.C.S. In one handsome octavo volume of 1008 pages, with 428 engravings. Cloth, \$6; leather, \$7.

— A SYSTEM OF SURGERY. With notes and additions by various American authors. Edited by JOHN H. PACKARD, M.D. In three very handsome 8vo. volumes containing 3137 double-columned pages, with 979 engravings and 13 lithographic plates. Per volume, cloth, \$6; leather, \$7; half Russia, \$7.50. *For sale by subscription only.*

HORNER (WILLIAM E.). SPECIAL ANATOMY AND HISTOLOGY. Eighth edition, revised and modified. In two large 8vo. volumes of 1007 pages, containing 320 engravings. Cloth, \$6.

HUDSON (A.). LECTURES ON THE STUDY OF FEVER. In one octavo volume of 308 pages. Cloth, \$2.50.

HUTCHINSON (JONATHAN). SYPHILIS. In one pocket-size 12mo. volume of 542 pages, with 8 chromo-lithographic plates. Cloth, \$2.25. See *Series of Clinical Manuals*, p. 13.

HYDE (JAMES NEVINS). A PRACTICAL TREATISE ON DISEASES OF THE SKIN. New (3d) edition, thoroughly revised. In one octavo volume of 802 pages, with 108 engravings and 9 colored plates. Cloth, \$5; leather, \$6. *Just ready.*

JACKSON (GEORGE THOMAS). THE READY-REFERENCE HANDBOOK OF DISEASES OF THE SKIN. In one 12mo. volume of 544 pages, with 50 engravings. \$2.75.

JAMIESON (W. ALLAN). DISEASES OF THE SKIN. Third edition. In one octavo volume of 656 pages, with 1 engraving and 9 double-page chromo-lithographic plates. Cloth, \$6.

JONES (C. HANDFIELD). CLINICAL OBSERVATIONS ON FUNCTIONAL NERVOUS DISORDERS. Second American edition. In one octavo volume of 340 pages. Cloth, \$3.25.

JULER (HENRY). A HANDBOOK OF OPHTHALMIC SCIENCE AND PRACTICE. New (2d) edition. In one octavo volume of 549 pages, with 201 engravings, 17 chromo-lithographic plates, test-types of Jaeger and Snellen, and Holmgren's Color-Blindness Test. Cloth, \$5.50; leather, \$6.50.

KING (A. F. A.). A MANUAL OF OBSTETRICS. Fifth edition. In one 12mo. volume of 446 pages, with 150 illustrations. Cloth, \$2.50.

KLEIN (E.). ELEMENTS OF HISTOLOGY. Fourth edition. In one pocket-size 12mo. volume of 376 pages, with 194 engravings. Cloth, \$1.75. See *Student's Series of Manuals*, p. 14.

LANDIS (HENRY G.). THE MANAGEMENT OF LABOR. In one handsome 12mo. volume of 329 pages, with 28 illus. Cloth, \$1.75.

LA ROCHE (R.). YELLOW FEVER. In two 8vo. volumes of 1468 pages. Cloth, \$7.

— PNEUMONIA. In one 8vo. volume of 490 pages. Cloth, \$3.

LAURENCE (J. Z.) AND MOON (ROBERT C.). A HANDY-BOOK OF OPHTHALMIC SURGERY. Second edition. In one octavo volume of 227 pages, with 66 engravings. Cloth, \$2.75.

LAWSON (GEORGE). INJURIES OF THE EYE, ORBIT AND EYE-LIDS. From the last English edition. In one handsome octavo volume of 404 pages, with 92 engravings. Cloth, \$3.50.

LEA (HENRY C.). CHAPTERS FROM THE RELIGIOUS HISTORY OF SPAIN; CENSORSHIP OF THE PRESS; MYSTICS AND ILLUMINATI; THE ENDEMONIADAS; EL SANTO NIÑO DE LA GUARDIA; BRIANDA DE BARDAXI. In one 12mo. volume of 522 pages. Cloth, \$2.50.

LEA (HENRY C.). FORMULARY OF THE PAPAL PENITENTIARY. In one 8vo. vol. of 221 pages, with frontispiece. Cloth, \$2.50.

— **SUPERSTITION AND FORCE; ESSAYS ON THE WAGER OF LAW, THE WAGER OF BATTLE, THE ORDEAL AND TORTURE.** Fourth edition, thoroughly revised. In one handsome royal 12mo. volume of 629 pages. Cloth, \$2.75.

— **STUDIES IN CHURCH HISTORY.** The Rise of the Temporal Power—Benefit of Clergy—Excommunication. New edition. In one handsome 12mo. volume of 605 pages. Cloth, \$2.50.

— **AN HISTORICAL SKETCH OF SACERDOTAL CELIBACY IN THE CHRISTIAN CHURCH.** Second edition. In one handsome octavo volume of 685 pages. Cloth, \$4.50.

LEDGER. THE MEDICAL NEWS PHYSICIAN'S LEDGER. Contains 300 pages ledger paper, ruled in approved style. Strongly bound with patent flexible back. Price, \$4.

LEE (HENRY) ON SYPHILIS. In one 8vo. volume of 246 pages. Cloth, \$2.25.

LEHMANN (C. G.). A MANUAL OF CHEMICAL PHYSIOLOGY. In one 8vo. volume of 327 pages, with 41 engravings. Cloth, \$2.25.

LEISHMAN (WILLIAM). A SYSTEM OF MIDWIFERY. Including the Diseases of Pregnancy and the Puerperal State. Fourth edition. In one octavo volume.

LOOMIS (ALFRED L.) AND THOMPSON (W. GILMAN). A SYSTEM OF PRACTICAL MEDICINE. In Treatises by Various Authors. In very handsome 8vo. volumes with illus. *Preparing.*

LUCAS (CLEMENT). DISEASES OF THE URETHRA. *Preparing.* See *Series of Clinical Manuals*, p. 13.

LUDLOW (J. L.). A MANUAL OF EXAMINATIONS UPON ANATOMY, PHYSIOLOGY, SURGERY, PRACTICE OF MEDICINE, OBSTETRICS, MATERIA MEDICA, CHEMISTRY, PHARMACY AND THERAPEUTICS. To which is added a Medical Formulary. Third edition. In one royal 12mo. volume of 816 pages, with 370 engravings. Cloth, \$3.25; leather, \$3.75.

LUFF (ARTHUR P.). MANUAL OF CHEMISTRY, for the use of Students of Medicine. In one 12mo. volume of 522 pages, with 36 engravings. Cloth, \$2. See *Student's Series of Manuals*, p. 14.

LYMAN (HENRY M.). THE PRACTICE OF MEDICINE. In one very handsome octavo volume of 925 pages, with 170 engravings. Cloth, \$4.75; leather, \$5.75.

LYONS (ROBERT D.). A TREATISE ON FEVER. In one octavo volume of 362 pages. Cloth, \$2.25.

MAISCH (JOHN M.). A MANUAL OF ORGANIC MATERIA MEDICA. Fifth edition. In one very handsome 12mo. volume of 544 pages, with 270 engravings. Cloth, \$3.

MANUALS. See *Student's Quiz Series*, p. 14, *Student's Series of Manuals*, p. 14, and *Series of Clinical Manuals*, p. 13.

MARSH (HOWARD). DISEASES OF THE JOINTS. In one 12mo. volume of 468 pages, with 64 engravings and a colored plate. Cloth, \$2. See *Series of Clinical Manuals*, p. 13.

MAY (C. H.). MANUAL OF THE DISEASES OF WOMEN. For the use of Students and Practitioners. Second edition, revised by L. S. RAY, M. D. In one 12mo. volume of 360 pages, with 31 engravings. Cloth, \$1.75.

MILLER (JAMES). PRINCIPLES OF SURGERY. Fourth American from the third Edinburgh edition. In one large octavo volume of 688 pages, with 240 engravings. Cloth, \$3.75.

— **THE PRACTICE OF SURGERY.** Fourth American from the last Edinburgh edition. In one large octavo volume of 682 pages, with 364 engravings. Cloth, \$3.75.

- MORRIS (HENRY).** SURGICAL DISEASES OF THE KIDNEY. In one 12mo. volume of 554 pages, with 40 engravings and 6 colored plates. Cloth, \$2.25. See *Series of Clinical Manuals*, p. 13.
- MORRIS (MALCOLM).** DISEASES OF THE SKIN. In one square 8vo. volume of 572 pages, with 19 chromo-lithographic figures and 17 engravings. Cloth, \$3.50. *Just ready.*
- MÜLLER (J.).** PRINCIPLES OF PHYSICS AND METEOROLOGY. In one large 8vo. vol. of 623 pages, with 538 cuts. Cloth, \$4.50.
- MUSSER (JOHN H.).** A PRACTICAL TREATISE ON MEDICAL DIAGNOSIS, for Students and Physicians. In one octavo volume of 873 pages, illustrated with 162 engravings and 2 colored plates. Cloth, \$5; leather, \$6. *Just ready.*
- NATIONAL DISPENSATORY.** See *Stillé, Maisch & Caspari*, p. 14.
- NATIONAL MEDICAL DICTIONARY.** See *Billings*, p. 3.
- NETTLESHIP (E.).** DISEASES OF THE EYE. Fourth American from fifth English edition. In one 12mo. volume of 504 pages, with 164 engravings, test-types and formulæ and color-blindness test. Cloth, \$2.
- NORRIS (WM. F.) AND OLIVER (CHAS. A.).** TEXT-BOOK OF OPHTHALMOLOGY. In one octavo volume of 641 pages, with 357 engravings and 5 colored plates. Cloth, \$5; leather, \$6.
- OWEN (EDMUND).** SURGICAL DISEASES OF CHILDREN. In one 12mo. volume of 525 pages, with 85 engravings and 4 colored plates. Cloth, \$2. See *Series of Clinical Manuals*, p. 13.
- PARRISH (EDWARD).** A TREATISE ON PHARMACY. With many Formulæ and Prescriptions. Fifth edition, enlarged and thoroughly revised by THOMAS S. WIEGAND, Ph. G. In one octavo volume of 1093 pages, with 257 engravings. Cloth, \$5; leather, \$6.
- PARRY (JOHN S.).** EXTRA-UTERINE PREGNANCY, ITS CLINICAL HISTORY, DIAGNOSIS, PROGNOSIS AND TREATMENT. In one octavo volume of 272 pages. Cloth, \$2.50.
- PARVIN (THEOPHILUS).** THE SCIENCE AND ART OF OBSTETRICS. Second edition. In one handsome 8vo. volume of 701 pages, with 239 engravings and a colored plate. Cloth, \$4.25; leather, \$5.25.
- PAVY (F. W.).** A TREATISE ON THE FUNCTION OF DIGESTION, ITS DISORDERS AND THEIR TREATMENT. From the second London edition. In one 8vo. volume of 238 pages. Cloth, \$2.
- PAYNE (JOSEPH FRANK).** A MANUAL OF GENERAL PATHOLOGY. Designed as an Introduction to the Practice of Medicine. In one octavo volume of 524 pages, with 153 engravings and 1 colored plate. Cloth, \$3.50.
- PEPPER'S SYSTEM OF MEDICINE.** See p. 2.
- PEPPER (A. J.).** FORENSIC MEDICINE. *In press.* See *Student's Series of Manuals*, p. 14.
- SURGICAL PATHOLOGY. In one 12mo. volume of 511 pages, with 81 engravings. Cloth, \$2. See *Student's Series of Manuals*, p. 14.
- PICK (T. PICKERING).** FRACTURES AND DISLOCATIONS. In one 12mo. volume of 530 pages, with 93 engravings. Cloth, \$2. See *Series of Clinical Manuals*, p. 13.
- PIRRIE (WILLIAM).** THE PRINCIPLES AND PRACTICE OF SURGERY. In one octavo volume of 780 pages, with 316 engravings. Cloth, \$3.75.

PLAYFAIR (W. S.). A TREATISE ON THE SCIENCE AND PRACTICE OF MIDWIFERY. Sixth American from the eighth English edition. Edited, with additions, by R. P. HARRIS, M. D. In one octavo volume of 697 pages, with 217 engravings and 5 plates. Cloth, \$4; leather, \$5. *Just ready.*

— THE SYSTEMATIC TREATMENT OF NERVE PROSTRATION AND HYSTERIA. In one 12mo. vol. of 97 pp. Cloth, \$1.

POLITZER (ADAM). A TEXT-BOOK OF THE DISEASES OF THE EAR AND ADJACENT ORGANS. Second American from the third German edition. Translated by OSCAR DODD, M. D., and edited by SIR WILLIAM DALBY, F. R. C. S. In one octavo volume of 748 pages, with 330 original engravings. Cloth, \$5.50. *Just ready.*

POWER (HENRY). HUMAN PHYSIOLOGY. Second edition. In one 12mo. volume of 396 pages, with 47 engravings. Cloth, \$1.50. See *Student's Series of Manuals*, p. 14.

PURDY (CHARLES W.). BRIGHT'S DISEASE AND ALLIED AFFECTIONS OF THE KIDNEY. In one octavo volume of 288 pages, with 18 engravings. Cloth, \$2.

PYE-SMITH (PHILIP H.). DISEASES OF THE SKIN. In one 12mo. vol. of 407 pp., with 28 illus., 18 of which are colored. Cloth, \$2.

QUIZ SERIES. See *Student's Quiz Series*, p. 14.

RALFE (CHARLES H.). CLINICAL CHEMISTRY. In one 12mo. volume of 314 pages, with 16 engravings. Cloth, \$1.50. See *Student's Series of Manuals*, p. 14.

RAMSBOTHAM (FRANCIS H.). THE PRINCIPLES AND PRACTICE OF OBSTETRIC MEDICINE AND SURGERY. In one imperial octavo volume of 640 pages, with 64 plates and numerous engravings in the text. Strongly bound in leather, \$7.

REICHERT (EDWARD T.). A TEXT-BOOK ON PHYSIOLOGY. In one handsome octavo volume of about 800 pages, richly illustrated. *Preparing.*

REMSEN (IRA). THE PRINCIPLES OF THEORETICAL CHEMISTRY. Fourth edition, thoroughly revised and much enlarged. In one 12mo. volume of 325 pages. Cloth, \$2.

REYNOLDS (J. RUSSELL). A SYSTEM OF MEDICINE. Edited, with notes and additions, by HENRY HARTSHORNE, M. D. In three large 8vo. vols., containing 3056 closely printed double-columned pages, with 317 engravings. Per volume, cloth, \$5; leather, \$6; very handsome half Russia, \$6.50. *For sale by subscription only.*

RICHARDSON (BENJAMIN WARD). PREVENTIVE MEDICINE. In one octavo volume of 729 pages. Cloth, \$4; leather, \$5.

ROBERTS (JOHN B.). THE PRINCIPLES AND PRACTICE OF MODERN SURGERY. In one octavo volume of 780 pages, with 501 engravings. Cloth, \$4.50; leather, \$5.50.

— THE COMPEND OF ANATOMY. For use in the Dissecting Room and in preparing for Examinations. In one 16mo. volume of 196 pages. Limp cloth, 75 cents.

ROBERTS (SIR WILLIAM). A PRACTICAL TREATISE ON URINARY AND RENAL DISEASES, INCLUDING URINARY DEPOSITS. Fourth American from the fourth London edition. In one very handsome 8vo. vol. of 609 pp., with 81 illus. Cloth, \$3.50.

ROBERTSON (J. MCGREGOR). PHYSIOLOGICAL PHYSICS. In one 12mo. volume of 537 pages, with 219 engravings. Cloth, \$2. See *Student's Series of Manuals*, p. 14.

ROSS (JAMES). A HANDBOOK OF THE DISEASES OF THE NERVOUS SYSTEM. In one handsome octavo volume of 726 pages, with 184 engravings. Cloth, \$4.50; leather, \$5.50.

SAVAGE (GEORGE H.). INSANITY AND ALLIED NEUROSES, PRACTICAL AND CLINICAL. In one 12mo. volume of 551 pages, with 18 typical engravings. Cloth, \$2. See *Series of Clinical Manuals*, p. 13.

SCHAFER (EDWARD A.). THE ESSENTIALS OF HISTOLOGY, DESCRIPTIVE AND PRACTICAL. For the use of Students. New (3d) and enlarged edition. In one handsome octavo volume of 311 pages, with 325 illustrations. Cloth, \$3.

SCHMITZ AND ZUMPT'S CLASSICAL SERIES.

ADVANCED LATIN EXERCISES. Cloth, 60 cents; half bound, 70 cents.

SCHMIDT'S ELEMENTARY LATIN EXERCISES. Cloth, 50 cents.

SALLUST. Cloth, 60 cents; half bound, 70 cents.

NEPOS. Cloth, 60 cents; half bound, 70 cents.

VIRGIL. Cloth, 85 cents; half bound, \$1.

CURTIUS. Cloth, 80 cents; half bound, 90 cents.

SCHOFIELD (ALFRED T.). ELEMENTARY PHYSIOLOGY FOR STUDENTS. In one 12mo. volume of 380 pages, with 227 engravings and 2 colored plates. Cloth, \$2.

SCHREIBER (JOSEPH). A MANUAL OF TREATMENT BY MASSAGE AND METHODOICAL MUSCLE EXERCISE. Translated by WALTER MENDELSON, M. D., of New York. In one handsome octavo volume of 274 pages, with 117 fine engravings.

SEILER (CARL). A HANDBOOK OF DIAGNOSIS AND TREATMENT OF DISEASES OF THE THROAT AND NASAL CAVITIES. Fourth edition. In one 12mo. volume of 414 pages, with 107 engravings, and 2 colored plates. Cloth, \$2.25.

SENN (NICHOLAS). SURGICAL BACTERIOLOGY. Second edition. In one octavo volume of 268 pages, with 13 plates, 10 of which are colored, and 9 engravings. Cloth, \$2.

SERIES OF CLINICAL MANUALS. A Series of Authoritative Monographs on Important Clinical Subjects, in 12mo. volumes of about 550 pages, well illustrated. The following volumes are now ready: BROADBENT on the Pulse, \$1.75; YEO on Food in Health and Disease, \$2; CARTER and FROST'S Ophthalmic Surgery, \$2.25; HUTCHINSON on Syphilis, \$2.25; MARSH on Diseases of the Joints, \$2; MORRIS on Surgical Diseases of the Kidney, \$2.25; OWEN on Surgical Diseases of Children, \$2; PICK on Fractures and Dislocations, \$2; BUTLIN on the Tongue, \$3.50; SAVAGE on Insanity and Allied Neuroses, \$2; and TREVES on Intestinal Obstruction, \$2. The following is in press: LUCAS on Diseases of the Urethra.

For separate notices, see under various authors' names.

SERIES OF STUDENT'S MANUALS. See next page.

SIMON (W.). MANUAL OF CHEMISTRY. A Guide to Lectures and Laboratory Work for Beginners in Chemistry. A Text-book specially adapted for Students of Pharmacy and Medicine. New (4th) edition. In one 8vo. volume of 490 pages, with 44 engravings and 7 plates showing colors of 56 tests. Cloth, \$3.25.

SLADE (D. D.). DIPHTHERIA; ITS NATURE AND TREATMENT. Second edition. In one royal 12mo. vol., 158 pp. Cloth, \$1.25.

SMITH (EDWARD). CONSUMPTION; ITS EARLY AND REMEDIABLE STAGES. In one 8vo. volume of 253 pp. Cloth, \$2.25.

SMITH (J. LEWIS). A TREATISE ON THE DISEASES OF INFANCY AND CHILDHOOD. Seventh edition, revised and enlarged. In one large 8vo. volume of 881 pages, with 51 illustrations. Cloth, \$4.50; leather, \$5.50.

SMITH (STEPHEN). OPERATIVE SURGERY. Second and thoroughly revised edition. In one octavo volume of 892 pages, with 1005 engravings. Cloth, \$4; leather, \$5.

STILLÉ (ALFRED). CHOLERA; ITS ORIGIN, HISTORY, CAUSATION, SYMPTOMS, LESIONS, PREVENTION AND TREATMENT. In one 12mo. volume of 163 pages, with a chart showing routes of previous epidemics. Cloth, \$1.25.

— **THERAPEUTICS AND MATERIA MEDICA.** Fourth and revised edition. In two octavo volumes, containing 1936 pages. Cloth, \$10; leather, \$12.

STILLÉ (ALFRED), MAISCH (JOHN M.) AND CASPARI (CHAS. JR.). THE NATIONAL DISPENSATORY: Containing the Natural History, Chemistry, Pharmacy, Actions and Uses of Medicines, including those recognized in the latest Pharmacopœias of the United States, Great Britain and Germany, with numerous references to the French Codex. New (5th) edition, revised and enlarged in accordance with the new U. S. Pharmacopœia, Seventh Decennial Revision. In one magnificent imperial octavo volume of 1910 pages, with 320 engravings. Cloth, \$7.25; leather, \$8. With ready reference Thumb-letter Index. Cloth, \$7.75; leather, \$8.50. *Just ready.*

STIMSON (LEWIS A.). A TREATISE ON FRACTURES AND DISLOCATIONS. In two handsome octavo volumes. Vol. I., FRACTURES, 582 pages, 360 engravings. Vol. II., DISLOCATIONS, 540 pages, 163 engravings. Complete work, cloth, \$5.50; leather, \$7.50. Either volume separately, cloth, \$3; leather, \$4.

— **A MANUAL OF OPERATIVE SURGERY.** Second edition. In one royal 12mo. volume of 503 pages, with 342 illus. Cloth, \$2.50.

STUDENT'S QUIZ SERIES. A New Series of Manuals in question and answer for Students and Practitioners, covering the essentials of medical science. Thirteen volumes, pocket size, convenient, authoritative, well illustrated, handsomely bound in limp cloth, and issued at a low price. 1. Anatomy (double number); 2. Physiology; 3. Chemistry and Physics; 4. Histology, Pathology and Bacteriology; 5. Materia Medica and Therapeutics; 6. Practice of Medicine; 7. Surgery (double number); 8. Genito-Urinary and Venereal Diseases; 9. Diseases of the Skin; 10. Diseases of the Eye, Ear, Throat and Nose; 11. Obstetrics; 12. Gynecology; 13. Diseases of Children. Price, \$1 each, except Nos. 1 and 7, *Anatomy and Surgery*, which being double numbers are priced at \$1.75 each. Full specimen circular on application to publishers.

STUDENT'S SERIES OF MANUALS. A Series of Fifteen Manuals by Eminent Teachers or Examiners. The volumes are pocket-size 12mos. of from 300–540 pages, profusely illustrated, and bound in red limp cloth. The following volumes may now be announced: HERMAN'S First Lines in Midwifery, \$1.25; LUFF'S Manual of Chemistry, \$2; BRUCE'S Materia Medica and Therapeutics (fifth edition), \$1.50; TREVES' Manual of Surgery (monographs by 33 leading surgeons), 3 volumes, per set, \$6; BELL'S Comparative Anatomy and Physiology, \$2; ROBERTSON'S Physiological Physics, \$2; GOULD'S Surgical Diagnosis, \$2; KLEIN'S Elements of Histology (4th edition), \$1.75; PEPPER'S Surgical Pathology, \$2; TREVES' Surgical Applied Anatomy, \$2; POWER'S Human Physiology (2d edition), \$1.50; RALFE'S Clinical Chemistry, \$1.50; and CLARKE and LOCKWOOD'S Dissector's Manual, \$1.50. The following is in press: PEPPER'S Forensic Medicine.

For separate notices, see under various authors' names.

STURGES (OCTAVIUS). AN INTRODUCTION TO THE STUDY OF CLINICAL MEDICINE. In one 12mo. volume. Cloth, \$1.25.

SUTTON (JOHN BLAND). SURGICAL DISEASES OF THE OVARIES AND FALLOPIAN TUBES. Including Abdominal Pregnancy. In one 12mo. volume of 513 pages, with 119 engravings and 5 colored plates. Cloth, \$3.

— **TUMORS, INNOCENT AND MALIGNANT.** Their Clinical Features and Appropriate Treatment. In one 8vo. vol. of 526 pp., with 250 engravings and 9 full-page plates. Cloth, \$4.50. *Just ready.*

TAIT (LAWSON). DISEASES OF WOMEN AND ABDOMINAL SURGERY. In two handsome octavo volumes. Vol. I. contains 546 pages and 3 plates. Cloth, \$3. Vol. II., *preparing.*

TANNER (THOMAS HAWKES). A MANUAL OF CLINICAL MEDICINE AND PHYSICAL DIAGNOSIS. Third American from the second revised English edition. Edited by TILBURY FOX, M. D. In one handsome 12mo. vol. of 362 pp., with illus. Cloth, \$1.50.

— ON THE SIGNS AND DISEASES OF PREGNANCY. From the second English edition. In one octavo volume of 490 pages, with 4 colored plates and 16 engravings. Cloth, \$4.25.

TAYLOR (ALFRED S.). MEDICAL JURISPRUDENCE. Eleventh American from the twelfth English edition, specially revised by CLARK BELL, Esq., of the N. Y. Bar. In one octavo volume of 787 pages, with 56 engravings. Cloth, \$4.50; leather, \$5.50.

— ON POISONS IN RELATION TO MEDICINE AND MEDICAL JURISPRUDENCE. Third American from the third London edition. In one 8vo. vol. of 788 pp., with 104 illus. Cloth, \$5.50; leath., \$6.50.

TAYLOR (ROBERT W.). A CLINICAL ATLAS OF VENEREAL AND SKIN DISEASES. Including Diagnosis, Prognosis and Treatment. In eight large folio parts, measuring 14 x 18 inches, and comprising 213 beautiful figures on 58 full-page chromo-lithographic plates, 85 fine engravings, and 425 pages of text. Complete work now ready. Price per part, sewed in heavy embossed paper, \$2.50. Bound in one volume, half Russia, \$27; half Turkey Morocco, \$28. *For sale by subscription only.* Address the publishers. Specimen plates by mail on receipt of ten cents.

— THE PATHOLOGY AND TREATMENT OF VENEREAL DISEASES. Being the sixth edition of BUMSTEAD and TAYLOR. In one very handsome 8vo. volume of about 900 pages, with about 150 engravings as well as chromo-lithographic plates. *Preparing.*

THOMAS (T. GAILLARD) AND MUNDÉ (PAUL F.). A PRACTICAL TREATISE ON THE DISEASES OF WOMEN. New (6th) edition, thoroughly revised by PAUL F. MUNDÉ, M. D. In one large and handsome octavo volume of 824 pages, with 347 engravings. Cloth, \$5; leather, \$6.

THOMPSON (SIR HENRY). CLINICAL LECTURES ON DISEASES OF THE URINARY ORGANS. Second and revised edition. In one octavo vol. of 203 pp., with 25 engravings. Cloth, \$2.25.

— THE PATHOLOGY AND TREATMENT OF STRICTURE OF THE URETHRA AND URINARY FISTULÆ. From the third English edition. In one octavo volume of 359 pages, with 47 engravings and 3 lithographic plates. Cloth, \$3.50.

TODD (ROBERT BENTLEY). CLINICAL LECTURES ON CERTAIN ACUTE DISEASES. In one 8vo. vol. of 320 pp., cloth, \$2.50.

TREVES (FREDERICK). OPERATIVE SURGERY. In two 8vo. vols. containing 1550 pp., with 422 illus. Cloth, \$9; leath., \$11.

— A MANUAL OF SURGERY. In Treatises by 33 leading surgeons. Three 12mo. volumes, containing 1866 pages, with 213 engravings. Price per set, \$6. See *Student's Series of Manuals*, p. 14.

— THE STUDENTS' HANDBOOK OF SURGICAL OPERATIONS. In one 12mo. vol. of 508 pp., with 94 illus. Cloth, \$2.50.

— SURGICAL APPLIED ANATOMY. In one 12mo. vol. of 540 pp., with 61 engravings. Cloth, \$2. See *Student's Series of Manuals*, p. 14.

— INTESTINAL OBSTRUCTION. In one 12mo. volume of 522 pages, with 60 illus. Cloth, \$2. See *Series of Clinical Manuals*, p. 13.

TUKE (DANIEL HACK). THE INFLUENCE OF THE MIND UPON THE BODY IN HEALTH AND DISEASE. Second edition. In one 8vo. volume of 467 pages, with 2 colored plates. Cloth, \$3.

VAUGHAN (VICTOR C.) AND NOVY (FREDERICK G.). PTOMAINES, LEUCOMAINES AND BACTERIAL PROTEIDS, or the Chemical Factors in the Causation of Disease. Second edition. In one 12mo. volume of 389 pages. Cloth, \$2.25.

VISITING LIST. THE MEDICAL NEWS VISITING LIST for 1894.

Four styles: Weekly (dated for 30 patients); Monthly (undated for 120 patients per month); Perpetual (undated for 30 patients each week); and Perpetual (undated for 60 patients each week). The 60-patient book consists of 256 pages of assorted blanks. The first three styles contain 32 pages of important data, thoroughly revised, and 176 pages of assorted blanks. Each in one volume, price, \$1.25. With thumb-letter index for quick use, 25 cents extra. Special rates to advance-paying subscribers to THE MEDICAL NEWS or THE AMERICAN JOURNAL OF THE MEDICAL SCIENCES, or both. See p. 1.

WALSHE (W. H.). PRACTICAL TREATISE ON THE DISEASES OF THE HEART AND GREAT VESSELS. Third American from the third revised London edition. In one 8vo. vol. of 420 pp. Cloth, \$3.

WATSON (THOMAS). LECTURES ON THE PRINCIPLES AND PRACTICE OF PHYSIC. A new American from the fifth and enlarged English edition, with additions by H. HARTSHORNE, M. D. In two large 8vo. vols. of 1840 pp., with 190 cuts. Cloth, \$9; leather, \$11.

WELLS (J. SOELBERG). A TREATISE ON THE DISEASES OF THE EYE. In one large and handsome octavo volume.

WEST (CHARLES). LECTURES ON THE DISEASES PECULIAR TO WOMEN. Third American from the third English edition. In one octavo volume of 543 pages. Cloth, \$3.75; leather, \$4.75.

— ON SOME DISORDERS OF THE NERVOUS SYSTEM IN CHILDHOOD. In one small 12mo. volume of 127 pages. Cloth, \$1.

WHARTON (HENRY R.). MINOR SURGERY AND BANDAGING. New (2d) edition. In one 12mo. volume of 529 pages, with 416 engravings, many of which are photographic. Cloth, \$3. *Just ready.*

WHITLA (WILLIAM). DICTIONARY OF TREATMENT, OR THERAPEUTIC INDEX. Including Medical and Surgical Therapeutics. In one square octavo volume of 917 pages. Cloth, \$4.

WILLIAMS (CHARLES J. B. AND C. T.). PULMONARY CONSUMPTION: ITS NATURE, VARIETIES AND TREATMENT. In one octavo volume of 303 pages. Cloth, \$2.50.

WILSON (ERASMUS). A SYSTEM OF HUMAN ANATOMY. A new and revised American from the last English edition. Illustrated with 397 engravings. In one octavo volume of 616 pages. Cloth, \$4; leather, \$5.

— THE STUDENT'S BOOK OF CUTANEOUS MEDICINE. In one 12mo. volume. Cloth, \$3.50.

WINCKEL ON PATHOLOGY AND TREATMENT OF CHILDBED. Translated by JAMES R. CHADWICK, A. M., M. D. With additions by the Author. In one octavo volume of 484 pages. Cloth, \$4.

WÖHLER'S OUTLINES OF ORGANIC CHEMISTRY. Translated from the eighth German edition, by IRA REMSEN, M. D. In one 12mo. volume of 550 pages. Cloth, \$3.

YEAR-BOOK OF TREATMENT FOR 1894. A Critical Review for Practitioners of Medicine and Surgery. In contributions by 24 well-known medical writers. 12mo., 500 pages. Cloth, \$1.50. In combination with THE MEDICAL NEWS and THE AMERICAN JOURNAL OF THE MEDICAL SCIENCES, 75 cents. See p. 1.

YEAR-BOOKS OF TREATMENT FOR 1891, 1892 and 1893, similar to above. Each, cloth, \$1.50.

YEAR-BOOKS OF TREATMENT FOR 1886 and 1887. Similar to above. 12mo., 320-341 pages. Limp cloth, \$1.25.

YEO (I. BURNEY). FOOD IN HEATH AND DISEASE. In one 12mo. vol. of 590 pp. Cloth, \$2. See *Series of Clinical Manuals*, p. 13.

— A MANUAL OF MEDICAL TREATMENT, OR CLINICAL THERAPEUTICS. Two volumes containing 1275 pages. Cloth, \$5.50.

YOUNG (JAMES K.). ORTHOPEDIC SURGERY. In one 12mo. volume of 475 pages, with 286 illustrations. Cloth, \$4; leather, \$5. *Just ready.*

LIBRARY OF CONGRESS



0 003 833 850 A

